

# Ideal Gas Thermodynamic Properties of Ethane and Propane

Cite as: Journal of Physical and Chemical Reference Data **2**, 427 (1973); <https://doi.org/10.1063/1.3253123>  
Published Online: 29 October 2009

J. Chao, R. C. Wilhoit, and B. J. Zwolinski



View Online



Export Citation

## ARTICLES YOU MAY BE INTERESTED IN

### Thermophysical Properties of Fluids. II. Methane, Ethane, Propane, Isobutane, and Normal Butane

Journal of Physical and Chemical Reference Data **16**, 577 (1987); <https://doi.org/10.1063/1.555785>

### Thermophysical Properties of Ethane

Journal of Physical and Chemical Reference Data **20**, 275 (1991); <https://doi.org/10.1063/1.555881>

### A Reference Equation of State for the Thermodynamic Properties of Ethane for Temperatures from the Melting Line to 675 K and Pressures up to 900 MPa

Journal of Physical and Chemical Reference Data **35**, 205 (2006); <https://doi.org/10.1063/1.1859286>



# Ideal Gas Thermodynamic Properties of Ethane and Propane

J. Chao, R. C. Wilhoit, and B. J. Zwolinski

Thermodynamics Research Center, Texas A&M University, College Station, Texas 77843

The thermodynamic properties ( $C_p^\circ$ ,  $S^\circ$ ,  $H^\circ-H_0^\circ$ ,  $(H^\circ-H_0^\circ)/T$ ,  $-(G^\circ-H_0^\circ)/T$ ,  $\Delta H_f^\circ$ ,  $\Delta G_f^\circ$  and  $\log K_f$ ) for ethane and propane in the ideal gaseous state in the temperature range from 0 to 1500 K and at 1 atm were calculated by statistical thermodynamic methods based on a rigid-rotor harmonic-oscillator model. The internal rotation contributions to thermodynamic functions were evaluated by using a partition function formed by summation of internal rotation energy levels. The calculated heat capacities and entropies compare favorably with available experimental data.

Key words: Critically evaluated data; enthalpy; enthalpy function; enthalpy of formation; entropy; equilibrium constant of formation; ethane; Gibbs energy function; Gibbs energy of formation; heat capacity; ideal gas thermodynamic properties; internal rotation; internal rotation barrier height; internal rotation energy levels; propane; torsional frequencies.

## Contents

	Page
1. Introduction .....	427
2. Ethane .....	427
3. Propane.....	428
4. Discussion.....	430
5. Acknowledgment.....	431
6. References.....	431

## List of Tables

Table 1.	Structural Data for Ethane
Table 2.	Vibrational Assignments for Ethane
Table 3.	Determinations of Torsional Frequency ( $\nu_4$ ) in Ethane.
Table 4.	Reported Potential Barriers to Internal Rotation in Ethane.
Table 5.	Molecular Constants for Ethane
Table 6.	Ideal Gas Thermodynamic Properties for Ethane.

Table 7.	Summary of Heat Capacity Measurements for Ethane.
Table 8.	Comparison of Calculated and Measured Heat Capacities for Ethane.
Table 9.	Structural Data for Propane
Table 10.	Vibrational Assignments for Propane
Table 11.	Summary of Reported Torsional Frequencies in Propane.
Table 12.	Reported Potential Barriers to Internal Rotation in Propane.
Table 13.	Comparison of Calculated $C_p^\circ$ and $S^\circ$ for Propane.
Table 14.	Molecular Constants for Propane
Table 15.	Ideal Gas Thermodynamic Properties for Propane.
Table 16.	Comparison of Calculated and Measured Heat Capacities for Propane.
Table 17.	Comparison of Calculated and API 44 Table 1952 Values of $C_p^\circ$ and $S^\circ$ for Ethane and Propane.

## 1. Introduction

Ethane and propane are two basic organic raw materials used extensively in chemical and petrochemical industries for manufacture of numerous important products. The thermodynamic properties for these two compounds have been investigated intensively by many researchers over a long period. In view of the availability of more reliable molecular and spectroscopic constants and of changes in the fundamental constants [1]<sup>1</sup>, the ideal gas chemical thermodynamic properties for ethane and propane have been re-evaluated in the temperature range from 0 to 1500 K and at 1 atm. The mass units used are  $C=12.011$  and  $H=1.008$  (1969 atomic weights).

The contribution of internal rotation of the methyl groups to the thermodynamic properties of ethane and propane was calculated by use of partition functions

formed by summation of the internal rotation energy levels for each  $\text{CH}_3$  rotor. These energy levels were generated using a procedure developed by Laane, et al. [32]. The calculated internal rotational contributions to thermodynamic properties were added to the contributions from the other degrees of freedom, less internal rotation of each methyl group, to yield the thermodynamic properties as shown in tables 6 and 15.<sup>2</sup>

The calculated heat capacities and entropies are compared with the available measured values in tables 8 and 16.

## 2. Ethane

Ethane ( $\text{C}_2\text{H}_6$ ) has been a subject of great interest to chemists and physicists for many years. Since the work of Kemp and Pitzer [2, 3] established the existence of a potential barrier hindering the free rotation of the  $\text{CH}_3$  group about the C-C bond in the  $\text{CH}_3\text{CH}_3$  molecule, a significant number of experimental as well as theoretical investigations have been devoted to the study of the

<sup>1</sup> Figures in brackets indicate literature references in section 6.

Copyright © 1973 by the U.S. Secretary of Commerce on behalf of the United States. This copyright will be assigned to the American Institute of Physics and the American Chemical Society, to whom all requests regarding reproduction should be addressed.

<sup>2</sup> All tables have been placed at the end of this paper.

molecular properties of ethane. These molecular data are needed for the calculation of the thermodynamic properties of ethane in the ideal gaseous state by statistical thermodynamic techniques. The method of selection of these values for calculation is described as follows.

The structural parameters of the ethane molecule have been elucidated from electron diffraction studies [4, 5], and infrared [6, 7] and Raman [8, 9] spectra. Because of its high symmetry ( $D_{3d}$ ), ethane has no permanent dipole moment and hence cannot be studied by microwave techniques. Four sets of typical results obtained by use of different experimental methods are presented in table 1. The values determined by Shaw, Lepard, and Welsh [9] from the rotational constants derived from the Raman spectrum were adopted for this work. Based on these data, we derived the three principal moments of inertia as:  $I_A = 10.4811 \times 10^{-40}$  and  $I_B = I_C = 42.2486 \times 10^{-40}$  g cm<sup>2</sup>. The product  $I_A I_B I_C$  was  $1.8708 \times 10^{-116}$  g<sup>3</sup> cm<sup>6</sup> and the reduced moment ( $I_r$ ) for internal rotation was calculated to be  $2.6203 \times 10^{-40}$  g cm<sup>2</sup>.

The vibrational frequencies of ethane have been investigated extensively by analyses of infrared [6, 7, 10–16] and Raman [8, 17–20] spectra. Recently, Shimanouchi [21] has reviewed the reported spectral data on ethane and selected a complete set of vibrational wavenumbers for this molecule as shown in table 2 (units in cm<sup>-1</sup>).<sup>3</sup> In the same table two important sets of vibrational assignments reported in the literature are listed. The values given by Smith [10] were evaluated from the infrared spectrum and those by Lepard, Shaw, and Welsh [19] were obtained from the Raman spectrum.

The hindered internal rotation of the two methyl groups in ethane has been a subject of intensive research since the 1930's. Because the torsional vibration is inactive in both the infrared and Raman spectra of ethane, spectroscopists were forced to deduce the torsional frequency ( $\nu_4$ ) from combination bands. Most determinations favor a value near 290 cm<sup>-1</sup> as shown in table 3. Based on low temperature heat capacity data, Wilson [29] and Kistiakowsky, et al. [26] derived  $\nu_4$  as 280 and 275 cm<sup>-1</sup>, respectively. Lambert and Salter [30] analyzed the ultrasonic relaxation data on ethane and evaluated the torsional vibration wavenumber to be 290 cm<sup>-1</sup>. Recently, the direct excitation of the torsional vibration in ethane was observed in a slow-neutron inelastic-scattering experiment by Strong and Brugger [31]. Theoretical curves fitted by them to the data indicated that the frequency of transition from the zero to first excited state of this oscillation was 277 cm<sup>-1</sup>, and a barrier height of 2.75 kcal mol<sup>-1</sup> was derived. It is known that selection rules often break down as the result of

intermolecular interactions or interaction among vibrations causing forbidden transitions to become possible. Based on this fact Weiss and Leroi [15] recorded the infrared spectrum of ethane in the region of the torsional fundamental and first overtone under high-pressure-path-length conditions. This was the first time that the torsional spectra of gaseous ethane have been observed directly in the infrared. The fundamental,  $\nu_4$  (0 $\rightarrow$ 1), was observed at 289 cm<sup>-1</sup>. From the above results, as listed in table 3, we adopted the value 289 cm<sup>-1</sup>, which was observed directly in the infrared by Weiss and Leroi [15], as the torsional wavenumber  $\nu_4$  for ethane for our calculation of the internal rotation contributions to thermodynamic properties.

Table 4 summarizes the reported potential barriers to internal rotation in ethane in the period 1932–1970. Except for the values of Eyring [28], Howard [27] and R. M. Pitzer [23], the barrier heights reported by the remaining investigators are in reasonable agreement. It should be emphasized that those reported  $V_0$ 's were derived based on different molecular structures (see table 1). Therefore direct comparison of the listed values is not appropriate, strictly speaking. Using the adopted  $\nu_4 = 289$  cm<sup>-1</sup>, we calculated  $V_0 = 2.96$  kcal mol<sup>-1</sup> for ethane.

By use of the selected molecular data, as shown in table 5, the ideal gas thermodynamic functions ( $H^\circ - H_0^\circ$ ,  $-(G^\circ - H_0^\circ)/T$ ,  $(H^\circ - H_0^\circ)/T$ ,  $S^\circ$ , and  $C_p^\circ$ ) for ethane in the temperature range 50–1500 K were evaluated by the statistical thermodynamic method based on a rigid-rotor harmonic-oscillator model. The internal rotation contributions to thermodynamic properties were calculated separately by using the internal rotation partition function obtained by summation of 87 energy levels (0 to 20120 cm<sup>-1</sup>) derived from the adopted potential function  $V = \frac{1}{2}V_0(1 - \cos 3\theta)$  where  $V_0 = 2.96$  kcal mol<sup>-1</sup> and  $\theta$  = angle of internal rotation [32]. The calculated first three energy levels (289, 545, and 549 cm<sup>-1</sup>) are in agreement with observed peaks (289, 544, and 547 cm<sup>-1</sup>) reported by Weiss and Leroi [15]. The values thus obtained were added to the contribution from other degrees of freedom (i.e. translation, rotation, vibration, and electronic) less internal rotation. The results are presented in table 6.

Based on the thermodynamic functions ( $H^\circ - H_0^\circ$ ) and  $-(G^\circ - H_0^\circ)/T$  for C (c) and H<sub>2</sub> (g) from [33], for C<sub>2</sub>H<sub>6</sub> (g) from this study, and  $\Delta H_f^\circ$  (298.15, C<sub>2</sub>H<sub>6</sub>, g) = -20.24 kcal mol<sup>-1</sup> [34], the values of  $\Delta H_f^\circ$ ,  $\Delta G_f^\circ$  and  $\log K_f$  in the temperatures 50–1500 K were derived and given in table 6.

### 3. Propane

Propane (C<sub>3</sub>H<sub>8</sub>) is the simplest hydrocarbon which has more than one methyl internal rotation in the molecule. Investigations on the internal rotation problem in propane have been pursued by chemists for decades. The solution of this problem may lead to a better understanding of internal rotations in more complex organic

<sup>3</sup>In keeping with the common convention in molecular spectroscopy, the fundamental frequencies and vibrational energies are frequently expressed in their wavenumber (cm<sup>-1</sup>) equivalents. The actual frequency in units of hertz may be obtained by multiplying the numbers in these tables by the speed of light expressed in centimeters per second. Energies are obtained by multiplying the frequencies by Planck's constant.

molecules. The research work on the determination of the torsional frequency and potential barrier to internal rotation of the two methyl groups in propane is briefly reviewed in this report. The thermodynamic properties of propane were evaluated by using the most recent molecular data. The procedures employed for the selection of the molecular constants and calculation of the functions are described as follows.

The molecular structure of propane has been investigated by electron diffraction [47,48] and microwave spectroscopy [49]. The molecular parameters obtained from these investigations are summarized in table 9. The values given by Lide [49] were adopted for calculation of the reduced moment ( $I_r$ ) for internal rotation of the  $\text{CH}_3$  group in propane as  $4.4202 \times 10^{-40}$  g cm<sup>2</sup>. For the reduced moment calculation, the internal rotations of the two  $\text{CH}_3$  rotors were assumed to be independent of each other. The product of the three principal moments of inertia was calculated from the ground-state rotational constants of reference [49].

The infrared spectrum of propane has been observed extensively by many researchers [50–58]. Bartholome [50], Kohlrausch and Köppl [59], and Elyashevich and Stepanov [51] have studied the Raman spectra of propane. By use of theoretical calculation, Takahashi [60], Schachtschneider and Snyder [61], Gayles, et al. [62], and Shimanouchi [21] assigned different vibrational wavenumbers for propane. Based on their excellent infrared absorption spectrum, Wu and Barker [57] made the most detailed assignment at that time. Later, Pitzer [63] extended their assignment and reported definite values (within 2%) for all frequencies in good agreement with spectral information and thermodynamic data for propane. The fundamental frequencies are listed in table 10 together with three other sets of vibrational assignments. The assignments by Schachtschneider and Snyder [61] were based on extensive normal coordinate calculations on the extended  $n$ -paraffins  $\text{C}_2\text{H}_6$  through  $n\text{-C}_{14}\text{H}_{30}$  and polyethylene, using a perturbation method which adjusted force constants for any number of molecules simultaneously. Gayles, et al. [62] assigned their vibrational wavenumbers based upon the observed band types and rotational structure, frequency shifts upon deuterium substitution, published assignments for the propanes and previous work on the characteristic normal modes of the methyl group [71, 72]. Recently, Shimanouchi [21] critically reviewed the reported spectra data and assigned a complete set of vibrational wavenumbers (see table 10). His reported values were adopted for evaluation of the thermodynamic properties for propane, except for the two torsional wavenumbers,  $\nu_{14}$  and  $\nu_{27}$ .

The two torsional frequencies in propane have been directly measured at  $217 \pm 8$  and  $265 \pm 8$  cm<sup>-1</sup> recently by Grant, et al. [74] with inelastically scattered neutrons using the "small k" method of neutron molecular spectroscopy (NMS). Not subject to the same selection rules as optical spectroscopy, NMS is capable of observing

vibrational transitions which are otherwise forbidden by molecular symmetry. The vibrational frequencies were interpreted by numerically solving the wave equation based upon a two-dimensional Fourier expansion of the potential energy. They obtained a  $V_3 = 3680 \pm 190$  cal mol<sup>-1</sup> for individual methyl-group librations and a  $V_6 = -280 \pm 130$  cal mol<sup>-1</sup> term which couples the two methyl tops. The microwave spectrum of propane in the ground torsional state has been observed by Lide [49], and he did not observe any splitting due to internal rotation. He concluded that the barrier height for internal rotation of a methyl group must be at least 2700 cal mol<sup>-1</sup>. This value is consistent with the other reported  $V_3$  barriers, as presented in table 12. Hirota, et al. [64] observed the microwave spectrum of propane in the first-excited states of the  $\text{CH}_3$  torsion. The top-top interaction and its effect on the rotational spectra were treated by perturbation theory. They reported that the rotational barrier is described by a threefold parameter  $V_3 = 3325 \pm 20$  cal mol<sup>-1</sup> and a coupling parameter  $V_6 = -170 \pm 20$  cal mol<sup>-1</sup>. The two methyl torsional frequencies were calculated as 216.1 and 270.7 cm<sup>-1</sup>.

Hoyland [65] reanalyzed the data of Hirota, et al. [64] and carried out quantum-mechanical calculations on various conformations of the propane molecule using a Gaussian basis set and the Hartree-Fock-Roothaan Formalism. He obtained  $V_3 = 3575 \pm 100$  cal mol<sup>-1</sup> and a coupling term  $V_6 = -310 \pm 40$  cal mol<sup>-1</sup>. These values imply  $\nu_{14} = 216.2$  and  $\nu_{27} = 268.1$  cm<sup>-1</sup>. By the high-pressure-long-path-length technique, Weiss and Leroi observed a weak peak at 264 cm<sup>-1</sup>, which they assigned to the  $B_2$  torsional frequency, and determined  $V_3 = 2945$  and  $V_6 = -283$  cal mol<sup>-1</sup> [74]. The reported methyl torsional frequencies and internal rotation potential barrier in propane are summarized in tables 11 and 12.

Using their values for the  $A_2$  and  $B_2$  torsional modes, Grant, et al. [74] obtained the first five propane torsional energy levels as: 0, 217.1, 265.5, 431.1, 465.5, and 523.7 cm<sup>-1</sup>. The higher energy states were not calculated because of the unknown tunneling which will perturb the higher states. Evaluation of additional Fourier constants in the potential-energy expansion must await additional spectroscopic data concerning higher vibrational levels. Since we cannot now obtain the complete set of propane torsional energy levels derived from the two-dimensional potential function including the coupling effect of the two methyl tops, the partition function was calculated on the assumption of two independent one-dimensional internal rotations with identical barrier heights.

The thermodynamic properties of propane listed in table 15 were calculated from the selected molecular parameters given in tables 10, 11, 12, and 14 for the rigid-rotor harmonic-oscillator model. The partition function for internal rotation was calculated from two identical sets of 113 levels from 0 to 20300 cm<sup>-1</sup> gen-

erated from a computer solution of the Schroedinger equation [32].

The values of  $\Delta H_f^\circ$ ,  $\Delta G_f^\circ$  and  $\log K_f$  in the temperature range from 0 to 1500 K were derived using  $(H^\circ - H_0^\circ)$  and  $-(G^\circ - H_0^\circ)/T$  for C (c) and H<sub>2</sub> (g) from [33], for C<sub>3</sub>H<sub>8</sub> (g) from this work, and  $\Delta H_f^\circ$  (298.15, C<sub>3</sub>H<sub>8</sub>, g) = -24.82 kcal mol<sup>-1</sup> from [34]. The results are shown in table 15.

#### 4. Discussion

Ethane is an important raw material used extensively in petrochemical industries for manufacture of various organic chemicals. The low temperature heat capacities (15.5–184.1 K), melting point (89.87 ± 0.1 K), and enthalpies of fusion (682.9 ± 0.7 cal mol<sup>-1</sup>) and vaporization (3514 ± 3.5 cal mol<sup>-1</sup>) were measured by Witt and Kemp [35] in 1937. They derived the third law entropy of ethane (g) at the boiling point (184.1 ± 0.1 K) to be 49.38 ± 0.15 cal K<sup>-1</sup> mol<sup>-1</sup>. Adopting a correction for gas imperfection as 0.16 cal K<sup>-1</sup> mol<sup>-1</sup>, they obtained the entropy of ideal gas at the boiling point as 49.54 cal K<sup>-1</sup> mol<sup>-1</sup>. Using a correction for the gas imperfection as 0.19 cal K<sup>-1</sup> mol<sup>-1</sup> [34], we recalculated the value of  $S^\circ$  (184.1, C<sub>2</sub>H<sub>6</sub>, g) to be 49.57 ± 0.15 cal K<sup>-1</sup> mol<sup>-1</sup>. The  $S^\circ$  value obtained by the present statistical thermodynamic calculation is 49.48 cal K<sup>-1</sup> mol<sup>-1</sup> which is in agreement with the third law entropy value within the experimental uncertainty.

The heat capacities of ethane (g) in the temperature range 92.4–603.2 K have been determined by many investigators; the most reliable of these studies are listed in table 7. Table 8 compares the selected experimental values with the calculated ones. In consideration of the precision of the experimental equipment available at that time, the agreement between the  $C_p^\circ$  values from this work and the measured ones as listed in table 8 are excellent. For consistency, we converted the reported real gas heat capacities for ethane to ideal gas heat capacities by use of the gas imperfection corrections from the API Research Project 44 Tables [34]. It is interesting to note that our gas imperfection corrections for the  $C_p$  data of Dailey and Felsing [46] are virtually the same as those used by the authors in 1943. The results in table 8 illustrate the good agreement of our calculated thermodynamic properties for ethane (g) with the available experimental measurements.

Kemp and Egan [67] measured the low temperature heat capacities, 13.29–231.04 K, for propane. They also determined enthalpies of fusion (842.2 ± 0.8 cal mol<sup>-1</sup>), and vaporization (4487 ± 4 cal mol<sup>-1</sup>) and the melting (85.45 ± 0.05 K) and boiling (231.04 ± 0.05 K) points. Based on these data they derived the entropies of the real and ideal gas at the boiling point as 60.29 ± 0.1 and 60.45 cal K<sup>-1</sup> mol<sup>-1</sup>, respectively. Using the value of the correction for the gas imperfection, 0.21 cal K<sup>-1</sup> mol<sup>-1</sup> from API 44 tables [34], instead of 0.16 cal K<sup>-1</sup> mol<sup>-1</sup> used by Kemp and Egan, we recalculated the entropy of propane in the ideal gaseous state at 1 atm

and 231.04 K to be 60.50 ± 0.12 cal K<sup>-1</sup> mol<sup>-1</sup> which is in excellent agreement with our entropy value of 60.50 ± 0.1 cal K<sup>-1</sup> mol<sup>-1</sup> obtained from statistical thermodynamic calculations. The entropy value calculated by Pitzer [63] was 60.46 ± 0.1 cal K<sup>-1</sup> mol<sup>-1</sup>. He also reported the entropy of propane at 180.0 K as 57.04 ± 0.1 cal K<sup>-1</sup> mol<sup>-1</sup> while our value is 57.11 cal K<sup>-1</sup> mol<sup>-1</sup>.

The heat capacities of propane (g) have been determined by Beccck (273–573 K) [42], Sage, et al. (294–444 K) [73], Kistiakowsky, et al. (148–258 K) [70], Kistiakowsky and Rice (272–369) [69], and Dailey and Felsing (344–693 K) [46]. For consistency, their reported  $C_p$  values were converted to those in the ideal gaseous state ( $C_p^\circ$ ) by using the values of the correction for the gas imperfection obtained from API 44 tables [34]. The results are compared with our calculated values as shown in table 16. The agreement between our values and those measured by Kistiakowsky, et al. [70], Kistiakowsky and Rice [69], and Dailey and Felsing [46] is reasonably good.

The calculated heat capacities and entropies for ethane and propane are compared with those given in the API 44 tables [34] in table 17.

The calculation of all the energy levels associated with the internal rotations of the two methyl groups in propane is still in an unsatisfactory state. In the normal coordinate treatment this torsional motion is described in terms of two symmetry coordinates which account for the two harmonic oscillations identified by  $\nu_{14}$  and  $\nu_{27}$  in table 10. The frequencies of these two oscillations, and their corresponding reduced moments of inertia, are different even when it is assumed that there are no cross terms in the potential energy for the internal rotations of the two groups. Grant, et al. [74], and others, have shown, moreover, that the energy levels are significantly perturbed by the effect of interaction terms in the potential energy. An exact set of energy levels is not necessarily required for a reliable statistical calculation of thermodynamic properties since the interactions tend to split levels in a roughly symmetric way. Thus the effect on the partition function will be small as long as the perturbation is much smaller than  $kT$ .

Table 13 shows a comparison of heat capacity and entropy calculated from several approximate sets of energy levels for internal rotation. In each case two sets of levels for a single internal rotor whose potential energy is given by the usual three-fold cosine function of rotational angle were used. In the first three lines, two identical sets, corresponding to the fundamental frequency listed, were used. The value of 210 cm<sup>-1</sup> is the average of  $\nu_{14}$  and  $\nu_{27}$  obtained by Schachtschneider and Snyder [61] while 241 cm<sup>-1</sup> is the average of the two values reported by Grant, et al. [74]. The results for 241 cm<sup>-1</sup> agree better with the experimental heat capacity and entropy than do those for 210 cm<sup>-1</sup>. The results obtained when two different sets of levels, corresponding to the  $\nu_{14}$  and  $\nu_{27}$  of Grant, et al., were used are given in the

fourth line. These do not differ from the third line by an experimentally significant amount. An additional series of calculations of thermodynamic properties in the range from 50 to 200 K was also made with about 12 internal rotational levels. Higher levels do not contribute at these temperatures. These sets of levels were obtained by combining the first five levels of Grant, et al., with higher levels approximated in various ways, including those corresponding to table 13. No significant differences were obtained in the results. Hirota, et al. [64] tried to neglect the top-top coupling and calculated the  $\nu(0 \rightarrow 1)$  transition as  $243.4 \text{ cm}^{-1}$  and  $V_0 = 3.325 \text{ kcal mol}^{-1}$  which are in reasonable agreement with our selected values.

### 5. Acknowledgment

This study was partially supported by the American Petroleum Institute Research Project 44 of the Thermodynamics Research Center and the Texas Engineering Experiment Station of Texas A&M University.

### 6. References

- [1] Taylor, B. N., Parker, W. H., and Langenberg, D. N., *Rev. Mod. Phys.* **41**, 375 (1969).
- [2] Kemp, J. D., and Pitzer, K. S., *J. Chem. Phys.* **4**, 749 (1936).
- [3] Kemp, J. D., and Pitzer, K. S., *J. Amer. Chem. Soc.* **59**, 276 (1937).
- [4] Hedberg, K., and Schomaker, V., *J. Amer. Chem. Soc.* **73**, 1482 (1951).
- [5] Almenningsen, A., and Bastiansen, O., *Acta Chem. Scand.* **9**, 815 (1955).
- [6] Hansen, G. E., and Dennisen, D. M., *J. Chem. Phys.* **20**, 313 (1952).
- [7] Allen, H. C., Jr., and Plyler, E. K., *J. Chem. Phys.* **31**, 1062 (1959).
- [8] Lepard, D. W., Sweeney, D. M. C., and Welsh, H. L., *Can. J. Phys.* **40**, 1567 (1962).
- [9] Shaw, D. E., Lepard, D. W., and Welsh, H. L., *J. Chem. Phys.* **42**, 3736 (1965).
- [10] Smith, L. G., *J. Chem. Phys.* **17**, 139 (1949).
- [11] Van Riet, R., *Bull. Roy. Acad. Belg.* **188** (1955).
- [12] Lafferty, W. J., and Plyler, E. K., *J. Chem. Phys.* **37**, 2688 (1962).
- [13] Lafferty, W. J., and Plyler, E. K., *J. Res. Nat. Bur. Stand.* **A67**, 225 (1963).
- [14] Lysne, P. C., and Meister, A. G., *J. Chem. Phys.* **48**, 918 (1968).
- [15] Weiss, S., and Leroy, G. E., *J. Chem. Phys.* **48**, 962 (1968).
- [16] Cole, A. R. H., Lafferty, W. J., and Thibault, R. J., *J. Mol. Spectrosc.* **29**, 365 (1969).
- [17] Romanko, J., Feldman, T., and Welsh, H. L., *Can. J. Phys.* **33**, 588 (1955).
- [18] Stoitcheff, B. P., *Can. J. Phys.* **40**, 358 (1962).
- [19] Lepard, D. W., Shaw, D. E., and Welsh, H. L., *Can. J. Phys.* **44**, 2353 (1966).
- [20] Shaw, D. E., and Welsh, H. L., *Can. J. Phys.* **45**, 3823 (1967).
- [21] Shimanouchi, T., *NSRDS-NBS*—39, 1972.
- [22] Durig, J. R., Craven, S. M., and Bragin, J., *J. Chem. Phys.* **53**, 38 (1970).
- [23] Pitzer, R. M., *J. Chem. Phys.* **47**, 965 (1967).
- [24] Lide, D. R., Jr., *J. Chem. Phys.* **29**, 1426 (1958).
- [25] Pitzer, K. S., *Discuss. Faraday Soc.* **10**, 66 (1951).
- [26] Kistiakowsky, G. B., Lacher, J. R., and Stitt, F., *J. Chem. Phys.* **7**, 289 (1939).
- [27] Howard, J. B., *J. Chem. Phys.* **5**, 451 (1937).
- [28] Eyring, H., *J. Amer. Chem. Soc.* **54**, 3191 (1932).
- [29] Wilson, E. B., Jr., *J. Chem. Phys.* **6**, 740 (1938).
- [30] Lambert, J. D., and Salter, R., *Proc. Roy. Soc. (London)* **253**, 277 (1959).
- [31] Strong, K. A., and Brugger, R. M., *J. Chem. Phys.* **47**, 421 (1967).
- [32] Lewis, J. D., Malloy, T. B., Jr., Chao, T. H., and Laane, J., *J. Mol. Struct.* **12**, 427 (1972).
- [33] Zwolinski, B. J., et al., "Selected Values of Properties of Chemical Compounds," Thermodynamics Research Center Data Project, Thermodynamics Research Center, Texas A&M University, College Station, Texas 77843 (loose-leaf data sheets, extant 1972).
- [34] Zwolinski, B. J., et al., "Selected Values of Properties of Hydrocarbons and Related Compounds," American Petroleum Institute Research Project 44, Thermodynamics Research Center, Texas A&M University, College Station, Texas 77843 (loose-leaf data sheets, extant 1972).
- [35] Witt, R. K., and Kemp, J. D., *J. Amer. Chem. Soc.* **59**, 273 (1937).
- [36] Scheel, K., and Heuse, W., *Ann. Phys. (4)* **37**, 79 (1912); **40**, 473 (1913).
- [37] Heuse, W., *Ann. Phys.* **59**, 86 (1919).
- [38] Partington, J. R., and Shilling, W. G., "The Specific Heats of Gases," Ernest Benn Limited, London, 1924.
- [39] Thayer, V. R., "with" Stegemann, G., *J. Phys. Chem.* **35**, 1505 (1931).
- [40] Eucken, A., and Parts, A., *Z. Phys. Chem.* **B20**, 184 (1933).
- [41] Eucken, A., and Weigert, K., *Z. Phys. Chem.* **B23**, 265 (1933).
- [42] Beeck, O., *J. Chem. Phys.* **4**, 680 (1936).
- [43] Kistiakowsky, G. B., and Nazmi, F., *J. Chem. Phys.* **6**, 18 (1938).
- [44] Kistiakowsky, G. B., Lacher, J. R., and Stitt, F., *J. Chem. Phys.* **7**, 289 (1939).
- [45] Kistiakowsky, G. B., and Rice, W. W., *J. Chem. Phys.* **7**, 281 (1939).
- [46] Dailey, B. P., "with" Felsing, W. A., *J. Amer. Chem. Soc.* **65**, 42 (1943).
- [47] Wierl, R., *Ann. Phys. (5)* **13**, 453 (1932).
- [48] Bauer, S. H., *J. Chem. Phys.* **4**, 406 (1936).
- [49] Lide, D. R., Jr., *J. Chem. Phys.* **33**, 1514 (1960).
- [50] Bartholome, E., *Z. Phys. Chem.* **B23**, 152 (1933).
- [51] Elyashevich, M. A., and Stepanov, B. I., *Dokl. Akad. Nauk SSSR* **32**, 481 (1941).
- [52] Gates, D. M., *J. Chem. Phys.* **17**, 393 (1949).
- [53] McMurry, H. L., Thornton, V., and Condon, F. E., *J. Chem. Phys.* **17**, 918 (1949).
- [54] McMurry, H. L., and Thornton, V., *J. Chem. Phys.* **19**, 1014 (1951).
- [55] Comeford, J. J., and Gould, J. H., *J. Mol. Spectrosc.* **5**, 474 (1960).
- [56] Snyder, R. G., and Schachtschneider, J. H., *Spectrochim. Acta* **19**, 85 (1963).
- [57] Wu, V. L., and Barker, E. F., *J. Chem. Phys.* **9**, 487 (1941).
- [58] Gayles, J. N., Jr., and King, W. T., *Spectrochim. Acta* **21**, 543 (1965).
- [59] Kohlrausch, K. W. F., and Köppl, F., *Z. Phys. Chem.* **B26**, 209 (1934).
- [60] Takahashi, H., *Nippon Kagaku Zasshi*, **82**, 1304 (1961).
- [61] Schachtschneider, J. H., and Snyder, R. G., *Spectrochim. Acta* **19**, 117 (1963).
- [62] Gayles, J. N., Jr., King, W. T., and Schachtschneider, J. H., *Spectrochim. Acta* **23A**, 703 (1967).
- [63] Pitzer, K. S., *J. Chem. Phys.* **12**, 310 (1944).
- [64] Hirota, E., Matsumura, C., and Morino, Y., *Bull. Chem. Soc. Jap.* **40**, 1124 (1967).
- [65] Hoyland, J. R., *J. Chem. Phys.* **49**, 1908 (1968).
- [66] Pitzer, K. S., *J. Chem. Phys.* **5**, 473 (1937).
- [67] Kemp, J. D., and Egan, C. J., *J. Amer. Chem. Soc.* **60**, 1521 (1938).

- [68] Pitzer, K. S., Chem. Rev. **27**, 39 (1940).  
 [69] Kistiakowsky, G. B., and Rice, W. W., J. Chem. Phys. **8**, 610 (1940).  
 [70] Kistiakowsky, G. B., Lacher, J. R., and Ransom, W. W., J. Chem. Phys. **8**, 970 (1940).  
 [71] King, W. T., and Crawford, B., Jr., J. Mol. Spectrosc. **5**, 421 (1960).  
 [72] King, W. T., and Crawford, B., Jr., J. Mol. Spectrosc. **8**, 58 (1962).  
 [73] Sage, B. H., Webster, D. C., and Lacey, W. N., Ind. Eng. Chem. **29**, 1309 (1937).  
 [74] Grant, D. M., Pugmire, R. J., Livingston, R. C., Strong, K. A., McMurtry, H. L., and Brugger, R. M., J. Chem. Phys. **52**, 4424 (1970).

TABLE 1. Structural data for ethane

Structural parameter <sup>a</sup>	Reference			
	Almenningen and Bastiansen (1955) <sup>b</sup>	Allen and Plyler (1959) <sup>c</sup>	Lepard, et al. (1962) <sup>d</sup>	Shaw, et al. (1965) <sup>e</sup>
$r(\text{C-H})$	1.107	(1.093) <sup>f</sup>	$1.0914 \pm 0.0003$	$1.095 \pm 0.002$
$r(\text{C-C})$	1.536	1.534	$1.5362 \pm 0.0005$	$1.534 \pm 0.002$
$\angle \text{HCH}$	109.3°	109.75	$108.01 \pm 0.06$	$107.8 \pm 0.2$

<sup>a</sup> Bond distances,  $10^{-8}$  cm; bond angle, degrees.<sup>b</sup> Reference [5]; electron diffraction method.<sup>c</sup> Reference [7]; infrared spectrum.<sup>d</sup> Reference [8]; Raman spectrum.<sup>e</sup> Reference [9]; Raman spectrum. These values were adopted.<sup>f</sup> Estimated value.TABLE 2. Vibrational assignments for ethane ( $\text{cm}^{-1}$ )

Mode no.	Symmetry species	Reference		
		Smith (1949) <sup>b</sup>	Lepard, et al. (1966) <sup>c</sup>	Shimanouchi (1972) <sup>a, d</sup>
$\nu_1$	$a_{1g}$	2955.1 (b) 2899.2 (a)	2953.7	2954
$\nu_2$	$a_{1g}$	1375	1388.4	1388
$\nu_3$	$a_{1g}$	993	994.8	995
$\nu_4$	$a_{1u}$	290	289	289
$\nu_5$	$a_{2u}$	2954 (b) 2895.7 (a)	2895.8	2896
$\nu_6$	$a_{2u}$	1379.14	1379.19	1379
$\nu_7$	$e_u$	2995.5	2995.5	2969
$\nu_8$	$e_u$	1472.2	1472.2	1468
$\nu_9$	$e_u$	821.52	821.60	1190
$\nu_{10}$	$e_g$	2963 (b) 2939.5 (a)	2968.69	2985
$\nu_{11}$	$e_g$	1460	1468.1	1469
$\nu_{12}$	$e_g$	1190	1190	822

<sup>a</sup> The reported values were adopted in this work.<sup>b</sup> Reference [10].<sup>c</sup> Reference [19].<sup>d</sup> Reference [21].TABLE 3. Determinations of the torsional frequency ( $\nu_4$ ) in ethane<sup>a</sup>

Experimental method	$\nu_4$ , $\text{cm}^{-1}$	Reference
Combination bands:		
Infrared.....	290	[10]
	290	[24]
Raman.....	289	[19]
	278.4	[17]
Low-temperature $C_p$ measurements.....	280	[29]
	275	[26]
Ultrasonic relaxation data.....	290	[30]
Slow-neutron inelastic scattering.....	$277 \pm 8$	[31]
Infrared spectrum.....	289	[15]

<sup>a</sup> This work adopted the value  $289 \text{ cm}^{-1}$  for the torsional wave-number  $\nu_4$ .TABLE 4. Reported potential barriers to internal rotation in ethane<sup>a</sup>

$V_0$ , kcal $\text{mol}^{-1}$	Investigator (year)	Reference
2.96	Durig, et al. (1970)	[22]
$2.928 \pm 0.025$	Weiss and Leroi (1968)	[15]
3.5	R. M. Pitzer (1967)	[23] <sup>b</sup>
3.03	D. R. Lide (1958)	[24]
$2.875 \pm 0.125$	K. S. Pitzer (1951)	[25]
2.75	Kistiakowsky, et al. (1939)	[26]
3.00	E. B. Wilson (1938)	[29]
3.15	Kemp and Pitzer (1937)	[2]
$\geq 2.00$	J. B. Howard (1937)	[27]
0.36	H. Eyring (1932)	[28]

<sup>a</sup> This work derived  $V_0 = 2.96 \text{ kcal mol}^{-1}$  based on the adopted value of torsional frequency =  $289 \text{ cm}^{-1}$ .<sup>b</sup> The reported value was theoretically calculated. In this paper a summary of other theoretically calculated  $V_0$  values is given and references quoted therein.

TABLE 5. Molecular constants for ethane

Molecular weight.....	30.070
Point group.....	D <sub>3d</sub>
Symmetry number.....	18
Ground state configuration.....	<sup>1</sup> A <sub>1g</sub>
Product of the three principal moments of inertia, g <sup>2</sup> cm <sup>6</sup> .....	1.8708 × 10 <sup>-116</sup>
Reduced moment, g cm <sup>2</sup> .....	2.6203 × 10 <sup>-40</sup>
Potential barrier height, kcal mol <sup>-1</sup> ....	2.96
Vibrational wavenumbers, cm <sup>-1</sup> .....	2954, 1388, 995, 2896, 1379, 2969 (2), 1468 (2), 1190 (2), 2985 (2), 1469 (2), 822 (2)

TABLE 6. Ideal gas thermodynamic properties for ethane

<i>T</i>	<i>H</i> <sup>°</sup> - <i>H</i> <sub>0</sub> <sup>°</sup>	$\frac{(C_p - H_0^{\circ})}{T}$	$\frac{(H^{\circ} - H_0^{\circ})}{T}$	<i>S</i> <sup>°</sup>	<i>C</i> <sub>p</sub> <sup>°</sup>	Δ <i>H</i> <i>f</i> <sup>°</sup>	Δ <i>G</i> <i>f</i> <sup>°</sup>	log <i>K</i> <i>f</i>
<i>K</i>	cal mol <sup>-1</sup>	cal K <sup>-1</sup> mol <sup>-1</sup>			kcal mol <sup>-1</sup>			
0	0	0	0	0	0	-16.50	-16.50	infinite
50	398	30.32	7.953	38.27	7.98	-17.62	-16.40	71.68
100	808	35.06	8.002	43.94	8.52	-18.00	-15.04	32.86
150	1252	39.19	8.348	47.53	9.24	-18.48	-13.45	19.60
200	1735	41.63	8.673	50.31	10.10	-19.04	-11.69	12.78
273.15	2534	44.42	9.276	53.70	11.84	-19.94	-8.88	7.10
298.15	2838	45.24	9.520	54.76	12.54	-20.24 <sup>a</sup>	-7.84	5.74
300	2862	45.30	9.538	54.84	12.60	-20.26	-7.76	5.65
400	4272	48.20	10.681	58.88	15.65	-21.43	-3.41	1.86
500	5989	50.72	11.978	62.70	18.64	-22.45	1.22	-0.53
600	7990	53.02	13.317	66.34	21.33	-23.30	6.03	-2.20
700	10245	55.17	14.635	69.81	23.71	-24.00	10.98	-3.43
800	12723	57.21	15.903	73.11	25.81	-24.56	16.01	-4.38
900	15398	59.15	17.109	76.26	27.67	-24.99	21.11	-5.13
1000	18249	61.02	18.249	79.26	29.30	-25.30	26.26	-5.74
1100	21252	62.81	19.320	82.13	30.73	-25.52	31.40	-6.24
1200	24390	64.53	20.325	84.86	31.99	-25.66	36.61	-6.67
1300	27644	66.20	21.265	87.46	33.08	-25.75	41.80	-7.03
1400	31002	67.80	22.144	89.95	34.04	-25.78	47.02	-7.34
1500	34449	69.36	22.966	92.33	34.87	-25.77	52.20	-7.61

1 cal = 4.184 J

<sup>a</sup> While this paper was in proof, Pittam and Pilcher reported a new value of -20.04 kcal mol<sup>-1</sup> for Δ*H**f*<sup>°</sup>(g, 298.15 K) ethane. J. Chem. Soc. Faraday Trans. 1 68, 2224 (1972).

TABLE 7. Summary of heat capacity measurements for ethane

Investigator	Number of data points	Temperature, K	Reference
Scheel and Heuse (1913).....	3	191.2-288.2	[36]
Heuse (1919).....	3	191.2-288.2	[37]
Parlington and Shilling (1924).....	3	283.2-373.2	[38]
Thayer and Stegeman (1931).....	4	275.4-337.6	[39]
Eucken and Parts (1933).....	7	189.1-373.5	[40]
Eucken and Weigert (1933).....	5	143 -280	[41]
Beeck (1936).....	4	273.2-573.2	[42]
Kistiakowsky and Nazmi (1938)...	2	143 -163	[43]
Kistiakowsky, et al. (1939).....	9	92.4-134.1	[44]
Kistiakowsky and Rice (1939).....	4	272.1-364.8	[45]
Dailey and Felsing (1943).....	7	347.7-603.2	[46]



TABLE 8. Comparison of calculated and measured heat capacities for ethane<sup>a</sup>

$T$ K	$C_p$	$C_p - C_p^\circ$	$C_p^\circ$		$C_p^\circ$ (calc.) - $C_p^\circ$ (exptl.)
			Experimental	Calculated	

## Dailey and Felsing (1943)

347.7	14.14	0.07	14.07	14.03	-0.04
359.8	14.49	0.06	14.43	14.40	-0.03
387.6	15.32	0.05	15.27	15.27	0.00
452.0	17.34	0.03	17.31	17.24	-0.07
520.6	19.16	0.02	19.14	19.22	+0.08
561.6	20.64	0.01	20.63	20.33	-0.30
603.2	21.63	0.01	21.62	21.41	-0.21

## Kistiakowsky and Rice (1939)

272.07	11.96	0.17	11.79	11.81	+0.02
302.70	12.82	0.12	12.70	12.67	-0.03
335.82	13.79	0.08	13.71	13.67	-0.04
364.78	14.64	0.06	14.58	14.56	-0.02

## Eucken and Parts (1933)

189.2	10.48	0.61	9.87	9.89	+0.02
209.3	10.68	0.46	10.22	10.29	+0.07
229.6	11.02	0.31	10.71	10.74	+0.03
249.9	11.48	0.23	11.25	11.23	-0.02
250.2	11.46	0.23	11.23	11.23	0.00
272.0	12.01	0.17	11.84	11.81	-0.03
292.0	12.57	0.16	12.41	12.36	-0.05
373.6	14.90	0.05	14.85	14.83	-0.02

## Heuse (1919)

191.2	10.44	0.60	9.84	9.93	+0.09
238.2	11.04	0.27	10.77	10.94	+0.17
288.2	12.40	0.13	12.27	12.26	-0.01

<sup>a</sup> Unit = cal K<sup>-1</sup> mol<sup>-1</sup>; 1 cal = 4.184 J.

TABLE 9. Structural data for propane

Structural parameter <sup>a</sup>	Reference		
	Wierl (1932) <sup>b</sup>	Bauer (1936) <sup>c</sup>	Lide (1960) <sup>d</sup>
$r(\text{C}-\text{C})$	$1.52 \pm 0.05$	$1.50 \pm 0.02$	$1.526 \pm 0.002$
$\angle \text{CCC}$	(109.47) <sup>e</sup>	114	$112.4 \pm 0.2$
$r(\text{C}-\text{H})$ , in $\text{CH}_2$	1.10	$1.08 \pm 0.02$	$1.096 \pm 0.002$
$\angle \text{HCH}$ , in $\text{CH}_2$	(109.47)		$106.1 \pm 0.2$
$r(\text{C}-\text{H})$ , in $\text{CH}_3$	1.10		$1.091 \pm 0.01$
$\angle \text{HCH}$ , in $\text{CH}_3$	(109.47)		$107.7 \pm 1.0$

<sup>a</sup> Bond distances, 10<sup>-8</sup> cm; bond angle, degrees.<sup>b</sup> Reference [47].<sup>c</sup> Reference [48].<sup>d</sup> Reference [49]; the reported values were adopted in this work.<sup>e</sup> Estimated value.

TABLE 10. Vibrational assignments for propane ( $\text{cm}^{-1}$ )

Mode no.	Symmetry species	Reference				
		Wu and Barker (1941) <sup>a</sup>	Pitzer (1944) <sup>b</sup>	Schachtschneider and Snyder (1963) <sup>c</sup>	Gayles and King (1965) <sup>d</sup>	Schimanouchi (1972) <sup>e</sup>
$\nu_1$	$a_1$	2966	2966	2966	2976.7	2977
$\nu_2$	$a_1$	2960	2960	2882	2961.7	2962
$\nu_3$	$a_1$	2914	2914	2856	2887.0	2887
$\nu_4$	$a_1$	1436	1460	1471	1476.0	1476
$\nu_5$	$a_1$	1468	1468	1445	1462.0	1462
$\nu_6$	$a_1$	1370	1370	1378	1391.9	1392
$\nu_7$	$a_1$	1149	1155	1151	1157.5	1158
$\nu_8$	$a_1$	867	868	870	869.3	869
$\nu_9$	$a_1$	375	375	382	369.2	369
$\nu_{10}$	$a_2$		2970	2964		2967
$\nu_{11}$	$a_2$	1436	1450	1459		1451
$\nu_{12}$	$a_2$	1361	1278	1279	(1292)	1278
$\nu_{13}$	$a_2$	870	940	903		940
$\nu_{14}$	$a_2$		(202)	200	(208)	216
$\nu_{15}$	$b_1$	2968	2968	2963	2968.2	2968
$\nu_{16}$	$b_1$	2942	2942	2882	2887.0	2887
$\nu_{17}$	$b_1$	1465	1470	1465	1464.0	1464
$\nu_{18}$	$b_1$	1375	1375	1367	1378.0	1378
$\nu_{19}$	$b_1$		1338	1342	1338.4	1338
$\nu_{20}$	$b_1$	1053	1053	1046	1053.8	1054
$\nu_{21}$	$b_1$	922	922	924	921.7	922
$\nu_{22}$	$b_2$	2980	2980	2965	2972.6	2973
$\nu_{23}$	$b_2$	2968	2968	2921	2968.2	2968
$\nu_{24}$	$b_2$	1443	1450	1464	1472.0	1472
$\nu_{25}$	$b_2$	1179	1179	1185	1191.5	1192
$\nu_{26}$	$b_2$	748	748	747	748.1	748
$\nu_{27}$	$b_2$		(283)	220	(223.0)	268

<sup>a</sup> Reference [57].<sup>b</sup> Reference [63].<sup>c</sup> Reference [61].<sup>d</sup> Reference [58].<sup>e</sup> Reference [21], the reported values adopted in this work except  $\nu_{14}$  and  $\nu_{27}$ .

TABLE 11. Summary of reported torsional frequencies in propane

Torsional frequency, $\text{cm}^{-1}$		Method of determination	Reference
$\nu_{14}$	$\nu_{27}$		
202	283	calculated	[63]
200	220	calculated	[61]
208	223	estimated from combination band	[58]
202	225	calculated	[62]
216.1	270.7	calculated	[64]
243	243	calculated	
216.2	268.1	calculated	[65]
217 $\pm$ 8	265 $\pm$ 8	directly measured	[74]
	264 <sup>a</sup>		
216	268	calculated	[21]
241	241	selected by this work	

<sup>a</sup> This transition was observed by Weiss and Leroy; quoted by Grant, et al. [74] as private communication.

TABLE 12. Reported potential barriers to internal rotation in propane

$V_o$	$V_a$	Investigator (year)	Reference
kcal mol <sup>-1</sup>			
3.40		Pitzer (1937)	[66]
3.30		Kemp and Egan (1938)	[67]
3.40		Pitzer (1940)	[68]
3.1 – 3.2		Kistiakowsky and Rice (1940)	[69]
3.30		Kistiakowsky, et al. (1940)	[70]
3.40		Pitzer (1944)	[63]
3.40		Pitzer (1951)	[25]
>2.70		Lide (1960)	[49]
3.325	– 0.170	Hirota, et al. (1967)	[64]
3.575	– 0.310	Hoyland (1968)	[65]
3.33		Durig, et al. (1970)	[22]
3.680	– 0.280	Grant, et al. (1970)	[74]
2.945	– 0.283	Weiss and Leroi <sup>a</sup>	
3.29		This work	

<sup>a</sup> Quoted by Grant, et al. [74] as private communication.

TABLE 13. Comparison of calculated  $C_p^\circ$  and  $S^\circ$  for propane<sup>a</sup>

Torsional frequency		Potential barrier	$C_p^\circ$			$S^\circ$		
$\nu_{14}$	$\nu_{27}$	$V_0$	298.15 K	500 K	1000 K	298.15 K	500 K	1000 K
210	210	2.55, 2.55	17.65	26.48	41.38	65.21	76.46	99.94
239	239	3.25, 3.25	17.60	26.89	41.71	64.61	75.96	99.73
241	241	3.29, 3.29	17.59	26.91	41.73	64.58	75.93	99.72
217	265	2.71, 3.94	17.54	26.84	41.75	64.60	75.92	99.70

<sup>a</sup> Units:  $\text{cm}^{-1}$  for torsional frequency;  $\text{kcal mol}^{-1}$  for potential barrier;  $\text{cal K}^{-1} \text{mol}^{-1}$  for  $C_p^\circ$  and  $S^\circ$ .

TABLE 14. Molecular constants for propane

Molecular weight.....	44.097
Point group.....	$C_{2v}$
Symmetry number.....	18
Ground state configuration.....	$1A_{1g}$
Product of the three principal moments of inertia, $I_A I_B I_C$ , $\text{g}^3 \text{cm}^6$ .....	$3.2148 \times 10^{-115}$
Reduced moment, $I_r$ , $\text{g cm}^2$ .....	$4.4202 \times 10^{-40}$
Potential barrier height, $\text{kcal mol}^{-1}$ .....	3.29
Vibrational wavenumber, $\text{cm}^{-1}$ .....	2977, 2962, 2887, 1476, 1462, 1392, 1158, 869, 369, 2967, 1451, 1278, 940, 2968, 2887, 1464, 1378, 1338, 1054, 922, 2973, 2968, 1472, 1192, 748

TABLE 15. Ideal gas thermodynamic properties for propane (g)

$T$	$H^\circ - H_0^\circ$	$\frac{(G^\circ - H_0^\circ)}{T}$	$\frac{(H^\circ - H_0^\circ)}{T}$	$S^\circ$	$C_p^\circ$	$\Delta H_f^\circ$	$\Delta G_f^\circ$	$\log K_f$
K	cal $\text{mol}^{-1}$	cal $\text{K}^{-1} \text{mol}^{-1}$			kcal $\text{mol}^{-1}$			
0	0	0	0	0	0	-19.50	-19.50	infinite
50	399	36.44	7.976	44.42	8.14	-21.11	-19.15	83.72
100	845	42.09	8.453	50.54	9.87	-21.73	-16.96	37.07
150	1385	45.66	9.231	54.89	11.66	-22.42	-14.43	21.02
200	2011	48.43	10.053	58.48	13.40	-23.21	-11.64	12.72
273.15	3098	51.75	11.341	63.09	16.43	-24.42	-7.24	5.79
298.15	3523	52.76	11.816	64.58	17.59	-24.82 <sup>a</sup>	-5.63	4.13
300	3555	52.83	11.852	64.68	17.67	-24.85	-5.51	4.01
400	5563	56.52	13.907	70.43	22.47	-26.36	1.18	-0.64
500	8036	59.85	16.072	75.93	26.91	-27.64	8.21	-3.59
600	10924	62.97	18.207	81.18	30.76	-28.68	15.48	-5.64
700	14171	65.94	20.245	86.18	34.10	-29.52	22.91	-7.15
800	17729	68.77	22.161	90.93	36.99	-30.16	30.45	-8.32
900	21558	71.48	23.953	95.43	39.52	-30.64	38.05	-9.24
1000	25623	74.09	25.623	99.72	41.73	-30.96	45.72	-9.99
1100	29895	76.61	27.178	103.79	43.66	-31.15	53.36	-10.60
1200	34348	79.04	28.623	107.66	45.35	-31.26	61.08	-11.12
1300	38958	81.38	29.967	111.35	46.81	-31.28	68.77	-11.56
1400	43704	83.65	31.217	114.86	48.09	-31.24	76.50	-11.94
1500	48570	85.84	32.380	118.22	49.21	-31.14	84.17	-12.26

1 cal = 4.184 J.

<sup>a</sup> While this paper was in proof, Pittam and Pilcher reported a new value of  $-25.02 \text{ kcal mol}^{-1}$  for  $\Delta H_f^\circ(\text{g}, 298.15 \text{ K})$  propane. J. Chem. Soc. Faraday Trans. 1 68, 2224 (1972).

TABLE 16. Comparison of calculated and measured heat capacities for propane (g)<sup>a</sup>

T K	$C_p$	$C_p - C_p^\circ$	$C_p^\circ$		$C_p^\circ$ (calc.) - $C_p^\circ$ (exptl.)
			Experimental	Calculated	
O. Beeck (1936)					
273.15	15.86	0.415	15.44	16.43	+0.99
373.15	20.26	0.125	20.14	21.19	+1.05
473.15	24.74	0.055	24.68	25.77	+1.09
573.15	28.71	0.026	28.68	29.78	+1.10
B. H. Sage, D. C. Webster and W. N. Lacey (1937)					
294.3	17.83	0.31	17.52	17.41	-0.11
310.9	18.31	0.25	18.06	18.19	+0.13
327.6	18.79	0.21	18.58	18.99	+0.41
344.3	19.28	0.17	19.11	19.80	+0.69
360.9	19.80	0.14	19.66	20.60	+0.94
377.6	20.32	0.12	20.20	21.40	+1.20
394.3	20.84	0.11	20.73	22.20	+1.47
410.9	21.36	0.09	21.27	22.98	+1.71
427.6	21.88	0.08	21.80	23.75	+1.95
444.3	22.40	0.06	22.34	24.50	+2.16
G. G. Kistiakowsky, J. R. Lacher and W. W. Ransom (1940)					
148.2			11.69	11.60	-0.09
157.8			12.04	11.93	-0.11
213.1			14.00	13.90	-0.10
219.0			14.17	14.13	-0.04
258.0			15.75	15.75	0.00
G. B. Kistiakowsky and W. W. Rice (1940)					
272.38	16.39	0.42	15.97	16.39	+0.42
300.37	17.73	0.25	17.48	17.69	+0.21
334.05	19.35	0.19	19.16	19.30	+0.14
368.55	20.97	0.13	20.84	20.97	+0.13
B. P. Dailey and W. A. Felsing (1943)					
343.7	19.80	0.17	19.63	19.77	+0.14
360.1	20.48	0.14	20.34	20.56	+0.22
387.8	21.72	0.11	21.61	21.89	+0.28
452.6	25.13	0.06	25.07	24.87	-0.20
521.0	27.93	0.04	27.89	27.77	-0.12
562.0	29.13	0.03	29.10	29.37	+0.27
603.3	30.78	0.02	30.76	30.88	+0.12
693.2	33.63	0.015	33.62	33.88	+0.26

<sup>a</sup> Unit = cal K<sup>-1</sup> mol<sup>-1</sup>; 1 cal = 4.184 J.TABLE 17. Comparison of calculated and API 44 tables 1952 values of  $C_p^\circ$  and  $S^\circ$  for ethane and propane<sup>a</sup>

T K	$C_p^\circ$ , cal K <sup>-1</sup> mol <sup>-1</sup>			$S^\circ$ , cal K <sup>-1</sup> mol <sup>-1</sup>		
	This work	API 44 1952 value	Difference	This work	API 44 1952 value	Difference
Ethane						
100	8.52	8.59	-0.07	43.94	43.94	0.00
298.15	12.54	12.58	-0.04	54.76	54.85	-0.09
700	23.71	23.72	-0.01	69.81	69.93	-0.12
1000	29.30	29.33	-0.03	79.26	79.39	-0.13
1500	34.88	34.90	-0.02	92.32	92.46	-0.14
Propane						
100	9.87	9.84	+0.03	50.54	50.52	+0.02
298.15	17.59	17.57	+0.02	64.58	64.51	+0.07
700	34.10	34.20	-0.10	86.18	86.17	+0.01
1000	41.73	41.83	-0.10	99.72	99.77	-0.05
1500	49.21	49.26	-0.05	118.22	118.29	-0.07

<sup>a</sup> Average percentage difference in  $C_p^\circ$  for ethane and propane, respectively: 100-700 K: -0.32 percent, -0.01 percent; 700-1500 K: -0.08 percent, -0.20 percent. Average percentage difference in  $S^\circ$  for ethane and propane, respectively: 100-700 K: -0.13 percent, +0.06 percent; 700-1500 K: -0.16 percent, -0.05 percent.