

Compilation and Evaluation of Solubility Data in the Mercury (I) Chloride-Water System

Cite as: Journal of Physical and Chemical Reference Data **9**, 1307 (1980); <https://doi.org/10.1063/1.555633>
Published Online: 15 October 2009

Y. Marcus



View Online



Export Citation

ARTICLES YOU MAY BE INTERESTED IN

Static Dielectric Constant of Water and Steam

Journal of Physical and Chemical Reference Data **9**, 1291 (1980); <https://doi.org/10.1063/1.555632>

The Solubility of Mercury and Some Sparingly Soluble Mercury Salts in Water and Aqueous Electrolyte Solutions

Journal of Physical and Chemical Reference Data **14**, 631 (1985); <https://doi.org/10.1063/1.555732>

The Solubility of Some Sparingly Soluble Salts of Zinc and Cadmium in Water and in Aqueous Electrolyte Solutions

Journal of Physical and Chemical Reference Data **21**, 941 (1992); <https://doi.org/10.1063/1.555909>

Where in the **world** is AIP Publishing?
Find out where we are exhibiting next



Compilation and Evaluation of Solubility Data in the Mercury (I) Chloride-Water System

Y. Marcus

Department of Inorganic and Analytical Chemistry, The Hebrew University, Jerusalem, Israel

The more than one dozen papers dealing with the solubility of mercury (I) chloride in water or in aqueous chloride solutions have been compiled in the format set by the IUPAC Solubility Data Project, and have been evaluated. Mercury (I) chloride dissolves in water, forming the following species: $\text{Hg}(\text{OH})_2$, HgCl_2 , HgOH^+ , HgCl^+ , Hg_2^{2+} and Hg_2OH^+ , in addition to H^+ and Cl^- . In excess chloride solutions it dissolves to give, mainly, HgCl_3^- and HgCl_4^{2-} . Thus, many homogeneous equilibria have to be considered beside the two heterogeneous ones: $\text{Hg}_2\text{Cl}_2(\text{s}) = \text{Hg}_2^{2+}(\text{aq}) + 2\text{Cl}^-(\text{aq})$ and $\text{Hg}_2^{2+}(\text{aq}) = \text{Hg}^{2+}(\text{aq}) + \text{Hg}(\ell)$, of which K_{s0} and $(K_r)^{-1}$, respectively, are the equilibrium constants. The papers in which the total solubility (sum of all the mercury containing aqueous species) and the solubility product (derived from e.m.f. data) are reported do not give as accurate and reliable quantities as are obtained from the appropriate standard electrode potentials. The following values are recommended as valid at 298.15 K: $\log(K_{s0}/\text{mol}^2\text{kg}^{-2}) = -17.844 \pm 0.017$, $d \log(K_{s0}/\text{mol}^2\text{kg}^{-2})/dT = (0.0622 \pm 0.0002) - (6.0 \pm 0.4) \times 10^{-4} (\text{K}^{-1})$, $\Delta G_{s0}^\circ = 101.86 \pm 0.10 \text{ kJ mol}^{-1}$, $\Delta S_{s0}^\circ = -12.7 \pm 0.9 \text{ JK}^{-1} \text{ mol}^{-1}$, $\Delta H_{s0}^\circ = 98.08 \pm 0.18 \text{ kJ mol}^{-1}$, $\Delta C_{p,s0}^\circ = -0.36 \pm 0.04 \text{ JK}^{-1} \text{ mol}^{-1}$ (this item, tentatively), and $c_{\text{Hg}} = (8.4 \pm 1.6) \times 10^{-6} \text{ mol dm}^{-3}$ (the total aqueous solubility).

Key words: Compilation of solubility data; disproportionation of mercury (I); electromotive force measurements; mercury (I) chloride; solubility; solubility product; standard electrode potentials; standard thermodynamic functions.

Contents

	Page		Page
1. Introduction	1307	5.3. Data of Sherrill (1903)	1317
2. Evaluation	1308	5.4. Data of Ley and Heimbucher (1904)	1318
2.1. Solubility Measurements	1308	5.5. Data of Kohlrausch (1908)	1319
2.2. E.m.f. Measurements	1309	5.6. Data of Herz (1911)	1320
2.3. Standard Electrode Potentials	1310	5.7. Data of Brodsky and Scherschewer (1926) ..	1321
2.4. Calculated Solubility	1311	5.8. Data of Eversole (1932)	1322
3. Recommended Values	1311	5.9. Data of Garrett, Noble and Miller (1942) ..	1323
3.1. Solubility Product Constant of Hg_2Cl_2	1311	5.10. Data of Law (1946)	1324
3.2. Standard Thermodynamic Functions	1312	5.11. Data of Jonsson, Qvarfort and Sillén (1947) ..	1325
3.3. The Aqueous Solubility of Hg_2Cl_2	1313	5.12. Data of Dry and Gledhill (1955)	1326
4. References	1313	5.13. Data of Galloway (1961)	1327
5. Compilation	1314	5.14. Data of Hansen, Izatt and Christensen	
5.1. Data of Behrend (1893, 1894)	1314	(1963)	1328
5.2. Data of Richards and Archibald (1902) ...	1315	5.15. Other data	1329

1. Introduction

The Subcommittee on Solubility Data of Commission V.6, Equilibrium Data, Analytical Chemistry Division, International Union of Pure and Applied Chemistry, (secretariat: Oxford OX4 3YF, UK), is currently undertaking an extensive project of compiling and evaluating solubility data existing in the literature. As a part of this project, systems involving the

solubility of solids in liquids are being treated, and it became necessary to provide prospective authors of compilations and evaluations of such systems with a reasonably complicated, well worked out system as a sample. This led to the commissioning of the author with selecting such a system and providing this sample compilation and evaluation.

The mercury (I) chloride-water system appears to fulfill the requirements of being quite well documented in the literature, of providing reasonably complicated side effects which must be, and can be, dealt with, yet being amenable to a sufficiently rigorous treatment, so as to provide final, recommended

© 1981 by the U.S. Secretary of Commerce on behalf of the United States. This copyright is assigned to the American Institute of Physics and the American Chemical Society.

values. It was therefore selected for presentation as a sample compilation and evaluation.


Three groups of papers have to be examined in order to obtain definite values for the solubility of mercury (I) in water. The first, refs. [1]–[7], deal with direct determinations of the solubility. The second, refs. [3], [8]–[14], deal with determination of the solubility product via e.m.f. measurements. The third, refs. [15], [18], [19], [21]–[28], pertains to papers in which the standard potentials of the calomel electrode and of the mercury/mercury (I) electrode are determined separately, without going on to calculate the solubility product. This last group of papers is not included in the compilation on solubilities. It turns out however, that it provides, over a temperature range, the best values of the solubility.

2. Evaluation

2.1. Solubility Measurements

Kohlrausch [1a]–[1c]¹ compared the specific conductivity of saturated mercury (I) chloride solutions at 0.5, 18.0, 24.6 and ~43 °C with that of 0.0005 mol dm⁻³ mercury (I) nitrate (stated as 0.001N HgNO₃), knowing its temperature coefficient for conductivity. The conductivity of the water used in the saturated solutions was probably [6] overcorrected for, while on the other hand, the contribution of the hydrolysis to the conductivity, which is overwhelming [6], was disregarded. Hence the value found for the solubility, $s = (5.9 \pm 3.0) \times 10^{-6}$ mol dm⁻³ at 24.6 °C, is fortuitously of the right magnitude.

Richards and Archibald [2] measured the total solubility of mercury in saturated solutions of Hg₂Cl₂ in aqueous HCl, NaCl, and BaCl₂ at 25 °C, where HgCl₂²⁺ is the main mercury species, by precipitation as HgS. The equilibrium constant for the reaction



(the solubility product for producing the 4-ligand complex of mercury (II), i.e., K_{s114}) is obtained from


$$\begin{aligned} \log K_e^* &= \lim_{c_{\text{Cl}} \rightarrow 0} [\log c_{\text{Hg}}/c_{\text{Cl}}^2] \\ &- \log (1 + \beta_0\beta_1^2c_{\text{Cl}}^2) + \log \gamma_{\text{HgCl}_2^2+} \gamma_{\text{Cl}^-}^2 \end{aligned} \quad (2)$$

The second term on the rhs is small at the experimental chloride concentrations c_{Cl} employed, hence activity coefficient corrections to the stability constants β_i of HgCl₂²⁺ can be neglected. The third term on the rhs, however, is of considerable size, and can be approximated as $-2Ac_{\text{Cl}}^{1/2}/(1 + 1.6c_{\text{Cl}}^{1/2})^{-1} + \Delta b_{\text{Cl}}$. The first term of this approximation is known, and any deviations from the modified Debye-Hückel behavior are included in the second term, and is extrapolated out at the limit. The main uncertainty enters here, since as the Δb_{Cl} term decreases on extrapolation, the second term on the rhs of 2 increases. The solubility data themselves cannot be extrapolated to zero

excess chloride concentration to give the solubility of Hg₂Cl₂ in water. However, the solubility product can be calculated from

$$\begin{aligned} \log K_{s0}^* &= \log K_e^* + \log K_r^* - \log \beta_4 \\ &- \log \gamma_{\text{uncomplexed NaCl in NaClO}_4} \end{aligned} \quad (3)$$

where K_r^* is the reproporationation constant for the equilibrium



The values adopted for the constants are $\log K_r^* = 1.94 \pm 0.01$ [15] and $\log (\beta_4(\text{HgCl}_2^2+)/\text{mol}^{-4} \text{dm}^{12}) = 15.07$ [16], the latter valid for 0.5 mol dm⁻³ NaClO₄ medium. The main errors in K_{s0}^* arise from those in K_e^* (± 12 to 15%) and in β_4 ($\pm 15\%$). The final value $K_{s0}^* = (1.86 \pm 0.37) \times 10^{-18}$ mol²dm⁻⁶ at 25 °C will be seen to be consistent with (but on the high side of) the value recommended further on.

Sherrell [3] presented one datum point, for the solubility of Hg₂Cl₂ in 1 mol dm⁻³ NaCl at 25 °C. A value of K_e^* (equil. (1)) could be estimated by using the same value of Δb applicable to the NaCl data of Richards and Archibald [2] evaluated above. This led by means of (3) to $K_{s0}^* = 1.2 \times 10^{-18}$ mol²dm⁻⁶, with a large margin of uncertainty, but still consistent with the recommended value.

Eversole and McLachlan [4] determined the solubility of Hg₂Cl₂ in dilute acids, HClO₄ and HNO₃ at 25 °C. The only soluble species of importance which needs to be considered is HgCl₂, since the excess acid represses the hydrolysis. In aqueous solutions of Hg₂Cl₂, a major species is soluble Hg(OH)₂ [6], and its non-formation in the dilute acid solutions decreases the solubility of mercury (I) chloride below what it is in water. The lack of sufficient data prevents the calculation of this solubility, but the solubility product can be calculated, from the total solubilities c_{Hg} in the acids HA:

$$\begin{aligned} c_{\text{Hg}} &= [\text{Hg}_2^{2+}] + [\text{Hg}_2\text{A}^+] + [\text{HgCl}^+] + [\text{HgCl}_2] \\ &= (K_{s0}^*/4)^{1/3} \gamma_{\pm \text{Hg}_2\text{Cl}_2}^{-1} [1 + \beta_{1A} [A^-] + 2 (K_{s0}^*/4)^{1/3} \beta_1 K_r^{-1}] \\ &\quad + K_{s0}^* K_r^{-1} \beta_2^* (\gamma_{\pm \text{HgCl}_2} / \gamma_{\pm \text{Hg}_2\text{Cl}_2})^3 \end{aligned} \quad (5)$$

The first term on the rhs of (5) is a relatively small (~20%) correction term, in which unity dominates over the other terms in the square brackets, and for which K_{s0}^* is obtained iteratively. Hence, activity coefficient corrections for β_{1A} (Hg₂ClO₄⁺ or Hg₂NO₃⁺) and β_1 (HgCl⁺) are unnecessary, while $\log \gamma_{\pm \text{Hg}_2\text{Cl}_2} = -2 \times 0.51 [A^-]^{1/2} / (1 + 1.6[A^-]^{1/2})$. The value of the solubility product is therefore:

$$K_{s0}^* = \frac{c_{\text{Hg}} - \text{the first term in the rhs of (5)}}{K_r^{-1} \beta_2^* (\gamma_{\pm \text{HgCl}_2} / \gamma_{\pm \text{Hg}_2\text{Cl}_2})^3} \quad (6)$$

It is a good approximation to equate the activity coefficient ratio of (the ionically dissociated parts of) Hg₂²⁺ · 2Cl⁻ and Hg₂²⁺ · 2Cl⁻ to unity, and in any case no systematic deviation of K_{s0}^* calculated from (6) with $[A^-]$ could be seen. The precision of K_{s0}^* depends on that of c_{Hg} while its accuracy depends

¹ Figures in brackets indicate literature references.

mainly on those of K_f and β_2° . The values adopted for these constants, $\log K_f = 1.94 \pm 0.01$ [15] and $\log (\beta_2^\circ/\text{mol}^{-2}\text{dm}^6) = 14.26 \pm 0.09$ [6] are responsible for the overall accuracy of the constant $\log K_{s0}^\circ = (1.84 \pm 0.37) \times 10^{-18} \text{ mol}^2\text{dm}^{-6}$ at 25 °C, again consistent with the recommended value, but a bit high.

The data of Garret et al. [5] can be treated in a manner similar to the one used on the data of Richards and Archibald [2], eq (2) and (3), but since there are much fewer data, no independent extrapolations could be made. Therefore, the Δb values obtained [2] for NaCl and CaCl_2 solutions were used for the calculation of K_s° . The final value is $(1.12 \pm 0.23) \times 10^{-18} \text{ mol}^2\text{dm}^{-6}$ at 25 °C, which is somewhat on the low side of the recommended value. (The fact that the data are on the mol kg^{-1} scale produces a small complication, which can be overcome by using the known densities of the salt solutions and converting to mol dm^{-3} .)

The work of Dry and Gledhill [6] has been very carefully done, and gives the best value for the solubility determined directly, $s_{\text{Hg}_2\text{Cl}_2} = (7.5 \pm 0.3) \times 10^{-6} \text{ mol dm}^{-3}$ at 25 °C. The method used, treating the filtered saturated solution with dilute HCl and a solution of dithizone in CCl_4 , should produce dependable data, when compared with photometric readings from known solutions of HgCl_2 . The value is within the limits of the recommended value below, although on the low side. However, the primary data were not disclosed, and no basis for the estimate [6] of the $\pm 4\%$ error was given by the authors. Therefore there is no good basis for preferring this lower value, in spite of its apparently higher precision.

The conductivity data of Dry and Gledhill [6] serve to establish the correctness of the value $[\text{H}^+] = (8.17 \pm 0.08) \times 10^{-6} \text{ mol dm}^{-3}$, obtained primarily from the measured pH of the solutions. This concentration of the acid produced by hydrolysis is a highly important quantity for the establishment of the recommended value of the solubility, see below. Since a value of K_{s0}° obtained from the work of previous authors [12] is used in the calculations [6] of the concentrations of the species, this work [6] cannot be used to obtain an independent value of the solubility product.


The work of Herz [7] should give a value of K_{s0}° at 25 °C in conjunction with the use of a value of $K_s^\circ(\text{HgO}(\text{s}) + \text{H}_2\text{O} = \text{Hg}^{2+} + 2\text{OH}^-)$, provided that solid Hg_2O disproportionates to $\text{HgO}(\text{s}) + \text{Hg}(\ell)$. The best value of $K_s^\circ(\text{HgO}) = 2.8 \times 10^{-26}$ [17], [18] (at 25 °C) however leads to a $K_{s0}^\circ(\text{Hg}_2\text{Cl}_2)$ value which is about two orders of magnitude too small, the same value obtained if the existence of $\text{Hg}_2\text{O}(\text{s})$ is accepted [10b]. This could be due to the sluggish establishment of equilibrium in the presence of two insoluble solids ($\text{Hg}_2\text{Cl}_2(\text{s})$ and $\text{Hg}_2\text{O}(\text{s})$ or $\text{HgO}(\text{s}) + \text{Hg}(\ell)$). The period during which the phases were equilibrated was not stated (it was only specified as "extended"), and the establishment of equilibrium was not demonstrated [7].

2.2. E.m.f. Measurements

Of the second group of papers dealing with determinations of K_{s0}° by means of e.m.f. measurements, Behrend's work [8] was pioneering, but of low precision. Since only one concentration of mercury (I) in its half cell and of KCl in the calomel


half cell were used, an extrapolation procedure is excluded, the liquid junction potential cannot be adequately estimated in spite of Behrend's later attempt, and only rough estimates of the activity coefficients can be made. The value obtained $2 \times 10^{-19} \text{ mol}^2\text{dm}^{-6}$ at 17 °C, is just an estimate of the order of magnitude. Similarly, Sherril's work [3] gives an estimate of $[\text{Hg}^{2+}]$ in a saturated Hg_2Cl_2 solution in 1 mol dm^{-3} NaCl, which leads to $K_{s0}^\circ = 0.44 \times 10^{-18}$ which is much too low. The estimate was, however, not supported by definite e.m.f. data on which it was purportedly based.

Ley and Heimbucher [9] provided fuller experimental information, but their data cannot either be used to extrapolate out liquid junction potentials. These were produced by their bridge electrolyte, $0.1 \text{ mol dm}^{-3} \text{ KNO}_3$, with the half cell solutions $0.05 \text{ mol dm}^{-3} \text{ Hg}_2(\text{ClO}_4)_2$ and 0.1 or $1.0 \text{ mol dm}^{-3} \text{ KCl}$. With the more dilute KCl, the junction potentials could be rather small, estimated at $\pm 0.01 \text{ V}$, leading to a possible error of a factor of two in $K_{s0}^\circ = 1.48 \times 10^{-18} \text{ mol}^2\text{dm}^{-6}$ at 20 °C. This value is about twice the recommended value at 20 °C. On the other hand, Brodsky's work [10] using the cell



permitted extrapolation to $c = 0$ (and less clearly to $c = 0$), thus eliminating the liquid junction potentials, which are probably small in any case, through the use of a saturated KNO_3 bridge. The values obtained at four temperatures are rather close to the recommended values at these temperatures, the interpolated value for 25 °C being $(1.07 \pm 0.13) \times 10^{-18} \text{ mol}^2\text{dm}^{-6}$, definitely on the low side.

An improved extrapolation method was provided by Law [11], but the ionic strength was not kept constant as $x \rightarrow 0$ in the cell




the extrapolation, hence, not eliminating sufficiently the liquid junction [14]. Law's data also result in an incorrect value of $E_{\text{Hg}/\text{Hg}_2}^{2+}$ (see below), which, in turn, leads to a much too high value of $K_{s0}^\circ = 5.1 \times 10^{-18} \text{ mol}^2\text{dm}^{-6}$ at 25 °C. Since Law's values were available only through secondary sources [14], [19], no further comments can be made on the discrepancy.

The work of Jonsson, Qvarfort and Sillen [12] and of Hansen, Izatt and Christensen [13] tried to overcome the liquid junction and activity coefficient problems by using a constant ionic medium of $0.50 \text{ mol dm}^{-3} \text{ NaClO}_4$ (of which 0.01 mol dm^{-3} [12], 0.10 mol dm^{-3} [13], were HClO_4 rather than NaClO_4). The value obtained at 25 °C in this medium [12], $K_{s0}^\circ = (1.32 \pm 0.03) \times 10^{-17} \text{ mol dm}^{-6}$, must be multiplied by

$\gamma_{\pm}^3(\text{Hg}_2^{2+} \cdot 2\text{Cl}^- \text{ trace in NaClO}_4 \text{ medium})$ to obtain K_{s0}° . For this quantity the value 0.09 was suggested, but the evaluator failed to trace its origin through the references provided [12], [16], [20]. Application of this correction leads to $K_{s0}^\circ = (1.19 \pm 0.03) 10^{-18} \text{ mol}^2 \text{dm}^{-6}$ at 25 °C. A somewhat different value of the correction term [6], 0.101 ± 0.001 , leads to $K_{s0}^\circ = (1.33 \pm 0.03) \times 10^{-18} \text{ mol}^2 \text{dm}^{-6}$, which is rather near the recommended value, but a bit low. Again no details of the derivation of this γ_{\pm}^3 were provided [6]. The values of K_{s0} for 0.50 mol dm⁻³ NaClO₄ medium at 7 and 40 °C are [13] 6.4×10^{-19} and $6.29 \times 10^{-17} \text{ mol}^2 \text{dm}^{-6}$, respectively. Conversion to K_{s0}° requires estimates of $\gamma_{\pm}^3(\text{Hg}_2^{2+} \cdot 2\text{Cl}^- \text{ trace in NaClO}_4 \text{ medium})$ at these temperatures. Acceptance of 0.10 ± 0.02 as valid for the range 7–40 °C, in lieu of any better values, leads to $K_{s0}^\circ = (6.5 \pm 1.3) \times 10^{-20}$ at 7 °C and $(6.9 \pm 1.4) \times 10^{-18}$ at 40 °C as the best estimates. These are about 30% lower than the recommended values.

No experimental details are provided in these studies [12], [13] to evaluate the results from the point of view of the purity of the reagents (e.g., contamination with bromide) or of the instrumentation used, although the former [12] contains enough details on the care with which the work was carried out to assure acceptability of the data on this account.

The work of Galloway [14] combined the merits of constant ionic strength media (for extrapolating away the effect of excess chloride concentrations) with the possibility to eliminate the activity coefficients by providing series of data at decreasing ionic strengths. The cell employed



was measured at 5 K intervals over the range 15 to 40 °C. At first extrapolation at constant m (and T) from $0.1 \leq x \leq 0.6$ to $x = 0$ was followed by adding an activity coefficient correction term, which carried the main burden of the second extrapolation from $0.01 \leq m \leq 0.05$ to $m = 0$. This device puts relatively small weights on this extrapolation on the one hand, and on inadequacies of the activity coefficient term on the other, leading to results of high validity. The final value of $K_{s0}^\circ = (1.49 \pm 0.05) \times 10^{-18} \text{ mol}^2 \text{kg}^{-2}$ (practically the same as $\text{mol}^2 \text{dm}^{-6}$) does not differ significantly from the recommended value $(1.43 \pm 0.06) \times 10^{-18} \text{ mol}^2 \text{kg}^{-2}$, both at 25 °C. For other temperatures there are increasing differences, but still within the combined uncertainties.

2.3. Standard Electrode Potentials

A third group of papers is now considered, in some of which the standard electrode potentials of the calomel electrode $E_{\text{Hg}/\text{Hg}_2\text{Cl}_2}^\circ$ and in others, those of the mercury/mercury(I) electrode, $E_{\text{Hg}/\text{Hg}_2^{2+}}^\circ$ were determined, for purposes other than the calculation of the solubility product of mercury(I) chloride.

These papers will not be reviewed here, since they have been reviewed quite adequately previously, and only the latest or "best" results will be presented here. The solubility constant can, obviously, be calculated from

$$\log K_{s0}^\circ = (E_{\text{Hg}/\text{Hg}_2\text{Cl}_2}^\circ - E_{\text{Hg}/\text{Hg}_2^{2+}}^\circ)/(RT/2F) \ln 10. \quad (10)$$

The most extensive and careful study of the calomel electrode has been made over the years by Ives and his coworkers [21a]–[21d]. Their work is well confirmed by those of Ahluwalia and Cobble [22] and others (see refs. [14], [15]). The value for 25 °C is established at $E_{\text{Hg}/\text{Hg}_2\text{Cl}_2}^\circ = 0.26818 \pm 0.00002$ V [24] and at other temperatures can be obtained from

$$\begin{aligned} E_{\text{Hg}/\text{Hg}_2\text{Cl}_2}^\circ (T)/V &= 0.26818 \pm 0.00002 \\ &- (2.99 \pm 0.03) 10^{-4} \Delta T - (3.1 \pm 0.3) 10^{-6} (\Delta T)^2 \end{aligned} \quad (11)$$


where $\Delta T = T/K - 298.15$, with an overall precision of ± 0.00004 V. The corresponding entropy term is $F(dE_{\text{Hg}/\text{Hg}_2\text{Cl}_2}^\circ/dT) = 151.37 - 0.6044 \Delta T \text{ JK}^{-1} \text{ mol}^{-1}$. At 298.15 K $\Delta S^\circ = -28.83 \text{ JK}^{-1} \text{ mol}^{-1}$, compared with the value suggested by Ahluwalia and Cobble [22] $-28.49 \text{ JK}^{-1} \text{ mol}^{-1}$.

The standard potential of the mercury/mercury(I) electrode has not been studied as extensively, and is not established to that degree of accuracy obtained for the calomel electrode. The thermodynamic functions of formation of $\text{Hg}_2^{2+}(\text{aq})$ were given in the latest NBS compilation [23] as $\Delta G_f^\circ = 153.55 \text{ kJ mol}^{-1}$ and $\Delta H_f^\circ = 172.4 \text{ kJ mol}^{-1}$ (the latter with one less significant digit than the former). These values lead to $E_{\text{Hg}/\text{Hg}_2^{2+}}^\circ(298.15 \text{ K}) = (0.79574 \pm 0.00022) \text{ V}$ and

$$\begin{aligned} E_{\text{Hg}/\text{Hg}_2^{2+}}^\circ (T/K)/V \\ = E_{\text{Hg}/\text{Hg}_2^{2+}}^\circ (298.15)/V - 3.27 \times 10^{-4} \Delta T, \end{aligned} \quad (12)$$

where $\Delta T = T/K - 298.15$, with a precision of ± 0.00019 V. The apparent precision given here is based merely on the apparent precision (\pm one unit of the last significant digit) of the values in the compilation [23]. Another set of $E_{\text{Hg}/\text{Hg}_2^{2+}}^\circ(T)$ data was obtained by Read [24], (quoted in ref. [14]), with 0.7956 V for $T = 298.15 \text{ K}$, $dE^\circ/dT = -2.97 \times 10^{-4} \text{ VK}^{-1}$ and an overall precision of ± 0.00014 V, in the range 15 to 45 °C. Most other workers used the old data of Linhart [25] corrected in one way or another, except for the more recent data of ElWakkad and Salem [26], of Bonner and Unietis [27], and of Schwarzenbach and Anderegg [28] who gave for $E_{\text{Hg}/\text{Hg}_2^{2+}}^\circ(298.15 \text{ K})$ 0.7960 \pm 0.0005 V, and 0.7966 \pm 0.0010 V, respectively. The latest review of these data, by Venderzee and Swanson [15] selected the value $(0.7960 \pm 0.0005) \text{ V}$ as the best for 298.15 K. As for the temperature coefficient, $dE^\circ/dT = -(3.25 \pm 0.06) \times 10^{-4} \text{ VK}^{-1}$ represents best both Read's [24] and the NBS [23] data.

The standard thermodynamic functions for the reaction



are obtained from

$$\Delta G_{s0}^{\circ} = 2F(E_{\text{Hg}/\text{Hg}_2\text{Cl}_2}^{\circ} - E_{\text{Hg}/\text{Hg}_2^{2+}}^{\circ}) \quad (14)$$

and from $\Delta S_{s0}^{\circ} = -(d\Delta G_{s0}^{\circ}/dT) = 2Fd(E_{\text{Hg}/\text{Hg}_2\text{Cl}_2}^{\circ} - E_{\text{Hg}/\text{Hg}_2^{2+}}^{\circ})/dT$. The value at 298.15 K of $\Delta G_{s0}^{\circ} = 101.86 \pm 0.10$ kJ mol⁻¹ is consistent with the other thermodynamic data [23]. However, $\Delta S_{s0}^{\circ}/\text{JK}^{-1} \text{ mol}^{-1} = 5.06 \pm 1.25 - (1.21 \pm 0.12) \times 10^{-3} \Delta T$ (with $\Delta T = T/\text{K} - 298.15$), a positive entropy change resulting from (11), (12) and (14) leads to $\Delta H_{s0}^{\circ} = 103.36 \pm 0.38$ kJ mol⁻¹ at 298.15 K. This differs considerably from the calorimetric value recently obtained by Vanderzee and Swanson [15], 98.08 ± 0.18 kJ mol⁻¹. Since the latter work is very reliable, the fault must be sought with the temperature dependence of either one of the E° values (or both) employed in (14), most probably in that of $E_{\text{Hg}/\text{Hg}_2^{2+}}^{\circ}$, which is less well established.

In fact, the enthalpy of formation of Hg_2^{2+} (aq) given by Vanderzee and Swanson [15], $\Delta H_f^{\circ} = 166.82 \pm 0.21$ kJ mol⁻¹ differs considerably from that in the NBS compilation [23], 172.4 kJ mol⁻¹ and should be more reliable. Use of the newer value leads to the relation, obtained from the 25 °C standard potential and the enthalpy of precipitation of calomel

$$E_{\text{Hg}/\text{Hg}_2^{2+}}^{\circ}(T)/V = 0.7960 \pm 0.0005 \\ - (2.30 \pm 0.04) \cdot 10^{-4} \Delta T \quad (15)$$

as the more reliable value.

2.4. Calculated Solubility

The solubility of mercury(I) chloride in water is not, as was assumed in the very early studies, $(K_{s0}^{\circ}/4)^{1/3}$, because of the disproportionation of Hg_2^{2+} to give Hg^{2+} and $\text{Hg}(\ell)$, the complexing of Hg^{2+} with chloride, its hydrolysis, and to a lesser extent the hydrolysis of Hg_2^{2+} . The following analysis is based on the work of Dry and Gledhill [6], who showed that the saturated solution contains the following mercury species: $\text{Hg}(\text{OH})_2$, HgCl_2 , HgOH^+ , HgCl^+ , Hg_2^{2+} and Hg_2OH^+ , in addition to H^+ and Cl^- ions. The total concentration of mercury in the solution can be expressed as

$$c_{\text{Hg}} - [\text{Hg}_2^{2+}] \left\{ K_{\text{R}}^{-1} \left[K_{\text{W}}^{\circ} \beta_{\text{H}_2\text{OH}}^{\circ} (\gamma_{\text{Hg}_2^{2+}}/\gamma_{\text{Hg}(\text{OH})_2} \gamma_{\text{H}^+}^2) [\text{H}^+]^{-2} \right. \right. \\ + \beta_{\text{H}_2\text{Cl}}^{\circ} (\gamma_{\text{Hg}_2^{2+}} \gamma_{\text{Cl}^-}^2/\gamma_{\text{HgCl}_2}) [\text{Cl}^-]^2 \\ + K_{\text{W}}^{\circ} \beta_{\text{H}_2\text{OH}}^{\circ} (\gamma_{\text{Hg}_2^{2+}}/\gamma_{\text{HgOH}^+} \gamma_{\text{H}^+}) [\text{H}^+]^{-1} \\ \left. \left. + \beta_{\text{H}_2\text{Cl}}^{\circ} (\gamma_{\text{Hg}_2^{2+}} \gamma_{\text{Cl}^-}/\gamma_{\text{HgCl}^+}) [\text{Cl}^-] \right] \right\} \\ + 2 + 2K_{\text{W}}^{\circ} \beta_{\text{H}_2\text{OH}}^{\circ} (\gamma_{\text{Hg}_2^{2+}}/\gamma_{\text{Hg}_2\text{OH}^+} \gamma_{\text{H}^+}) [\text{H}^+]^{-1} \quad (16)$$

In this expression, the activity coefficients will be calculated according to the modified Debye-Hückel equation

$$\log \gamma_i = -0.51z_i^2 I^{1/2}/(1 + 1.6I^{1/2}) \quad (17)$$

where I , the ionic strength will be put equal to $[\text{H}^+]$, and z_i is the charge of the i -th ion. Since the ionic strength is very low, $I = [\text{H}^+] \approx 8.2 \times 10^{-6}$ mol dm⁻³, the activity coefficient corrections are quite small, and any deviation of (17) from the true behaviour leads to negligible errors. Other equations which must be considered are

$$[\text{Hg}_2^{2+}] [\text{Cl}^-]^2 = K_{s0}^{\circ} \gamma_{\text{Hg}_2^{2+}}^{-1} \gamma_{\text{Cl}^-}^{-2} \quad (18)$$

$$[\text{Cl}^-] = 2(c_{\text{Hg}} - [\text{HgCl}_2] - \frac{1}{2}[\text{HgCl}^+]) \\ \approx 2(c_{\text{Hg}} - [\text{HgCl}_2]) \quad (19)$$

The approximation in (19) is permissible since, as will be found, $[\text{HgCl}_2]/c_{\text{Hg}} \approx 0.37$, but $[\text{HgCl}^+]/c_{\text{Hg}} \approx 0.01$. The hydrogen ion concentration, due to hydrolysis, obtained by pH measurements and confirmed by conductivity [6], is taken as $[\text{H}^+] = (8.17 \pm 0.08) \times 10^{-6}$ mol dm⁻³ at 25 °C. The following values of the constants appearing in (16) and valid for 25 °C will be used: $K_{\text{R}}^{\circ}(\text{Hg}_2^{2+} + \text{Hg}(\ell) = \text{Hg}_2^{2+}) = 10^{1.944 \pm 0.008}$ [15], $K_{\text{W}}^{\circ} \beta_{\text{H}_2\text{OH}}^{\circ}(\text{Hg}_2^{2+} + 2\text{H}_2\text{O} = \text{Hg}(\text{OH})_2 + 2\text{H}^+) = 10^{-5.65 \pm 0.12}$ [6], $\beta_{\text{H}_2\text{Cl}}^{\circ}(\text{Hg}_2^{2+} + 2\text{Cl}^- = \text{HgCl}_2) = 10^{19.26 \pm 0.09}$ [6], $K_{\text{W}}^{\circ} \beta_{\text{H}_2\text{OH}}^{\circ}(\text{Hg}_2^{2+} + \text{H}_2\text{O} = \text{HgOH}^+ + \text{H}^+) = 10^{-3.09 \pm 0.20}$ [6], $\beta_{\text{H}_2\text{Cl}}^{\circ}(\text{Hg}_2^{2+} + \text{Cl}^- = \text{HgCl}^+) = 10^{7.34 \pm 0.27}$ [6], $K_{\text{W}}^{\circ} \beta_{\text{H}_2\text{OH}}^{\circ}(\text{Hg}_2^{2+} + \text{H}_2\text{O} = \text{Hg}_2\text{OH}^+ + \text{H}^+) = 10^{-4.4 \pm 0.4}$ [6] and for $K_{s0}^{\circ}(\text{Hg}_2\text{Cl}_2(\text{s}) = \text{Hg}_2^{2+} + 2\text{Cl}^-) = 10^{-17.844 \pm 0.017}$ will be used.

The values quoted were estimated [6] as valid for the ionic strength of the saturated solution, from corrections applied to published values [16] valid for 0.50 mol dm⁻³ NaClO₄ medium. With these values of the constants, the value of $[\text{H}^+]$ and eq (16), (18) and (19), the following implicit equation is obtained (valid for 25 °C):

$$c_{\text{Hg}}^3 (1 - (3.09 \pm 0.65)/10^6 c_{\text{Hg}})^2 = (4.25 \pm 0.05) \\ \times 10^{-21} \{ 10^{7.63 \pm 0.27} c_{\text{Hg}} (1 - (3.09 \pm 0.65)/10^6 c_{\text{Hg}}) \\ + 10^{14.85 \pm 0.09} c_{\text{Hg}}^2 (1 - (3.09 \pm 0.65)/10^6 c_{\text{Hg}})^2 \\ + (3.43 \pm 1.06) \times 10^4 \}. \quad (20)$$

This was solved iteratively to give the value for the solubility

$$c_{\text{Hg}} = (8.4 \pm 1.0) \times 10^{-6} \text{ mol dm}^{-3} \quad (21)$$

as the solubility of mercury(I) chloride in water at 25 °C. This value is consistent with the solubility measured directly [6], $(7.5 \pm 0.3) \times 10^{-6}$ mol dm⁻³, as discussed above.

3. Recommended Values

3.1. Solubility Product Constant of Hg_2Cl_2

The above analysis and eq (10), (11) and (15) yield for the




Figure 1. Values of pK_{s0}^* ($= -\log K_{s0}^*$) recalculated from the authors' data compared with the "recommended" values (continuous line).

solubility product of mercury(I) in water in the range 5 to 45 °C

$$\log(K_{s0}^*/\text{mol}^2\text{kg}^{-2}) = -17.844 \pm 0.017 + (0.0622 \pm 0.0002)\Delta T - (3.0 \pm 0.2) \times 10^{-4} (\Delta T)^2 \quad (22)$$

where $\Delta T = T/\text{K} - 298.15$. The first term on the rhs represents the value for 25 °C

$$K_{s0}^*/\text{mol}^2\text{kg}^{-2}(298.15 \text{ K}) = (1.43_3 \pm 0.05_6) \times 10^{-18} \quad (23)$$

the relative error being $\pm 3.9\%$. The value in $(\text{mol dm}^{-3})^2$ units is 0.6% lower, i.e., insignificantly different. The values of $\log K_{s0}^*$ at different temperatures are plotted in figure 1, and compared there with values obtained by several authors.

3.2. Standard Thermodynamic Functions

There are several fixed quantities, selected above as reliable, to which the thermodynamic functions must conform. These are $E_{\text{Hg}/\text{Hg}_2\text{Cl}_2}^\circ(298.15 \text{ K}) = 0.26818 \pm 0.00002 \text{ V}$, $\Delta S_{298.15}^\circ(\text{Hg}/\text{Hg}_2\text{Cl}_2) = -28.83 \text{ JK}^{-1} \text{ mol}^{-1}$, $E_{\text{Hg}/\text{Hg}_2^{2+}}^\circ(298.15 \text{ K}) = 0.7960 \pm 0.0005 \text{ V}$ and $\Delta H_{298.15}^\circ(\text{Hg}_2^{2+}(\text{aq}) + 2\text{Cl}^-(\text{aq}) = \text{Hg}_2\text{Cl}_2(\text{s})) = -98.08 \pm 0.18 \text{ kJ mol}^{-1}$. These lead to the following functions for reaction (13)

$$\begin{aligned} \Delta G_{s0}^\circ/\text{kJ mol}^{-1} &= 101.86 \pm 0.10 + (0.0127 \pm 0.0009)\Delta T + (0.60 \pm 0.06) \\ &\times 10^{-3} (\Delta T)^2 = -151.4 \\ &+ 0.345 T + 0.60 \times 10^{-3} T^2 \end{aligned} \quad (24)$$

$$\begin{aligned} \Delta S_{s0}^\circ/\text{JK}^{-1}\text{mol}^{-1} &= -12.7 \pm 0.9 - (1.20 \pm 0.12)\Delta T = 345 - 1.20 T \end{aligned} \quad (25)$$

$$\begin{aligned} \Delta H_{s0}^\circ/\text{kJ mol}^{-1} &= 98.08 \pm 0.18 - (0.358 \pm 0.036)\Delta T \\ &- (0.60 \pm 0.06) \times 10^{-3} (\Delta T)^2 \\ &= 151.4 - 0.60 \times 10^{-3} T^2 \end{aligned} \quad (26)$$

$$\begin{aligned} \Delta C_{p,s0}^\circ/\text{JK}^{-1}\text{mol}^{-1} &= -(0.358 \pm 0.036) \\ &- (1.20 \pm 0.12) \times 10^{-3} \Delta T \\ &= -1.20 \times 10^{-3} T \\ &\text{(tentative value)} \end{aligned} \quad (27)$$

These functions have the following values at 25 °C:

$$\begin{aligned}\Delta G_{\text{a0}}^{\circ}(298.15) &= 101.86 \pm 0.10 \text{ kJ mol}^{-1}, \\ \Delta S_{\text{a0}}^{\circ}(298.15) &= -12.70 \pm 0.9 \text{ JK}^{-1} \text{ mol}^{-1}, \\ \Delta H_{\text{a0}}^{\circ}(298.15) &= 98.08 \pm 0.18 \text{ kJ mol}^{-1} \\ \Delta C_{p,\text{a0}}^{\circ}(298.15) &= 0.36 \pm 0.04 \text{ JK}^{-1} \text{ mol}^{-1} \\ &\quad \text{(tentative value)}\end{aligned}\quad (28)$$

The uncertainty about $\Delta C_{p,\text{a0}}^{\circ}$ is due to the ignorance of $\Delta C_{p,\text{f}}^{\circ}(\text{Hg}_2^{2+}(\text{aq}))$, so that it is based solely on the second derivative of $E_{\text{Hg}/\text{Hg}_2\text{Cl}_2}^{\circ}$. The standard entropy change vanishes at 14.4 °C (according to Galloway at 19.0 °C [14]).

3.3. The Aqueous Solubility of Hg_2Cl_2

The aqueous solubility of mercury(I) chloride at 25 °C is given by eq (16) to (20) as

$$c_{\text{Hg}} = (8.4 \pm 1.0) \times 10^{-6} \text{ mol dm}^{-3} \quad (21)$$

The values at other temperatures cannot be calculated, since the temperature coefficients of the various constants and of the pH of the saturated solutions are unknown.

Acknowledgements

Acknowledgements. The help of Prof. H.L. Clever, in providing copies of hard to obtain literature sources, is gratefully acknowledged. Useful remarks and encouragement have been obtained from Profs. L.G. Hepler, D.N. Hume, A.S. Kertes, and G.H. Nancollas and Dr. M. Salomon.

4. References

- [1a] Kohlrausch, F., and Rosc, F., *Z. physik. Chem.* **12**, 234 (1893).
- [1b] Kohlrausch, F., *Z. physik. Chem.* **44**, 217 (1903).
- [1c] Kohlrausch, F., *Z. physik. Chem.* **64**, 129 (1908).
- [2] Richards, T.W., and Archibald, E.H., *Z. phys. Chem.* **40**, 385 (1902).
- [3] Sherrill, M.S., *Z. physik. Chem.* **43**, 705 (1903).
- [4] Eversole, W.G., and McLachlan, R.W., *J. Am. Chem. Soc.* **54**, 864 (1932).
- [5] Garrett, A.B., Noble, N. V., and Miller, S., *J. Chem. Educ.* **19**, 485 (1942).
- [6] Dry, M.E., and Gledhill, J.A., *Trans. Faraday Soc.* **51**, 1119 (1955).
- [7] Herz, W., *Z. anorg. Chem.* **70**, 170 (1911).
- [8] Behrend, R., *Z. physik. Chem.* **11**, 466 (1893); **15**, 498 (1894).
- [9] Ley, H., and Heimbucher, C., *Z. Elektrochem.* **10**, 301 (1904).
- [10a] Brodsky, A.E., and Scherschewer, J.M., *Z. Elektrochem.* **32**, 1 (1926).
- [10b] Brodsky, A.E., *Z. Elektrochem.* **35**, 833 (1929).
- [11] Law, J.T., M.S. Thesis, Univ. New Zealand, 1946.
- [12] Jonsson, J., Qvarfort, I., and Sillén, L.G., *Acta Chem. Scand.* **1**, 46 (1947).
- [13] Hansen, I.D., Izatt, R.M., and Christensen J.J., *Inorg. Chem.* **2**, 1243 (1963).
- [14] Galloway, W.D., M.S. Thesis, Univ. New Zealand, 1961.
- [15] Vanderzee, C.E., and Swanson, J.A., *J. Chem. Thermod.* **6**, 827 (1974).
- [16] Sillén, L.G., *Acta Chem. Scand.* **3**, 529 (1949).
- [17] Feitknecht, W., and Schindler, P., *Pure Appl. Chem.* **6**, 130 (1963).
- [18] Hepler, L.G., and Olofsson, G., *Chem. Rev.* **75**, 585 (1975).
- [19] Berecki, C., Biedermann, G., and Sillén, L.G., Report to (precursor of) Commission V.6, Analytical Division of IUPAC, 1953.
- [20] Qvarfort, I., and Sillén, L.G., *Acta Chem. Scand.* **3**, 517 (1949); Lindgren, B., Jonsson, A., and Sillén, L.G., *Acta Chem. Scand.* **1**, 479 (1947).
- [21a] Hills, G.J., and Ives, D.J.G., *J. Chem. Soc.* 318 (1951).
- [21b] Das, S.N., and Ives, D.J.G., *J. Chem. Soc.* 1619 (1962).
- [21c] Gupta, S.R., Hills, G.J., and Ives, D.J.G., *Trans. Faraday Soc.* **59**, 1874 (1963).
- [21d] Ives, D.J.G., and Prasad, D., *J. Chem. Soc. B* 1649 (1970).
- [22] Ahluwalia, J.C., and Coblle, P.W., *J. Am. Chem. Soc.* **86**, 5381 (1964).
- [23] Wagman, D.D., Evans, W.H., Parker, V.B., Halow, L., Bailey, S.M., and Schumm, R.H., *U.S. Natl. Bureau Stand., Tech. Note* 270-4 (1969).
- [24] Read, M.S. Thesis, Univ. New Zealand, 1960.
- [25] Linhart, G.A., *J. Am. Chem. Soc.* **38**, 2356 (1916).
- [26] ElWakkad, S.E.S., and Salem, T.M., *J. Phys. Chem.* **54**, 1371 (1950).
- [27] Bonner, O.D., and Unietis, F.A., *J. Am. Chem. Soc.* **75**, 5111 (1953).
- [28] Schwarzenbach, G., and Anderegg, G., *Helv. Chim. Acta* **37**, 1289 (1954).

5. Compilation

5.1. Data of Behrend (1893, 1894)

COMPONENTS: 1. Mercury(I) chloride, Hg_2Cl_2 (10112-91-1) 2. Potassium chloride, KCl (7447-40-7) 3. Water, H_2O (7732-18-5)	ORIGINAL MEASUREMENTS: Behrend, R., Z. Phys. Chem. (1893) <u>11</u> , 466; (1894) <u>15</u> , 498.
VARIABLES: One temperature One KCl concentration	PREPARED BY: Y. Marcus, January 1978
EXPERIMENTAL VALUES: <p>E.m.f. E at room temperature (17°C) of cell: $\text{Hg}/\text{Hg}_2(\text{NO}_3)_2$ 0.05M, HNO_3 (?M)/0.1M $\text{KNO}_3/\text{Hg}_2\text{Cl}_2(\text{s})$, 0.1M KCl/Hg found to be 0.356V and 0.360V in two experiments. Final value adopted is 0.358V. This value has to be multiplied by 1.092 to give 0.391V, as stated by the author in the correction published by him in the second source quoted (see below). From this is derived (Y.M.)</p> $\log K_{\text{SO}}^\circ(\text{Hg}_2\text{Cl}_2, 17^\circ\text{C}) = -17.10 - (E_j/V)/0.02879$ <p>A realistic estimate for E_j is $0.05 \pm 0.05\text{V}$, hence $\log K_{\text{SO}}^\circ = -17.1 - (1.7 \pm 1.0) = -18.8 \pm 1.0$ at 290K.</p>	
AUXILIARY INFORMATION	
METHOD: $\log K_{\text{SO}}^\circ(\text{Hg}_2\text{Cl}_2, 17^\circ\text{C}) = -(2/0.05757) \{ (E/V) - (E_j/V) \} - \log(c_{\text{Hg}_2(\text{NO}_3)_2} \cdot c_{\text{KCl}}^2) - \log(\gamma_{\text{Hg}_2(\text{NO}_3)_2} \cdot \gamma_{\text{KCl}}^2 \cdot \gamma_{\text{KNO}_3}^{-2})$ $= -2(0.391 - E_j/V)/0.05757 - \log(0.05 \cdot 0.1^2) - \log(0.55 \cdot 0.77^2/0.74^2)$ $= -13.58 - (E_j/V)/0.02879 - 3.30 - 0.22$ $= -17.10 - (E_j/V)/0.02879$ <p>$\gamma_{\text{Hg}_2(\text{NO}_3)_2}$ estimated from value for $\text{Ca}(\text{NO}_3)_2^1$, value for that, KCl and KNO_3 from ref. 2.</p> <p>Unthermostated cell, using an electrometer, a resistance box and a Leclanché standard cell, constant to ca. ± 3 mV over two months.</p>	SOURCE AND PURITY OF MATERIALS: Not stated ESTIMATED ERROR: From range of $E(\pm 0.002\text{V})$, uncertainty of temperature ($\pm 2^\circ\text{C}$) and activity coefficient product ($\pm 10\%$), error of $\log K_s$ is ± 0.22 . REFERENCES: 1. Bonner, O.D., Unietis, F., J. Am. Chem. Soc. (1953) <u>75</u> , 5111. 2. Robinson, R.A., Stokes, R.H., Electrolyte Solutions, Butterworth, London, 2nd Ed., 1959.

5.2. Data of Richards and Archibald (1902)

COMPONENTS:		ORIGINAL MEASUREMENTS:	
1. Mercury(I) chloride, Hg ₂ Cl ₂ (10112-91-1)		Richards, T.W., Archibald, E.H., Z. phys. Chem. (1902) 40, 385.	
2. Hydrochloric Acid, HCl (7647-01-0)			
3. Water, H ₂ O (7732-18-5)			
VARIABLES:		PREPARED BY:	
HCl concentration One temperature		Y. Marcus, January 1978	
EXPERIMENTAL VALUES:			
Solubility of Hg ₂ Cl ₂ in aqueous HCl at 25°C. Composition of the solutions:			
c _{HCl} /mol dm ⁻³	c _{Hg, total} /g dm ⁻³	10 ⁴ c _{Hg, total} /mol dm ⁻³	
0.83	0.034	1.69	
0.83	0.034	1.69	
1.00	0.048	2.39	
1.00	0.048	2.39	
2.50	0.206	10.27	
2.50	0.208	10.37	
4.15	0.400	19.94	
4.15	0.398	19.84	
5.48	0.548	27.32	
5.48	0.548	27.32	
(Remarks: c _{Hg, total} calculated by compiler. Data were also presented for 7.00, 7.30, 8.31 and 10.00 mol dm ⁻³ HCl, but not used for calculations below.)			
log K _e ^o (Hg ₂ Cl ₂ (s) + 2Cl ⁻ = Hg(l) + HgCl ₄ ²⁻) = -3.93±0.01 (calculated by YM)			
log K _{so} ^o (Hg ₂ Cl ₂ (s) = Hg ₂ ²⁺ + 2Cl ⁻) = -17.72±0.08 (calculated by YM with additional data ¹⁻³).			
AUXILIARY INFORMATION			
METHOD:		SOURCE AND PURITY OF MATERIALS:	
log K _e ^o = log(c _{Hg, total} /c _{HCl} ²) - - log(1+β ₃ β ₄ ⁻¹ c _{HCl} ⁻¹ + β ₂ β ₄ ⁻¹ c _{HCl} ⁻²) _{ref. 2} - 2AI ^{1/2} /(1+BI ^{1/2}) + ΔbI; A = 0.51, B = 1.6, I = c _{HCl} . Least squares calc. gave log K _e ^o = -3.93 and Δb = -0.146. log K _{so} ^o = log K _e ^o + log K _r (Hg ₂ ²⁺ +Hg(l) = Hg ₂ ²⁺) _{ref. 1} - log β ₄ (Hg ₂ ²⁺ +4Cl ⁻ = HgCl ₄ ²⁻) _{ref. 2} - 4 log y(trace NaCl in 0.5M NaClO ₄) _{ref. 3} = = -3.93+1.94-15.07-0.66 = -17.72. Excess HgCl ₂ shaken for >7 hr with 0.1 g Hg(l) and 50 ml HCl solution in glass vessel provided with purified rubber stopper in a thermostated bath at 25.00±0.05°C. Total Hg in equilibrium solutions determined gravimetrically as HgS precipitated by H ₂ S, washed and dried at 100°C.		Hg ₂ Cl ₂ sublimed at low temperature, the source shown to be immaterial. HCl was redistilled c.p. acid.	
		ESTIMATED ERROR: log K _e ^o : ±0.05 from least squares fitting log K _{so} ^o : ±0.06 on log β ₄ , ±0.01 on log y _{NaCl} and ±0.02 on log K _r yield total ±0.08	
		REFERENCES: 1. Hietanen, S., Sillén, L.G., Ark. Kemi (1956) 10, 103. 2. Sillén, L.G., Acta Chem. Scand. (1949) 3 539. 3. Estimated by YM from Robinson, R.A., Stokes, R.H., Electrolyte Solutions, Butterworth, London, 2nd ed., 1959, and Langer, R.D., J. Phys. Chem. (1965) 69, 3992.	

COMPONENTS:		ORIGINAL MEASUREMENTS:			
1. Mercury(I) chloride, Hg ₂ Cl ₂ (10112-91-1)		Richards, T.W., Archibald, E.H.,			
2a. Sodium chloride, NaCl (7647-14-5)		Z. phys. Chem. (1902), 385.			
2b. Calcium chloride, CaCl ₂ (10043-52-4)					
2c. Barium chloride, BaCl ₂ (10361-56-3)					
3. Water, H ₂ O (7732-18-5)					
VARIABLES:		PREPARED BY:			
NaCl, CaCl ₂ or BaCl ₂ concentrations		Y. Marcus, January 1978			
One temperature					
EXPERIMENTAL VALUES:					
Solubility of Hg ₂ Cl ₂ in aqueous chlorides at 25°C. Composition of the solutions:					
c_{NaCl}	$10^4 c_{\text{Hg, total}}$	c_{CaCl_2}	$10^4 c_{\text{Hg, total}}$	c_{BaCl_2}	$10^4 c_{\text{Hg, total}}$
mol dm ⁻³	mol dm ⁻³	mol dm ⁻³	mol dm ⁻³	mol dm ⁻³	mol dm ⁻³
1.00	2.04	0.36	1.10	0.50	2.19
2.00	6.43	0.50	1.64	0.75	4.39
2.50	9.67	1.00	4.04	1.00	5.33
3.80	18.9	1.25	5.88	1.50	11.5
5.00	32.1	1.76	11.5		
Remarks: $c_{\text{Hg, total}}$ calculated by compiler from $c_{\text{Hg, total}}/\text{g dm}^{-3}$ data, which are averages of two experiments at each concentration. Data also presented for 2.32 and 2.93 mol dm ⁻³ CaCl ₂ , but not used for calculations below. Salt concentrations presented as equivalents dm ⁻³ , recalculated here.					
$\log K_e^\circ(\text{Hg}_2\text{Cl}_2(\text{s}) + 2\text{Cl}^- = \text{Hg}(\text{l}) + \text{HgCl}_4^{2-}) = -3.98$ for NaCl solutions					
$= -3.94$ for CaCl ₂ solutions					
$= -3.93$ for BaCl ₂ solutions					
av. = -3.95±0.06					
$\log K_{\text{SO}}^\circ(\text{Hg}_2\text{Cl}_2(\text{s}) = \text{Hg}_2^{2+} + 2\text{Cl}^-) = -17.74 \pm 0.09$ on the average for the three series.					
AUXILIARY INFORMATION					
METHOD:		SOURCE AND PURITY OF MATERIALS:			
$\log K_e^\circ = \log(c_{\text{Hg, total}}/c_{\text{MCl}}^2) - \log(1 + \beta_3 \beta_4^{-1} c_{\text{MCl}}^{-1} + \beta_2 \beta_4^{-1} c_{\text{MCl}}^{-2})_{\text{ref. 2}} - 2A\gamma^{1/2}/(1+B\gamma^{1/2}) + \Delta bT$; $A=0.51$, $B=1.6$, $M=\text{Na}, \frac{1}{2}\text{Ca}$ or $\frac{1}{2}\text{Ba}$. Least squares calc. gave $\log K_e^\circ$ values given and $\Delta b=0.160$ for NaCl, 0.198 for CaCl ₂ and 0.136 for BaCl ₂ .		Hg_2Cl_2 sublimed at low temperature, the source shown to be immaterial. NaCl precipitated from solution by HCl, then recrystallized, CaCl ₂ prepared from pure Ca(NO ₃) ₂ , converted to CaCO ₃ and to CaCl ₂ and then recrystallized.			
$\log K_{\text{SO}}^\circ = \log K_e^\circ + \log K_r(\text{Hg}_2^{2+} + \text{Hg}(\text{l}) = \text{Hg}_2^{2+})_{\text{ref. 1}} - \log \beta_4(\text{Hg}_2^{2+} + 4\text{Cl}^- = \text{HgCl}_4^{2-})_{\text{ref. 2}} - 4 \log \gamma(\text{trace NaCl in 0.5M NaClO}_4)_{\text{ref. 3}} = 3.95 + 1.94 - 15.07 - 0.66 = -17.74 \pm 0.008$.		ESTIMATED ERROR: $\log K_e^\circ: \pm 0.06$ from least squares fitting. $\log K_{\text{SO}}^\circ: \pm 0.06$ on $\log \beta_4$, ± 0.01 on $\log \gamma_{\text{NaCl}}$ and ± 0.02 on $\log K_r$ yield ± 0.09 .			
Excess Hg ₂ Cl ₂ shaken for >7 hr with 0.1 g Hg(l) and 50 ml aqueous NaCl, CaCl ₂ or BaCl ₂ in glass vessel provided with purified rubber stopper in thermostated bath at 25.00±0.05°C. Total Hg in equilibrium solutions determined gravimetrically as HgS precipitated by H ₂ S, washed and dried at 100°C.		REFERENCES:			
		1. Hietanen, S., Sillén, L.G., Ark. Kemi (1956) 10, 103.			
		2. Sillén, L.G., Acta Chem. Scand. (1949) 3, 539.			
		3. Estimated by Y.M. from Robinson, R.A., Stokes, R.H., Electrolyte Solutions, Butterworth, London, 2nd Ed., 1959.			

5.3. Data of Sherrill (1903)

COMPONENTS: 1. Mercury(I) chloride, Hg_2Cl_2 (10112-91-1) 2. Sodium chloride, NaCl (7647-14-5) 3. Water, H_2O (7732-18-5)	ORIGINAL MEASUREMENTS: Sherrill, M.S., Z. physik. Chem. (1903) 43, 705.
VARIABLES: One NaCl concentration One temperature	PREPARED BY: Y. Marcus, January 1978
EXPERIMENTAL VALUES: (a) Total solubility of $\text{Hg}_2\text{Cl}_2(\text{s})$ in aqueous NaCl , 1 mol dm^{-3} , at 25°C is 1.51×10^{-4} mol dm^{-3} . (b) Concentration of species $[\text{Hg}_2^{2+}]$ in aqueous NaCl , 1 mol dm^{-3} , at equilibrium with $\text{Hg}_2\text{Cl}_2(\text{s})$ at 25°C is 5.3×10^{-20} mol dm^{-3} . From datum (b), $K_s(\text{Hg}_2^{2+} + \text{Hg}(\ell) = \text{Hg}_2^{2+}) = 120^1$ and the assumed degree of dissociation of the NaCl of 0.75, ¹ the author obtained $\log K_s(\text{Hg}_2\text{Cl}_2(\text{s}) = \text{Hg}_2^{2+} + 2\text{Cl}^-) = \log 5.3 \times 10^{-20} + \log 120 + 2 \log(1 \times 0.75) = -17.45$ Calculated (Y.M.) from datum (b), $K_s(\text{Hg}_2^{2+} + \text{Hg}(\ell) = \text{Hg}_2^{2+}) = 88 \pm 4^2$ and the activity coefficient of calcium chloride ³ in 1 mol dm^{-3} sodium chloride ⁴ $\log K_s^0(\text{Hg}_2\text{Cl}_2(\text{s}) = \text{Hg}_2^{2+} + 2\text{Cl}^-) = \log 5.3 \times 10^{-20} + \log 88 + 2 \log 1 + 3(-0.344) = -18.36$ Calculated (Y.M.) from datum (a): $\log K_e(\text{Hg}_2\text{Cl}_2(\text{s}) + 2\text{Cl}^- = \text{Hg}(\ell) + \text{HgCl}_4^{2-}) = -4.1$ $\log K_s^0 = \log K_e + \log K_T^{\text{ref.2}} - \log \beta_4(\text{Hg}^{2+} + 4\text{Cl}^- = \text{HgCl}_4^{2-})^{\text{ref.5}} - 4 \log \gamma_{\text{NaCl}} = -4.1 + 1.94 - 15.07 - 0.66 = -17.9$	
AUXILIARY INFORMATION	
METHOD: (a) This datum appears in Table 18 of the paper; the accompanying text does not make it clear that it is the author's own determination, rather than Richards and Archibald's ⁶ , but the value does not appear in these authors' paper. (b) This datum apparently obtained from E.m.f. measurements, but these have not been specified. E.m.f. measured by compensation method using capillary electrometer as null instrument. Cells, not described, kept in thermostated bath at 25° . Total mercury in solution determined by precipitation as HgS gravimetrically with no details given.	SOURCE AND PURITY OF MATERIALS: Pure commercial salts (Kahlbaum), not further purified. ESTIMATED ERROR: Large, since experimental details are not given. REFERENCES: 1. Abel, E., Z. anorg. Chem. (1901) 26, 361. 2. Hietanen, S., Sillén, L.G. Ark. Kem. (1956) 10, 103. 3. Bonner, O.D., Unietis, F., J. Am. Chem. Soc. (1953) 75, 5111. 4. Lanier, R.D., J. Phys. Chem. (1965) 69, 3992. 5. Sillén, L.G., Acta. Chem. Scand. (1949) 3, 539. 6. Richards, T.W., Archibald, E.H., Z. physik. Chem. (1907) 49, 385.

5.4. Data of Ley and Heimbucher (1904)

COMPONENTS: 1. Mercury (I) chloride, Hg ₂ Cl ₂ (10112-91-1) 2. Potassium chloride, KCl (7447-40-7) 3. Water, H ₂ O (7732-18-5)	ORIGINAL MEASUREMENTS: Ley, H., Heimbucher, C., Z. Elektrochem. (1904) <u>10</u> , 301.										
VARIABLES: Two KCl concentrations One temperature	PREPARED BY: Y. Marcus, January 1978										
EXPERIMENTAL VALUES: The e.m.f. <i>E</i> of the following cell determined at 20°C Hg/Hg ₂ (ClO ₄) ₂ 0.05M/0.1M KNO ₃ /Hg ₂ Cl ₂ (s), cM KCl/Hg (M = mol dm ⁻³). Found <table><tr><td><i>c</i> = 0.1M</td><td><i>E</i>/V = 0.4193</td><td>0.4197</td><td>0.4200</td><td>0.4199</td></tr><tr><td><i>c</i> = 1.0M</td><td><i>E</i>/V = 0.4721</td><td>0.4717</td><td></td><td></td></tr></table> Authors took [Hg ₂ ²⁺] = 0.047M in 0.05M Hg ₂ (ClO ₄) ₂ (the rest being hydrolyzed species), 0.73 as its degree of dissociation. For 0.1M and 1.0M KCl the degrees of dissociation used were 0.86 and 0.75 respectively. For (RT/2 <i>F</i>)ln 10 the authors took 0.0295V. For the two KCl concentrations they calculated $\log K_{\text{SO}}^{\circ}(\text{Hg}_2\text{Cl}_2(\text{s}) = \text{Hg}_2^{2+} + 2\text{Cl}^-) = -17.83 \text{ (} c = 0.1\text{M) and } -17.95 \text{ (} c = 1.0\text{M)}.$ Calculated from the data (YM), avoiding rounding-off errors, using the accepted value of (RT/2 <i>F</i>)ln 10 at 20°C (0.02908V), and the authors' estimate of "degree of dissociation" and hydrolysis $\log K_{\text{SO}}^{\circ} = -18.03(c = 0.1\text{M}) \text{ and } -17.95 \text{ (} c = 1.0\text{M)}.$ No correction for any liquid junction potentials can, however, be applied.		<i>c</i> = 0.1M	<i>E</i> /V = 0.4193	0.4197	0.4200	0.4199	<i>c</i> = 1.0M	<i>E</i> /V = 0.4721	0.4717		
<i>c</i> = 0.1M	<i>E</i> /V = 0.4193	0.4197	0.4200	0.4199							
<i>c</i> = 1.0M	<i>E</i> /V = 0.4721	0.4717									
AUXILIARY INFORMATION											
METHOD: E.m.f. measured by compensation method. Degree of hydrolysis by comparing rate of inversion of sucrose by 0.05M Hg ₂ (ClO ₄) ₂ with that by (1/250)M HClO ₄ . Degree of dissociation by comparing conductivity of 0.05M Hg ₂ (ClO ₄) ₂ with that of 0.05M Ba(ClO ₄) ₂ in an unspecified manner. Capillary electrometer used as null detector in e.m.f. measurements.	SOURCE AND PURITY OF MATERIALS: Hg ₂ (ClO ₄) ₂ obtained by dissolving Hg ₂ O (prepared by precipitating purest (Merck) Hg ₂ (NO ₃) ₂ with NaOH) in 2M HClO ₄ , and recrystallizing from water. This was then dissolved in conductivity water. Excess acid remaining <0.08%. No sources for other materials given. ESTIMATED ERROR: Main error in assumed activity coefficients (authors' "degree of dissociation"), ±0.08 units in log <i>K</i> _S (after rounding off errors corrected). REFERENCES:										

5.5. Data of Kohlrausch (1908)

COMPONENTS:		ORIGINAL MEASUREMENTS:	
1. Mercury(I) chloride, Hg_2Cl_2 [10112-91-1] 2. Water, H_2O [7732-18-5]		Kohlrausch, F., Z. physik. Chem. (1908) 64, 129.	
VARIABLES:		PREPARED BY:	
Four temperatures		Y. Marcus, January 1978	
EXPERIMENTAL VALUES:			
Solubilities s_{Hg} given in mg mercury(I) chloride per dm^{-2} of saturated solution:			
$t/^{\circ}\text{C}$	0.5	18.0	24.6 (43)
$s_{\text{Hg}}/\text{mg dm}^{-3}$	1.4	2.1*	2.8 7
$10^6 s_{\text{Hg}}/\text{mol dm}^{-3**}$	3.0	4.4	5.9 15
* In the earliest study ¹ , the estimated solubility at this temperature was 3.1 mg dm^{-3} .			
** Calculated by YM.			
AUXILIARY INFORMATION			
METHOD:		SOURCE AND PURITY OF MATERIALS:	
The conductivity of saturated solutions of Hg_2Cl_2 compared with that of $5 \times 10^{-4} \text{ mol dm}^{-3}$ $\text{Hg}_2(\text{NO}_3)_2$, taking into account its temperature coefficient and the expansibility of the solutions. Hydrolysis recognized as contributing to the conductivity, but not corrected for. The apparatus and procedure were described in earlier work ^{1,2} . Material as finely divided powder suspended in conductivity water as often as needed until constant conductivity achieved. Daylight was excluded. Conductivity of the water employed subtracted from that of the saturated solution.		Described in earlier work ^{1,2} . Hg_2Cl_2 precipitated from aqueous solution of $\text{Hg}_2(\text{NO}_3)_2 \cdot x\text{H}_2\text{O}$ with excess HCl . Some re-examined after 2.5 years' storage.	
		ESTIMATED ERROR:	
		Author's estimate: data could be 50% wrong.	
		REFERENCES:	
		1. Kohlrausch, F., Rose, F., Z. physik. Chem. (1893) 12, 234. 2. Kohlrausch, F., Rose, F., Z. physik. Chem. (1903) 44, 197.	

5.6. Data of Herz (1911)

COMPONENTS: 1. Mercury(I) chloride, Hg_2Cl_2 (10112-91-1) 2. Potassium hydroxide, KOH (1310-58-3) 3. Potassium chloride, KCl (7447-40-7) 4. Water, H_2O (7732-18-5)	ORIGINAL MEASUREMENTS: Herz, W., Z. Anorg. Chem. (1911) <u>70</u> , 170.								
VARIABLES: One temperature, varying KOH and KCl concentrations.	PREPARED BY: Y. Marcus, January 1978								
EXPERIMENTAL VALUES: The composition of the aqueous salts in equilibrium with a mixture of solid Hg_2Cl_2 and Hg_2O ($= \text{Hg}(\text{l}) + \text{HgO} ?$) ¹ determined at 25°C: <table><tr><td>$c_{\text{KOH}}/\text{mol dm}^{-3}$</td><td>0.038</td><td>0.018</td><td>0.009</td></tr><tr><td>$c_{\text{KCl}}/\text{mol dm}^{-3}$</td><td>2.177</td><td>1.087</td><td>0.543</td></tr></table> Calculated (YM): $\log K_s(\text{Hg}_2\text{Cl}_2(\text{s}) = \text{Hg}_2^{2+} + 2\text{Cl}^-) = \log (c_{\text{KCl}}/c_{\text{KOH}})^2 + \log K(\text{HgO}(\text{s}) = \text{Hg}^{2+} + 2\text{OH}^-) + \log K(\text{Hg}^{2+} + \text{Hg}(\text{l}) = \text{Hg}_2^{2+}) + \log (y_{\pm\text{KCl}}/y_{\pm\text{KOH}})^2 = 3.61 \pm 0.04 - 25.4$ (ref. 2) + 1.94 (ref. 3) - 0.03 (ref. 4) = -19.9 ± 0.1 . Alternatively, if existence of solid Hg_2O is accepted, with the solubility product $K_s(\text{Hg}_2\text{O}(\text{s}) = \text{Hg}_2^{2+} + 2\text{OH}^-) = 1.8 \times 10^{-24}$ (ref. 5), then $\log K_s(\text{Hg}_2\text{Cl}_2(\text{s})) = \log K_s(\text{Hg}_2\text{O}(\text{s})) + \log (c_{\text{KCl}}/c_{\text{KOH}})^2 + \log (y_{\pm\text{KCl}}/y_{\pm\text{KOH}})^2 =$ $= -23.74 + 3.61 \pm 0.04 - 0.03 = -20.16 \pm 0.05$.		$c_{\text{KOH}}/\text{mol dm}^{-3}$	0.038	0.018	0.009	$c_{\text{KCl}}/\text{mol dm}^{-3}$	2.177	1.087	0.543
$c_{\text{KOH}}/\text{mol dm}^{-3}$	0.038	0.018	0.009						
$c_{\text{KCl}}/\text{mol dm}^{-3}$	2.177	1.087	0.543						
AUXILIARY INFORMATION									
METHOD: Excess solid Hg_2Cl_2 shaken at 25°C for an extended period (unspecified) with aqueous KOH. The remaining base titrated with standardized acid to the phenolphthalein endpoint. The KCl content of the equilibrium solution obtained by difference. Solids assumed to be Hg_2Cl_2 and Hg_2O , but latter probably ¹ disproportionates.	SOURCE AND PURITY OF MATERIALS: Not given ESTIMATED ERROR: Main error arises from ignorance of the nature of the solid " Hg_2O ," hence from K_s of Hg_2O . The concentrations of KOH are imprecise, contributing to the error. REFERENCES: 1. Sillén, L.G., Martell, A.E., Stability Constants, Chem. Soc. Spec. Publ. 17, London, 1964, p. 64. 2. Feitknecht, W., Schindler, P., Pure Appl. Chem. (1963) 6, 130. 3. Hietanen, S., Sillén, L.G., Arkiv Kemi (1956) 10, 103. 4. Harned, H.S., Hamer, W.J., J. Am. Chem. Soc. (1937) 59, 1890. 5. Brodsky, A.E., Z. Electrochem. (1929) 35, 833.								

5.7. Data of Brodsky and Scherschewer (1926)

COMPONENTS:

1. Mercury(I) chloride, Hg_2Cl_2 (10112-91-1)
2. Potassium chloride, KCl (7447-40-7)
3. Water, H_2O (7732-18-5)

ORIGINAL MEASUREMENTS:

Brodsky, A.E., Scherschewer, J.M., Z. Elektrochem. (1926) 32, 1.

VARIABLES:

Temperature
 KCl concentration
 $\text{Hg}_2(\text{NO}_3)_2$ concentration

PREPARED BY:

Y. Marcus, January 1978

EXPERIMENTAL VALUES:

The e.m.f. of the following cell determined ($M = \text{mol dm}^{-3}$)
 $\text{Hg}/\text{Hg}_2(\text{NO}_3)_2, c_0 M, \text{HNO}_3, 0.005M^1/\text{KNO}_3 \text{ satd.}/\text{Hg}_2\text{Cl}_2(s), \text{KCl}cM/\text{Hg}$

$c_0/\text{mol dm}^{-3}$	$t/^{\circ}\text{C}$	10.8	14.9				19.2		26.5	
	$c/\text{mol dm}^{-3}$	0.01	0.1	0.01	0.1	0.5	1.0	0.1	1.0	0.1
		$10^4 E/V$ (Int. volts corrected to volts)								
0.0466		3756	4133	3754	4124	4475	4664	4103		4089
0.0233		3524	4080	3514	4067	4424	4608	4054	4609	4032
0.01165		3463	4023	3465	4002	4366	4555	4000		3977
0.00583		3399	3959	3397	3949	4307	4491	3934	4488	3914
0.00291		3275	3866	3324	3879	4233	4426	3881		3833
0.00146		3256	3825	3256	3807	4164	4350	3813	4435	3749
0.000728		3185	3751	3146	3700	4055	4243	3781	4360	3667
0.000364		3077	3627	3062	3611	3958	4158	4328		
$\log K_{s0}^*(\text{Hg}_2\text{Cl}_2(s) = \text{Hg}_2^{2+} + 2\text{Cl}^-)$		-18.72 \pm 0.09		-18.54 \pm 0.05			-18.33 \pm 0.05		-17.88 \pm 0.05	

* Calculated by Y.M.

AUXILIARY INFORMATION

METHOD:

$\log K_{s0}^* = \lim_{c_0 \rightarrow 0} [(-2FE/RT \ln 10) + \log c_0^2 y_{\pm\text{KCl}}^2]$
where $y_{\pm\text{KCl}}$ is activity coefficient of KCl at temp. T and molar concentration c . Authors' values of F and $y_{\pm\text{KCl}}^1$ replaced by modern values. Extrapolation to $c_0 = 0$ intended to eliminate $y_{\text{K}^+}/y_{\text{Hg}_2^{2+}} y_{\text{Cl}^-}$.
E.m.f. measured by compensation method with calibrated Weston cell standard, a mirror galvanometer as null instrument, and half-cells which were intercompared for consistency. No thermostat was used, but e.m.f.'s and temperatures steady for 4 hrs at least.

SOURCE AND PURITY OF MATERIALS:

$\text{Hg}_2(\text{NO}_3)_2$ freshly prepared from salt of undisclosed source.

ESTIMATED ERROR:

Errors given for $\log K_{s0}^*$ are the nonsystematic errors from the relationship of the measured E , c_0 , c and T , and the extrapolation. The systematic error inherent in the latter is less than ± 0.1 .

REFERENCES:

1. Brodsky, A.E., Z. Elektrochem. (1929) 35, 833.

5.8. Data of Eversole (1932)

COMPONENTS:	ORIGINAL MEASUREMENTS:
1. Mercury(I) chloride, Hg ₂ Cl ₂ (10112-91-1)	Eversole, W.G., McLachlan, R.W., J. Am. Chem. Soc. (1932) <u>52</u> , 864.
2. Perchloric acid, HClO ₄ (7601-90-3)	
3. Nitric acid, HNO ₃ (7697-37-2)	
4. Water, H ₂ O (7732-18-5)	
VARIABLES:	PREPARED BY:
One temperature, several concentrations of HClO ₄ or HNO ₃ .	Y. Marcus, January 1978
EXPERIMENTAL VALUES:	
Concentration of soluble mercury species in equilibrium with solid Hg ₂ Cl ₂ at 25.0°C in solutions of the acids	
c _{HClO₄} /mol dm ⁻³ :	0.0005 0.005 0.01 0.1 0.2
10 ⁶ c _{Hg(total)} /mol dm ⁻³ :	4.7 4.8 5.0 5.5 6.3
c _{HNO₃} /mol dm ⁻³ :	0.005 0.01 0.1
10 ⁶ c _{Hg(total)} /mol dm ⁻³ :	5.2 5.6 5.8
Calculated (Y.M.): The contributions of the species HgNO ₃ ⁺ , HgOH ⁺ , Hg(OH) ₂ are negligible, those of Hg ₂ ClO ₄ ⁺ , Hg ₂ NO ₃ ⁺ and HgCl ⁺ are small ¹⁻³ , so that Hg ₂ ²⁺ and HgCl ₂ are the main species. Thus to a good approximation	
c _{Hg} = (K _{SO} ^o /4) ^{1/3} y _{Hg₂Cl₂} ⁻¹ {1+β ₁ (A) [A ⁻] + 2(K _{SO} ^o /4) ^{1/3} β ₁ K _r ⁻¹ } + K _{SO} ^o K _r ⁻¹ β ₂ (y _{HgCl₂} /y _{Hg₂Cl₂}) ³	
Following estimates were used: y _{Hg₂Cl₂} ≈ y _{HgCl₂} = 10 ^{-3√I/(1+1.6√I)} ; I = c _{acid} ; β ₁ (ClO ₄ ⁻) = 0.9 mol dm ⁻³ ; β ₁ (NO ₃ ⁻) = 2.25 mol dm ⁻³ ; β ₁ (HgCl ⁺) = 10 ^{7.34} ; β ₂ (HgCl ₂) = 10 ^{14.26} ; K _r = 10 ^{1.94} . These gave for all acid concentrations by a short iteration the consistent value log K _{SO} ^o = -17.74±0.02.	
AUXILIARY INFORMATION	
METHOD:	SOURCE AND PURITY OF MATERIALS:
Increasing volumes of very dilute (10 ⁻⁴ mol dm ⁻³) aqueous Hg ₂ (ClO ₄) ₂ and KCl added together with excess water, until incipient precipitation could be observed by Tyndall effect. Equilibrium tested by cooling 0.2°C, then reheating to 25.0°C, with no change in results. Solubility given as that solution which is just short of showing Tyndall effect. Solutions agitated for >4 hrs at 25°C regulated to ±0.02°C and carefully guarded against dust.	H ₂ O redistilled from KMnO ₄ ; KCl, c.p. twice recryst. from redist. H ₂ O; Hg ₂ (ClO ₄) ₂ prepared by authors ⁴ ; HNO ₃ , c.p., twice redistilled and nitrous gases expelled; HClO ₄ distilled in vacuum.
Sealed pyrex glass vessels used. A home-made Tyndallometer was employed, in conjunction with a Zeiss Pulfrich gradation photometer.	ESTIMATED ERROR: Error in solubility data ca. 2%, to which error in K _{SO} ^o is proportional. Main uncertainties are in log K _r (±0.02) and log β ₂ (±0.09), contributing ±0.10 systematic error in log K _{SO} ^o .
	REFERENCES:
	1. Hietanen, S., Sillén, L.G., Arkiv Kemi (1956), <u>10</u> , 103.
	2. Hietanen, S., Sillén, L.G., Acta Chem. Scand. (1952) <u>6</u> , 747.
	3. Sillén, L.G., Acta Chem. Scand. (1949) <u>3</u> , 539.
	4. Popoff, S., J. Am. Chem. Soc. (1931) <u>53</u> , 1195.

5.9. Data of Garrett, Noble and Miller (1942)

COMPONENTS:		ORIGINAL MEASUREMENTS:		
1. Mercury(I) chloride, Hg ₂ Cl ₂ (10112-91-1)		Garrett, A.B., Noble, M.V., Miller, S.,		
2a. Sodium chloride, NaCl (7647-14-5)		J. Chem. Educ. (1942) 19, 485.		
2b. Calcium chloride, CaCl ₂ (10043-52-4)				
3. Water (7732-18-5)				
VARIABLES:		PREPARED BY:		
One temperature, several NaCl or CaCl ₂ concentrations		Y. Marcus, January 1978		
EXPERIMENTAL VALUES:				
1. The solubility of Hg ₂ Cl ₂ in water at 25°C is 5.9×10 ⁻⁶ mol (kg H ₂ O) ⁻¹ .				
2. The solubility of Hg ₂ Cl ₂ in aqueous salt solution at 25°C is				
	m/mol kg ⁻¹	c*/mol dm ⁻³	10 ⁶ m _{Hg,total} /mol kg ⁻¹	10 ⁶ c _{Hg,total} */mol dm ⁻³
NaCl	0.101	0.100	8.7	8.6
	1.02	1.00	88.4	86.7
	2.12	2.03 ₄	185.	177.
CaCl ₂	0.374	0.370	48.5	48.0
	0.528	0.522	74.2	73.4
	1.05	1.02	170.	165.
* Calculated (Y.M.). Calculated also log K _{SO} ^o = lim _{m_{Cl⁻}→0} [log(c _{Hg,total} /c _{Cl⁻} ²)				
- log(1 + β ₃ β ₄ ⁻¹ c _{Cl⁻} ⁻¹ + β ₂ β ₄ ⁻¹ c _{Cl⁻} ⁻²) - 2A ^{1/2} /(1 + B ^{1/2})] + log K _r (Hg ²⁺ + Hg(l)) = Hg ₂ ²⁺)				
- log β ₄ - 4 log γ _± (trace NaCl in 0.5M NaClO ₄) ^{ref.3} = -17.95±0.08.				
AUXILIARY INFORMATION				
METHOD:		SOURCE AND PURITY OF MATERIALS:		
β _i (Hg ²⁺ + iCl ⁻ = HgCl _i ²⁻ⁱ) = 10 ^{13.22} for i=2, 10 ^{14.07} for i=3, 10 ^{15.07} for i=4, valid for 0.5M (mol dm ⁻³) NaClO ₄ . ¹ A = 0.51 and B = 1.6, I = ionic strength, is arbitrary form of activity coefficient term, the rest of which is taken into account by the extrapolation. K(Hg ²⁺ + Hg(l) = Hg ₂ ²⁺) = 88 (ref. 2).		No details given.		
No details are given of the apparatus and procedure.		ESTIMATED ERROR: Mainly from activity coefficient terms, necessitating the extrapolation, and uncertainties in the constants		
REFERENCES: 1. Sill��n, L.G., Acta Chem. Scand. (1949) 3. 539.				
2. Hietanen, S., Sill��n, L.G., Arkiv Kemi (1956) 10, 103.				
3. Estimated (YM) from Robinson, R.A., Hiteck, R.H., Electrolyte Solutions, Butterworth, London, 2nd Ed., 1959.				

5.10. Data of Law (1946)

COMPONENTS: 1. Mercury(I) chloride, Hg_2Cl_2 (10112-91-1) 2. Sodium chloride, NaCl (7647-14-5) 3. Perchloric acid, HClO_4 (7601-90-3) 4. Water, H_2O (7732-18-5)	ORIGINAL MEASUREMENTS: Law, J.T., M.S. Thesis, University of New Zealand, 1946.
VARIABLES: Temperatures, between 15 and 45°C; concentrations of NaCl and HClO_4 .	PREPARED BY: Y. Marcus, January 1978
EXPERIMENTAL VALUES: E.m.f.'s of the cell $\text{Hg}/\text{Hg}_2\text{Cl}_2(\text{s}), \text{NaCl} (1.5m)m, \text{HClO}_4 (m-mx)m/\text{HClO}_4 (m+0.5mx)m/\text{Hg}_2(\text{ClO}_4)_2 (0.5mx)m, \text{HClO}_4 (m-mx)m/\text{Hg}$ measured at 5K intervals between $288.15 \leq T/K \leq 318.15$ for various concentrations in the ranges $0.02 \leq m \leq 0.05$ and $0.2 \leq x \leq 0.6$. The results were summarized as follows: 1. $\lim_{m \rightarrow 0} [\lim_{x \rightarrow 0, m \text{ const.}} (E + (RT/2F) \ln(4.5mx))] = 0.5108V_{\text{int}} = 0.5110V$ for $T/K=298.15$ (ref. 1) 2. $\Delta G^\circ(\text{Hg}_2\text{Cl}_2(\text{s}) = \text{Hg}_2^{2+} + 2\text{Cl}^-)/\text{kJ mol}^{-1} = 42.947 + 0.3243T - 4.588T^2$ (ref. 2) $\Delta H^\circ/\text{kJ mol}^{-1} = 42.947 + 4.588T^2$ (ref. 2) $\Delta S^\circ/\text{JK}^{-1}\text{mol}^{-1} = -324.3 + 0.9177T$ (ref. 2) 3. $E^\circ(\text{Hg}/\text{Hg}_2^{2+}) = 0.7789V$ for $T/K = 298.15$ (ref. 2) Calculated (Y.M.) from 1. $\log K_{\text{SO}}^\circ(\text{Hg}_2\text{Cl}_2(\text{s})) = -0.5110/(RT \ln 10/2F) = -17.275$; from 2. $\log K_{\text{SO}}^\circ = -\Delta G^\circ/RT \ln 10 = -98849.5/5708 = -17.318$; from 3. $\log K_{\text{SO}}^\circ = (-0.7789 + E^\circ(\text{Hg}/\text{Hg}_2\text{Cl}_2))/ (RT \ln 10/2F) = (-0.7789 + 0.2680)/0.02958 = -17.272$, all at 298.15K	
AUXILIARY INFORMATION	
METHOD: Information as to apparatus/procedure is not available to evaluator, who saw only quotations of the work in refs. 1 and 2.	SOURCE AND PURITY OF MATERIALS: Information is not available to evaluator, who saw only quotations of the work in refs. 1 and 2.
ESTIMATED ERROR:	
REFERENCES: 1. Berecki, C., Biedermann, G., Sillén, L.G. Report to Commission V.6 of IUPAC (rather, its precursor), 1953. 2. Galloway, W.J., M.S. Thesis, University of New Zealand, 1961.	

5.11. Data of Jonsson, Qvarfort and Sillén (1947)

COMPONENTS: 1. Mercury(I) chloride, Hg_2Cl_2 (10112-91-1) 2. Sodium chloride, NaCl (7647-14-5) 3. Sodium perchlorate, NaClO_4 (7601-89-0) 4. Water, H_2O (7732-18-5)	ORIGINAL MEASUREMENTS: Jonsson, A., Qvarfort, I., Sillén, L.G., Acta Chem. Scand. (1947) 1, 461.
VARIABLES: One temperature, constant ionic strength, variable NaCl concentration.	PREPARED BY: Y. Marcus, January 1978.
EXPERIMENTAL VALUES: Measured at 25°C: 1) E.m.f. $E/V = E_1^0/V + (RT/2F)\ln(b/\text{mol dm}^{-3})$ of cell $\text{Hg}, \text{Hg}_2\text{Cl}_2(\text{s}), \text{NaCl } 4\text{M}/\text{NaClO}_4 \text{ } 0.5\text{M}/\text{Hg}_2(\text{ClO}_4)_2 \text{ } b\text{M}, \text{HClO}_4 \text{ } 0.010\text{M}, \text{NaClO}_4 \text{ } 0.490\text{-}3b \text{ M}/\text{Hg}$ $(E_1^0 = \text{authors' "millimolar potential"} E_{10} + (6RT/F)\ln 10; \text{M} \equiv \text{mol dm}^{-3})$. 2) E.m.f. $E/V = E_2^0/V - (RT/F)\ln(c/\text{mol dm}^{-3})$ of cell $\text{Hg}, \text{Hg}_2\text{Cl}_2(\text{s}), \text{NaCl } 4\text{M}/\text{NaClO}_4 \text{ } 0.5\text{M}/\text{Hg}_2\text{Cl}_2(\text{s}), \text{NaCl } c\text{M}, \text{HClO}_4 \text{ } 0.010\text{M}, \text{NaClO}_4 \text{ } 0.490\text{-}c\text{M}/\text{Hg}$ $(E_2^0 = \text{authors' "millimolar potential"} E_{1X} + (3RT/F)\ln 10)$ For one experiment, b was specified as $4.83 \times 10^{-3}/(1+0.01v/\text{cm}^3)\text{M}$, and $c = 0.05(v/\text{cm}^3)/(100 + v/\text{cm}^3)\text{M}$. For v see under Method. Six values of $\Delta E^0 - E_2^0 - E_1^0$ obtained over a period of a year are reported: $\Delta E^0/V = -0.4990, -0.4994, -0.4995, -0.4995, -0.4993, -0.4995$ average: -0.4993 ± 0.0003 . Hence $\log K_{\text{SO}}^1 = \Delta E^0/(RT/2F)\ln 10 = -16.88 \pm 0.01$ for 0.5M NaClO_4 medium. Correction for activity coefficients ² is -1.05 , hence $\log K_s = -17.93$.	
AUXILIARY INFORMATION	
METHOD: 100cm^3 of a solution containing $b\text{M}$ $\text{Hg}_2(\text{ClO}_4)_2 + 0.008b \text{ M Hg}(\text{ClO}_4)_2 + 0.010\text{M}$ $\text{HClO}_4 + 0.490 - 3b \text{ M NaClO}_4$, connected via a 0.5 M NaClO_4 bridge to the $\text{Hg}, \text{Hg}_2\text{Cl}_2(\text{s})$, 4M NaCl reference electrode, were titrated with $v\text{cm}^3$ of a solution containing $c\text{M NaCl} +$ $0.010 \text{ M HClO}_4 + 0.490 - c\text{M NaClO}_4$. Care was taken to exclude O_2 from the solutions by bubbling purified and water vapor presaturated nitrogen. Salt bridge designed to minimize diffusion of contaminating ions. E.m.f. measured to $\pm 0.1 \text{ mV}$ by Radiometer potentiometer, calibrated with Weston cell, or with Raps or Jensen compensators and Multiflex galvanometer to $\pm 0.05 \text{ mV}$. Temperature kept at $25.0 \pm 0.1^\circ\text{C}$ of all parts of cell and buret.	SOURCE AND PURITY OF MATERIALS: NaClO_4 thrice recrystallized from 70% ethanol and dried. NaCl freed from bromide ³ . Hg twice distilled. $\text{Hg}_2(\text{ClO}_4)_2$ prepared ⁴ from Hg, HgO (Merck) and HClO_4 (Schering) analytical reagents. Hg_2Cl_2 was formed during titrations, but no description given of that salt in the (left hand) reference electrode. ESTIMATED ERROR: Error in $\log K_{\text{SO}}^1$ is ± 0.01 , from $\pm 0.3\text{mV}$ error in ΔE^0 . REFERENCES: 1. Sillén, L.G., Svensk Kem. Tidskr. (1946) 58, 52. 2. Sillén, L.G., Acta Chem. Scand. (1949) 3, 539; Qvarfort, I., Sillén, L.G., Acta Chem. Scand. (1949) 3, 517. 3. Guntelberg, E., Dissertation, Univ. Copenhagen (1938). 4. Pugh, W., J. Chem. Soc. (1937), 1824.

5.12. Data of Dry and Gledhill (1955)

COMPONENTS: 1. Mercury(I) chloride, Hg_2Cl_2 (10112-91-1) 2. Water, H_2O (7732-18-5)	ORIGINAL MEASUREMENTS: Dry, M.E., Gledhill, J.A., Trans. Faraday Soc. (1955) <u>51</u> , 1119.
VARIABLES: One temperature	PREPARED BY: Y. Marcus, January 1978
EXPERIMENTAL VALUES: 1. Conductivity of a saturated aqueous Hg_2Cl_2 solution at 25°C is $350.8 \pm 1.0 \mu\text{Sm}^{-1}$. 2. The pH of a saturated aqueous Hg_2Cl_2 solution at 25°C is 5.085 ± 0.010 . 3. The total concentration of soluble mercury species in a saturated aqueous Hg_2Cl_2 solution at 25°C is $(7.5 \pm 0.3) \times 10^{-6} \text{ mol dm}^{-3}$. In an iterative procedure, utilizing equilibrium constants ¹⁻³ and estimates of activity coefficient terms, authors estimated following concentrations, in $10^{-6} \text{ mol dm}^{-3}$ to be present in the saturated solution: H^+ 8.17 ± 0.08 ; Cl^- 8.40 ± 0.14 ; $\text{Hg}(\text{OH})_2$ 4.03 ± 0.10 ; HgCl_2 3.3 ± 0.4 ; Hg_2OH^+ 0.10 ; HgCl^+ 0.027 ; Hg_2^{2+} 0.0190 ± 0.0001 ; HgOH^+ 0.014 . These values sum up to the total concentration and are consistent with the pH and the conductivity (which is 99.5% due to H^+ and Cl^-). The ionic strength is $8.4 \times 10^{-5} \text{ mol dm}^{-3}$, hence: $\log K_{\text{SO}}^\circ (\text{Hg}_2\text{Cl}_2(\text{s}) = \text{Hg}_2^{2+} + 2\text{Cl}^-) = \log(1.9 \times 10^{-8}) + 2 \log(8.4 \times 10^{-6}) + 3 \log \gamma_{\text{Hg}_2\text{Cl}_2} =$ $= -17.873 - 0.009 = -17.882.$	
AUXILIARY INFORMATION	
METHOD: Conductivity data extrapolated to zero time to correct for ion exchange with vessel walls, and to zero bubbling rate of nitrogen. Total mercury determined with dithizone ⁴ at 490 nm in supernatant solutions filtered through a sintered glass filter, made 0.06 mol dm^{-3} in HCl . Treatment with Cl_2 and boiling gave same results as without treatment. Dithizone applied in equal volume of CCl_4 , and spectrophotometric readings compared with those from known amounts of HgCl_2 , similarly treated. The pH measured with a glass electrode pH-meter, N_2 bubbled to remove dissolved CO_2 , instrument standardized at pH=4.005 for 0.05 mol dm^{-3} potassium hydrogen phthalate. A Cenco-Sheard "photometer" used for the mercury-dithizone spectrophotometric determinations. Results were independent from addition of liquid Hg to the samples for all measurements.	SOURCE AND PURITY OF MATERIALS: Hg_2Cl_2 was precipitated in the cold from pure $\text{Hg}_2(\text{NO}_3)_2$ or $\text{Hg}_2(\text{ClO}_4)_2$ with KCl solutions at equivalent concentrations, digested and washed 50 times with conductivity water. Latter had conductivities between 1.1 and $2.4 \mu\text{Sm}^{-1}$. ESTIMATED ERROR: No data provided to evaluate the reported random error of 4% of the total solubility, but procedure reported should eliminate systematic errors. REFERENCES: 1. Sillén, L.G., Acta Chem. Scand. (1949) <u>3</u> , 539. 2. Hietanen, S., Sillén, L.G., Acta Chem. Scand. (1952) <u>6</u> , 747. 3. Forsling, W., Hietanen, S., Sillén, L.G., Acta Chem. Scand. (1952) <u>6</u> , 901. 4. Fischer, H., Leopoldi, G., Z. anal. Chem. (1935) <u>103</u> , 241.

5.13. Data of Galloway (1961)

COMPONENTS: 1. Mercury(I) chloride, Hg_2Cl_2 (10112-91-1) 2. Barium chloride, BaCl_2 (10361-37-2) 3. Perchloric acid, HClO_4 (7601-90-3) 4. Water, H_2O (7732-18-5)	ORIGINAL MEASUREMENTS: Galloway, W.J., M.S. Thesis, Univ. of New Zealand, 1961.																
VARIABLES: Seven temperatures, five ionic strengths	PREPARED BY: Y. Marcus, January 1978																
EXPERIMENTAL VALUES: Measured and reported e.m.f. E of the cell $\text{Hg}, \text{Hg}_2\text{Cl}_2/\text{BaCl}_2 (1/2)x\text{mm}, \text{HClO}_4 (1-x)\text{mm}/\text{Ba}(\text{ClO}_4)_2 (1/3)x\text{mm}, \text{HClO}_4 (1-x)\text{mm}/\text{Hg}_2(\text{ClO}_4)_2 (1/3)x\text{mm}, \text{HClO}_4 (1-x)\text{mm}/\text{Hg}_2\text{Cl}_2, \text{Hg}$ at seven temperatures at intervals of 5°C from 15 to 45°C , varying the concentrations in the range $0.005 \leq m$ (five values) ≤ 0.05 and $0.1 \leq x$ (four values) ≤ 0.6 . Calculated and reported the double limit $\lim_{m \rightarrow 0} [\lim_{x \rightarrow 0} (E - (3RT/2F) \ln(xm/3) - (RT/F) \ln 2) + 3RTAm^{1/2} / (1+m^{1/2})] = -(RT/2F) \ln K_{\text{so}}^\circ$ <table><tr><td>$t/^\circ\text{C}$</td><td>15</td><td>20</td><td>25</td><td>30</td><td>35</td><td>40</td><td>45</td></tr><tr><td>$-(RT/2F) \ln K_{\text{so}}^\circ / V$</td><td>0.5273</td><td>0.5272</td><td>0.5273</td><td>0.5277</td><td>0.5280</td><td>0.5284</td><td>0.5291</td></tr></table> Expressed this as the polynomial $-(RT/2F) \ln K_{\text{so}}^\circ / V = 1.3587 - 7.59 \times 10^{-3} T + 2.273 \times 10^{-5} T^2 - 2.222 \times 10^{-8} T^3$ from least squares fitting. Hence obtained expressions for the standard thermodynamic functions for the process $\text{Hg}_2\text{Cl}_2(\text{s}) = \text{Hg}_2^{2+}(\text{aq}) + 2\text{Cl}^-(\text{aq})$ as a function of T , and specifically for $T = 298.15\text{K}$: $\log K_{\text{so}}^\circ = -17.827$, $\Delta G_{\text{so}}^\circ = 101.77 \text{ kJ mol}^{-1}$, $\Delta H_{\text{so}}^\circ = 99.65 \text{ kJ mol}^{-1}$ and $\Delta S_{\text{so}}^\circ = -7.10 \text{ JK}^{-1} \text{ mol}^{-1}$		$t/^\circ\text{C}$	15	20	25	30	35	40	45	$-(RT/2F) \ln K_{\text{so}}^\circ / V$	0.5273	0.5272	0.5273	0.5277	0.5280	0.5284	0.5291
$t/^\circ\text{C}$	15	20	25	30	35	40	45										
$-(RT/2F) \ln K_{\text{so}}^\circ / V$	0.5273	0.5272	0.5273	0.5277	0.5280	0.5284	0.5291										
AUXILIARY INFORMATION																	
METHOD: Extrapolation procedure for E° of the cell removes first the effect of the chloride concentration at a constant ionic strength, m , and then extrapolates to zero ionic strength a function partly corrected for activity coefficients. The points at $m=0.005\text{m}$ were disregarded in this extrapolation (to avoid hydrolysis effects). Other activity coefficient expressions, within reason, lead to similar results. A Cambridge Instrument potentiometer, with calibrated Weston std. cell (1.01861V), Galvanometer readable to 0.2 mA, permitting precision of $\pm 0.1\text{mV}$ (at $45^\circ\text{C} \pm 0.2\text{mV}$). Temperature regulated to $\pm 0.02^\circ\text{C}$. Electrodes prepared according to Hilles and Ives ¹ . Deoxygenated N_2 bubbled through solutions or kept above them for removal of O_2 .	SOURCE AND PURITY OF MATERIALS: Analytical reagent HClO_4 , BaCl_2 , HCl , $\text{Ba}(\text{OH})_2$; prepared solutions of $\text{Hg}_2(\text{ClO}_4)_2$ by dissolving HgO in HClO_4 , recrystallizing twice from dilute HClO_4 , and reducing with Hg under CO_2 atmosphere, and keeping it with CO_2 and Hg . Prepared Hg_2Cl_2 by anodic dissolution of Hg in dilute HCl , washing with HCl and H_2O and drying. $\text{Ba}(\text{ClO}_4)_2$ prepared by dissolving $\text{Ba}(\text{OH})_2$ in HClO_4 . ESTIMATED ERROR: The precision of $\log K_{\text{so}}^\circ$ is ± 0.003 , while the accuracy is within ± 0.014 . REFERENCES: 1. Hilles, G.J., Ives, D.J.G., J. Chem. Soc. (1951) 154, 311.																

5.14. Data of Hansen, Izatt and Christensen (1963)

COMPONENTS: 1. Mercury(I) chloride, Hg_2Cl_2 (10112-91-1) 2. Sodium chloride, NaCl (7647-14-5) 3. Sodium perchlorate, NaClO_4 (7601-89-0) 4. Water, H_2O (7732-18-5)	ORIGINAL MEASUREMENTS: Hansen, L.D., Izatt, R.M., Christensen, J.J. Inorg. Chem. (1963) <u>2</u> , 1243.
VARIABLES: Two temperatures, constant ionic strength, variable NaCl concentration.	PREPARED BY: Y. Marcus, January 1978
EXPERIMENTAL VALUES: Measured e.m.f. of cells ($M \equiv \text{mol dm}^{-3}$) $\text{Hg, Hg}_2\text{Cl}_2(\text{s})/\text{NaClO}_4$ 0.40M, HClO_4 0.1M/ $\text{Hg}_2(\text{ClO}_4)_2$ bM, NaClO_4 (0.40-3b)M, HClO_4 0.10M/ Hg $\text{Hg, Hg}_2\text{Cl}_2(\text{s})/\text{NaClO}_4$ 0.40M, HClO_4 0.1M/ $\text{Hg}_2\text{Cl}_2(\text{s})$, NaCl cM, NaClO_4 (0.40-c)M, HClO_4 0.10M/ Hg No primary data reported, only final results of calculations: $\log K'_{\text{SO}} = -18.19$ at 7°C $\log K'_{\text{SO}} = -16.16$ at 40°C valid for 0.40M NaClO_4 + 0.10M HClO_4 medium. Measurements and calculations followed closely those of Jonsson, Qvarfort and Sillén ¹ . If estimate of -1.05 for activity coefficient term for 0.49M NaClO_4 + 0.01M HClO_4 at 25°C is taken to be valid also for 0.40M NaClO_4 + 0.10M HClO_4 medium at 7 and 40°C , then $\log K^\circ_{\text{SO}} = -19.24$ at 7°C $\log K^\circ_{\text{SO}} = -17.21$ at 40°C	
AUXILIARY INFORMATION	
METHOD: Procedure of measurement similar to that in ref. 1, except that E° for $\text{Hg}/\text{Hg}_2^{2+}$ ob- tained from the appropriate concentration cell, rather than from titration, while E° for $\text{Hg}/\text{Hg}_2\text{Cl}_2-2\text{Cl}^-$ obtained from titration of freshly precipitated Hg_2Cl_2 with excess NaCl solution.	SOURCE AND PURITY OF MATERIALS: Solutions of HClO_4 , NaClO_4 and NaCl pre- pared from commercial reagents of highest purity. $\text{Hg}_2(\text{ClO}_4)_2$ solutions prepared by equilibrating Hg , HgO and HClO_4 and fil- tration. ESTIMATED ERROR: Precision given by authors on $\log K'_{\text{SO}}$ is ± 0.007 . Precision for $\log K^\circ_{\text{SO}}$, due to uncertainty in activity coefficient term is ± 0.08 . REFERENCES: 1. Jonsson, A., Qvarfort, I., Sillén, L.G., Acta Chem. Scand. (1947) <u>1</u> , 461.

5.15. Other data

COMPONENTS: 1. Mercury(I) chloride, Hg ₂ Cl ₂ (10112-91-1) 2. Water, H ₂ O (7732-18-5)	ORIGINAL MEASUREMENTS: Unknown ¹								
VARIABLES: Three temperatures	PREPARED BY: Y. Marcus, January 1978								
EXPERIMENTAL VALUES: The solubility of Hg ₂ Cl ₂ in water given as <table><tr><td>t/°C</td><td>0</td><td>20</td><td>40</td></tr><tr><td>10⁶s_{Hg}/mol kg⁻¹</td><td>3.0</td><td>4.9</td><td>12.7</td></tr></table> The solubility product is reported as 2×10 ⁻¹⁸ mol ² kg ⁻² at 25°C		t/°C	0	20	40	10 ⁶ s _{Hg} /mol kg ⁻¹	3.0	4.9	12.7
t/°C	0	20	40						
10 ⁶ s _{Hg} /mol kg ⁻¹	3.0	4.9	12.7						
AUXILIARY INFORMATION									
METHOD: The report appears in a compilation ¹ which does not give its sources.	SOURCE AND PURITY OF MATERIALS:								
	ESTIMATED ERROR:								
	REFERENCES: 1. Freier, R.K., Aqueous Solutions, de Gruyter, Berlin (1976), p. 68.								