

Triplet-Triplet Absorption Spectra of Organic Molecules in Condensed Phases

Cite as: Journal of Physical and Chemical Reference Data **15**, 1 (1986); <https://doi.org/10.1063/1.555770>
Published Online: 15 October 2009

Ian Carmichael, and Gordon L. Hug



View Online



Export Citation

ARTICLES YOU MAY BE INTERESTED IN

Extinction Coefficients of Triplet-Triplet Absorption Spectra of Organic Molecules in Condensed Phases: A Least-Squares Analysis

Journal of Physical and Chemical Reference Data **16**, 239 (1987); <https://doi.org/10.1063/1.555782>

Triplet-Singlet Transitions in Organic Molecules. Lifetime Measurements of the Triplet State
The Journal of Chemical Physics **17**, 905 (1949); <https://doi.org/10.1063/1.1747085>

A Theory for the Triplet-Triplet Absorption Spectra of Porphyrins

The Journal of Chemical Physics **33**, 1523 (1960); <https://doi.org/10.1063/1.1731436>

Where in the **world** is AIP Publishing?
Find out where we are exhibiting next



Triplet-Triplet Absorption Spectra of Organic Molecules in Condensed Phases

Ian Carmichael and Gordon L. Hug

Radiation Chemistry Data Center, Radiation Laboratory, University of Notre Dame, Notre Dame, IN 46556, U.S.A.

We present a compilation of spectral parameters associated with triplet-triplet absorption of organic molecules in condensed media. The wavelengths of maximum absorbance and the corresponding extinction coefficients, where known, have been critically evaluated. Other data, for example, lifetimes, energies and energy transfer rates, relevant to the triplet states of these molecules are included by way of comments but have not been subjected to a similar scrutiny. Work in the gas phase has been omitted; as have theoretical studies. We provide an introduction to triplet state processes in solution and solids, developing the conceptual background and offering an historical perspective on the detection and measurement of triplet state absorption. Techniques employed to populate the triplet state are reviewed and the various approaches to the estimation of the extinction coefficient of triplet-triplet absorption are critically discussed. A statistical analysis of the available data is presented and recommendations for a hierarchical choice of extinction coefficients are made. Data collection is expected to be complete through the end of 1984. Compound name, molecular formula and author indexes are appended.

Key words: condensed phase; data compilation; extinction coefficients; glass, organic; lifetimes; photophysical processes; review; solution; triplet states; triplet-triplet absorption.

Contents

1. Introduction	2	3.1.5. Special Problems in Flash Photolysis	11
1.1. Conceptual Background	3	3.1.6. Summary of Advantages and Disadvantages	12
1.2. Historical Perspective	4	3.2. Singlet Depletion (SD) Method	12
2. Methods for Populating and Detecting the Triplet State	5	3.2.1. Description of Method	12
2.1. Generation of Triplets by Pulse Radiolysis	6	3.2.2. Reconstruction of True Transient Absorbance	13
2.1.1. Theoretical Concepts	6	3.2.3. Corrections for Triplet Decay...	13
2.1.2. Special Considerations for Pulse Radiolysis	6	3.2.4. Summary of Advantages and Disadvantages	13
2.2. Generation of Triplets by Photolysis....	7	3.3. Total Depletion (TD) Method	14
2.2.1. Theoretical Concepts	7	3.3.1. Description of Method	14
2.2.2. Special Considerations for Photolysis	7	3.3.2. A Three-state Excitation- deactivation Model	14
2.3. Miscellaneous Methods	9	3.3.3. Corrections for Triplet Decay...	15
3. Methods for Measuring Extinction Coefficients	9	3.3.4. Summary of Advantages and Disadvantages	15
3.1. Energy Transfer (ET) Method	9	3.4. Relative Actinometry (RA) Method	15
3.1.1. Description of Method	9	3.4.1. Description of Method	15
3.1.2. Kinetic Corrections	9	3.4.2. Corrections for Singlet Depletion	15
3.1.3. Uncertainty in Probability of Transfer	10	3.4.3. Corrections for Triplet Decay...	15
3.1.4. Special Problems in Radiolytic Excitation	10	3.4.4. Corrections for Laser Excitation.	16
		3.4.5. Extensions	16
		3.4.6. Summary of Advantages and Disadvantages	16
		3.5. Intensity Variation (IV) Method	16
		3.5.1. Description of Method	16

©1986 by the U.S. Secretary of Commerce on behalf of the United States. This copyright is assigned to the American Institute of Physics and the American Chemical Society.
Reprints available from ACS; see Reprints List at back of issue.

3.5.2. Summary of Advantages and Disadvantages	17	5. Arrangement of the Data Table	28
3.6. Kinetic Method (KM)	17	5.1. List of Abbreviations	29
3.6.1. Description of Method	17	5.2. List of Symbols	29
3.6.2. An Alternative Derivation of the Kinetic Method	17	6. Acknowledgments	30
3.6.3. Summary of Advantages and Disadvantages	19	7. References to Text	30
3.7. Partial Saturation (SM) Method	19	8. Table 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases	33
3.7.1. Description of Method	19	9. References to Table 6	184
3.7.2. An Alternate Model for Photoexcitation	19	10. Indexes to Table 6	205
3.7.3. Summary of Advantages and Disadvantages	20	10.1. Compound Name Index	205
3.8. Miscellaneous Methods for Measuring Extinction Coefficients	20	10.2. Compound Formula Index	228
3.8.1. Electron Spin Resonance (ESR)	20	10.3. Author Index	240
3.8.2. Other Comparative Techniques (HAT,ELT,RF)	21		
3.8.3. Pulsed Intensity Variation (PIV)	21		
4. Evaluation Procedure	22		
4.1. General Methodology	22		
4.2. A Statistical Analysis	23		
4.2.1. Statistical Test of Solvent and Temperature Effects	23		
4.2.2. Mean ϵ and Confidence Intervals	25		
4.2.3. Discussion of Statistical Results	26		
4.3. Tentative Standard ϵ 's	27		

List of Tables

1. Excitation methods per 5-year period.	5
2. Extinction methods per 5-year period.	5
3. t -tests of solvent effects in anthracene.	24
4. ANOVA table for anthracene.	24
5. Average extinction coefficients.	26
6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases	33

List of Figures

1. Rate processes connecting electronic energy levels.	4
---	---

1. Introduction

In 1971, as part of the classic volumes on organic molecular photophysics edited by the late J.B. Birks, Labhart and Heinzelmann contributed a review¹ which contained an extensive compilation of spectral data relevant to the absorption of light by the excited triplet state of organic molecules in condensed phases. The energies of absorption maxima in both fluid solution and in various glasses and solids were recorded together with the corresponding extinction coefficients where known. This compilation was neither limited to a specific class of compounds nor to a particular experimental method as had been the case with previous reviews^{2,3}. Data on the polarization properties of triplet-triplet absorption transitions in rigid matrices were also listed, and a bibliography of theoretical calculations on and assignments of triplet spectra was given. A short discussion of all techniques then currently in use for both transient excitation and extinction coefficient determination was provided and the importance of correct identification of the absorbing species was highlighted.

Some years later Bensasson and Land produced a review⁴ on the physical properties of excited states which also contained a compilation of extinction coefficients of triplet-triplet absorption maxima for a range of organic molecules, some of biological importance, in solution. This review was limited to those measurements made by

the comparative energy transfer technique introduced by Land⁵ and subsequently developed together with several coworkers. Some previous data obtained under similar experimental conditions, but by different approaches, were also listed by way of comparison. These reviews have served to illustrate the ubiquitousness and importance of the triplet state in a wide variety of photochemical and photophysical phenomena, under conditions ranging from steady illumination to pulse radiolysis.

With the availability of commercial laser flash photolysis equipment and nanosecond pulse radiolysis facilities and the subsequent explosion of time-resolved studies monitored by light absorption, the need for routine identification of transients has become crucial. One of the main transients observed in organic nonaqueous solution in such experiments is the triplet state. Whether this state is itself under investigation, or whether the purpose of study is to eliminate or implicate the triplet state as a precursor for a longer-lived species such as a free radical, a compilation of spectral characteristics of triplet-triplet absorption is of great value.

The quantification of such photophysical processes as intersystem crossing requires the knowledge of the triplet state concentration. One method to obtain this information relies on the existence of a dependable estimate of the extinction coefficient for triplet-triplet absorption⁶. This then can lead to the determination of the quantum yield for triplet formation and, subsequently, if the appropriate

lifetime data, ideally measured under closely similar experimental conditions, is available, to the calculation of the rate of intersystem crossing. However, while the qualitative shapes of the transient absorption spectra are relatively routinely measured, though often unsatisfactorily reported⁷, the determination of reliable extinction coefficients is both more difficult and tedious. Clearly an evaluation of the existing extinction data is in order.


For these reasons we have undertaken the compilation and critical evaluation of information relative to the triplet-triplet absorption spectra of organic molecules in condensed phases. In the course of laying the foundations of this work we have published three bibliographies⁸⁻¹⁰ citing the relevant works in the literature up through the end of 1984. These lists were produced with the assistance of our local bibliographic data base¹¹, and it is our intention to provide regular updates to this series whenever the number of appropriate references accumulated is sufficient. From these bibliographic lists we have culled the data presented in the main body of this work, the table of the wavelengths of maximum absorbance (λ_{\max}) and extinction coefficients (ϵ_{\max}), where determined. Attention has been paid to the critical evaluation of these data as discussed below. In the comments associated with each entry in the table are found numerous other relevant data such as triplet state lifetimes, energies and energy transfer rate constants. No effort has been expended on this data to provide a comparable evaluation of these numbers which must thus be regarded as simply the authors' reported values. The subject of this work is then triplet-triplet absorption spectra of organic molecules in condensed media, i.e. solution, glass, liquid and solid. Gas phase studies are not included. For completeness, and to enhance the usefulness of the present document we have also scanned the literature for the period rather thoroughly covered in the work of Labhart and Heinzelmann. However we have addressed neither the compilation nor the evaluation of theoretical calculations on the triplet-triplet absorption process and provide no update to that section of their review.

We begin with a conceptual introduction to the photophysical processes involved in triplet-triplet absorption, followed by an historical perspective on the discovery and characterization of the triplet state. In Sec. 2. we provide a discussion of the methods commonly used for populating the excited triplet state focusing on the differences between photolysis and radiolysis and the problems and advantages inherent in each technique. The various methods which have been used for the estimation of the extinction coefficient of triplet-triplet absorption are critically analyzed in Sec. 3. Seven principal approaches are outlined, numerous required corrections and precautions are pointed out, possible extensions are illustrated and each technique is evaluated in the light of its advantages or disadvantages. Several less used methods are catalogued in Sec. 3.8. A discussion of the data evaluation procedure follows in Sec. 4. The arrangement of the principal data table, Table 6, is presented in Sec. 5. and lists of abbreviations and symbols used are given.

Since this compilation derives from part of a series of computer accessible numerical data bases under construction at the Radiation Chemistry Data Center, we are able to include, in an automatic fashion, a number of indexes which we hope will enhance the usefulness of this work. We have provided a compound name index, a molecular formula index and an author index.





1.1. Conceptual Background

Triplet states can be generated in several ways. We will discuss the direct intramolecular photoproduction of triplet states in the process of introducing the general photophysical scheme, saving until later photosensitization and pulse radiolysis techniques, where the triplet energy is transferred intermolecularly from the triplet manifold of another solute or the solvent, respectively. When an organic molecule, M, is excited, it is most likely to go to an excited singlet state,



where k_{ex} is the intensity dependent excitation rate. The excited singlet state, ${}^1M^*$, will then decay via several independent processes, again both intermolecular and intramolecular. Intramolecularly it can undergo photochemistry, with rate constant k_{pc} , intersystem crossing, k_{isc} , internal conversion, k_{ic} , or fluorescence, k_f .

These decay channels are represented by the following four processes:



In the usual kinetic scheme these processes are independent, and the singlet state decays exponentially with a rate constant,

$$k_s = k_{pc} + k_{isc} + k_{ic} + k_f, \quad (1.1)$$

which is equal to the inverse of the measured fluorescence lifetime. Triplet states are thus generated directly by the intersystem crossing process, 1.3. The quantum yield, Φ_T for production of the triplet state in this scheme is simply given by

$$\Phi_T = k_{isc}/k_s. \quad (1.2)$$

Schematically, these elementary processes may be summarized in the energy level diagram displayed in Fig. 1.

Depending on the frequency of the incident radiation, the initial state achieved in the absorption event, 1.1, may lie in the vibrational manifold associated with the first electronically excited state (S_1) or that of a higher singlet (S_n). Rapid radiationless deactivation (vibrational relaxation and internal conversion) soon (rate constant $10^{11} - 10^{13} \text{ s}^{-1}$) leads to population of S_1 (Kasha's Rule¹²),





FIG. 1. Rate processes connecting electronic energy levels.

though in exceptional cases radiative emission from S_2 has been observed¹³. These processes are of course in competition with intersystem crossing and photochemistry from the higher excited singlet states, but because of the high rates of internal conversion, this competition is generally ineffective.

Figure 1 also provides an illustration of the decay channels available to the triplet state (again within the above simplified model). Triplet states once formed can undergo similar processes to those of the singlet state. The intramolecular processes are again photochemistry, with rate constant k'_{pc} , (reverse) intersystem crossing, k'_{isc} , and phosphorescence, k_p .



The intramolecular decay is usually observed to be exponential with a decay rate constant of

$$k_T = k'_{pc} + k'_{isc} + k_p. \quad (1.3)$$

Phosphorescence is almost always absent in fluid media because of collision-induced intersystem crossing.


Energy transfer from the singlet manifold, process 1.3, is, of course, spin-forbidden, but the presence of several small interactions, both intramolecular and intermolecular, serve to make the process weakly allowed. Such radiationless transitions may lead to an excited vibrational level of the lowest triplet state (T_1) or that of a higher electronic origin (T_n). Rapid radiationless deactivation (vibrational relaxation and internal conversion) within the triplet manifold soon (rate constant $\sim 10^{11} \text{ s}^{-1}$) leads to population of the lowest electronic state, T_1 , though in exceptional cases triplet-triplet fluorescence has been observed¹⁴. Again these processes compete with

(reverse) intersystem crossing and photochemistry and again this competition is generally ineffective. In addition, once thermal equilibration has occurred, a further deactivation channel is open, provided the necessary thermal energy is available.



This process leads to a type of delayed fluorescence.

In the presence of radiation of a suitable frequency, the relaxed triplet excited state, ${}^3M^*$, can again absorb photons populating higher triplet states with characteristic probabilities,



This process, then, is triplet-triplet absorption, $T_n \leftarrow T_1$ in terms of the symbols used above, the subject of the present compilation.

1.2. Historical Perspective

The discovery of the nature of the "phosphorescent state" of photoexcited organic molecules as the triplet and its detection in absorption have led to the development of triplet-triplet absorption spectroscopy as an important tool in quantitative photochemistry and in the study of photophysical and, more recently, radiation chemical processes in general. But it was really with the study of phosphorescence that the history of triplet states of organic molecules began. The origin of phosphorescence was a research puzzle until Lewis and Kasha ascribed the phosphorescing state to an excited state with a total electronic spin of 1, namely a triplet excited state¹⁵. Throughout the early history of the phenomena implicating the triplet state (up to and including the Lewis and Kasha work and indeed for many years thereafter) phosphorescence spectra and decay characteristics remained the primary method of studying these transient states. However even before the nature of the phosphorescing state was assigned to the triplet state, Lewis and co-workers had recorded that state in absorption¹⁶. Such early work employed high intensity continuous light sources and was limited by the lifetime of the metastable state to viscous media.

By maintaining a boric acid glass at low temperature and monitoring the bleaching of the ground state absorbance, Lewis et al. were able to estimate the extinction coefficient of the phosphorescing state of fluorescein relative to that of the ground state. This was the first use, in the study of the triplet state, of the method we designate singlet depletion, employed here in specially favourable circumstances since distinct spectral regions of transient and ground state absorption could be found. Some years after the clarification of the states participant in triplet-triplet absorption, again using a photostationary technique, McClure¹⁷ recorded transient spectra of a number of aromatic molecules in EPA glass at liquid nitrogen temperature. To deduce the extinction coefficients of the observed transitions he introduced the new method of intensity variation.

A big breakthrough in the absorption studies of triplet states came when Porter and Windsor¹⁸ applied the then recently developed technique of flash photolysis^{19,20}. The introduction of this excitation source allowed the facile detection of triplet-triplet absorption in fluid media, thus extending considerably not only the range of materials and solvents open to investigation but also the time scale available for observation. Studies of triplet states in fluid solution on the microsecond time range, which is generally inaccessible to phosphorescence monitoring, now became possible. Porter and Windsor²¹ were able to obtain an absorption spectrum for anthracene in liquid hexane and used the singlet depletion criterion to estimate extinction coefficients far in excess of McClure's values. (Improvements in the intensity variation technique later confirmed that the original measurements by this method were indeed in error.)

The introduction of the technique of electron pulse radiolysis²²⁻²⁵ and its use to produce triplet states of organic molecules by Dainton et al.²⁶ provided access to a wider range of compounds by removing the limitation formerly imposed by low quantum yields of intersystem crossing. It was, however, four years before Land⁵ was able to report extinction coefficients using this excitation source, through the development of an estimation method involving triplet energy transfer. (See Sec. 3.1. for details).

Finally the late 1960's brought the introduction of pulsed laser techniques²⁷ to the study of triplet states. Further developments in this equipment, namely the application of Q-switching, together with advances in the associated hardware for photoelectric detection eventually brought flash photolysis to the nanosecond time scale²⁸. Shorter lived transients such as the carbonyl triplets²⁹ in fluid solution could now be studied. Picosecond pulse photolysis techniques³⁰, made possible through the phenomenon of mode-locking, have also been used to study triplets^{31,32} on the picosecond time-scale. The much wider availability of commercial nanosecond laser flash photolysis equipment, together with the inherent lower cost of associated detection apparatus, make the nanosecond work much more extensive and the derived data more reliable.

Based on the number of entries corresponding to each of the four principal excitation techniques, Table 1 summarizes their introduction and development over five year spans. However we expect that the numbers in the 80-84 row will be underestimated due to unavoidable timelags in our collection procedure. (Abbreviations are explained in appended list).

Roughly parallel to the development of new and faster techniques for the population of the triplet state has been the growth of the associated methods for estimating the extinction coefficient of the transient population. Singlet depletion and intensity variation initially deployed in photostationary experiments, as mentioned above, have been extended as needed to accommodate the use of pulsed sources. The kinetic method has been used with modulated excitation sources, while relative actinometry

and energy transfer techniques were also popularized with the introduction of flash photolysis and pulse radiolysis. Several other comparative techniques based on electron and hydrogen atom transfer have also been developed and exploited mainly with the help of laser facilities. Sec. 3. presents a critical discussion of these methods, and here we trace the growth in the application of these methods to the study of the triplet state in Table 2 below. This table lists the number of entries in the data base per 5-year period for several popular methods. (Again an explanation of the symbols used can be found in the lists appended to Sec. 5.)

TABLE 1. Excitation methods per 5-year period

Period	LP	FP	PR	PS	Total ^a
50-54	0	19	0	32	51
55-59	0	75	0	9	84
60-64	0	88	4	11	104
65-69	23	166	63	193	452
70-74	121	222	80	45	569
75-79	443	218	75	26	795
80-84	768	174	67	32	1046
Total	1355	962	289	350	3103

^aHorizontal totals reflect the presence of entries assigned to methods other than those specifically indicated in the vertical columns.

TABLE 2. Extinction methods per 5-year period

Period	ET	SD	TD	RA	KM	IV	Total ^a
50-54	0	3	0	0	0	3	6
55-59	0	17	0	0	0	0	17
60-64	0	11	0	0	0	0	11
65-69	28	54	11	3	12	22	173
70-74	73	45	19	8	31	0	185
75-79	93	49	37	28	7	1	255
80-84	96	75	68	28	13	1	320
Total	290	225	135	67	63	27	972

^aHorizontal totals again reflect the presence of entries assigned to methods other than those specifically indicated in the vertical columns.

2. Methods for Populating and Detecting the Triplet State

There are two basic methods for achieving substantial triplet state populations: photolysis with light in the visible or, more commonly, ultraviolet spectral regions and radiolysis with ionizing radiation, usually high energy electrons. In fluid solution, where the triplet state lifetime is on the order of microseconds, both techniques are generally applied in a pulsed form. In low temperature glasses, where triplets can endure for up to about 30 seconds, steady irradiation can be usefully employed.

Experimental setups for the detection of triplet-triplet absorption spectra differ greatly, but there are at least two elements common to almost all of these experiments. First of all every experiment has an "exciting" source to produce triplet state species. Second, each experiment has a

"monitoring" light source to probe their absorbance. In laser photolysis setups the laser often provides both monitoring and exciting light when used with suitable frequency multiplication and light path adjustment. Additionally, if a complete spectrum is desired, the laser pulse can be used to generate a continuum as a monitoring source.

Details of the detection equipment will not be given. We mention only the following historical development. In the earliest flash photolysis systems the transient spectrum was photographed, and the changes in optical density laboriously converted from densitometer measurements of the exposure intensity. With the growth of efforts to discern the mechanism of the various rate processes involved, kinetic spectroscopy using photomultipliers became popular. Here the decay of the transient is monitored at a particular wavelength. When this pattern has been recorded at several observational wavelengths, the entire spectrum may be judiciously reconstructed. More recently the introduction of vidicon, and later reticon, detectors has offered a way to fulfil both of the above functions simultaneously.


We discuss the two primary excitation sources, starting with pulse radiolysis, focusing on the special features of each method, and mention some problems encountered in the respective detection process.

2.1. Generation of Triplets by Pulse Radiolysis

In the technique of pulse radiolysis the sample is irradiated by short bursts of ionizing radiation. Most commonly, pulses of high energy (1–20 MeV) electrons extracted from a linear accelerator or Van de Graaff generator are employed. Initially, the usable pulse width was of the order of a few microseconds, but steady improvement in equipment design soon made the nanosecond regime attainable.

2.1.1. Theoretical Concepts



The deposition of energy from these injected electrons does not discriminate among the chemical components of the systems irradiated. Thus in a dilute solution, which is almost always employed in the study of triplet-triplet absorption, the radiation is absorbed almost exclusively by the solvent, S, which is consequently ionized,




to give solvent radical cations and electrons. The subsequent mechanism of solute excitation depends to a large extent on the properties of the solvent.

In polar solvents, such as water and the lower alcohols, the ionized electron is rapidly thermalized and readily solvated. Thus on the nanosecond time scale the reactive species are the various oxidizing and reducing radicals and radical ions formed from the solvent. These intermediates are also stabilized by solvation. Correspondingly, solute radical ions (and other redox or radical addition products) are predominantly produced by pulse radiolysis in polar solvents.


If the solvent lacks the driving force for electron and (radical) ion solvation, by virtue of its nonpolar character, then radical recombination will promptly occur. In room temperature hexane, for example, the electrons are largely (90%) captured by their parent cations, the particles being unable to escape from their mutual Coulombic attraction. While this geminate recombination is complete within a few nanoseconds, those electrons which do escape recombine homogeneously over a much longer microsecond timescale. With the addition of an electron (or hole) acceptor, A, these processes can be intercepted to produce solute radical ions.



These species then recombine to give both singlet and triplet solute excited states.




In general the solvent radical cation can also participate in this neutralization process,




Further triplets may then be formed by intersystem crossing from the singlet manifold.


A further complication ensues in the above scheme if the solvent itself possesses long-lived excited states, for example benzene (but note that the triplet state lifetime in neat benzene is strongly dependent on the purity of the liquid³³). Electron attachment



is followed by solvent radical ion recombination to give solvent excited states.



Energy transfer may then occur from the excited states of the solvent molecules to those of the solute in a spin-selective fashion.



Again intersystem crossing may augment the solute triplet yield.

The above simplified picture has been debated in the literature for several years. Clearly each individual solvent must be carefully examined to determine the extent to which a given process contributes to the overall triplet yield. Within these limitations the scheme presented serves as an extremely useful guideline.

2.1.2. Special Considerations for Pulse Radiolysis

The absorption intensity of a transient created by such pulses of electrons depends both on the dose per pulse and the G value (100 eV yield) for that transient in addition to its intrinsic extinction coefficient at the monitoring wavelength. For a typical triplet with G = 2 say, to obtain an optical density of 0.1 at a wavelength where the ex-

tion coefficient is on the order of $10^4 \text{ L mol}^{-1} \text{ cm}^{-1}$ over a path length of 1 cm, requires a dose per pulse in excess of 2.5 Gy (1 Gy \equiv 1 gray = 1 J kg^{-1}).

Van de Graaff generators with pulse widths on the order of a few nanoseconds can produce bursts of 3 MeV electrons to give a maximum dose of about 20 Gy. Care must be taken in quantitative work to ensure a sufficiently low dark charge to beam charge ratio when operating under conditions of high dose per pulse or high pulse repetition rate.

Single-pulse generators such as the Febetron eliminate the dark current problem completely and can deliver doses of up to 5000 Gy in a 25 ns pulse of 2 MeV electrons. Problems arise from the lack of penetration depth of such electrons. If they are stopped within the sample severe space charge distortions of the dose distribution result³⁴. In addition spurious optical effects on the microsecond time scale have been observed³⁵ in typical experimental geometries, presumably due to pressure waves generated by the radiation source.

The incremental acceleration of electron by repeated traversal of microwave fields in a linear accelerator has proved to be the most versatile source of ionizing radiation for transient spectroscopy. Coupled with powerful computerized signal averaging techniques to reduce noise, the characteristics of reproducible $\sim 5 \text{ ns}$ pulses of $\sim 20 \text{ MeV}$ electrons delivering a dose per pulse of $\sim 100 \text{ Gy}$ has made this the method of choice for irradiation of fluid samples. Certain problems specific to optical detection of intermediates created by nanosecond pulse radiolysis have been summarized by Hunt et al.³⁶. More recently, modern technology has enabled the resolution of the fine-structure components of linear accelerator pulses, pushing the domain of excitation by this method into the picosecond time range. This is much in analogy to the development of mode-locked lasers, though quantitative work on this timescale still remains problematic due to instabilities in the integrated yield arising from pulse to pulse jitter in both the electron beam and the analyzing light.

Electron linear accelerators produce their own source of analyzing light. Broad band (ultraviolet to infrared) Cerenkov radiation, emitted by electrons traveling above the speed of light pertinent to the solvent under study can be used for the detection of transients. A stroboscopic pulse radiolysis system capable of picosecond time resolution which implements this light as a probe has been reported³⁷.

The principal advantage of the technique of pulse radiolysis then lies in the possibility of populating the excited triplet state of molecules which have intrinsically low efficiency for intersystem crossing. Note however that to study species with short excited state lifetimes, substantial concentrations ($\sim 0.1 \text{ mol L}^{-1}$) of solute are required. For the simple reason of solubility this problem can set a limitation on the usefulness of the technique.

Extensive reviews of the development of this field and the instrumentation required have been provided by Dorfman³⁸ and Hunt³⁹.

2.2. Generation of Triplets by Photolysis

The sources of light employed to excite molecules to the triplet state are quite varied. Intense steady-state sources have been a popular excitation source for samples in glasses; long triplet state lifetimes allow large steady-state concentrations of triplets to build up. Conventional flash lamps have provided most of the triplet-triplet absorption spectra recorded up to the middle 1970's. Their short, but intense, pulses make them ideal for studying spectra and kinetics on the μs time scale, which is in the range of the lifetimes of many triplets in fluid solution. In order to study shorter lived triplets such as those of aromatics ketones, it is most convenient to use Q-switched lasers²⁹ to get into the nanosecond time range. Recently mode-locked lasers have been used to study triplets (including rise times) in the picosecond time range⁴⁰. Non-pulsed lasers (continuous wave lasers) have also been used in much the same fashion as conventional steady-state light sources.

2.2.1. Theoretical Concepts

Triplet states are generated in two ways by photons. In the first method, exemplified in Sec. 1.1. of the above introduction, the compound under study, irradiated within its ground state absorption band, absorbs the photon as in process 1.1, going first to a singlet excited state. The excited molecule then undergoes intersystem crossing, process 1.3, to reach the triplet state.

Should the efficiency of this process be low, another technique is available⁴¹. In this method another molecule, M_1 with a suitable intense ground state absorption and high quantum yield for subsequent triplet state formation is added to the solution (usually in excess). This species intercepts the light energy and transfers it via the intermediary of a triplet state to the triplet state of the compound under study, M_2 during collisions.



At least in dilute solution, excited singlet states are usually too short lived to participate in an analogous singlet-singlet energy transfer, though several examples have been documented⁴². This photosensitization technique will generally be successful provided the triplet energy of the sensitizer (donor) is at least 8–12 kJ mol^{-1} greater than that of the acceptor^{43,44}. Given a suitable energy difference as a driving force, it is usually assumed that triplet-triplet energy transfer proceeds with unit efficiency. However this assumption is often erroneous and detailed kinetic analysis is then required to produce valid estimates of the triplet state concentration of acceptor molecules formed and its temporal evolution. Such corrections are discussed in Sec. 3.1.2. If the acceptor itself absorbs the incident radiation further adjustments must be made to the estimated extinction coefficients.

2.2.2. Special Considerations for Photolysis

Steady-state light sources have often been used with a modulator. In some cases the modulation of the light

source has been used to enhance the signal/noise ratio of the signal⁴⁵, enabling the study of very low triplet concentrations. This is particularly effective when combined with a phase-sensitive detection system⁴⁶. In other cases a modulated light source has been used in connection with the kinetic method to measure extinction coefficients^{47,48}. Modulated moderate intensity light sources are also useful if the sample under study is susceptible to photo-decomposition by more intense pulse techniques⁴⁹.

The advent of laser flash photolysis with relatively monochromatic pulses greatly enhanced the accuracy to which the total energy absorbed by a species in solution could be measured. Previously, conventional flash techniques with intense but polychromatic sources, had made this estimation difficult. This feature enabled, in principle, substantial improvements in the results derived from both the pulsed intensity variation and relative actinometry methods discussed below (Secs. 3.8.3. and 3.4. respectively).

The monitoring light sources that measure the spectrum of the transients have not been nearly as varied in type as have been the exciting sources. They are usually a fairly weak source, for instance of the arc-lamp type. Sometimes the sources have been pulsed to increase their intensity or modulated as part of phase-sensitive detection systems⁴⁶.

A more salient feature of analyzing beams is their spatial arrangement with reference to the path of the exciting light. Both in-line (front-face) and crossed beam (right-angle) geometries are popular. This arrangement of the exciting and monitoring beams can be of crucial importance when using photons to generate the triplet states. The concentration of triplets is central to the determination of extinction coefficients, but this concentration varies along the exciting beam according to Beer's Law. When right-angle monitoring is used, the particular region sampled, along the exciting beam, is one in which the concentration is uniform along the direction of the monitoring beam. Thus kinetics (especially second-order kinetics) can be measured accurately, but spectra are more difficult to obtain since the region sampled will not in general be near the front face of the cell where the concentration of transients is greatest.

In-line monitoring of the triplets has the disadvantage that the observation is along a gradient in the concentration. Unless the concentration of the triplets is kept low, for example with the optical density of triplet absorption < 0.05 ,⁵⁰ the kinetics can be severely distorted when looking at second-order reactions⁵¹. In-line monitoring, however, has an advantage in that it is possible to easily access the region of high concentration of triplets near the surface of the cell on which the exciting light is incident. On the other hand the usual way of calculating concentration from the optical density, in regions where the ground state does not absorb,

$$\Delta OD = \epsilon_i [^3M^*] \ell, \quad (2.1)$$

will not be meaningful because of the variation in $[^3M^*]$ with the distance across the cell along the exciting light

beam. An early work by Ross⁵² provided an estimate of the effect of such nonuniform triplet concentrations on the absorption spectrum recorded during steady-state experiments in glasses. In that study it was concluded that while the general features of the spectrum should be preserved, band to band distortion was not ruled out. Further corrections are discussed in Sec. 3.

There are indeed several other possible sources of spectral distortion. If there is substantial singlet depletion, i.e. removal of ground state molecules in the path of excitation, the resulting difference spectra can offer only a limited guide to triplet-triplet absorption maxima in the vicinity of regions of negative ΔOD . On the other hand, this effect can be very useful because it can often give a quantitative indication of the amount of ground state population that has been converted into triplets. This is the basis for the method of measuring extinction coefficients of triplet-triplet transitions discussed in Sec. 3.2. However due to possible distortions, while we retain difference spectra that are clearly removed from the singlet depletion region in this compilation, we do not report maxima of spectra near the singlet depletion region of uncorrected spectra.

Light scattering of the monitoring beam can cause distortions in the form of the spectral shape and can also shift the position of the spectra in some cases. The intensity of scattered light varies as $1/\lambda^4$ and so becomes a problem in the ultraviolet. In addition conventional monitoring sources usually have very low output in the ultraviolet, and thus they are susceptible to light scattering from the more intense visible regions. The effect on the spectra can be seen by the following example. The experimental optical density is given by

$$\Delta OD_{\text{exp}} = \log_{10} \{ (I_0 + I_s) / (I + I_s) \}, \quad (2.2)$$

where I_0 is the incident light which is measured before the pulse, I_s is the intensity of the visible scattered light that is measured along with I_0 , I is the true intensity of the transmitted light through the solution containing the transient, and I_s is the excess scattered light from the visible that is measured along with I . ΔOD_{exp} is the experimentally measured optical density which is just the equal to the logarithm (to the base ten) of the total intensity measured before the pulse divided by the intensity at some specific time during the decay of the transient. The true optical density is given by

$$\Delta OD_{\text{true}} = \log_{10} (I_0/I). \quad (2.3)$$

ΔOD_{true} can be written in terms of ΔOD_{exp} as

$$\Delta OD_{\text{true}} = -\log_{10} \{ (1 + I_s/I_0) \exp(-2.3\Delta OD_{\text{exp}}) - (I_s/I_0) \}. \quad (2.4)$$

If we consider an example in which scattered light constitutes 50% of the incident and also the transmitted beams, then $(I_s/I_0) = 1 = (I_s/I)$. If we also assume $\Delta OD_{\text{exp}} = 0.01$, then $\Delta OD_{\text{true}} \sim 0.02$. For a second example take the same numbers except for $(I_s/I_0) = 0.996$. This would mean that not as much light is being scattered from the visible while the transient is present. $\Delta OD_{\text{true}} \sim 0.018$

in this case. These examples illustrate the general behavior of excess scattered light on the true spectrum. This effect is often seen on the "blue" side of absorption peaks in the ultraviolet, namely the "blue" side of the uncorrected spectra is suppressed.

Distortions of the spectra can also arise from improper overlap of the analyzing and exciting beams. Proper overlap means that the analyzing beam should go only through the region of the cell that is excited and that the cell should be excited across the total width of the cell in right angle apparatus. In a right-angle setup, if the cell is not excited over the whole width, then the optical path is not the same as the path through the excited region of the cell⁵³. This does not lead to a distortion, but it leads to an incorrect concentration of the transient. The distortion of the spectra due to incorrect overlap comes when the analyzing beam samples regions of the cell that contain no excited molecules. In effect this extra light is scattered light, and the equations that Bazin and Ebbesen⁵³ derive for the relationship between the true optical density and the experimental optical density are very similar to Eq. (2.4) above for scattered light. Such distortions can lead to large errors in measuring extinction coefficients.

Novel sources of nonuniform concentration of transients are hot spots generated in a sample photolyzed by a pulse of laser light (T. W. Ebbesen and M. Bazin, unpublished data). The equations derived by Ebbesen and Bazin in this extension are similar to those reported in their earlier work⁵³ and indicate the possibility of large effects because lasers can cause very severe non-uniformities in transient concentration.

Extensive reviews of the development of pulsed photolytic techniques and the instrumentation involved have been provided by Porter and West⁵⁴ and West⁵⁵.

2.3. Miscellaneous Methods

While there exist other methods for the formation of the triplet states of organic molecules, for example enzymolysis⁵⁶ and thermolysis⁵⁷, we know of no reports concerning the estimation of the extinction coefficients of triplet-triplet absorption in these ways and of none in which new spectra, otherwise unobserved, have been recorded.

3. Methods for Measuring Extinction Coefficients

We discuss seven principal techniques currently in use for the determination of the extinction coefficient for triplet-triplet absorption. Special problems, extensions of the basic method, and advantages and disadvantages of each approach are documented. Several features (e.g. required corrections) are common to more than one technique. To avoid repetition these items are detailed only on their first occurrence. The various methods are presented approximately in order of number of entries in our database, energy transfer being the most frequently adopted approach. Miscellaneous methods are collected in Sec. 3.8.

3.1. Energy Transfer (ET) Method

3.1.1. Description of Method

The energy transfer method is currently one of the most widespread methods of measuring extinction coefficients. It was originally used for measuring extinction coefficients of triplet states generated by pulse radiolysis⁵ and later extended to include population by pulsed laser photolysis⁵⁸. Energy transfer methods have been criticized because of uncertainties in the probability of transfer. However over the past few years this method for measuring triplet state extinction coefficients has been improved and can now yield reliable results when used carefully.

The basic idea of the method is quite simple⁴. Two compounds are placed in a cell. One compound, R, has a triplet state whose extinction coefficient is well-known. The other compound, T, is the compound whose extinction coefficient is to be determined. In the ideal experiment, one of the triplet states (either R's or T's) is initially populated, and the other triplet state is not populated. The compound with the initially populated triplet is chosen so that it can act as the triplet donor of the pair of triplet states. It does not matter whether R or T acts as the triplet donor. After the donor triplet is populated it can then transfer its energy to the acceptor triplet. This is done by collisional quenching involving the exchange interaction⁵⁹. In the ideal experiment where the triplet lifetimes of the isolated molecules are infinite on the time scale of the quenching experiment, every donor triplet molecule produces exactly one acceptor triplet molecule. The "one donor triplet yielding one acceptor triplet molecule" is the heart of the method because it allows one to get a handle on the persistent problem of measuring extinction coefficients of transients, namely determining the concentration of the transient. So in the ideal experiment the initial concentration of the donor triplet is equal to the final concentration of the acceptor triplet and so




$$\epsilon_T^* = \epsilon_R^* (\Delta OD_T / \Delta OD_R). \quad (3.1)$$

In this equation ϵ_T^* is the unknown extinction coefficient, and ϵ_R^* is the reference extinction coefficient and the superscript "*" refers to the excited state. The meaning of the ΔOD 's depends on whether T or R is functioning as the donor. The ΔOD of the compound serving as the donor is set equal to the ΔOD extrapolated back to time zero, and the ΔOD of the compound serving as the acceptor is set equal to the ΔOD at infinite time.

3.1.2. Kinetic Corrections

Corrections need to be made for several things when the conditions are not as described above. One problem occurs because the probability for transfer from donor, D, to acceptor, A, is not always unity. In addition to populating the acceptor triplet, the donor triplet can decay by other paths. In well degassed solutions, the most likely path is by unimolecular, radiationless decay to the ground state. The kinetic scheme is often modelled by the following reactions:

Scheme I



Ignoring the last process, the probability of transfer, P_{tr} , is

$$P_{\text{tr}} = k_{\text{et}}[{}^1\text{A}] / (k_{\text{et}}[{}^1\text{A}] + k_{\text{D}}). \quad (3.2)$$

The concentration of the acceptor is reduced by this factor. So an equation analogous to Eq. (3.1) can be written with

$$\epsilon_{\text{A}}^* = \epsilon_{\text{D}}^* (\Delta\text{OD}_{\text{A}} / \Delta\text{OD}_{\text{D}}) / P_{\text{tr}} \quad (3.3)$$

In order to make this correction one has to know k_{D} and $k_{\text{et}}[{}^1\text{A}]$. The unimolecular decay rate constant of the donor, k_{D} , is just the rate constant of the decay of the donor triplet measured in the absence of the acceptor, but with all other conditions similar to those in the actual energy transfer experiment. If the acceptor is then added to the cell, the combination $k_{\text{D}} + k_{\text{et}}[{}^1\text{A}]$ can then be measured as an effective first-order decay rate constant of the donor triplet. This effective first-order rate constant is the sum of the donor's unimolecular decay rate constant and the pseudo first-order, energy transfer rate constant, $k_{\text{et}}[{}^1\text{A}]$. With the measurement of $k_{\text{D}} + k_{\text{et}}[{}^1\text{A}]$ and k_{D} , P_{tr} can be calculated using Eq. (3.2).

Two precautions are often taken when applying this kinetic correction factor. First, care is taken to eliminate second-order decay of the donor triplets. These processes could complicate the kinetic scheme. Second, $k_{\text{D}} + k_{\text{et}}[{}^1\text{A}]$ is often measured by checking both the decay rate constant of the donor absorption and also the inverse rise time of the acceptor triplet. If the kinetic scheme is as simple as the model assumed above, these two rate constants should be the same. If they are not, complicating factors have probably entered the analysis.

The rise time of the acceptor is often obscured by the decay of the acceptor. For instance, the rise and decay kinetics of the acceptor in Scheme I is given by

$$[{}^3\text{A}^*] = W \times \{\exp(-k_{\text{A}}t) - \exp(-k_{\text{et}}[{}^1\text{A}]t - k_{\text{D}}t)\}, \quad (3.4)$$

where

$$W = [{}^3\text{D}^*]_0 k_{\text{et}}[{}^1\text{A}] / (k_{\text{D}} + k_{\text{et}}[{}^1\text{A}] - k_{\text{A}}), \quad (3.5)$$

and where $[{}^3\text{D}^*]_0$ is the initial triplet concentration of the donor. If k_{A} is not small compared to the argument of the second exponential in Eq. (3.4), then the growth of A will be complicated by the first term in Eq. (3.4).

When k_{A} is not small compared to $k_{\text{et}}[{}^1\text{A}] + k_{\text{D}}$, a modification to Eq. (3.3) is often used⁶⁰. The maximum of the decay curve represented by Eq. (3.4) is the one unique feature of the curve. The time at which this point is reached is found by differentiating Eq. (3.4) and setting the derivative to zero. This time is given by

$$t_{\text{max}} = \frac{\ln\{k_{\text{A}}/(k_{\text{et}}[{}^1\text{A}] + k_{\text{D}})\}}{k_{\text{A}} - k_{\text{et}}[{}^1\text{A}] - k_{\text{D}}}. \quad (3.6)$$

This time, t_{max} , and the measured $\Delta\text{OD}_{\text{A}}(t_{\text{max}})$ can be used to find the $\Delta\text{OD}_{\text{A}}$ needed for Eq. (3.3) in the following manner. First one can calculate $[{}^3\text{A}^*]$ at t_{max} for the Eqs. (3.4,3.5). Second one can calculate $[{}^3\text{A}^*]$ as $t \rightarrow \infty$ under the Eqs. (3.4,3.5) for $k_{\text{A}} = 0$, that is no decay of ${}^3\text{A}^*$. Taking the ratio of these two results gives a simple correction factor, and the final result is

$$\Delta\text{OD}_{\text{A}} = \Delta\text{OD}_{\text{A}}(t_{\text{max}}) \times \exp(k_{\text{A}}t_{\text{max}}). \quad (3.7)$$

This is the $\Delta\text{OD}_{\text{A}}$ that is to be used in Eq. (3.3) to obtain the sought for ϵ_{A}^* . In order to do the kinetic correction in Eq. (3.7), an extra rate constant, k_{A} , must be obtained in addition to the rate constants needed to calculate P_{tr} . If $[{}^3\text{A}^*]$ cannot be directly populated, then k_{A} must be obtained from a fit to the two-exponential decay given by Eq. (3.4).

3.1.3. Uncertainty in Probability of Transfer

Even after the two kinetic corrections are made, there still remains an uncertainty in the energy transfer method. This uncertainty involves the probability of transfer. With the two kinetic corrections already discussed, account can be taken of the unimolecular decay of both the donor, Eq. (3.3), and the acceptor, Eqs. (3.3,3.7). Nothing can be done in this scheme if the acceptor deactivates the donor without the acceptor itself being excited.



Eq. (3.2) would have to be modified by the pseudo first-order rate constant, $k_{\text{DA}}[{}^1\text{A}]$ to read

$$P_{\text{tr}} = k_{\text{et}}[{}^1\text{A}] / (k_{\text{DA}}[{}^1\text{A}] + k_{\text{et}}[{}^1\text{A}] + k_{\text{D}}). \quad (3.8)$$

The concentration dependence of this process and the qualitative kinetics are the same as that in Scheme I alone, and, on a routine basis, it is not easy to check whether this type of deactivation of ${}^3\text{D}^*$ is occurring.

3.1.4. Special Problems in Radiolytic Excitation

There are special problems with both photoexcitation and radiolytic excitation methods. In pulse radiolysis, benzophenone has been the most widely used triplet donor. The benzophenone triplet can either transfer its energy to the acceptor via reaction 3.2 in Scheme I or it can abstract a hydrogen from the solvent and form a ketyl radical.



By varying the concentration of the acceptor, the amount of benzophenone ketyl radical will also vary. The absorption spectrum of the ketyl radical is well-known⁵, and variations in its spectrum reflect the competition between reactions 3.2 and 3.5.

The first problem with the radiolytic method arises out of one of its strengths. One of the attractive features of the method is that the extinction coefficient of the ketyl radical can be independently and reliably measured by pulse radiolysis. As a general rule it is true that measurements of extinction coefficients in aqueous solution of radicals will

be more reliable when measured by pulse radiolysis than by flash photolysis. This is due mainly to the existence of reliable G values of the radicals in aqueous solution and also due to the fact that in radiolysis, uniform concentrations of the transients can be generated throughout the cell. The problem arises precisely because the well established extinction coefficient of the ketyl radical is known in water; whereas the radiolytic generation of triplets can be done only in hydrocarbon solvents where radiolytic yields are not at all well established. The procedure that is used is that the extinction coefficient of the ketyl radical in the hydrocarbon solvent is calculated from the extinction coefficient in water by assuming that the oscillator strength is independent of solvent⁶¹.

The second special problem with the pulse radiolysis is that there is part of the ketyl radical population that cannot be scavenged. The unscavengeable fraction of ketyl radicals can be quite large. In applications it is assumed that the unscavengeable fraction is constant, independent of the acceptor concentration.⁴

The third special problem with the radiolysis method is that ketyl radicals themselves are often reactive with the acceptors. This is not a problem with hydrocarbons, but with quinones and acridines the method cannot be used⁴.

There are pulse radiolytic methods, of course, which do not involve the competitive step of ketyl radical formation. In these methods, a potential triplet donor, for example biphenyl, is present at roughly 100 times the acceptor concentration. The donor triplets are formed by energy transfer from the solvent triplets. This pulse radiolysis technique is more analogous to the flash photolytic method in that it does not require competition between two reactive paths with the subsequent formation of two transients which may or may not interfere with one another.

3.1.5. Special Problems in Flash Photolysis

The complications unique to the photolytic method are bound up with the production of nonuniform distributions of transients according to Beer's Law. Using the differential form of the law, namely that $\Delta I = -2.3\epsilon_D[I] \Delta x$, one can see that the number of photons absorbed in a volume of unit cross section and of depth Δx (during unit time) is equal in magnitude to $\text{abs}(\Delta I)$. To convert this number, $\text{abs}(\Delta I)$, of excited molecules into a concentration that varies with x across the cell, one must divide the number of excited molecules by the volume (1 unit cross section $\times \Delta x$) containing them. In addition one needs to multiply by the duration of the exciting pulse, Γ_p , and by the triplet quantum yield, Φ_T , which results in

$$[^3D^*](x) = 2303 \epsilon_D [^1D] I_p(x) \Gamma_p \Phi_T(D). \quad (3.9)$$

In Eq. (3.9), $I_p(x)$ is the intensity of the photolyzing light at x , $[^1D]$ is the concentration of the absorbing species whose extinction coefficient at the exciting frequency is ϵ_D . When the x -dependence of $I_p(x)$ is displayed, using the integrated form of Beer's Law, $I = I_0 \exp\{-2.3\epsilon_D [^1D]x\}$, Eq. (3.9) becomes

$$[^3D^*](x) = 2303 \epsilon_D [^1D] I_p^0 \exp(-2.3\epsilon_D [^1D] x) \times \Gamma_p \Phi_T(D), \quad (3.10)$$

where I_p^0 is the incident intensity of the photolyzing light. When $[^1D]$ and ϵ_D are small enough, ie.

$$\epsilon_D [^1D] x \approx 0.05, \quad (3.11)$$

then the exponential in Eq. (3.10) is approximately unity and $[^3D^*](x)$ is roughly constant across the cell⁵⁰. This is the usual technique to obtain uniform concentration distributions in flash photolysis. However, this technique often fails when the absorbance of the resulting transient is low because then a larger concentration of donor is needed than is allowed by condition (3.11).

When no light is absorbed by the acceptor, the non-uniform distribution does not present a problem even at concentrations where Eq. (3.11) is no longer valid. The following discussion is for a linear arrangement of the monitoring and photolyzing light. Placing the excited state concentration given by Eq. (3.10) into the differential form of Beer's Law for the monitoring light intensity, I_m , the effective optical density,

$$OD = \log_{10}(I_m^0/I_m), \quad (3.12)$$

is found by integration to give

$$OD(\lambda_m) = \epsilon_D^*(\lambda_m) \left\{ \frac{[^3D^*]_{x=0} - [^3D^*]_{x=\ell}}{\epsilon_D(\lambda_p)[^1D]\ell} \right\} \ell, \quad (3.13)$$

where quantities without asterisks refer to the ground state, and ℓ is the optical path length of the cell. Even though Eq. (3.13) does not look immediately like the conventional formula for an optical density, namely $OD = \epsilon_D^* [^3D^*] \ell$, it can be understood as such if the quantity in curly brackets is interpreted as an effective concentration. From Eq. (3.13), it can be seen that the effective concentration, in front-face experiments, is equal to the difference between the true triplet concentrations at the front and back faces of the cell divided by the optical density of the solution at the exciting wavelength, λ_p . Note that this interpretation of an effective concentration keeps intact the important relationships between extinction coefficient and optical density, namely that they are proportional to each other and that the λ_m -dependence of the right-hand side of Eq. (3.13) is only in the factor ϵ_D^* . (In most of the equations in this work we will use the traditional form for the optical density, but it should be kept in mind that for front-face alignment the concentration of the transient refers to an effective concentration defined by the expression in curly brackets in Eq. (3.13)).

The complications arise when the acceptor starts to absorb the exciting light directly. Even when the quantum yield of direct triplet formation is zero for the acceptor, problems can enter. Because of the problem of extrapolating the donor absorbance accurately back to zero time when an acceptor is present, the initial optical density can be taken from a measurement before the acceptor is added to the cell. Then an inner filter effect⁵⁰ can occur just as is common in emission spectroscopy⁶². The

basic equations for numbers of the excited state species now become

$$[^3A^*](x) = 2303 P_{tr} \Phi_T(D) \Gamma_p \epsilon_D [^1D] \times I_p^0 \exp\{-2.3(\epsilon_D [^1D] + \epsilon_A [^1A])x\} \quad (3.14)$$

and Eq. (3.10). There is no inner filter effect with only the donor in the cell, Eq. (3.10); so after integrating the differential form of Beer's Law using separately Eq. (3.10) and Eq. (3.14) and dividing the two resulting equations, one obtains Eq. (3.15) which is a modification to Eq. (3.3).

$$\epsilon_A^*(\lambda_m^A) = C_{inner} \epsilon_D^*(\lambda_m^D) (\Delta OD_A / \Delta OD_D) / P_{tr} \quad (3.15)$$

where

$$C_{inner} = N_{inner} / D_{inner} \quad (3.16)$$

$$N_{inner} = \{ \epsilon_D(\lambda_p) [^1D] \ell \} \times (1 - \exp\{-2.3\epsilon_D(\lambda_p) [^1D] \ell - 2.3\epsilon_A(\lambda_p) [^1A] \ell\}), \quad (3.17)$$

$$D_{inner} = \{ \epsilon_D(\lambda_p) [^1D] \ell + \epsilon_A(\lambda_p) [^1A] \ell \} \times (1 - \exp\{-2.3\epsilon_D(\lambda_p) [^1D] \ell\}), \quad (3.18)$$

and λ_p is the exciting wavelength. The terms $\epsilon_D(\lambda_p) [^1D] \ell$ and $\epsilon_D(\lambda_p) [^1D] \ell + \epsilon_A(\lambda_p) [^1A] \ell$ are easily accessible to measurement because they are just the usual optical densities of the unexcited solutions at the exciting wavelength and thus can be measured with an ordinary absorption spectrometer. This correction is not necessary when the absorbance of the donor is measured in the presence of the acceptor because inner filter effects will be the same for both the donor and acceptor triplet, and both will cancel. Even when the absorbance of the donor is measured separately, as long as the concentration of the donor is such that

$$\epsilon_D(\lambda_p) [^1D] \ell \gg \epsilon_A(\lambda_p) [^1A] \ell, \quad (3.19)$$

C_{inner} will approach unity and Eq. (3.3) will hold.

When the quantum yield for direct triplet formation in the acceptor is not zero, the corrections for the acceptor absorbing the exciting light are even more complicated. In order to present the results simply, we assume that both the donor and acceptor absorbances are made with both species present, eliminating the correction of type containing C_{inner} in Eq. (3.15). With this qualification, the basic concentration equations, for the case where the acceptor absorbs and forms triplets directly, are

$$[^3A^*](x) = 2303 P_{tr} \Phi_T(D) \epsilon_D [^1D] \Gamma_p I_p^0 \times \exp\{-2.3(\epsilon_D [^1D] + \epsilon_A [^1A])x\} + 2303 \Phi_T(A) \epsilon_A [^1A] \Gamma_p I_p^0 \times \exp\{-2.3(\epsilon_D [^1D] + \epsilon_A [^1A])x\} \quad (3.20)$$

and

$$[^3D^*](x) = 2303 \Phi_T(D) \epsilon_D [^1D] \Gamma_p \times I_p^0 \exp\{-2.3(\epsilon_D [^1D] + \epsilon_A [^1A])x\}. \quad (3.21)$$

Using these equations with Beer's Law gives the equation

$$\epsilon_A^* = \epsilon_D^* (\Delta OD_A / \Delta OD_D) \times \frac{\Phi_T(D) \epsilon_D [^1D]}{P_{tr} \Phi_T(D) \epsilon_D [^1D] + \Phi_T(A) \epsilon_A [^1A]} \quad (3.22)$$

which is the analog of Eqs. (3.3,3.15).

3.1.6. Summary of Advantages and Disadvantages

In summary, the energy transfer method has an advantage over the singlet depletion method in that it does not depend on overlaps of the ground-singlet and triplet-triplet absorption spectra. In addition using the pulse radiolysis excitation method, independent reference extinction coefficients can be obtained because of accurate dosimetry in radiolysis experiments. However the accuracy of the method is only as good as the reference extinction coefficients. The disadvantages and limitations of the energy transfer method have been discussed above. The most troublesome problem is the uncertainty in the probability of transfer. This is mainly manifested in the process of collisional deactivation without energy transfer, namely reaction 3.4.

3.2. Singlet Depletion (SD) Method

3.2.1. Description of Method

If the measurement of the change in OD due to the population of the triplet state can be extended into a spectral region in which the starting material (singlet) absorbs, then another technique is generally available for the estimation of the extinction coefficient of triplet-triplet absorption. This is the method of singlet depletion.

The precision of the technique can be considerably enhanced if the accessible region is one in which the underlying singlet absorption is sharply structured, as will be seen below.

Assuming that all the molecules which are pumped out of the starting ground state, S_0 either return to S_0 or populate the lowest triplet state, T_1 within the time scale of the observations, then the change in optical density may be simply expressed as

$$\Delta OD = (\epsilon_T^* - \epsilon_S) [^3M^*] \ell, \quad (3.23)$$

where $[^3M^*]$ is the concentration of the triplet state at the time of observation and ℓ is the optical path length of the monitoring beam in the sample (which can usually be determined with reasonable precision).

From the resulting difference spectrum (ΔOD as a function of observational wavelength), the triplet-triplet spectrum may be reconstructed in a number of ways which are documented in Sec. 3.2.2.

It is often assumed that a region of the conventional ground state absorption spectrum ($S_n \leftarrow S_0$) can be located (and accessed for excitation) where the triplets

formed on excitation will not absorb. Thus if the change in optical density, ΔOD_s , due to singlet depletion at this wavelength, λ_1 , is measured, then

$$\Delta OD_s(\lambda_1) = -\epsilon_s [^3M^*] \ell, \quad (3.24)$$

assuming, of course, all find their way to the lowest excited triplet state, T_1 . The triplet extinction follows simply by using the value of $[^3M^*]$ from Eq. (3.24) and by observing the change in absorbance, ΔOD_T , at some further wavelength, λ_2 , where the ground state molecule is known not to absorb.

$$\Delta OD_T(\lambda_2) = \epsilon_T [^3M^*] \ell. \quad (3.25)$$

If it is assumed that the nascent triplet state does not absorb in the spectral region where the singlet depletion is monitored, then an upper limit to ϵ_T will be derived. In many cases this assumption will be unjustified and the more complicated procedures documented below must be adopted to recover correct quantitative estimates of $[^3M^*]$ and thus ϵ_T .

3.2.2. Reconstruction of True Triplet-triplet Absorbance

Labhart⁴⁵ recasts equation (3.23) to give the optical density, OD_T , due to triplet-triplet absorption as

$$OD_T \equiv \epsilon_T [^3M^*] \ell = \Delta OD + \epsilon_s [^3M^*] \ell. \quad (3.26)$$

It is then possible, by trial and error, to find a value of $[^3M^*]$ such that OD_T shows no evidence of the peaks in the original singlet absorption spectrum. If $[^3M^*]$ is chosen to be too large, then the peaks of the predicted triplet spectrum will coincide with those of the underlying singlet. Too small a value for $[^3M^*]$ aligns the peaks of the triplet spectrum with troughs in the singlet absorbance. Upper and lower limits for $[^3M^*]$ are thus readily established.

Dawson⁶³ assumes that the triplet extinction coefficient did not change between the wavelength of the peak, λ_1 of a strong singlet absorption and an adjacent trough, λ_2 . This gives an equation for $[^3M^*]$ in terms of the ΔOD s and the ground state extinction only.

$$\ell [^3M^*] = \frac{\Delta OD(\lambda_1) - \Delta OD(\lambda_2)}{\epsilon_s(\lambda_2) - \epsilon_s(\lambda_1)} \quad (3.27)$$

The resulting value for the concentration of triplets may be checked, in a similar fashion through the region of overlapping singlet absorption. Bowers and Porter⁶⁴ had previously derived an equation similar to Eq. (3.27). In that work the wavelengths, λ_1 and λ_2 , were chosen to bracket a strong singlet absorption and the value for ϵ_T was again assumed constant in this range. Unfortunately their Eq. 1 contains a misprint with extinction coefficients instead of ΔOD 's in the denominator.

By utilizing the change in OD at both troughs neighboring a strong singlet peak, and at the peak itself, Hadley

and Keller⁶⁵ obtain three simultaneous equations for $[^3M^*]$ and the triplet extinction at the three wavelengths in question. These are

$$\Delta OD(\lambda_i) = \{\epsilon_T(\lambda_i) - \epsilon_s(\lambda_i)\} [^3M^*] \ell; i=1,2,3. \quad (3.28)$$

Assuming that the triplet extinction changes linearly over the spanned wavelength range provides a fourth relation,

$$\frac{\epsilon_T(\lambda_1) - \epsilon_T(\lambda_2)}{\epsilon_T(\lambda_3) - \epsilon_T(\lambda_2)} = (\lambda_1 - \lambda_2) / (\lambda_3 - \lambda_2), \quad (3.29)$$

which is sufficient to allow solution for $[^3M^*]$. An alternative choice for the observation wavelengths is made by Pavlopoulos⁶⁶. Assuming that the triplet extinction coefficient varies linearly

$$\epsilon_T(\lambda) = A + B \lambda \quad (3.30)$$

over the region between the onset of a ground state absorption band, λ_a , and the isosbestic point in the difference spectrum, λ_c , and recording the change in optical density at these wavelengths and also at some intermediate wavelength, λ_b , again leads to a set of equations which may be solved for the triplet concentration. Noting that, at the isosbestic point,

$$\epsilon_T(\lambda_c) = \epsilon_s(\lambda_c) \quad (3.31)$$

these equations may be written as

$$\Delta OD(\lambda_a) = (A + B \lambda_a) [^3M^*] \ell, \quad (3.32)$$

$$\Delta OD(\lambda_b) = \{(A + B \lambda_b - \epsilon_s(\lambda_b))\} [^3M^*] \ell, \quad (3.33)$$

and

$$\epsilon_s(\lambda_c) = A + B \lambda_c, \quad (3.34)$$

and A, B and $[^3M^*]$ may be recovered.

The success of these methods obviously depends on the existence of accessible spectral regions where $S_n \leftarrow S_0$ and $T_n \leftarrow T_1$ transitions overlap, which is often found to be the case.

3.2.3. Corrections for Triplet Decay

As long as no (irreversible) photochemistry has occurred upon excitation, no corrections for triplet decay are required. In a pulsed laser experiment, after the excited singlet has decayed, the original population will be distributed between the ground state and the triplet state only. Changes in OD at the depletion and observation wavelength should thus simply be measured after equivalent time delays.

3.2.4. Summary of Advantages and Disadvantages

Obviously, if other channels are open (eg. photochemical) and can be quantified, appropriate corrections can be made — but these complications would greatly

reduce the usefulness of this method and detract from its principal advantage, its simplicity. This simplicity rests in the fact that no detailed knowledge is required concerning the rate constants of the various participating elementary radiationless processes.

Though this method is free from kinetic considerations one practical weakness remains. In the singlet depletion region no precise knowledge of the triplet absorption profile can be gained without recourse to one of the approximation schemes discussed above. Only at an isosbestic point in the difference spectrum can a certain value of the triplet extinction coefficient be given. The values at other wavelengths are critically dependent on the functional form for $\epsilon(\lambda)$ chosen above.

3.3. Total Depletion (TD) Method

3.3.1. Description of Method

It is often assumed that by simply increasing the intensity of a pulse of exciting radiation complete conversion of a suitably small ground state concentration to the triplet manifold is possible, provided, of course, that the quantum yield of intersystem crossing is not negligibly small. Thus $[^3M^*]$ is simply set equal to $[^1M]$.

Kinetic considerations based on a simple two-state model for the population of the triplet state may be used to justify the assumption of complete conversion⁶⁷. In this model the intermediacy of the excited singlet state is neglected and the excitation process is modelled by two kinetic equations,

$$\frac{d[^1M]}{dt} = -2303\epsilon_s I_p(t) \Phi_T [^1M] \quad (3.35)$$

and

$$\frac{d[^3M^*]}{dt} = +2303\epsilon_s I_p(t) \Phi_T [^1M]. \quad (3.36)$$

The excitation rate, k_{ex} has been explicitly defined as

$$k_{ex} = 2303 \epsilon_s I_p(t) \quad (3.37)$$

to clearly exhibit its intensity dependence, ϵ_s is the extinction coefficient of the ground state at the photolyzing wavelength and optically thin samples are assumed. Solving these equations for $[^3M^*]$ with the initial conditions $[^1M] = [^1M]_0$ and $[^3M^*] = 0$ gives

$$[^3M^*] = [^1M]_0 (1 - \exp\{-2303\epsilon_s I_p \Phi_T t\}) \quad (3.38)$$

for the concentration of triplet at time, t , after a pulse of constant intensity, I_p , is turned on. In the limit of infinite excitation intensity the exponential vanishes and Eq. (3.38) collapses to give the initial concentration of the ground state. Notice that the triplet state is not allowed to decay in this model.

If it is assumed that the ground state is totally depleted then a lower limit can be set on the extinction estimation. While this assumption may indeed be valid for systems

studied by conventional flash photolysis^{18,68} it does not hold, in general, when laser pulses of a few (or a few tens of) nanoseconds duration are employed⁶⁹. Under such conditions kinetic limitations become important and the participation of S_1 must be accounted for. Below we formulate a model which adequately encompasses such features.

3.3.2. A Three-state Excitation-deactivation Model

To describe the detailed time development of the population of an excited state, Processes 1.1–1.8 may be modelled by a set of three coupled kinetic equations

$$\frac{d[^1M]}{dt} = -k_{ex}[^1M] + (k_f + k_{ic})[^1M^*] + k_T[^3M^*], \quad (3.39)$$

$$\frac{d[^1M^*]}{dt} = k_{ex}[^1M] - k_s[^1M^*], \quad (3.40)$$

and

$$\frac{d[^3M^*]}{dt} = k_{isc}[^1M^*] - k_T[^3M^*]. \quad (3.41)$$

Note that loss by photochemical processes has been explicitly neglected. For several choices of pulse shape these equations may be solved analytically. For example, for a square pulse of width, Γ_p , the triplet concentration at some time, t , after the pulse, is given by⁶⁹

$$[^3M^*]_t = [^3M^*]_{\Gamma_p} \exp\{-(t - \Gamma_p)k_T\} + \{k_{isc}/(k_s - k_T)\} [^1M^*]_{\Gamma_p} \times (\exp\{-(t - \Gamma_p)k_T\} - \exp\{-(t - \Gamma_p)k_s\}), \quad (3.42)$$

where $[^3M^*]_{\Gamma_p}$ and $[^1M^*]_{\Gamma_p}$ are the end-of-pulse values of the triplet and excited singlet state concentrations respectively. In the limit of an infinitely short (δ -function) pulse, as the excitation intensity grows toward infinity and when the sampling time, t , is sufficiently long that the excited singlet has decayed then

$$[^3M^*]_t / [^1M]_0 = \Phi_T. \quad (3.43)$$

However, for a finite pulse width there will exist a finite end-of-pulse concentration of both T_1 and S_1 regardless of the excitation intensity. Indeed, in the limit of infinite excitation rate (or, equivalently, intensity) we find⁶⁹

$$[^3M^*]_t = [^1M]_0 \frac{k_{isc}}{(k_T + k_{isc})} \times \{ (1 - \exp\{-(k_T + k_{isc})\Gamma_p\}) \exp\{-k_T(t - \Gamma_p)\} + (k_T + k_{isc}) \exp\{-(k_T + k_{isc})\Gamma_p\} \} \times \frac{\exp\{-(t - \Gamma_p)k_T\} - \exp\{-(t - \Gamma_p)k_s\}}{k_s - k_T}. \quad (3.44)$$

Only in the limit of sufficiently fast singlet decay (or sufficiently long pulse duration) can complete conversion be attained. For a small number of systems examined

previously (polycyclic aromatic hydrocarbons) we find⁶⁹ that 95% conversion can be expected only when

$$\tau_s \leq \Phi_T \Gamma_p / 2. \quad (3.45)$$

If this inequality is not satisfied for the system under study then a systematic error will be introduced into the extinction coefficient of triplet-triplet transitions if the concentration of the absorbing triplet is simply set equal to that of the starting ground state. The method will afford, in general, only a lower limit of the true extinction coefficient.

3.3.3. Corrections for Triplet Decay

In a pulsed laser experiment the change in optical density at the wavelength of observation will be a function of time. Usually these changes are extrapolated back to "zero" time (i.e. immediately after the pulse) by assuming a simple single exponential decay of the triplet absorbance. This point should of course be checked and for this purpose the initial ground state concentration should generally be kept as low as possible. Complications in the decay curve due to triplet-triplet annihilation may thus be avoided.

3.3.4. Summary of Advantages and Disadvantages

The principal advantage of the total depletion method is that it offers a simple direct estimate of the triplet concentration and hence of the triplet-triplet extinction coefficient.

Empirically, the approach to total depletion is inferred from a saturation in the ΔOD at some monitoring wavelength as the excitation intensity rises. The assumption then is that the asymptote toward infinite intensity represents complete conversion. Unfortunately, especially in laser photolysis, many other sources of this apparent saturation are possible. Multiphotonic processes⁷⁰ including biphotonic ionization^{71,72} and excited state absorption⁷¹⁻⁷³ have been shown to produce a similar behavior. In order to reduce the chance of interference from such spurious phenomena a partial saturation method has been developed⁷¹ which is discussed below in Sec. 3.7. However even when these and other complications such as photochemical decomposition can be avoided one basic problem still remains. The desired conversion to the triplet state may be kinetically unattainable within a typical pulse duration⁶⁹, although the observed ΔOD may still appear to saturate.

3.4. Relative Actinometry (RA) Method

3.4.1. Description of Method

The basic principle of the technique defined here as relative actinometry is simple⁷⁴. A comparison is made between the change in optical density produced in some solvent, S_T at some wavelength, λ_1 upon excitation of the triplet, T for which the extinction coefficient, ϵ_T^* is required and that change produced under comparable conditions (but in a separate sample) in some reference triplet,

R of known extinction coefficient, ϵ_R^* at some wavelength, λ_2 (not necessarily equal to λ_1) in another solvent, S_R . The concentrations of the ground state precursors of T and R are adjusted, based on a knowledge of their respective absorption spectra so that the same number of excitation photons are absorbed by both sample and reference.

If ΔOD_R is the observed change in optical density due to the formation of the triplet state, R and ΔOD_T that change for T , then, if $\Phi_T(R)$ is the (known) quantum yield for intersystem crossing in the reference compound in the same solvent, S_R in which the extinction coefficient ϵ_R^* was determined (under the same conditions of excitation), then the extinction coefficient of the unknown triplet state is given by

$$\epsilon_T^*(\lambda_1) = \frac{\Delta OD_T \Phi_T(R)}{\Delta OD_R \Phi_T(T)} \epsilon_R^*(\lambda_2), \quad (3.46)$$

where $\Phi_T(T)$ is the quantum yield for intersystem crossing leading to the triplet state of unknown extinction (and again measured in the appropriate solvent, S_T under similar excitation conditions).

3.4.2. Corrections for Singlet Depletion

As written, Eq. (3.46) requires that the ground state depletion (in OD) is negligible at the respective wavelengths of observation. Thus either the starting singlets do not absorb appreciably at λ_1 and λ_2 respectively or the excitation intensity is kept low. This latter factor weighs significantly in the use of a laser source in this technique. Considerable care must be taken to ensure that all the excitation beam is channeled into the appropriate photochemical processes and none is lost to $S_n \leftarrow S_1$ (biphotonic) absorption for instance, if this process does not exclusively lead to population of the triplet manifold.

If such singlet depletion is, in fact, a problem, it will be evidenced by the presence of curvature in a ΔOD vs intensity plot. Restricting the excitation intensity to the initial linear region is then recommended. The extrapolation of ΔOD 's observed in a nonlinear regime is problematical since the curvature may also be due to other effects.

Often the respective singlet ground states also absorb the monitoring radiation. The extinction coefficients appearing in Eq. (3.46) may be altered to take this feature into account to give

$$\{\epsilon_T^*(\lambda_1) - \epsilon_S^T(\lambda_1)\} = \frac{\Delta OD_T \Phi_T(R)}{\Delta OD_R \Phi_T(T)} \{\epsilon_R^*(\lambda_2) - \epsilon_S^R(\lambda_2)\}, \quad (3.47)$$

where ϵ_S^T is the extinction coefficient of the ground state absorption ($S_n \leftarrow S_0$), eventually leading to the triplet, T , at the indicated wavelength. The extinction coefficient, ϵ_S^R , is similarly defined for the reference compound.

3.4.3. Corrections for Triplet Decay

Comments entirely similar to those presented in Sec. 3.3.3., applicable to the technique of total depletion are again relevant to the method of relative actinometry.

3.4.4. Corrections for Laser Excitation

The achievement of an equal initial excitation population in both sample and reference is usually performed by matching starting ODs in a separate "conventional" spectrometer. If a laser is then employed to populate the respective transients, questions of bandpass differences should be addressed. These are, in general, difficult to answer. However, to be most suitable the relevant photo-physical properties (such as the spectral band shapes) of both sample and reference should be comparable.

3.4.5. Extensions

Obviously, if the initial ODs are different but known then the differential excitation probability may be accounted for by the inclusion of a simple multiplicative factor in either of Eq. (3.46) or Eq. (3.47), whichever is appropriate to the experimental situation.

There is also no need to restrict the reference to another triplet-triplet absorption. The benzophenone ketyl radical, for which formation quantum yields are known, has also provided a useful actinometer.

Recently an interesting extension of the relative actinometry method has been presented⁷⁵ which removes the need for a knowledge of the quantum yields for triplet formation. The triplet populations formed in both the sample and the reference cells are scavenged by biacetyl and the intensities of the resulting phosphorescence are compared. The extinction coefficients are then related by

$$\epsilon_T(\lambda_1) = \frac{\Delta OD_T I_p(R)}{\Delta OD_R I_p(T)} \epsilon_R(\lambda_2). \quad (3.48)$$

The ratio, $I_p(R)/I_p(T)$ of the phosphorescence intensities then takes the place of that involving the triplet yields, though assumptions as to the extent of conversion of individual triplets must be confirmed or kinetic considerations cannot be obviated.

3.4.6. Summary of Advantages and Disadvantages

The principal disadvantage in the technique of relative actinometry as outlined above is the required knowledge of the respective quantum yields for intersystem crossing. Indeed, the method is most commonly used to determine these yields given estimates of the triplet-triplet extinction coefficients⁷⁶. If the quantum yields are known (and not inferred from those measured or estimated in disparate experimental circumstances as is unfortunately common), if the ΔOD can be satisfactorily extrapolated and if the appropriate precautions are taken when singlet depletion is a complicating factor, then this technique should yield extinction coefficients of comparable accuracy of those derived from the other methods documented here.

3.5. Intensity Variation (IV) Method

3.5.1. Description of Method

The intensity variation method was one of the first techniques to be used to measure extinction coefficients of triplet states¹⁷. The initial application gave results which

turned out to be several orders of magnitude too low. Later applications of the method^{77,78}, however gave values that were in accord with those results obtained with different techniques.

The derivation of the method is again based on the kinetic equations for Processes 1.1–1.8, given by Eqs. (3.40) and (3.41). However, conventionally, a steady-state approximation for both singlet and triplet excited states is made and the depletion of the ground state is specifically neglected leading to

$$[^3M^*]/[^1M] = 2303 \tau_T \epsilon_S I_p(x) \Phi_T. \quad (3.49)$$

Here τ_T is the measured triplet lifetime, for example the measured phosphorescence lifetime in glassy solutions and the concentrations are measured at some volume element in the cell at distance, x , along the optical path, from the point of incidence of the photolyzing light. Φ_T is the probability of forming the triplet state from the lowest excited singlet. To obtain extinction information then, it is thus necessary to know the quantum yield for intersystem crossing, Φ_T .

At the time of the inception of this technique, the triplet yield could be related to experimentally measured values in several ways¹⁷. In particular, it could be related to the ratio, R , of the fluorescence to phosphorescence quantum yields and also to the total luminescence quantum yield, Φ . If internal conversion takes place from the singlet state but no intersystem crossing occurs from the triplet to the ground state then

$$\Phi_T = \Phi/(R + 1). \quad (3.50)$$

If the reverse is true, then

$$\Phi_T = (R + 1 - R\Phi)/(R + 1). \quad (3.51)$$

If only intersystem crossing from the singlet state occurs, then

$$\Phi_T = 1/(R + 1) = 1 - \Phi_f. \quad (3.52)$$

Currently many other approaches are in use for determining Φ_T . These methods have been comprehensively reviewed most recently by Wilkinson⁷⁹.

Using Beer's Law for the intensity variation with x , one gets, for an in-line arrangement of the monitoring and photolyzing light, in a manner analogous to the pulsed excitation results of Eqs. (3.9,3.10) that

$$OD = \epsilon_T^* \Phi_T I_p^0 (1 - \exp[-2.3\epsilon_S [^1M] l]) \times 10^3. \quad (3.53)$$

In this equation, everything is assumed known except for the extinction coefficient of the triplet. The intensity, I_p^0 , is varied, and from the slope of the plot of OD vs I_p^0 , one can calculate the extinction coefficient.

The derivation given above follows closely that of Keller and Hadley⁷⁷. They take Eq. (3.53) one step further and assume that the exponential makes no contribution because of a large OD of the ground state at the exciting wavelength. However this is not likely to be the usual case because OD's are usually of the order of unity or lower; in which case the exponential cannot be ignored.

McClure¹⁷ used a modification of this derivation in which the initial concentration, $[^1M]_0$ appeared instead of the concentration of ground state, $[^1M]$ in the steady-state mixture as the denominator of the lefthand side of Eq. (3.49). This lead to a more complicated expression for the conversion to the triplet state. In particular, the intensity dependence varied as $1/(1/I + \text{constant})$. This equation reduces to the one given above when the concentration of triplet is low and when the ground state is not depleted significantly in the steady-state. However because of this dependence, McClure chose to plot $1/OD$ vs. $1/I_p^0$. In effect this means that he was emphasizing the OD at infinite I_p^0 , whereas Keller and Hadley were able to extrapolate their OD vs I_p^0 plots easily to zero I_p^0 which is in conformity to the assumptions of the method.

3.5.2. Summary of Advantages and Disadvantages

This method has been applied to glassy systems with weak steady-state illumination. Under these conditions this method is at its best because neither the energy-transfer nor the singlet-depletion methods can be used.

Beyond these conditions, the method becomes cumbersome. Like some of the other methods photochemistry can cause problems. However in addition one needs a very complete knowledge of the photophysics of the molecule before one can use the method. First of all one must decide which of the three cases to use for Φ_T in Eqs. (3.50–3.52). Then the triplet lifetime and the ratio, R , of fluorescence and phosphorescence yields must be known under conditions of the experiment. That is the minimum information that must be available in addition to the intensity variation measurements. For some of the forms of Φ_T , the absolute luminescence yield must also be known. Finally if the method is to be used, the $1/OD$ vs. $1/I_p^0$ is more susceptible to extrapolation errors than plotting OD vs. I_p^0 .

3.6. Kinetic Method (KM)

3.6.1. Description of Method

Another approach employed to estimate the triplet extinction coefficient, closely related to the intensity variation technique of Sec. 3.5., is termed here the kinetic method. This method was developed originally in conjunction with phosphorescence studies⁸⁰, and later, independently, applied to triplet-triplet absorption spectroscopy^{47,48,81}. The basis of this technique also lies in Eqs. (3.40–3.41). It is convenient for the present derivation to rewrite these as

$$\frac{d[^1M^*]}{dt} = k_{ex} \{[^1M]_0 - [^1M^*] - [^3M^*]\} - k_s [^1M^*], \quad (3.54)$$

and

$$\frac{d[^3M^*]}{dt} = k_{isc} [^1M^*] - k_T [^3M^*], \quad (3.55)$$

where k_s and k_T are defined in Eqs. (1.1) and (1.3) respectively and $[^1M]_0$ is the initial concentration of solute.

In the photostationary state it is assumed that $d[^1M^*]/dt = 0$ which implies

$$[^1M^*] = k_{ex} \{[^1M]_0 - [^3M^*]\} / (k_{ex} + k_s). \quad (3.56)$$

Then, from equation (3.55) and using the fact that $[^3M^*]_0 = 0$,

$$[^3M^*] = \gamma [^1M]_0 \frac{1 - \exp\{-(\gamma + k_T)t\}}{\gamma + k_T}, \quad (3.57)$$

where the (excitation intensity dependent) constant γ is given by

$$\gamma = \frac{k_{ex} k_{isc}}{k_{ex} + k_s}. \quad (3.58)$$

At long times, i.e. $t \gg (\gamma + k_T)^{-1}$ equation (3.57) may be reduced to

$$[^3M^*]_{\infty} = \frac{\gamma [^1M]_0}{\gamma + k_T}. \quad (3.59)$$

Defining the (excitation flux dependent) rise time of the triplet-triplet absorbance, τ_r as

$$\tau_r = \frac{k_s + k_{ex}}{k_{ex}(k_T + k_{isc}) + k_T k_s} \quad (3.60)$$

and the triplet decay time as

$$\tau_T = 1/k_T, \quad (3.61)$$

then equation (3.59) for the photostationary triplet concentration may be recast as

$$[^3M^*]_{\infty} = [^1M]_0 (1 - \tau_r / \tau_T). \quad (3.62)$$

Thus determination of the growth time of triplet absorption and its decay time after cessation of excitation allows the straightforward estimation of the triplet concentration.

Assuming that the monitoring beam does not significantly deplete the ground singlet state (through transitions to higher singlet states for example) then the relation

$$OD_T = \epsilon_T^* [^3M^*]_{\infty} \ell \quad (3.63)$$

is valid and ϵ_T^* may be estimated.

3.6.2. An Alternative Derivation of the Kinetic Method

As opposed to the technique of intensity variation, the kinetic method is inherently a nonstationary approach. It is thus not clear that the traditional method^{47,48,80} of solving equations (3.54) and (3.55) using the steady state approximation, as described in Sec. 3.6.1., is valid. In fact it turns out that this approximation is unnecessary and leads to an unsatisfactory conceptualization. Thus the following is a more elaborate derivation than is usually given. From this derivation it can be easily seen how the usual equations arise and under what conditions the usual equations apply.

The scheme and mechanism are identical to the intensity variation method. However the kinetic method is a time-dependent method. The photolyzing light is turned on as a very long square pulse. Initially the triplet signal builds up after the pulse is switched on. This build-up is

taken to be exponential with a characteristic time, τ_r . When the light of the pulse is turned off at the end of the square pulse, the triplet decays with a lifetime of τ_T , given by Eq. (3.61). These two lifetimes can easily be measured as indicated above, the derivation of the kinetic method relates them to the fraction of triplets in a steady-state mixture that has evolved during the duration of the photolyzing pulse.

A consistent derivation of Eq. (3.57), with a rise time given by Eq. (3.60), starts with a solution of the Eqs. (3.54, 3.55) for the initial conditions $[^3M^*]_0 = [^1M^*]_0 = 0$. The general solutions are given by

$$[^1M^*] = -G_- \exp\{m_+ t\} + G_+ \exp\{m_- t\} + \alpha \quad (3.64)$$

$$[^3M^*] = -Z_+ G_- \exp\{m_+ t\} + Z_- G_+ \exp\{m_- t\} + \beta, \quad (3.65)$$

where

$$m_{\pm} = -(k_{ex} + k_S + k_T)/2 \pm 1/2 \sqrt{(k_{ex} + k_S + k_T)^2 - 4k_{ex}k_{isc}}, \quad (3.66)$$

$$G_{\pm} = \frac{\alpha(k_{ex} + k_S + m_{\pm}) - k_{ex}\beta}{m_- - m_+}, \quad (3.67)$$

$$Z_{\pm} = (k_{ex} + k_S + m_{\pm})/k_{ex}, \quad (3.68)$$

$$k_S = k_f + k_{isc} + k_{ic}, \quad (3.69)$$

$$\alpha = k_T k_{ex} [^1M]_0 / (k_{ex} k_T + k_{ex} k_{isc} + k_S k_T), \quad (3.70)$$

and

$$\beta = \alpha k_{isc} / k_T. \quad (3.71)$$

It can be seen immediately that this derivation gives results that are quite different in general from the results of the standard derivation of Sec. 3.6.1. The results given by Eqs. (3.64–3.71) hold for the duration of the square pulse, given the conditions of no triplets and excited singlets at the beginning of the pulse.

In order to make further progress in reducing the double exponential behavior of the general results in Eqs. (3.65, 3.66) to the single exponential rise time expression of Eq. (3.60), it is necessary to make several approximations. The first step is to remove the radical in Eq. (3.66) by making an appropriate expansion. There are several ways in which this can be done. The most convenient way is to rewrite the radical as

$$\sqrt{(k_{ex} + k_S + k_T)^2 - 4(k_{ex}k_T + k_{ex}k_{isc} + k_Sk_T)}, \quad (3.72)$$

to then extract a factor of $(k_{ex} + k_S + k_T)$, and finally to expand the remaining radical, $\sqrt{[1 - Y]}$, where

$$Y = \frac{4(k_{ex}k_T + k_{ex}k_{isc} + k_Sk_T)}{(k_{ex} + k_S + k_T)^2}. \quad (3.73)$$

The expansion is allowed if $Y < 1$. Y must be ≤ 1 or the radical will be imaginary. After making the expansion,

$$m_+ \sim -\frac{k_{ex} + k_S}{(k_{ex} + k_S + k_T)\tau_r} + O(1) \quad (3.74)$$

and, as always,

$$m_- = -(k_{ex} + k_S + k_T) - m_+, \quad (3.75)$$

where τ_r is defined by Eq. (3.60). From Eqs. (3.74, 3.75), it can be seen that if

$$(k_{ex} + k_S) \gg k_T, \quad (3.76)$$

then

$$m_+ \sim -\tau_r^{-1} \quad (3.77)$$

and

$$m_- \sim -(k_{ex} + k_S + k_T) + \tau_r^{-1}. \quad (3.78)$$

The value of m_+ is exactly what is required of one of the exponentials in Eq. (3.65) if the simplified theory of Sec. 3.6.1. is to be obtained. Furthermore, since m_- contains the measured fluorescence rate constant, one expects it to be too fast to see on the time resolution of most instrumentation conventionally employed in the kinetic method. Mathematically, this empirical requirement that $m_- \gg m_+$ can be written as

$$(k_{ex} + k_S) \gg \tau_r^{-1} \quad (3.79)$$

since one can neglect k_T in Eq. (3.78) for m_- by the condition in Eq. (3.76). So ignoring the second exponential in Eq. (3.65), Eq. (3.65) can be written as

$$[^3M^*] \sim \beta \{1 - (Z_+ G_- / \beta) \exp(m_+ t)\}. \quad (3.80)$$

If conditions can be found such that the coefficient of the exponential, $Z_+ G_- / \beta$, can be shown to be one, then under those conditions Eq. (3.57) will hold. Using the definitions of Z_+ and G_- and applying the conditions in Eqs. (3.76, 3.79) to the expression for $Z_+ G_- / \beta$ becomes

$$Z_+ G_- / \beta = \{k_{ex} - (\tau_r^{-1} - k_T)k_T / k_{isc}\} / k_{ex}. \quad (3.81)$$

This does reduce to one if

$$k_{ex} \gg (\tau_r^{-1} - k_T)k_T / k_{isc}. \quad (3.82)$$

Thus Eq. (3.65) reduces to the traditional result of Eq. (3.57) under the approximations in Eqs. (3.76, 3.79, 3.82).

Even though the standard result holds, the standard justification does not necessarily hold. Applying the three approximations in Eqs. (3.76, 3.79, 3.82) to the ratio of G_- , the coefficient of $\exp\{m_+ t\}$ in Eq. (3.65) for $[^1M^*]$, to α , the time-independent part of $[^1M^*]$ in the same equation, gives

$$G_- / \alpha = \{k_{ex} / (k_{ex} + k_S)\} \times (k_{isc} / k_T). \quad (3.83)$$

Only if this expression is small does the steady-state condition for $[^1M^*]$ hold. The second factor on the r.h.s of Eq. (3.83) is equal to β/α by Eq. (3.71), and β/α is the ratio of the time-independent concentrations of the triplet to singlet. This ratio is not likely to be small in a glass where k_T may be quite slow and where k_{isc} will still have to be large enough compared to the natural fluorescence lifetime or no triplet will be able to form. The only way for

the time-dependent part of $[^1M^*]$ to be small would be for the first factor on the r.h.s. of Eq. (3.83) to be small. This means that the excitation rate constant, k_{ex} , must be very small compared to the decay rate constant for the singlet state, k_s .

3.6.3. Summary of Advantages and Disadvantages

The parallel derivations of the intensity variation technique in Sec. 3.5.1. and the traditional approach to the kinetic method outlined in Sec. 3.6.1. suggest the close kinship of these two methods⁸¹. Both are usually applied to studies in glassy systems because of the inherent mechanical limitations of rotating sector excitation devices which require long triplet lifetimes. The kinetic method has some advantage over the intensity variation technique in that not as many assumptions need be made about the photophysics involved. Rather this information is obtained directly.

3.7. Partial Saturation (SM) Method

3.7.1. Description of the Method

Recently a method⁷¹ was developed that makes use of the large photon fluxes available with lasers. It is different from the total depletion method, which was discussed in Sec. 3.3., but the final form of the descriptive equations is very similar to the expressions employed in that approach. Originally the technique was devised for a situation other than triplet-triplet absorption, but the method has been applied in this field in the last few years.

The basic equation of the method is derived from a two-state model of the excitation process, where the two states considered are the ground state and the triplet state. In the partial saturation method, the ground state may become significantly depleted, and its kinetic behavior can no longer be ignored. The two kinetic equations for the excitation process in this model are given by Eqs. (3.35) and (3.36). Combining the solution given in Eq. (3.38) for the initial conditions $[^1M] = [^1M]_0$ and $[^3M^*] = 0$ with the expression for optical density change given by Eq. (3.23) gives

$$\Delta OD = a(1 - \exp\{-bI_p\}), \quad (3.84)$$

where

$$a = (\epsilon_T^* - \epsilon_S)[^1M]_0 \ell \quad (3.85)$$

and

$$b = 2303\epsilon_S t \Phi_T. \quad (3.86)$$

We will refer to Eq. (3.84) as Lachish's equation for identification purposes. The intermediate excited singlet does not appear in the equations even though it is the state that is initially populated by the radiation. Eq. (3.84) is used by fitting the data to the two-parameter functional form. The excitation intensities must be high enough so that the plot of ΔOD vs. I_p reaches a nonlinear region, a partial saturation region, or the method will not work. When the exponential in eq. (3.84) is expanded in low I_p , the leading term is linear in I_p with a coefficient proportional to $(a \times$

b). There is no constant term, so a and b cannot be determined separately, and thus the ϵ_T^* also cannot be determined independently of Φ_T .

Even though one must use fairly high intensities to reach the nonlinear region, the intensities are lower than those needed to obtain total depletion. Herein lies the advantage of the method over the total depletion method whenever photochemistry is involved.

3.7.2. An Alternate Model for Photoexcitation

In order to see under what conditions the singlet can be ignored in a kinetic model of excitation, we solve a more complete model for the excitation process⁸². The equations in this model will apply only while a square pulse of intensity I_p is on, and the decay of the triplet state will be ignored, as was also done in the two-state model described in Sec. 3.7.1. Such a model for excitation can be written as the following rate equations:

$$\frac{d[^1M]}{dt} = (k_f + k_{ic})[^1M^*] - k_{ex}[^1M], \quad (3.87)$$

$$\frac{d[^1M^*]}{dt} - k_{ex}[^1M] - k_s[^1M^*], \quad (3.88)$$

and

$$\frac{d[^3M^*]}{dt} = k_{isc}[^1M^*]. \quad (3.89)$$

Notice that these equations are simply Eqs. (3.39–3.41) with $k_T=0$ (chosen to correspond with the two-state model above).

The solution to these kinetic equations, using the same initial conditions as in Sec. 3.7.1., together with $[^1M^*]_0=0$, is

$$[^1M] = \Xi \{ (\lambda_2 - k_{isc}) \exp\{-\lambda_2 t\} / \lambda_2 + (k_{isc} - \lambda_3) \exp\{-\lambda_3 t\} / \lambda_3 \}, \quad (3.90)$$

$$[^1M^*] = \Xi \{ -\exp\{-\lambda_2 t\} + \exp\{-\lambda_3 t\} \}, \quad (3.91)$$

and

$$[^3M^*] = k_{isc} \Xi \{ \exp\{-\lambda_2 t\} / \lambda_2 - \exp\{-\lambda_3 t\} / \lambda_3 + (\lambda_2 - \lambda_3) / \lambda_2 \lambda_3 \}. \quad (3.92)$$

The definitions of the symbols used in Eqs. (3.90), (3.91), and (3.92) are

$$\Xi = k_{ex}[^1M]_0 / (\lambda_2 - \lambda_3), \quad (3.93)$$

$$\lambda_2 = (X + Y) / 2, \quad (3.94)$$

$$\lambda_3 = (X - Y) / 2, \quad (3.95)$$

where

$$X = k_{ex} + k_f + k_{ic} + k_{isc} = k_{ex} + k_s \quad (3.96)$$

and

$$Y = \sqrt{X^2 - 4k_{isc}k_{ex}}. \quad (3.97)$$

In order to relate this to Lachish's equation, one can expand Eq. (3.92) in a systematic fashion. This is done by rewriting Eq. (3.97) as

$$Y = X \sqrt{1 - 4k_{isc}k_{ex}/X^2} \quad (3.98)$$

and expanding everything as a power series in

$$\gamma = k_{isc}k_{ex}/X^2. \quad (3.99)$$

The series is valid if

$$k_{isc}k_{ex} < (k_{ex} + k_s)^2. \quad (3.100)$$

The lowest order term in such an expansion is

$$[^3M^*] \sim [^1M]_0(1 - \exp\{-\lambda_3 t\}) + O(\gamma). \quad (3.101)$$

This is almost in the form of Lachish's equation. Using the inequality of Eq. (3.100)

$$\lambda_3 \sim k_{ex}k_{isc}/(k_{ex} + k_s). \quad (3.102)$$

If the further assertion that

$$k_{ex} \ll k_s \quad (3.103)$$

holds, then

$$\lambda_3 \sim k_{ex}k_{isc}/k_s = k_{ex}\Phi_T \quad (3.104)$$

and Eq. (3.101) finally reduces to

$$[^3M^*] = [^1M]_0(1 - \exp\{-k_{ex}\Phi_T t\}). \quad (3.105)$$

Since k_{ex} is proportional to I_p by Eq. (3.37), Eq. (3.105) is equivalent to Lachish's equation.

Thus in order for the two-state model for exciting the triplet state to be valid the condition in Eq. (3.103) must hold. This means that k_{ex} must be in general much smaller than 10^8 s^{-1} . Two examples of photon densities can be used to illustrate whether this is a reasonable expectation or not. In one report⁶⁷ the onset of saturation occurred with pulses of $1.2 \times 10^{-8} \text{ einsteins/cm}^2$, and in another experiment⁷² nonlinearity was observed at 0.2 MW/cm^2 . Taking a moderate absorbance of the ground state of $\epsilon_s = 5000 \text{ L mol}^{-1} \text{ cm}^{-1}$, these two measurements translate into k_{ex} 's of $1.4 \times 10^7 \text{ s}^{-1}$ and $6.7 \times 10^7 \text{ s}^{-1}$, respectively. However the photon intensities in these two experiments are too large for the condition in Eq. (3.103) to be well-satisfied if the singlet lifetime is 10^{-8} s . So in such cases, the a and b parameters obtained by fitting the Lachish equation to the experimental data may not be related to the experimental quantities that they are supposed to represent in Eqs. (3.85) and (3.86).

3.7.3. Summary of Advantages and Disadvantages

The partial saturation method is basically a curve-fitting technique. In order to get extinction coefficients from the Lachish formula, high enough intensities must be reached for the ΔOD vs I_p curves to exhibit nonlinearities. However if the power densities and ground state ϵ_s 's are such that k_{ex} is of the order of k_s , then the more exact formula Eq. (3.92) must apply, but it is not simply related to ϵ_T^* . If the adverse condition of $k_{ex} \sim k_s$ applies, but the formula $\Delta OD = a(1 - \exp\{-bI\})$ is

used anyway, it is still likely that some fit will be obtained since the exact equation looks, in form, something like the Lachish equation. However the meaning of the extinction coefficients obtained by this method will be in doubt.

If this method can be used in an intensity region where the Lachish formula is valid, then the method has distinct advantages over the total depletion method. The advantages lie in that one can operate in some intermediately intense laser excitation regions that may not be plagued with multiphotonic effects such as intensity quenching due to excitation by the laser of the excited states,⁷³ multiphotonic ionization or photochemistry via the excited states,^{72,83,84} or other true multiphotonic processes.⁷⁰

The problem of course is whether a suitable intermediate excitation intensity range does exist for the molecular system under study. This will depend on the laser intensity, I_p , the ground state extinction coefficient, ϵ_s , and the singlet state lifetime, k_s^{-1} ; all of these are components of the condition in Eq. (3.103). If the intensity is too small, no extinction coefficients can be obtained by the method since ΔOD vs I_p must exhibit some nonlinear behavior. However this nonlinear behavior must be due only to saturation, not to the nonlinear processes listed in the last paragraph. Furthermore the intensity must be low enough so that $k_{ex} \ll k_s$ or the simple formula will not apply.

3.8. Miscellaneous Methods for Measuring Extinction Coefficients

3.8.1. Electron Spin Resonance (ESR)

Most of the methods used to measure triplet extinction coefficients try to avoid the difficult problem of measuring the triplet concentration directly. On the other hand the ESR method tackles this problem directly. The method makes use of the property that molecules in the triplet state are paramagnetic and thus have an esr signal⁸⁵.

Brinen and coworkers^{86,87} and Alfimov et al.⁸⁸ have developed this method. The former workers make use of an equation⁸⁹ which relates the concentration of triplets, N_T , which has selection rule $\Delta m = 2$ to the concentration of a free radical, N_r , which has selection rule $\Delta m = 1$. These concentrations are related by the following formula

$$N_T = N_r(15/8) \{(\hbar\nu)^2/(D^2 + 3E^2)\} (I_T/I_r), \quad (3.106)$$

where I_T and I_r are the integrated microwave absorptions of the triplet and radical, respectively, $\hbar\nu$ is the microwave energy, and D and E are the zero-field splitting parameters of the triplet. The factor containing D and E can be evaluated using the method of Kubo and Tomita⁹⁰ which relates it to the first moment, μ_1 , of the triplet esr spectrum by the formula,

$$\mu_1 = (4\pi/7)(D^2 + 3E^2)\nu/(\hbar\nu)^2. \quad (3.107)$$

The advantage of this method is that it attempts to determine the triplet concentration directly. A disadvantage is that the Eq. (3.106) is derived for a uniform concentration of triplets, but the concentration from optical

excitation is governed by Beer's Law. One way to minimize the effects of this last problem is to use small concentrations. Then the variable concentration represented by Eq. (3.10) can be expanded to keep only the first term, and the concentration is roughly constant⁵⁰.

3.8.2. Other Comparative Techniques (ELT,HAT,RF)

In the energy transfer technique there is the transfer of triplet energy from (to) a donor (acceptor) compound with known extinction coefficient to (from) the acceptor (donor) whose extinction coefficient is sought. Particularly with the development of nanosecond pulsed laser and electron sources the possibility of quantifying other photochemical quenching has been realized. Thus the extinction coefficient of the triplet state of 5-nitro-2-furoic acid has been estimated relative to the radical cations of various amines from which it has abstracted an electron⁹¹ (ELT). Hydrogen atom abstraction (HAT) by ketone triplets leads to the corresponding ketyl radicals for which extinction data is often independently obtainable. Photoreduction of benzophenone by lactams has been used in this fashion to evaluate the triplet properties of the ketone⁹². These approaches assume of course that some estimate of the amount of conversion of the triplet state can be obtained and that appropriate corrections can be made for the absorbance accompanying the production of the associated ionic and radical species.

The basis of another comparative method (RF) rests in the observations of Lavalette et al.⁹³ They have noted that the oscillator strength of the principal triplet-triplet absorption peaks in a series of aromatic hydrocarbons remains roughly constant for a given molecule in a range of solvents. Apart from a (usually small) refractive index correction, this consistency is expected on theoretical grounds, and the result serves as a basis for another useful comparative technique.

To estimate the oscillator strength, f , associated with a particular triplet-triplet absorption band, the area under the absorption profile expressed as a function of wavenumber must be measured. Note also that the extinction coefficient must first be known. Thus if we integrate the identity of Eq. (3.26)

$$\int OD(\nu) d\nu = \int \epsilon_T^* [^3M^*] \ell, \quad (3.108)$$

$$= \alpha_n f [^3M^*] \ell, \quad (3.109)$$

where the constant of proportionality, α_n , contains only fundamental constants and the solvent refractive index.

Using Eq. (3.26) once more to eliminate the unknown product, $[^3M^*] \ell$, gives an equation relating the oscillator strength to the area under the experimentally determined OD curve scaled to the maximum value of the extinction coefficient for the band.

$$\frac{\epsilon_T^*(\nu_{\max})}{OD(\nu_{\max})} \int OD(\nu) d\nu = \alpha_n f. \quad (3.110)$$

If the empirical OD curve is normalized to unit maximum absorbance and the enclosed area termed Σ , then

$$\alpha_n f = \epsilon_T^*(\nu_{\max}) \Sigma. \quad (3.111)$$

Thus for some compound in a certain solvent, if either the oscillator strength is known or if the extinction coefficient is available *and* an appropriate area can be measured, then simply repeating this area measurement in subsequent solvents allows the determination of the corresponding extinction coefficients.

We note again that strictly this procedure ignores the variation in refractive index in a set of solvents. A number of experimental difficulties also arise. Often the complete spectral shape for a given electronic band cannot be detected. Thus some extrapolation of the observed OD is usually necessary. Often, the assumption of "triangular" bands is made, and the area is approximated as the product of the height and halfwidth, δ , (i.e. full width at half maximum). In terms of the area, Σ , defined above this simplification amounts to a direct comparison of half-widths.

$$\alpha_n f = \epsilon_T^*(\nu_{\max}) \delta. \quad (3.112)$$

Preferably, some extrapolation of the observed OD should be attempted. Commonly the bands are assumed to be symmetrical around their maxima (as a function of wavenumber). Obviously, the OD observed must be strictly attributable to the unknown triplet state or else corrections such as those documented in the foregoing sections must be assessed. Also the electronic nature of the band must remain reasonably intact in the disparate solvating environments.

3.8.3. Pulsed Intensity Variation

The formulation of the intensity variation technique presented in Sec. 3.4. above, with its steady-state assumptions, is adequate under conditions of steady illumination⁷⁷ or conventional flash photolysis⁷⁸. With nanosecond pulsed excitation, however, a time-dependent solution of the corresponding kinetic equations must be sought. Such a solution, for low intensity, is possible for a square pulse within the context of the simple two-state model given in Eqs. (3.35–3.38) above. If the pulse is of width, Γ_p and the excitation rate, k_{ex} , is as in Eq. (3.37), then we find⁸²

$$[^3M^*]_{\Gamma_p} / [^1M]_0 \sim \Gamma_p \Phi_T k_{ex} - (\Gamma_p \Phi_T k_{ex})^2 / 2. \quad (3.113)$$

Triplet decay within the pulse has been specifically neglected. For a fixed pulse width a linear variation of ΔOD with pulse intensity is thus expected at low intensities. The slope of this line is proportional to the product of the triplet yield and the required extinction coefficient⁹⁴. This approach has recently been adopted⁹⁵ to suggest a refinement of the triplet extinction coefficient for benzophenone, a commonly used reference value.

Solution of the three-state model given in Eqs. (3.39–3.41) again with the assumption of no triplet decay during the pulse leads to a more complex limiting behavior. The end-of-pulse triplet concentration is given by⁸²

$$[^3M^*]_{\text{r.p.}}/[^1M]_0 \sim \Gamma_p \Phi_T k_{\text{ex}} + k_{\text{ex}} \Phi_T \exp(-k_s \Gamma_p)/k_s + O(k_{\text{ex}}^2). \quad (3.114)$$

It thus appears that even at low intensities a further requirement must be met to achieve correspondence with the simple result presented above; the rate constant for singlet deactivation must be sufficiently fast to reduce the contribution of the second term in Eq. (3.114). Otherwise a false estimate of the extinction coefficient will be obtained. However, empirically, the change in optical density upon laser photolysis is recorded after the induced fluorescence has subsided. Thus a more suitable comparison with the two-state result of Eq. (3.113) is provided by the long-time solution of the three-state approach which is⁸²

$$[^3M^*]_{\infty}/[^1M]_0 \sim \Gamma_p \Phi_T k_{\text{ex}} - (\Gamma_p \Phi_T k_{\text{ex}})^2/2 + k_{\text{ex}}^2 (\Phi_T^2 - \Phi_T) \{ \Gamma_p/k_s + \exp(-k_s \Gamma_p)/k_s^2 \}. \quad (3.115)$$

This expression differs from the commonly used limiting form of Eq. (3.113) only in terms of second order in k_{ex} . Thus, provided no curvature is observed in plots of ΔOD against intensity, the simple relation

$$[^3M^*]/[^1M]_0 \sim \Gamma_p \Phi_T k_{\text{ex}} \quad (3.116)$$

will be valid.

The principal disadvantage of this technique is, of course, that Φ_T must be known. This problem is weighed against the fact that at low intensities complications such as photochemistry can be alleviated.

4. Evaluation Procedure

4.1. General Methodology


The above discussions of the problems encountered in the application of the various types of excitation sources in Sec. 2. and, in particular, of the advantages and disadvantages inherent in the methods employed to determine the extinction coefficients of triplet-triplet absorption bands listed in Sec. 3., serve as a framework for a scientific evaluation of the wavelength maxima and extinction coefficient data presented in Table 6.

Thus we have pointed out various precautions and corrections required in each method and have outlined their respective domains of applicability. It was initially intended to develop an automatic evaluation procedure based on a quality indicator derived from a list of requirements which, ideally, would have to be satisfied to ensure sound application of the experimental extinction techniques. Unfortunately, in the course of extracting data in the primary literature, we regularly encountered inadequate reporting of experimental methodology. This defi-

ciency should hopefully be remedied in the future if the guidelines for the presentation of photochemical and photophysical data issued by IUPAC^{96,97} are routinely followed. For the purposes of the present compilation we have had to judge the data firstly on a case by case basis as they were converted for entry into our database and save any global comparison until this compilation was near completion. This procedure represents, then, a first step in our evaluation.

Many decisions and interpretations had to be made merely to get the data into our structured data file. Insufficient information was often present in a given work to fulfil even the minimum requirements for tabulation. In this case similar articles by the same authors could often be found and some extrapolation made to glean the necessary data for entry into the table. If this extrapolation was very great, it was documented in the comments column. A most common inadequacy was failure to mention the solvent system used.

A problem central to the kinetic spectroscopy of transients is the identification of the absorbing species. We indicate the method(s) adopted, where these were reported, in the comments for each entry in the main data table. Often, however, comparison with an earlier spectrum of the same molecule or a closely similar compound is implied (if not stated), and this comment has not been included. Of course we are careful to note or eliminate bands which have been interpreted or proven by later authors to be due to other species. Oxygen quenching, the susceptibility of the transient to aeration or oxygenation of the ambient medium, has been most widely quoted as an identification method. Unfortunately the decay rates of many transient species are reduced in the presence of oxygen either through photochemical quenching or energy transfer. For triplet states, the most likely process is thought to be⁹⁸



Another popular technique involves the sensitization of (by) the transient by (of) well characterized triplet state species. In this method it is implicitly assumed that only $^3D^* \rightarrow ^3A^*$ transitions are effective in transferring energy to (from) the triplet states of known acceptors (donors). However, both $^3D^* \rightarrow ^1A^*$ energy transfer⁹⁹ and $^1D^* \rightarrow ^3A^*$ energy transfer¹⁰⁰ have been reported, though the latter is quite rare. These two approaches are most reliable when some estimate of the quenching rates involved is given or when multiple references have been used, for example in an effort to bracket the triplet energy of the unknown sample.

If the absorption can be followed in a glassy matrix, and if the observed transient lifetime matches the decay of phosphorescence, then a triplet state can be confidently assigned. This technique is particularly useful carried out at a number of distinct wavelengths. Corresponding decay times would thus eliminate the possibility of multiple transients. Again in glassy samples, if the transient can be detected by ESR techniques, then the presence of $\Delta m =$

2 transitions in the ESR spectrum provides convincing evidence of its triplet nature. Another interesting technique involves an additional monitoring of a fluorescence signal. If the square of the transient absorbance can be correlated with the intensity of delayed luminescence arising from triplet-triplet annihilation, then a triplet state source is likely⁹⁴.

Simply observing a first-order decay of the absorption signal on a time scale appropriate for triplet disappearance under the conditions used, has been cited by some authors as evidence for triplet state participation. Radicals often vanish through second order processes and can be longer lived. But this method is obviously not free from criticism. Over the course of this work we have formed the impression that serious identification strategies are only attempted when the assignment is either novel or questionable.

Another example of a problem encountered in the data extraction phase is the precise chemical nature of the triplet species, for instance the state of ionization especially when the solvent is water. This problem was addressed by collecting pK_a 's (or pK_b 's) of the triplet conjugate acids (or conjugate bases), by including these in the comments and then by analyzing which triplet species should be present at the pH specified.

As we have stated previously, there exists a good deal of agreement among the values reported by various workers, using different excitation methods for the wavelength maxima of the triplet-triplet absorption spectrum of a particular compound in a particular environment. Solvent shifts in λ_{max} are also consistently documented in general. An exception however is found in the values of λ_{max} derived from uncorrected difference spectra. As a difference spectrum approaches the singlet depletion, SD, region, ΔOD drops precipitously. This effect can add spurious maxima and/or greatly shift maxima that occur in or near the SD region. For this reason we have attempted to avoid quoting maxima near the SD region which have been clearly obtained directly from difference spectra. Also we have not attempted to locate such peaks on published difference spectra. Indeed we have often simply omitted mention of original articles which record spectral data derived solely from such difference spectra. This criterion, unfortunately, means that we have excluded more data relevant to such molecules as porphyrins and chlorophylls than was our original intent. Most reports of triplet-triplet absorption in these compounds contain only uncorrected difference spectra and, in addition, such spectra occur almost entirely in the SD region.

Unfortunately, the relative reproducibility revealed in the various measurements of λ_{max} is not present in the corresponding estimations of the extinction coefficient, ϵ_{max} . Data for the same molecule in different solvents are of course subject to a wide scatter. However, data for the same solvent/molecule system, but obtained by different measurement techniques, are still inconsistent. Two of the principal methods employed to determine the extinction coefficient of triplet-triplet absorption, the energy trans-

fer method, ET, (see Sec. 3.1.) and that of relative actinometry, RA, (see Sec. 3.4.) as well as a number of the less widespread approaches (see Sec. 3.8.2.) are comparative techniques. Ideally the data derived from these methods should form a network within which questions of internal consistency both within a given technique and across a range of approaches could be answered. A number of problems remain which currently prohibit the achievement of such a goal. Usually authors mentioned the standards that they used for reference in some detail. However quite often the authors would quote a standard that was measured in a totally different solvent. The reference ϵ may have been measured in a similar solvent, but this does not automatically guarantee that it will be unchanged in the solvent that the authors found convenient to study. A good deal of work has been done to document⁹⁵ that this apparent neglect of solvent effects underlies much of the apparent discrepancies between ϵ_{max} measured in different laboratories (even if this determination is performed by the same technique). In fact it was shown that the use of oscillator strengths would serve to be a better guide to relating the absorption spectrum of a solute in one solvent with its spectrum in another solvent⁹⁵.

This empirical rule that oscillator strengths are relatively independent of solvent has a theoretical justification. Most of the well-studied spectra are allowed transitions; thus solvent effects, being first- or second-order perturbations, will not contribute a very large fraction to the transition matrix elements, which are directly related to oscillator strengths and not to ϵ 's. In order to deal with this problem for this compilation, we made a special effort to report, in the comments for comparative methods, both the reference ϵ used in addition to the solvent which was used to get that particular reference ϵ and the relevant oscillator strengths if given.

In summary, then, roughly 30% of the entries in the Table contain extinction information. Of these, about 40% are relative measurements which can be checked for internal consistency. Data derived from other techniques lie outside of this network, and their quality must be judged accordingly.

4.2. A Statistical Analysis

4.2.1. Statistical Test of Solvent and Temperature Effects

As a further step in a quantitative evaluation of the triplet-triplet extinction coefficients, we undertook a short statistical analysis of the data. One of the first things that strikes an observer of the extinction coefficient measurements is that there is a large variation between the values derived for most compounds for which multiple determinations exist. The question arises as to whether these variations are so large as to swamp more systematic scatter arising for example in solvent effects. Some progress¹⁰¹ has already been made in evaluating photophysical and photochemical data by ignoring solvent effects and by treating all the variation between measurements as random.

To judge whether or not the systematic variations can be ignored, we ran a series of statistical tests on one of the key compounds. The approach chosen was a *t* test which gauges the validity of the hypothesis that two sample distributions had the same mean. The test assumes that the variances of the two sample distributions are the same, but unknown. The procedure is described by Ostle and Mensing.¹⁰² A test statistic, *T*, is used in the method, and *T* is defined by

$$T = (\langle X_1 \rangle - \langle X_2 \rangle) / \sqrt{\{S^2(1/n_1 + 1/n_2)\}}, \quad (4.1)$$

where *S*² is the estimate of the common variance given by

$$S^2 = \{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2\} / (n_1 + n_2 - 1) \quad (4.2)$$

and where $\langle X_i \rangle$, *n_i*, and *s_i* are the mean, size, and sample standard deviation, respectively, of the *i*th sample. The critical region of the test is given by

$$\text{abs}(T) \geq t_{(1-\alpha/2)(n_1+n_2-2)} \quad (4.3)$$

t is obtained from tables of Student's *t*-distribution, and $\alpha \times 100\%$ is called the level of significance of the test. The level of significance is the probability that one will reject a true hypothesis for the given test procedure. Traditionally, the acceptance or rejection of hypotheses is based on a 5% level of significance, or even on a 1% level if it is really important to avoid rejecting a true hypothesis. The test is done by first calculating *T*, from Eq. (4.1) and then checking to see whether the calculated *T* falls in the region defined in Eq. (4.3) for the chosen level of significance. If *T* falls in this region, then the hypothesis is rejected.

Five such *t*-tests were run on the anthracene data using 31 measurements. The 2800 value was discarded because of a systematic error in the measurement, and another value was discarded from these initial tests because no solvent conditions were given. Five separate tests were run, but tests were made on three general areas. First the data was tested to see if the measurements in polar environments and nonpolar environments had different means. Second it was tested whether measurements at low-temperature and room temperature had the same means. Third it was tested whether measurements in benzene vs other nonpolar solvents were different.

In all of the five specific tests for anthracene, the hypotheses that "the means are equal" in all cases could not be rejected at the 5% level of significance. In fact the observed values of the test statistics were just significant at the "levels of significance" ranging from 14 to 30%, see Table 3. These values, being in the range of 14 to 30%, are well above the traditional levels of 5% or 1%. This implies that if one chooses to reject the hypotheses that "the means are equal", then one runs quite a large risk, 14 to 30%, of rejecting a true hypothesis.

The simple sequence of *t*-tests just described is only one way to test for the equality of a series of means. It can lead to several problems; one of which is the problem of transitivity of the means. This problem is that if one tests means two-by-two it is possible to find no significant differences from the tests even when the differences are

present. This could happen if not all possible pairs are tested. However if all pairs of means are tested two-by-two then it becomes more likely that a mistake will enter in the form of a rejection of a true hypothesis.¹⁰³

In order to circumvent the problems of doing a large number of two-by-two tests, but yet being able to test all the means, one can use an *F*-test along the lines of a technique called Analysis of Variance (ANOVA). The specific test we chose is described on pp. 123-5 of Ostle and Mensing¹⁰² with the ANOVA for an unequal number of experimental units described¹⁰² on pp. 295-7. The hypothesis to be considered is that "the means of all the groups are equal" as opposed to the alternate hypothesis that "at least one of the group means is different than the other means." In the ANOVA technique one compares the variation *within* the groups to variations *among* the groups. If the variation among the groups is significant relative to that within the group, then the hypothesis that the means are equal is rejected.

For the ANOVA, we chose five groups related to solvent and temperature among the anthracene measurements. The five groups were polar-RT, polar-LT, benzene-RT, cyclohexane-RT, and nonpolar-RT. RT and LT refer to room temperature and to low temperature, respectively. The nonpolar group did not contain any benzene or cyclohexane data. With this five-level classification of the same 31 measurements of anthracene as in the *t*-tests, we obtained an *F*-ratio of about 1. This indicated that the hypothesis was not to be rejected at the usual levels of significance. However since there may be extreme values that are significantly affecting this result, we used Chauvenet's criterion (see Sec. 4.2.2.) on the data to check for such outliers. The value of 118000 failed and was rejected. With this revision of the data groups we performed the ANOVA again and obtained the traditional ANOVA table given in Table 4.

TABLE 3. *t* tests of solvent effects in anthracene

Dual comparisons	No. in samples	<i>t</i> test statistic	<i>f</i> ^a /%
Polar / Nonpolar (RT ^b)	9/16	1.06	30
Polar (RT) / Polar (LT ^c)	9/6	1.61	14
All (RT) / All (LT)	25/6	1.21	24
Benzene / other Nonpolar	6/10	1.09	30
Benzene / Cyclohexane	6/5	1.17	28

^a Comparison is statistically significant at *f*%.

^b Room temperature.

^c Low temperature (usually 77 K).

TABLE 4. ANOVA table for anthracene ^a

Source of Variation	Degrees of Freedom	Sum of Squares / $\times 10^9$	Mean Square / $\times 10^8$	F Ratio
Mean	1	111	1110	—
Among Groups	4	1.11	2.78	1.15
Within Groups	25	6.05	2.42	—
Total	30	111	—	—

^a Excluding outlier.

The F-ratio of 1.15 was compared to an F -distribution with 4 and 25 degrees of freedom and was found to be not significant at the traditional levels. The F-ratio would have to be about 3 before one could reject the equality of the group means at a 5% level.

Both the t -tests and the ANOVA have limitations. They complement each other in the sense that the ANOVA assumes that all of the groups have the same variance, but each t -test makes a separate assumption about the equality of the variances of the means that it is testing. The ANOVA is more limited in this sense, but it has the advantage that it tests the means only once as discussed above.

One feature that is shared by both tests is the assumption of an underlying normal distribution. It is relatively easy to test whether the large distribution of 30 measurements used in the ANOVA is normal or not. The procedure we chose was the Kolmogorov-Smirnov test, described in a textbook by DeGroot¹⁰⁴ on pp. 465-8. The largest difference, D_n^* , between the empirical distribution function and the distribution from a standard normal distribution was 0.10 for the data used in the ANOVA above. Following through the rest of the procedures for the Kolmogorov-Smirnov test, we found a tail region of over 90%, which is very far from the 5% or 1% tails traditionally used in rejecting the hypothesis that the distribution is normal.

Another assumption of the statistical models underlying the t -tests and the ANOVA is that the variances of the normal distributions of the groups being tested are equal. One can also test this assumption if one uses an F -test. The F statistic chosen¹⁰² is

$$F = s_1^2/s_2^2 \quad (4.4)$$

where the rejection region of the test is

$$F \geq F_{(1-\alpha/2)(n_1-1, n_2-1)} \quad (4.5)$$

or

$$F \leq F_{(\alpha/2)(n_1-1, n_2-1)} \quad (4.6)$$

As before s_i and n_i are the sample standard deviation and size of the i th sample, respectively, and the F 's on the righthand sides of Eqs. (4.5, 4.6) are taken from tables of the F -distribution with n_1-1 and n_2-1 degrees of freedom. The variances of the five pairs of groups in Table 3 were tested using this test, and none of the five tests showed any statistically significant result.

In summary the t -tests and the ANOVA both failed to detect differences between the means of the groups of anthracene measurements based on solvent and temperature. This does not imply that the measurements all truly come from the same distribution and that there are no solvent or temperature effects. But it does indicate that, given the data in the literature, we cannot reject the hypothesis that *there is no statistically significant difference between the means of the various solvent-temperature groups.*

4.2.2. Mean ϵ and Confidence Intervals

Based on the results of the statistical tests of solvent and temperature effects on anthracene, we proceeded with the statistical analysis by ignoring these effects. We studied the 32 compounds that had five or more measurements at a particular wavelength. The number five was chosen because it allows reasonable statistics to be done, such as using Chauvenet's criterion.

Since the samples were taken from a distribution whose mean and variance are both unknown, we used the t factor to calculate the confidence interval.¹⁰² The assumption made here is that the sample is distributed normally with unknown mean and variance. The 95% confidence interval was calculated from

$$CI = t_{0.975(N-1)} \times s / \sqrt{N}, \quad (4.7)$$

where s is the sample standard deviation and N is the number of measurements. The values of $t_{0.975(N-1)}$ were taken from tables of the t -distribution. The calculated means of the samples of measurements on the various compounds are given in Table 5 along with the calculated 95% confidence intervals.

In an attempt to critically evaluate the data, we chose Chauvenet's criterion to examine the individual measurements on each of the 32 compounds. The guiding principle¹⁰⁵ of Chauvenet's criterion is that individual measurements should not inordinately influence the calculation of the mean of a sample. If an individual measurement has a much larger (or smaller) deviation from the mean than any other measurements, then it will pull the value of the mean toward it to a much greater extent than other measurements pull the mean toward themselves. Measurements that do affect the mean in such an unwanted fashion can be identified and eliminated from the sample if desired. The explicit form is such that if the deviation from the mean, d_i , is greater than q , where q is given by

$$1 - \{1/s\sqrt{2\pi}\} \times \int_{-q}^{+q} \exp\{-y^2/(2s^2)\} dy = 1/(2N), \quad (4.8)$$

then the i th measurement can be eliminated from the calculation of the mean. This is Chauvenet's criterion.

Chauvenet's criterion gives a more flexible procedure than simply discarding measurements that are 2 (or some other fixed number of) standard deviations away from the mean.¹⁰¹ For the same sample standard deviation, s , a sample with $N=5$ would throw away measurements that had $d_i \geq 1.68 \times s$. However, a sample with $N=30$ would not throw away a measurement unless $d_i \geq 2.39 \times s$. This implies that the criterion is much more tolerant of large deviations in large samples than it is in small samples, which is just another way of saying that the mean of a small sample is more sensitive to a large deviation than is the mean of a large sample.

TABLE 5. Average extinction coefficients

Compound Name (λ_{\max}/nm) ^a	No.	$\langle \epsilon \rangle$ /L mol ⁻¹ cm ⁻¹	95% confidence interval	$\epsilon(\text{benzene})^b$ /L mol ⁻¹ cm ⁻¹	$\epsilon(\text{cyclohexane})^b$ /L mol ⁻¹ cm ⁻¹
1. Acridine (440)	12	23900	± 3250 ($\pm 14\%$)	24300	31500
2. Anthracene (430)	31	61000	± 5670 ($\pm 9\%$)	45500	64700
3. Benz[a]anthracene (490)	10	23300	± 4190 ($\pm 18\%$)	20500	28800
4. Benzo[a]coronene (570)	5	22300	± 4170 ($\pm 19\%$)	—	—
5. Benzophenone (525)	17	7640	± 1050 ($\pm 14\%$)	7630	—
6. Biphenyl (360)	8	36600	± 6300 ($\pm 17\%$)	27100	42800
7. Carbazole (425)	6	14000	± 3110 ($\pm 22\%$)	—	—
8. 3-Carbethoxypsoralen (450) ^c	7	4620	± 1870 ($\pm 40\%$)	—	—
9. β -apo-14'-Carotenal (480)	4	120000	± 9840 ($\pm 8\%$)	—	—
10. Chloranil (510)	5	6810	± 1170 ($\pm 17\%$)	—	—
11. Chrysene (580)	7	30000	± 13700 ($\pm 46\%$)	—	—
12. Dibenz[a,h]anthracene (580)	6	21600	± 10000 ($\pm 46\%$)	—	—
13. 9,10-Diphenylanthracene (445)	6	16500	± 2230 ($\pm 13\%$)	—	—
14. DPB (390)	5 ^d	52600	± 8890 ($\pm 17\%$)	—	—
15. DPH (420)	7	112000	± 5390 ($\pm 5\%$)	—	—
16. DPO (440)	7	189000	± 22900 ($\pm 12\%$)	—	—
17. Duroquinone (490)	10	6180	± 1330 ($\pm 21\%$)	6950	5330
18. Methylene Blue ⁺ (420)	4	10800	± 3760 ($\pm 35\%$)	—	—
19. Methylene Blue ⁺ , protonated (370)	4	13800	± 1910 ($\pm 14\%$)	—	—
20. C ₁₇ -aldehyde (440)	7 ^d	59600	± 6640 ($\pm 11\%$)	—	—
21. Naphthalene (415)	16	22500	± 5230 ($\pm 23\%$)	13200	24500
22. Naphthalene-d ₈ (415)	6	23500	± 11700 ($\pm 50\%$)	—	—
23. 5-Nitro-2-furoic acid (490)	5 ^d	20700	± 1550 ($\pm 7\%$)	—	—
24. Phenanthrene (490)	10	26900	± 6500 ($\pm 24\%$)	15700	25200
25. Pyrene (415)	5	36400	± 14700 ($\pm 40\%$)	20900	30400
26. 1-Pyrenecarboxaldehyde (440)	10 ^d	18400	± 311 ($\pm 2\%$)	—	—
27. 11- <i>cis</i> -Retinal (450)	6	49700	± 20300 ($\pm 41\%$)	—	—
28. <i>all-trans</i> -Retinal (450)	14	71000	± 3420 ($\pm 5\%$)	—	—
29. TMPD (620)	5	15300	± 6610 ($\pm 43\%$)	—	12200
30. Triphenylene (430)	9	12700	± 4120 ($\pm 32\%$)	—	—
31. Ru(bpy) ₃ ²⁺ (370)	5	27600	± 1830 ($\pm 7\%$)	—	—
32. Zinc(II) phthalocyanine (480)	5	33300	± 15400 ($\pm 46\%$)	—	—

^a Nominal wavelength maximum (large environmental and instrumental shifts are often present).^b Data from R. Bensasson and E.J. Land, Trans. Faraday Soc. 67, 1904 (1971).^c 450 nm not necessarily a maximum.^d Average computed from one paper only. Scatter in the data for these compounds is likely due to real solvent and temperature effects.

The measurements that do not satisfy Chauvenet's criterion for the 32 compounds studied are marked in Table 6. These values are not included in the averages listed in Table 5. There are two other type of measurements that we have excluded from the averages in Table 5 and have marked in Table 6.

The first includes measurements that are commonly acknowledged to contain systematic errors. In this class we have marked only values from some of the first papers to measure extinction coefficients of triplet-triplet absorption. These values are uniformly too small by an order of magnitude or so.

The second type of measurement that we have excluded from averages in Table 5 are measurements in mixed crystals, neat liquids and micelles. These measurements are also uniformly low compared to other measurements. In this case however, we believe the values are low for a good physical reason. It seems likely that there are intermolecular interactions between the chromo-

phores that are causing the hypochromic effect. This intermolecular effect is well studied in DNA (see Chase and Rhodes¹⁰⁶ for a theoretical review). A similar effect is still likely in mixed crystals. Depending on the host, intermolecular communication between the guest molecules is possible.

4.2.3. Discussion of Statistical Results

The above statistical analysis can be summarized as follows: Based on the statistical tests on the anthracene data, we found no statistically significant solvent or temperature effects. Furthermore the data for anthracene appear to follow a normal distribution. Generalizing these two statistical results to all the compounds, we calculated means and 95% confidence intervals for those compounds with more than five measurements. In addition we chose again Chauvenet's criterion to tag outliers in the data.

However, as is well known from other varieties of molecular spectroscopy, solvent and temperature effects

do in fact exist. The question then arises as to how the statistical results obtained above should be viewed.

To support the view that solvent and temperature effects are significant in this data, one only needs to look at Table 6 and see how the extinction coefficients often go up, particularly in glasses at low temperature. So the data do in fact show systematic variations due to solvent and temperature. Furthermore the compounds tagged with superscript "d" in Table 5 are reported in single papers and should therefore give some indication of the solvent effect because systematic errors due to different laboratories are "blocked out", in the statistical sense. Finally the widely used standards⁶¹ listed in the last two columns of Table 5 show a significant deviation between cyclohexane and benzene which might not be expected to be so large because of the similarity of the physical properties of these two solvents.

The conflict between the lack of statistically significant solvent effects and the appearance of them in the several forms as listed above can be rationalized, of course. The resolution of the conflict most likely lies in the random experimental errors in the difficult measurements and in the various systematic errors between laboratories and methods. These errors are large enough so that the discrepancies between measurements within solvent groups are comparable to or larger than fluctuations among solvent categories. A quantitative test of this was given above in the ANOVA procedure on the anthracene data.

The sources of these systematic errors due to methods have already been discussed extensively in Sec. 3. For convenience we list some of the more prominent conclusions here again. Singlet depletion (SD) gives an upper bound due to underlying triplet absorption. Total depletion (TD) gives a lower bound due to lower triplet concentrations either from competing photophysical (multiphotonic) or photochemical processes or from insufficient laser pulse length. Results from the energy transfer (ET) technique can be either high or low due to lack of unit probability of energy transfer (high when the unknown is the triplet donor and low when the unknown is the triplet acceptor).

The systematic errors due to variations between laboratories may partially be accounted for by the systematic errors incurred by the choice of method, but an additional source is to be found among the various procedures adopted in different laboratories. Some of the more prominent of these include improper overlap of monitoring beam with exciting beam,⁵³ monochromator slits too wide for sharp spectra (particularly important for the aromatic hydrocarbons, leading extinction coefficients being too low),⁵⁰ different standard extinction coefficients for relative methods like energy transfer and relative actinometry, improper calibration of monochromator (making these standards even more tenuous in relative methods and throwing off the singlet wavelength where the standard ϵ is measured in singlet depletion measurements), and insufficient number of wavelength measurements to clearly define the absorption peaks.⁷ Many of the commonly used reference extinction coefficients

are unfortunately for compounds with particularly sharp spectra such as anthracene.

4.3. Tentative Standard ϵ 's

Taking into account that experimental errors, both methodological and instrumental, seem to induce variations in the data that are greater than, or of the same order as, the solvent effects, it is tempting to use the average extinction coefficients in Table 5 as standards. We feel it would be premature to do this at this time. However lacking further analysis, we would recommend that these values represent some of the best ϵ 's that are available at this time.

Until further analysis can be performed, we recommend that a hierarchy of values be used. First we would still recommend the Bensasson-Land values⁶¹ in the last two columns of Table 5 if the solvents used are benzene or cyclohexane. The only exception to this is that 7220 L mol⁻¹ cm⁻¹ should be used for benzophenone in benzene.⁹⁵ For these measurements great care was taken to make these values stand as bench marks. Our only reservation is that these measurements were made using primarily the energy transfer method and are susceptible to its weaknesses. The measurements are also susceptible to the usual instrumental errors, but great care seems to have been taken to avoid the instrumental pitfalls in these two works.

On the next level of the hierarchy of values, we tentatively recommend the average values in Table 5. These values are not equally recommended and should only be used in a nested hierarchy. The hierarchy we recommend can be used as follows: First, for compounds that have values in one of the last two columns of Table 5, use the average ϵ 's only for solvents that are not benzene or cyclohexane. Second, use the average values of ϵ for all other compounds with exceptions being made for the compounds with confidence intervals exceeding $\pm 30\%$. Third, average ϵ 's for compounds with large confidence intervals of over $\pm 30\%$ should not be used as a standard in a relative measurement of an unknown extinction coefficient (or quantum yield) unless no other compound with a more reliable standard can be used.

At this point it should be noted that the average ϵ 's with their accompanying confidence intervals are not always consistent with the values in the last two columns of Table 5. This is true even for the compounds with confidence intervals of less than $\pm 30\%$. The most notable example is the Bensasson-Land value⁶¹ for anthracene in benzene which falls outside the computed 95% confidence interval. The suggested hierarchy is set up to favor the Bensasson-Land values in this case.

Of the 10 compounds in Table 5 with confidence intervals exceeding $\pm 30\%$, most have small sample sizes. The sources of the large confidence intervals vary somewhat among the compounds. Over half of these compounds are aromatic hydrocarbons with sharp peaks that sharpen even further at low temperature. With the small sample sizes, our assumptions of normality and comparable experimental variations vs. solvent variations may not hold

for these compounds. For example in the case of chrysene there are only 7 measurements. Of these, three are measurements in low temperature glasses and have $\epsilon > 36000 \text{ L mol}^{-1} \text{ cm}^{-1}$. Another of the 7 measurements was an early¹⁰⁷ measurement that seems unusually low at $8800 \text{ L mol}^{-1} \text{ cm}^{-1}$. For chrysene Chauvenet's criterion does not help because of the particular scatter of the data. Furthermore there seems to be a well-founded physical reason for the scatter, namely sharper spectra at lower temperature, violating our assumption that temperature effects are comparable with experimental errors.

Another of the 10 compounds with larger than $\pm 30\%$ confidence intervals that is not well represented by the average ϵ 's is 11-*cis*-retinal. Like the aromatic hydrocarbons, it has a large confidence interval. In addition some workers¹⁰⁸ believe in a common triplet state for some polyene trans and cis isomers. If this were so in the all-trans and 11-*cis* isomers of retinal, then 11-*cis*-retinal should have an $\epsilon \sim 70000 \text{ L mol}^{-1} \text{ cm}^{-1}$. This points to the two values¹⁰⁹ of $27000 \text{ L mol}^{-1} \text{ cm}^{-1}$ as either suspiciously low or as showing unusual solvent effects.

A word should be said about the decision to cut the recommended ϵ 's to compounds with $\pm 30\%$ confidence intervals. First of all, $\pm 30\%$ is roughly on the most pessimistic limits of reported "error" limits by the authors of papers reporting extinction coefficients. In fact rarely are error bars stated as being more than $\pm 25\%$. However after spending much time evaluating the data, we tend to favor the more pessimistic estimates. In addition the average ϵ 's with confidence limits of less than $\pm 30\%$ tend to fall close to or between the Bensasson-Land values in the last two columns of Table 5.

As mentioned above the recommended hierarchy of values is only tentative and is contingent on further work. It was hoped that further evaluation could be done using the principle that oscillator strengths do not change with solvent. Although this approach is promising, it is difficult to carry out on a large scale because one must have complete spectra in each solvent. Another approach that seems more feasible is a global fit¹⁰¹ using the recommended values obtained to correct all the ϵ 's obtained from relative measurements. Using a consistent set of values as standards for the relative measurements should in itself be an improvement over the present analysis.

5. Arrangement of the Data Table

Table 6, the data table, contains over 3000 entries. These are grouped under the name of the compound to which they pertain. The compounds themselves are numbered and listed alphabetically, with secondary attention being paid to numerical substituent locations. Common names are employed and we rely on the use of a compound name index, which contains both synonyms and inverted names, and a molecular formula index to help locate specific data.

Multiple entries which occur for a particular compound are further ordered alphabetically by the solvent

name or abbreviation. Final questions of tabular precedence are settled by appeal to the serial number accorded to each paper upon acquisition for our local bibliographic database.¹¹

In addition to specifying the ambient medium, the "solvent" column reports, in parentheses, the temperature at which the measurement was carried out. If no temperature is shown then room temperature may be assumed unless otherwise noted in the comments.

The slash (/) in the "method" column separates the excitation method from the extinction determination technique. An explanation of the abbreviations used for each is given below. Occasionally the only reported data has been obtained as an average of unquoted primary values from different methods. In this case both methods are included and separated by an ampersand.

The third column contains the heart of the compilation: the wavelength(s), λ_{max} of maximum triplet-triplet absorbance together with the corresponding extinction coefficient(s), ϵ_{max} , where estimated. Multiple entries are presented in order of decreasing wavelength (i.e. in increasing energy). We highlight the following use of two superscripts:

^a implies that data have been obtained by computer-assisted digitization¹¹⁰ from a spectrum in the cited reference,

^b implies that the (wavelength, extinction coefficient) pair are not necessarily related to a spectral peak.

Other superscripts are used as indicated in the footnotes.

The fourth column, the "comments" column, collects information from a number of separate items in our database. In the table this construction is evidenced by the presence of semicolons (;) delimitating the items. Primary comments comprise an identification technique if reported. This is indicated by a preceding ‡. If this symbol is absent then no attempt to verify the nature of the transient, other than by similarity to a previous measurement, is mentioned in the article from which the data have been taken. Other comments, such as relative intensities in multiply peaked spectra, if stated or derived from a figure in the original work are included next. When a comparative method has been used to determine the extinction coefficient, the data relative to which the estimation has been made (if reported) are present in a set form giving both wavelength and extinction coefficient of the standard employed, as well as the solvent in which these values were obtained. Details of any additional assumptions, such as triplet yields in the relative actinometry technique, which were made by the original authors to enable the comparison, are also included. Further specification of, for example, the solvent system, e.g. ratio of mixture, etc. are then given together with any other information relevant to the evaluation of the data in the wavelength-extinction column.

The remainder of the comment is constructed in a fixed format from entries for triplet state lifetimes (μs), pH of solution, half-life of transient (if second order processes

are present and unsubtracted), oscillator strength of reported transition, rate constant for energy transfer to (or from) the species indicated earlier in the comment and rise-time of the triplet-triplet absorption. We note here again that these associated data, while relevant to the study and characterization of triplet states in general, have not been subjected to a procedure of critical evaluation.

The last column contains the serial number which may be used to locate the reference in the list provided subsequent to the table.

5.1. Lists of Abbreviations

We have made use of the following abbreviations in the body of the Table. These are mainly confined to the "Method" column though some do appear in the "Comments" column.

- a. Triplet State Population Method
- | | |
|-----|---|
| CWL | Continuous wave laser excitation |
| FP | Flash photolysis |
| LP | Laser photolysis (pulsed) |
| MOD | Modulated excitation (e.g. rotating sector) |
| PR | Pulse radiolysis |
| PS | Photostationary |
| -ET | (coupled with energy transfer) |
- b. Extinction Coefficient Estimation Method
- | | |
|-----|-----------------------------------|
| COM | See comments |
| ELT | Electron transfer |
| ESR | Electron spin resonance intensity |
| ET | Energy transfer |
| HAT | Hydrogen atom transfer |
| IV | Intensity variation |
| KM | Kinetic method |
| PIV | Pulsed intensity variation |
| RA | Relative actinometry |
| RF | Constant oscillator strength |
| SD | Singlet depletion |
| SM | Partial saturation |
| TD | Total depletion |

In addition, to conserve space, a number of solvents and mixtures have been abbreviated. These are mainly confined to the "Solvent" column and are listed below.

- | | |
|-------------------|---|
| AOT | Di-(2-ethylhexyl) sodium sulfosuccinate |
| BuOH | Butanol (isomer unspecified) |
| 1-BuOH | 1-Butanol |
| <i>tert</i> -BuOH | <i>tert</i> -Butyl alcohol |
| DFMeOH | Di(trifluoromethyl)methanol |
| CTAB | Cetyltrimethylammonium bromide |
| CTAC | Cetyltrimethylammonium chloride |
| DDDAB | Didodecyldimethylammonium bromide |
| DMF | Dimethyl formamide |
| DMSO | Dimethyl sulfoxide |
| DODAC | Diocetadecyldimethylammonium chloride |
| DTB | Dodecyltrimethylammonium bromide |
| EPA | Diethyl ether/Isopentane/Absolute ethanol (5:5:2) |
| EtOH | Ethanol |

- | | |
|-------------------|--|
| Et ₂ O | Diethyl ether |
| MCH | Methylcyclohexane |
| MeOH | Methanol |
| 3-MH | 3-Methylhexane |
| 3-MP | 2-Methylpentane |
| MTHF | Methyltetrahydrofuran (isomer unspecified) |
| 2-MTHF | 2-Methyltetrahydrofuran |
| PFMCH | Perfluoromethylcyclohexane |
| PMMA | Poly(methylmethacrylate) |
| PPFO | Potassium perfluorooctylsulfonate |
| PrOH | Propanol (isomer unspecified) |
| 1-PrOH | 1-Propanol |
| 2-PrOH | 2-Propanol |
| PVA | Poly(vinyl alcohol) |
| SDS | Sodium dodecyl sulfate surfactant |
| SHS | Sodium hexadecyl sulfate surfactant |
| SPFO | Sodium perfluorooctanoate |
| THF | Tetrahydrofuran |

5.2. Lists of Symbols

The following symbols, used in the body of the text are collected here for convenience.

- a. Conversion probabilities
- | | |
|-------------|---|
| Φ | total luminescence yield |
| Φ_f | quantum yield of fluorescence |
| Φ_T | quantum yield for triplet state formation |
| $\Phi_T(A)$ | quantum yield for triplet state formation in triplet energy acceptor, A |
| $\Phi_T(D)$ | quantum yield for triplet state formation in triplet energy donor, D |
| $\Phi_T(R)$ | quantum yield for triplet state formation in reference compound, R |
| $\Phi_T(T)$ | quantum yield for triplet state formation in unknown compound, T |
| P_{ir} | probability of triplet-triplet energy transfer |
- b. Extinction coefficients
- | | |
|----------------|--|
| ϵ_A | extinction coefficient for ground (singlet) state of energy acceptor, A |
| ϵ_A^* | extinction coefficient for excited (triplet) state of A |
| ϵ_D | extinction coefficient for ground (singlet) state of energy donor, D |
| ϵ_D^* | extinction coefficient for excited (triplet) state of D |
| ϵ_R^* | extinction coefficient for excited (triplet) state of reference triplet, R |
| ϵ_S | extinction coefficient for ground singlet state, S, of arbitrary molecule |
| ϵ_T^* | extinction coefficient for excited (triplet) state of unknown triplet, T |
- c. Intensities
- | | |
|----------|---|
| $I_p(R)$ | intensity of sensitized phosphorescence from reference compound |
| $I_p(T)$ | intensity of sensitized phosphorescence from unknown compound |
| I | true transmitted light intensity |

I_0	incident light intensity
I_m	intensity of monitoring light
I_m^0	incident intensity of monitoring light
I_p	intensity of photolyzing light
I_p^0	incident intensity of photolyzing light
I_s	scattered light intensity in absence of excitation
I_s'	scattered light intensity in presence of excitation

d. Miscellaneous

A_r	integrated microwave absorbance of reference radical
A_T	integrated microwave absorbance of triplet state
ℓ	optical path length (of monitoring beam)
x	measure of distance on along monitoring path
τ_T	triplet state lifetime
τ_r	(excitation rate dependent) rise time of triplet-triplet absorption
$[M]$	concentration of M
$[M]_0$	concentration of M at time zero
$[M]_\infty$	concentration of M at long times
$[M]_{x=0}$	concentration of M at front face of cell
$[M]_{x=\ell}$	concentration of M at rear of cell

e. Molecular species

1A	singlet ground state of energy acceptor, A
$^1A^*$	singlet excited state of A
$^2A^-$	radical anion of A
$^3A^*$	triplet excited state of A
1D	singlet ground state of energy donor, D
$^1D^*$	singlet excited state of D
$^3D^*$	triplet excited state of D
1M	singlet ground state of arbitrary molecule, M
$^1M^*$	singlet excited state of M
$^3M^*$	triplet excited state of M
$^3M^{**}$	higher triplet excited state of M, reached by TTA
1S	singlet ground state of solvent molecule, S
$^1S^*$	singlet excited state of S
$^2S^+$	radical cation of S
$^2S^-$	radical anion of S
$^3S^*$	triplet excited state of S

f. Molecular states

S_0	molecular (singlet) ground state
S_1	lowest excited singlet state
S_n	higher excited singlet state
T_1	lowest triplet (excited) state
T_n	higher excited triplet state

g. Optical densities

OD	optical density
OD _T	OD ascribed to triplet state absorbance
ΔOD	change in OD on excitation
ΔOD_A	change in OD of acceptor (triplet) extrapolated to long times

ΔOD_D	change in OD of donor (triplet) extrapolated to time zero
ΔOD_S	change in OD due to isolated singlet decay
ΔOD_T	change in OD due to isolated triplet growth

h. Rate constants

k_A	rate constant for acceptor triplet state deactivation
k_D	rate constant for donor triplet state deactivation
k_{DA}	rate constant for ET without acceptor excitation
k_{et}	rate constant for triplet energy transfer
k_{ex}	rate constant for ground state photo-excitation
k'_{ex}	rate constant for excited state photo-excitation
k_{ic}	rate constant for internal conversion in the singlet manifold
k_{isc}	rate constant for singlet to triplet inter-system crossing
k'_{isc}	rate constant for triplet to singlet inter-system crossing
k''_{isc}	rate constant for T_1 to S_1 intersystem crossing
k_p	rate constant for phosphorescence
k_{pc}	rate constant for photochemistry from S_1
k'_{pc}	rate constant for photochemistry from T_1
k_S	rate constant for singlet state deactivation
k_T	rate constant for triplet state deactivation

6. Acknowledgments

The Radiation Chemistry Data Center is supported jointly by the National Bureau of Standards, Office of Standard Reference Data and by the Office of Basic Energy Sciences of the Department of Energy. This is Radiation Laboratory Document No. NDRL-2690.

The authors thank Dr. W. P. Helman for the initial development of the software required to construct the data table from our numerical database and to derive the indexes. Dr. A. B. Ross is also thanked for her continuing interest in and support of this work. The authors would finally like to thank Drs. R. F. Ganey and S. Arndt for several instructive conversations on statistical analysis of data.

7. References to Text

- ¹H. Labhart and W. Heinzelmann, "Triplet-Triplet Absorption Spectra of Organic Molecules," in *Organic Molecular Photophysics*, Vol. 1, J. B. Birks, Editor (Wiley, New York, 1973) p. 297.
- ²J. B. Birks, *Photophysics of Aromatic Molecules* (Wiley-Interscience, London, 1970).
- ³A. K. Chibisov, *Russ. Chem. Rev.* **39**, 891 (1970).
- ⁴R. Bensasson and E. J. Land, "Physical Properties of Excited States: A General Method for Measuring Triplet-Triplet Extinction Coefficients, Singlet-Triplet Intersystem Crossing Efficiencies, and Related Parameters," in *Photochemical and Photobiological Reviews*, Vol. 3, K. C. Smith, Editor (Plenum Press, New York, 1978) p. 163.
- ⁵E. J. Land, *Proc. R. Soc. London, Ser. A* **305**, 457 (1968).
- ⁶W. Heinzelmann and H. Labhart, *Chem. Phys. Lett.* **4**, 20 (1969).

- ⁷J. C. Scaiano, *J. Photochem.* **18**, 395 (1982).
- ⁸I. Carmichael and G. L. Hug, *Radiat. Phys. Chem.* **20**, 119 (1982).
- ⁹I. Carmichael and G. L. Hug, *Radiat. Phys. Chem.* **20**, 179 (1982).
- ¹⁰I. Carmichael and G. L. Hug, *Radiat. Phys. Chem.* **26**, 229 (1985).
- ¹¹W. P. Helman and A. B. Ross, *Radiat. Phys. Chem.* **16**, 425 (1980).
- ¹²M. Kasha, *Discuss. Faraday Soc.* (9), 14 (1950).
- ¹³J. B. Birks, *Chem. Phys. Lett.* **17**, 370 (1972).
- ¹⁴G. D. Gillespie and E. C. Lim, *J. Chem. Phys.* **65**, 2022 (1976).
- ¹⁵G. N. Lewis and M. Kasha, *J. Am. Chem. Soc.* **66**, 2100 (1944).
- ¹⁶G. N. Lewis, D. Lipkin, and T. T. Magel, *J. Am. Chem. Soc.* **63**, 3005 (1941).
- ¹⁷D. S. McClure, *J. Chem. Phys.* **19**, 670 (1951).
- ¹⁸G. Porter and M. W. Windsor, *J. Chem. Phys.* **21**, 2088 (1953).
- ¹⁹R. G. W. Norrish and G. Porter, *Nature* **164**, 658 (1949).
- ²⁰G. Porter, *Proc. R. Soc. London, Ser. A* **200**, 284 (1950).
- ²¹G. Porter and M. W. Windsor, *Discuss. Faraday Soc.* (17), 178 (1954).
- ²²M. S. Matheson and L. M. Dorfman, *J. Chem. Phys.* **32**, 1870 (1960).
- ²³R. L. McCarthy and A. MacLachlan, *Trans. Faraday Soc.* **56**, 1187 (1960).
- ²⁴J. P. Keene, *Nature* **188**, 843 (1960).
- ²⁵J. W. Boag and R. E. Steel, *British Empire Cancer Campaign Report 38, Part II*, 1960, p. 251.
- ²⁶F. S. Dainton, T. J. Kemp, G. A. Salmon, and J. P. Keene, *Nature* **203**, 1050 (1964).
- ²⁷R. M. Danziger, K. H. Bar-Eli, and K. Weiss, *J. Phys. Chem.* **71**, 2633 (1967).
- ²⁸R. McNeil, J. T. Richards, and J. K. Thomas, *J. Phys. Chem.* **74**, 2290 (1970).
- ²⁹G. Porter and M. R. Topp, *Proc. R. Soc. London, Ser. A* **315**, 163 (1970).
- ³⁰P. M. Rentzepis, *Science* **169**, 239 (1970).
- ³¹N. Mataga, and N. Nakashima, *Spectrosc. Lett.* **8**, 275 (1975).
- ³²R. M. Hochstrasser, D. L. Narva, and A. C. Nelson, *Chem. Phys. Lett.* **43**, 15 (1976).
- ³³R. V. Bensasson, J. T. Richards, and J. K. Thomas, *Chem. Phys. Lett.* **9**, 13 (1971).
- ³⁴C. Willis, O. A. Miller, A. E. Rothwell, and A. W. Boyd, *Radiat. Res.* **35**, 428 (1968).
- ³⁵D. C. Walker and S. C. Wallace, *Chem. Phys. Lett.* **6**, 111 (1970).
- ³⁶J. W. Hunt, C. L. Greenstock, and M. J. Bronskill, *Int. J. Radiat. Phys. Chem.* **4**, 87 (1972).
- ³⁷J. E. Aldrich, P. Foldvary, J. W. Hunt, W. B. Taylor, and R. K. Wolff, *Rev. Sci. Instrum.* **43**, 991 (1972).
- ³⁸L. M. Dorfman, "Pulse Radiolysis," in *Techniques in Chemistry*, 3rd Edition, Part II, Vol. 6, G. G. Hammes, Editor (Wiley-Interscience, New York, 1974) p. 463.
- ³⁹J. W. Hunt, "Early Events in Radiation Chemistry," in *Advances in Radiation Chemistry*, Vol. 5, M. Burton and J. L. Magee, Editors, (Wiley-Interscience, New York, 1976) p. 185.
- ⁴⁰V. Sundstrom, P. M. Rentzepis, and E. C. Lim, *J. Chem. Phys.* **66**, 4287 (1977).
- ⁴¹M. Chessin, R. Livingston, and T. G. Truscott, *Trans. Faraday Soc.* **62**, 1519 (1966).
- ⁴²J. T. Dubois and M. Cox, *J. Chem. Phys.* **38**, 2536 (1963).
- ⁴³H. L. J. Baekstroem and K. Sandros, *Acta Chem. Scand.* **12**, 823 (1958).
- ⁴⁴G. Porter and F. Wilkinson, *Trans. Faraday Soc.* **57**, 1686 (1961).
- ⁴⁵H. Labhart, *Helv. Chim. Acta* **47**, 2279 (1964).
- ⁴⁶U. B. Ranalder, H. Kaenzig, and U. P. Wild, *J. Photochem.* **4**, 97 (1975).
- ⁴⁷M. V. Alfimov, I. G. Batekha, Y. B. Shek, and V. I. Gerko, *Spectrochim. Acta* **27**, 329 (1971).
- ⁴⁸D. Lavalette, *C. R. Seances Acad. Sci., Ser. B* **266**, 279 (1968).
- ⁴⁹T. F. Hunter and A. I. Younis, *J. Chem. Soc., Faraday Trans. 1* **75**, 550 (1979).
- ⁵⁰C. A. Parker, *Photoluminescence of Solutions* (Elsevier, Amsterdam, 1968).
- ⁵¹J. W. Boag, *Trans. Faraday Society* **64**, 677 (1968).
- ⁵²I. G. Ross, *J. Opt. Soc. Am.* **44**, 40 (1954).
- ⁵³M. Bazin and T. W. Ebbesen, *Photochem. Photobiol.* **37**, 675 (1983).
- ⁵⁴G. Porter and M. A. West, "Flash Photolysis" in *Techniques in Chemistry*, 3rd Edition, Part II, Vol. 6, G. G. Hammes, Editor (Wiley-Interscience, New York, 1974) p. 367.
- ⁵⁵M. A. West, *Photochemistry (Specialist Periodical Reports, Chem. Soc., London)* **8**, 3 (1977).
- ⁵⁶G. Cilento, *Acc. Chem. Res.* **13**, 225 (1980).
- ⁵⁷N. J. Turro, P. Lechtken, G. Schuster, J. Orell, H. C. Steinmetzer, and W. Adam, *J. Am. Chem. Soc.* **96**, 1627 (1974).
- ⁵⁸J. T. Richards and J. K. Thomas, *Trans. Faraday Soc.* **66**, 621 (1970).
- ⁵⁹D. L. Dexter, *J. Chem. Phys.* **21**, 836 (1953).
- ⁶⁰C. Capellos and B. H. J. Bielski, *Kinetic Systems* (Krieger, Huntington, N.Y., 1980).
- ⁶¹R. Bensasson and E. J. Land, *Trans. Faraday Soc.* **67**, 1904 (1971).
- ⁶²T. Takemura, P. K. Das, G. Hug, and R. S. Becker, *J. Am. Chem. Soc.* **100**, 2626 (1978).
- ⁶³W. R. Dawson, *J. Opt. Soc. Am.* **58**, 222 (1968).
- ⁶⁴P. G. Bowers and G. Porter, *Proc. R. Soc. London, Ser. A* **299**, 348 (1967).
- ⁶⁵S. G. Hadley and R. A. Keller, *J. Phys. Chem.* **73**, 4351 (1969).
- ⁶⁶T. G. Pavlopoulos, *J. Opt. Soc. Am.* **63**, 180 (1973).
- ⁶⁷R. Bensasson, C. R. Goldschmidt, E. J. Land, and T. G. Truscott, *Photochem. Photobiol.* **28**, 277 (1978).
- ⁶⁸H. Linschitz and K. Sarkanen, *J. Am. Chem. Soc.* **80**, 4826 (1958).
- ⁶⁹I. Carmichael and G. L. Hug, *J. Phys. Chem.* **89**, 4036 (1985).
- ⁷⁰M. R. Topp, *Appl. Spectrosc. Rev.* **14**, 1 (1978).
- ⁷¹U. Lachish, A. Shafferman, and G. Stein, *J. Chem. Phys.* **64**, 4205 (1976).
- ⁷²M. M. Fisher, B. Veyret, and K. Weiss, *Chem. Phys. Lett.* **28**, 60 (1974).
- ⁷³S. Speiser, R. van der Werf, J. Kommandeur, *Chem. Phys.* **1**, 297 (1973).
- ⁷⁴H. Lutz, E. Breheret, and L. Lindqvist, *J. Phys. Chem.* **77**, 1758 (1973).
- ⁷⁵A. Corval, P. Jardon, and R. Gautron, *J. Chim. Phys.-Chim. Biol.* **79**, 335 (1982).
- ⁷⁶B. Amand and R. Bensasson, *Chem. Phys. Lett.* **34**, 44 (1975).
- ⁷⁷R. A. Keller and S. G. Hadley, *J. Chem. Phys.* **42**, 2382 (1965).
- ⁷⁸W. G. Herkstroeter and D. S. McClure, *J. Am. Chem. Soc.* **90**, 4522 (1968).
- ⁷⁹F. Wilkinson, "Triplet Quantum Yields and Singlet-Triplet Inter-system Crossing," in *Organic Molecular Photophysics*, Vol. 2, J. B. Birks, Editor (Wiley, New York, 1975) p. 95.
- ⁸⁰V. A. Smirnov and M. V. Alfimov, *Kinetica Kataliz.* **7**, 583 (1966).
- ⁸¹D. Lavalette, *J. Chim. Phys.* **66**, 1853 (1969).
- ⁸²G. L. Hug and I. Carmichael, *J. Photochem.* **31**, 179 (1985).
- ⁸³M. S. Grodowski, B. Veyret, and K. Weiss, *Photochem. Photobiol.* **26**, 341 (1977).
- ⁸⁴B. Veyret, S. G. Davis, M. Yoshida, and K. Weiss, *J. Am. Chem. Soc.* **100**, 3283 (1978).
- ⁸⁵C. A. Hutchinson, Jr., and B. W. Magnum, *J. Chem. Phys.* **29**, 952 (1958).
- ⁸⁶J. S. Brinen, *J. Chem. Phys.* **49**, 586 (1968).
- ⁸⁷J. S. Brinen and M. K. Orloff, *J. Chem. Phys.* **51**, 527 (1969).
- ⁸⁸M. V. Alfimov, N. Y. Buben, A. I. Pristupa, and V. N. Shamshev, *Opt. Spectrosc.* **20**, 232 (1966).
- ⁸⁹I. V. Aleksandrov and K. K. Pukhov, *Opt. Spektrosk.* **17**, 944 (1964).
- ⁹⁰R. Kubo and K. Tomita, *J. Phys. Soc. Jpn.*, **9**, 888 (1954).
- ⁹¹T. J. Kemp and L. J. A. Martins, *J. Chem. Soc., Faraday Trans. 1* **77**, 1425 (1981).
- ⁹²R. V. Bensasson and J.-C. Gramain, *J. Chem. Soc., Faraday Trans. 1* **76**, 1801 (1980).
- ⁹³D. Lavalette, R. Bensasson, B. Amand, and E. J. Land, *Chem. Phys. Lett.* **10**, 331 (1971).
- ⁹⁴L. J. Andrews, A. Derouledé, and H. Linschitz, *J. Phys. Chem.* **82**, 2304 (1978).
- ⁹⁵J. K. Hurley, N. Sinai, and H. Linschitz, *Photochem. Photobiol.* **38**, 9 (1983).
- ⁹⁶A. A. Lamola and M. S. Wrighton, *Pure Appl. Chem.* **54**, 1251 (1982).
- ⁹⁷A. A. Lamola and M. S. Wrighton, *Pure Appl. Chem.* **56**, 939 (1984).
- ⁹⁸B. Stevens and B. E. Algar, *Ann. N.Y. Acad. Sci.* **171**, 50 (1970).
- ⁹⁹R. G. Bennett, R. P. Schwenker, and R. E. Kellogg, *J. Chem. Phys.* **41**, 3040 (1964).

- ¹⁰⁰V. L. Ermolaev and E. B. Sveshnikova, *Opt. Spectrosc.* **28**, 324 (1970).
- ¹⁰¹F. Wilkinson and J. G. Brummer, *J. Phys. Chem. Ref. Data* **10**, 809 (1981).
- ¹⁰²B. Ostle and R. W. Mensing, *Statistics in Research*, 3rd. edition (The Iowa State Univ. Press, Ames, 1975).
- ¹⁰³*SAS User's Guide: Statistics*, 1982 edition (SAS Institute, Cary, NC, 1982).
- ¹⁰⁴M. H. DeGroot, *Probability and Statistics* (Addison-Wesley, Reading, MA, 1975).
- ¹⁰⁵A. G. Worthing and J. Geffner, *Treatment of Experimental Data* (Wiley, New York, 1943).
- ¹⁰⁶W. Rhodes and M. Chase, *Rev. Modern Physics* **39**, 348 (1967).
- ¹⁰⁷G. Porter and M. W. Windsor, *Proc. R. Soc. London, Ser. A* **245**, 238 (1958).
- ¹⁰⁸H. Hamaguchi, H. Okamoto, and M. Tasumi, *Chem. Lett.* (4), 549 (1984).
- ¹⁰⁹R. Bensasson and E. J. Land, *Nouv. J. Chim.* **2**, 503 (1978).
- ¹¹⁰G. L. Hug, NSRDS-NBS 69 (National Bureau of Standards, Washington, DC, 1981).

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
1. Acenaphthene						
	Benzene	LP-ET	422		†Triplet ET from acetophenone	82A292
	EtOH (293 K)	FP	433			68E098
	EtOH (77 K)	MOD/KM	430, 6000 ± 1500			737055
2. Acenaphthylene, <i>cis</i> -photodimer						
	Benzene (293 K)	LP-ET	455		†Oxygen quenching and triplet ET from benzo-phenone and acetophenone; $\tau_T = 2.2 \mu\text{s}$	82A292
3. Acenaphthylene, <i>trans</i> -photodimer						
	Benzene (293 K)	LP-ET	455		†Triplet ET from acetophenone; $\tau_T = 0.12 \mu\text{s}$	82A292
4. 1-Acenaphthyl-1-phenylethylene						
	Benzene	LP-ET	502 ^a 424 ^a		†Triplet ET from xanthone and oxygen quenching; $\tau_T = 0.087 \mu\text{s}$	84B007
5. 1'-Acetonaphthone						
	Water/ <i>tert</i> -BuOH	FP	510 ^a		Solvent mixture contains "1-5%" <i>tert</i> -BuOH for solubility; very broad diffuse band; pH ~6	767189
6. 2'-Acetonaphthone						
	Benzene	PR/ET	430, 10500		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700 \text{ L mol}^{-1} \text{ cm}^{-1}$); ϵ obtained from a simultaneous least squares fit of data from several compounds making use of cyclohexane to benzene ϵ_{\max} ratios of 1.83 for naphthalene and 1.45 for anthracene	71E360
	Heptane	LP	424 404 384		Most intense peak at 424 nm	78A268
	Water/ <i>tert</i> -BuOH	FP	440 ^a		Solvent mixture contains "1-5%" <i>tert</i> -BuOH for solubility; shoulder at 420 nm; pH ~6	767189
7. Acetone						
	Acetonitrile (296 K)	FP	~302		†Phosphorescence decay; $\tau_T = 47 \pm 6 \mu\text{s}$	717489
	Liquid paraffin	FP	324		Solvent viscosity was 0.19 N·s/m ²	58E001
	Methylene chloride	LP	240		†Phosphorescence decay and oxygen quenching; $\tau_T = 6.3 \mu\text{s}$	84B051
	Water	LP/ET	300 ^b , 600 ± 100		ϵ relative to sodium 1,5-naphthalenedisulfonate in water ($\epsilon_{445} = 9900 \text{ L mol}^{-1} \text{ cm}^{-1}$); kinetic correction made in ET measurement; $\tau_T = 5.0 \mu\text{s}$; $k_{et} = 3.6 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	82B045
8. Acetophenone						
	? (81 K)	FP/SD	330, 1300		†Phosphorescence decay; glass was either 1:3:3 1-methylPrOH to Et ₂ O to isooctane or 1:9 MCH to mixture of EtOH and MeOH; $\tau_T = (1 \pm 0.1) \times 10^4 \mu\text{s}$	77B022
	Acetonitrile	LP/RA	320 ^a , 12600 ^a		ϵ relative to benzophenone in cyclohexane ($\epsilon_{533} = 7630 \text{ L mol}^{-1} \text{ cm}^{-1}$, assuming no solvent effect from benzene), and taking all the triplet yields to be unity	737198
	Cyclohexane	LP/ET	449 ^a , 1800 ^a 406 ^a , 2100 ^a		†Oxygen quenching, triplet ET to naphthalene, and quenching by piperylene; ϵ relative to naphthalene in cyclohexane ($\epsilon_{412.5} = 22600 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 0.23 \mu\text{s}$	717179
	Cyclohexane	LP/RA	441 ^a , 1900 ^a 401 ^a , 2100 ^a		ϵ relative to benzophenone in cyclohexane ($\epsilon_{533} = 7630 \text{ L mol}^{-1} \text{ cm}^{-1}$, assuming no solvent effect from benzene), and taking all the triplet yields to be unity; there was a more intense maximum <300 nm	737198

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	EtOH	LP/ET&RA	323 ^a , 6300 ^a		ϵ relative to naphthalene in cyclohexane ($\epsilon_{412.5} = 22600$ L mol ⁻¹ cm ⁻¹), assuming no solvent effects, for ET method; ϵ relative to benzophenone in benzene ($\epsilon_{532.5} = 10300$ L mol ⁻¹ cm ⁻¹), assuming no solvent effects and taking $\Phi_T(\text{acetophenone}) = 1 = \Phi_T(\text{benzophenone})$, and ϵ relative to benz[a]anthracene in cyclohexane ($\epsilon_{480} = 25100$ L mol ⁻¹ cm ⁻¹), assuming no solvent effects and taking $\Phi_T(\text{acetophenone}) = 1$ and $\Phi_T(\text{benz[a]anthracene}) = 0.82$, for RA method; there was a shoulder at $\sim 495^\circ$ nm; $\tau_T = 0.14$ μ s	717179
	Liquid paraffin	FP	281.5 265		Solvent viscosity was 0.03 N·s/m ²	58E001
	Water	FP	335		$\tau_T = \sim 100$ μ s	727098
	Water	LP/RA	337 ^a , 13100 ^a		ϵ relative to benzophenone in cyclohexane ($\epsilon_{533} = 7630$ L mol ⁻¹ cm ⁻¹), assuming no solvent effect from benzene), and taking all the triplet yields to be unity	737198
9.	Acetophenone, conjugate acid Sulfuric acid	LP	359 ^a		5 mol L ⁻¹ H ₂ SO ₄ ; solvent was 4:1 water to acetonitrile; pK_a 0.63 \pm 0.07	84E456
10.	2-Acetoxy-2-methyl-1-phenyl-1-propanone Benzene	LP	440 410 330		\dagger Phosphorescence decay, oxygen quenching (1.6×10^9 L mol ⁻¹ s ⁻¹); relative intensities (3:3:10); $\tau_T = 1.9 \pm 0.2$ μ s	80E642
	Cyclohexane	LP	440 410 330		\dagger Phosphorescence decay, oxygen quenching; relative intensities (3:3:10); $\tau_T = 1.9 \pm 0.2$ μ s	80E642
11.	Acetylacetone EtOH/MeOH (118 K)	FP	480		Solvent was 3:1 EtOH to MeOH	68D005
12.	1-Acetylanthracene EtOH (93 K)	PS	500			66B001
13.	9-Acetylanthracene ? (81 K)	FP/SD	426, 20000		\dagger Phosphorescence decay; glass was either 1:3:3 1-methylPrOH to Et ₂ O to isooctane or 1:9 MCH to mixture of EtOH and MeOH; $\tau_T = (2.8 \pm 0.3) \times 10^4$ μ s	77B022
	Acetonitrile	LP	421 ^a		Delay 320 ps	84B154
	EPA (77 K)	FP	429		$\tau_T = 2.95 \times 10^4$ μ s	82E338
	Toluene	LP	410		\dagger Oxygen quenching; rise time of 0.02 ± 0.004 ns	777635
14.	4-Acetylbiphenyl ? (81 K)	FP/SD	435, 130000		\dagger Phosphorescence decay; glass was either 1:3:3 1-methylPrOH to Et ₂ O to isooctane or 1:9 MCH to mixture of EtOH and MeOH; $\tau_T = (1.7 \pm 0.2) \times 10^4$ μ s	77B022
15.	3-Acetyl-9,10-epoxy-9,10-dihydrophenanthrene 1,2-Dichloroethane	I.P	395		\dagger Oxygen quenching	79A177
16.	N-(2-Acetylphenyl)acetamide EtOH	LP	430		\dagger Oxygen quenching (5.6×10^8 L mol ⁻¹ s ⁻¹); $\tau_T = 10$ μ s	78E308
	Hexane	LP/TD	450, 8900		$\tau_T = 8.5$ μ s; oscillator strength = 0.18	78E308
	Water	LP	430		\dagger Oxygen quenching (2.2×10^9 L mol ⁻¹ s ⁻¹); ionic strength 0.01 mol L ⁻¹ ; $\tau_T = 6.5$ μ s; pH 7.5	78E308
17.	N-(2-Acetylphenyl)formamide EtOH	LP/RF	450, 9000		\dagger Oxygen quenching (2.2×10^9 L mol ⁻¹ s ⁻¹); ϵ relative to N-(2-acetylphenyl)formamide in hexane ($\epsilon_{450} = 8100$ L mol ⁻¹ cm ⁻¹) assuming oscillator strength independent of solvent; $\tau_T = 10$ μ s	78E308

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Hexane	LP/TD	450, 6500			78E308
	Hexane	PR-ET/RA	450, 8100		†Triplet ET from benzene; ϵ relative to naphthalene in hexane ($\epsilon_{\max} = 24500$ L mol ⁻¹ cm ⁻¹) assuming same ET rate from benzene; $\tau_T = 7.7$ μ s; oscillator strength = 0.18	78E308
	Water	LP	450		†Oxygen quenching (3×10^9 L mol ⁻¹ s ⁻¹); ionic strength 0.01 mol L ⁻¹ ; $\tau_T = 4.1$ μ s; pH 7.5	78E308
	Water	LP	450		$\tau_T = 7$ μ s; pH 6.5	78E308
18.	<i>N</i> -(2-Acetylphenyl)- <i>N</i> -methylacetamide					
	EtOH	LP/RA	430, <1100		ϵ relative to naphthalene in hexane ($\Phi_T = 0.8$, $\epsilon_{\max} = 24500$ L mol ⁻¹ cm ⁻¹) taking $\Phi_T = 0.6$ for <i>N</i> -(2-Acetylphenyl)- <i>N</i> -methylacetamide in EtOH; $\tau_T = 0.2$ μ s	78E308
	Hexane	LP	420		$\tau_T = 0.2$ μ s	78E308
	Hexane	PR-ET/RA	420, 1200		†Triplet ET from benzene; ϵ relative to naphthalene in hexane ($\epsilon_{\max} = 24500$ L mol ⁻¹ cm ⁻¹) assuming same ET rate from benzene; $\tau_T = 0.2$ μ s; oscillator strength = 0.02	78E308
	Water	LP	430		†Oxygen quenching (6×10^8 L mol ⁻¹ s ⁻¹); ionic strength 0.01 mol L ⁻¹ ; $\tau_T = 7$ μ s; pH 7.5	78E308
19.	<i>N</i> -(2-Acetylphenyl)- <i>N</i> -methylformamide					
	1-ProH	LP	445		$\tau_T = 0.3$ μ s	78E308
	EtOH	LP/RA	445, <1100		ϵ relative to naphthalene in hexane ($\Phi_T = 0.8$, $\epsilon_{\max} = 24500$ L mol ⁻¹ cm ⁻¹) taking $\Phi_T = 0.6$ for <i>N</i> -(2-Acetylphenyl)- <i>N</i> -methylformamide in EtOH; $\tau_T = 0.6$ μ s	78E308
	Hexane	LP	420		$\tau_T = 0.2$ μ s	78E308
	Hexane	PR-ET/RA	420, 850		†Triplet ET from benzene; ϵ relative to naphthalene in hexane ($\epsilon_{\max} = 24500$ L mol ⁻¹ cm ⁻¹) assuming same ET rate from benzene; $\tau_T = 0.2$ μ s; oscillator strength = 0.02	78E308
	Water	LP	445		†Oxygen quenching (5.2×10^8 L mol ⁻¹ s ⁻¹); ionic strength 0.01 mol L ⁻¹ ; $\tau_T = 8.8$ μ s; pH 7.5	78E308
20.	3-Acetyltriptycene					
	Cyclohexane (289.5 K)	LP	550			83E483
21.	<i>N</i> -Acetyl-L-tryptophanamide					
	Water	LP	460		Buffered; pH 7.0	81A232
22.	Acridan					
	EtOH (93 K)	PS	520		Shoulder at 580 nm	69E214
23.	Acridine					
	2-ProH	LP	983 \pm 5			83E392
			440 \pm 1			
	Acetonitrile	LP	978 \pm 2			83E392
			434 \pm 1			
	Argon (13 K)	CWL	425.1		Infrared vibrational structure also observed	79B043
	Benzene	FP/TD	520*, 2000*		Uncertain correction of estimated peaks for SD; $\tau_T = 10^6$ μ s	677498
			440, 25000			
			435, 12000*			
			375*, 3000*			
			355*, 10000*			
			340*, 7000*			
	Benzene	FP/SD	440, 80000*		Delay 93 μ s	677259
	Benzene	PR/ET	440, 24300		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700$ L mol ⁻¹ cm ⁻¹); ϵ obtained from a simultaneous least squares fit of data from several compounds making use of cyclohexane to benzene ϵ_{\max} ratios of 1.83 for naphthalene and 1.45 for anthracene	71E360

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Benzene	FP/TD	442, 27000			776258
	Benzene (296 K)	LP	990		Shoulder at 415 nm	81E147
			870 ^a			
			781 ^a			
			699 ^a			
			518 ^a			
			440			
	Benzene	LP	980 ± 2			83E392
			442 ± 1			
	CCl ₄	LP	981 ± 2			83E392
			442 ± 1			
	Cyclohexane	PR/ET	432.5, 26300		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361.3} = 35400$ L mol ⁻¹ cm ⁻¹); author reported mean of 2 measurements (this one and another with a different ref. compd.) as 28800 ± 7200 L mol ⁻¹ cm ⁻¹	680727
	Cyclohexane	PR/ET	432.5, 31300		ϵ relative to naphthalene in cyclohexane ($\epsilon_{412.5} = 22600$ L mol ⁻¹ cm ⁻¹); author reported mean of 2 measurements (this one and another with a different ref. compd.) as 28800 ± 7200 L mol ⁻¹ cm ⁻¹	680727
	Cyclohexane	PR/ET	432.5, 31500		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700$ L mol ⁻¹ cm ⁻¹); reference ϵ obtained by starting from $\epsilon_{\max} = 3220$ L mol ⁻¹ cm ⁻¹ for this ketyl radical in water and assuming the f of the ketyl radical is independent of solvent; final ϵ obtained from a simultaneous least squares fit to data from several compounds	71E360
	EtOH	FP	440		Broad absorption < 290 nm also assigned to triplet; a band at 520 nm, which had been assigned to the triplet in water [61E008], was seen to decay at a different rate and was not assigned to the triplet in EtOH	66A003
	EtOH	FP/ET	440, 18500		Triplet ET from eosin and proflavine; ϵ relative to eosin in EtOH ($\epsilon_{580} = 9400$ L mol ⁻¹ cm ⁻¹) and proflavine in EtOH ($\epsilon_{550} = 11000$ L mol ⁻¹ cm ⁻¹)	716235
	EtOH	FP	985		Most intense peak at 440 nm	776227
			870			
			778			
			440			
	EtOH	LP/ET	440, 22500		ϵ relative to Methylene Blue cation ($\epsilon_{860} = 27000$ L mol ⁻¹ cm ⁻¹); solvent and temperature assumed; $\tau_T = 14 \mu s$	78E394
	Hexane	LP	435		145 ps delay	79B007
	Hexane	LP	980 ± 2			83E392
			430 ± 1			
	Hexane	LP	433		Delay 320 ps	84B154
			407			
	Isooctane	LP	430		A band was also observed at 511 ^a nm, but a similar band at 530 nm in MeOH was assigned to a precursor of the triplet state; rise time of 0.01 ns	767261
			408 ^a			
	MeOH	LP	983 ± 5			83E392
			438 ± 1			
	PMMA (77 K)	PS	440		Band seen at 480 nm and was attributed to a photo-product	70E291
			420			
	PMMA (293 K)	FP	438		$\tau_T = 1.9 \times 10^4 \mu s$	70E291
			418			
	PMMA	FP	980		Shoulder at 424 nm; 441 nm peak 10 times more intense than any other peak	82E561
			863 ^a			
			776 ^a			
			702 ^a			
			480 ^a			
			441 ^a			
	PVA	FP	980 ^a		Most intense peak at 450 nm; $\tau_T = 23.8 \times 10^3 \mu s$	776227
			870 ^a			
			780 ^a			
			450 ^a			

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Water	FP	443		$pK_b = 8.4$; a peak reported at 520 nm was later [66A003] shown to decay at a different rate than the 440 nm peak in EtOH; pH Basic	61E008
	Water (296 K)	FP-ET/ET	443, 14500		†Triplet ET from naphthalene-1,5-disulfonate; ϵ relative to proflavine conjugate acid in water ($\epsilon_{549} = 5700$ L mol ⁻¹ cm ⁻¹); $\tau_T = \sim 450$ μ s; pH 10.86	80B057
	Water	FP/TD	443 353 ^b , 7000		ϵ method assumed from text; pH 12.0	80F001
	Water (296 K)	FP/ET	880 443, 20000 \pm 2000		ϵ relative to proflavine in water ($\epsilon_{549} = 8000$ L mol ⁻¹ cm ⁻¹); pH Basic	81E147
	Water	LP	855 ^a 746 ^a 694 ^a 510 ^a 441 ^a		Most intense peak at 441 nm; pH 12.7	81E552
	Water (296 K)	FP/TD	880, 1800 ^a 753 ^a , 1100 ^a 689 ^a , 730 ^a 657 ^a , 470 ^a 521 ^a , 1500 ^a 443, 19400 \pm 1400		Shoulder at 420 nm; pH Basic	81E147
	Water	LP	983 \pm 5 862 ^a 775 ^a 658 ^a 443 \pm 1		pH \sim 13	83E392
	<i>tert</i> -BuOH	FP/TD	436, 26000			776258
24.	Acridine- <i>d</i> ₈ Benzene Fluorene- <i>d</i> ₈ (240 K)	FP/TD MOD	442, 26000 450		Single crystal	776258 83F334
25.	Acridine Orange, conjugate monoacid EtOH EtOH/Et ₂ O (90 K)	FP/TD PS/SD	490, 135000 1270, 54000 ^a 1050 ^a , 16000 ^a 645, 10500 ^a 540, 10000 ^a 390, 4000 ^a 285, 57000 ^a 245, 18000 ^a 205, 16000 ^a		pH Acidic Solvent was 2:1 EtOH to Et ₂ O	727073 57B001
	Glycerol (293 K)	FP	670 ^a 620 ^a 580 ^a		Solvent has added glucose	776171
	MeOH	FP/TD	610, 9400 560, 9200		$pK_a(\text{MeOH}) = 11.2$; $\tau_T = 105$ μ s; pH Acidic	79E219
	PMMA (77 K)	PS	1208 ^a 654 ^a 549 ^a		Relative intensities (10:3:3); shoulders at 1090 ^a and 1002 ^a nm; solvent contains 2-chloroethanol; pH 4.0	69B007
	Water	FP/TD	530, 9500 480, 14000 280, 35000		$\tau_T = \sim 3000$ μ s; pII 3	747187
26.	Acridine Orange, free base EtOH MeOH	FP/TD FP/TD	410, 42000 550, 7800 410, 39900		pH Basic $\tau_T = 285$ μ s; pH Basic	727073 79E219
	PMMA (193 K)	PS	530		Shoulders at 630 and 545 nm	59B001
	PMMA (77 K)	PS	1210 1100 1020 650			65B004

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Water	FP/TD	550, 7000 410, 37000 270, 40000		†Triplet ET to acridine; neutral form; an absorption was observable out to 900 nm; $\tau_T = 3.3 \times 10^3 \mu s$; pH 12	71B002
27.	Acridine Yellow, conjugate monoacid					
	Boric acid (77 K)	PS	1000 620			65B004
	Glycerol (293 K)	FP	660 ^a 550 ^a		Solvent has added glucose	776171
	PMMA (77 K)	PS	1000 ^a		Shoulders at 575 ^a and 532 ^a nm; solvent contains 2-chloroethanol; pH 4.0	69B007
28.	Acridine, conjugate acid					
	Water (296 K)	FP-ET/ET	490, 5500		†Triplet ET from naphthalene-1,5-disulfonate and to Methylene Blue; ϵ relative to proflavine conjugate acid in water ($\epsilon_{549} = 5700 \text{ L mol}^{-1} \text{ cm}^{-1}$); $pK_a = 5.6$; $\tau_T = 670 \mu s$; pH 1.0	80B057
	Water (296 K)	LP-ET/TD	770 490, 7700 \pm 800		†Triplet ET from disodium naphthalene-1,5-disulfonate; pH Acidic	81E147
29.	9(10H)-Acridone					
	Benzene	LP/ET	620, 37800		†Triplet ET to anthracene, oxygen quenching; ϵ relative to anthracene in benzene ($\epsilon_{430} = 45500 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 20 \mu s$	766377
	EtOH (300 K)	FP	580 310			81E649
	Pyridine	LP	605 ^a		†Triplet ET to anthracene; H-bonded species; $\tau_T = 3.3 \mu s$	766377
30.	Acriflavine cation					
	EPA (77 K)	PS	640 540		†Phosphorescence decay; EPA was 8:3:5 Et ₂ O to isopentane to EtOH; "tryptaflavine"	63F021
	Glycerol (293 K)	FP	620 ^a 550 ^a		Solvent has added glucose	776171
	MeOH	FP/TD	620, 8600		Peak at 490 nm obscured by singlet absorption (authors); $\tau_T = 110 \mu s$; pH Basic	79E219
	PMMA (193 K)	PS	555 530		Shoulders at 650 and 505 nm	59B001
	PMMA (77 K)	PS	1300 1190 1080 940 800 650			65B004
	PMMA (77 K)	PS	1060 ^a 679 ^a 541 ^a		Relative intensities (10:3:3); shoulder at 952 ^a nm; solvent contains 2-chloroethanol; pH 4.0	69B007
31.	Alloxazine					
	EtOH	LP/SD	500, 13000 438 ^a		†Phosphorescence decay at 77 K; 438 nm peak may be distorted by SD; $\tau_T = 13 \mu s$	737439
	Water	LP	550 420		†Oxygen quenching; $\tau_T = 9 \mu s$; pH 2	737439
	Water	LP	560 420		†Oxygen quenching; $\tau_T = 9 \mu s$; pH 6	737439
	Water	LP/RA	550, 7000 400 ^a , 12600 ^a 275 ^a , 28000 ^a		†Oxygen quenching; energy transfer to oxygen; lifetime measured at 560 nm; ϵ relative to anthracene in cyclohexane ($\Phi_T = 0.71$, $\epsilon_{423} = 64700 \text{ L mol}^{-1} \text{ cm}^{-1}$) and using $\Phi_T = 0.31$ for alloxazine in water; shoulder around 360 nm; $\tau_T = 9.1 \mu s$; pH 2.2; $k_{et} = (1.7 \pm 0.2) \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	777617

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
32.	Amiloride Water	LP/ET	400, 6300		†Triplet ET from benzophenone; pH was not specified and the pK_a of the triplet was not determined so the state of protonation of the triplet is uncertain; ϵ relative to benzophenone in water ($\epsilon_{520} = 5800$ L mol ⁻¹ cm ⁻¹); $\tau_T = 40$ μ s	84A227
33.	9-Aminoacridine Water	LP	885 ^a 826 ^a 794 ^a 476 ^a		Most intense peak at 476 nm; pH 12.7	81E552
34.	1-Amino-4-anilinoanthraquinone Benzene	PR-ET/ET	550, 27000		†Triplet ET from biphenyl; ϵ relative to biphenyl in benzene assuming ground state dimerization; $\tau_T = 8.3$ μ s; $E_T = 123 - 151$ kJ mol ⁻¹ ; $k_{et} = 3.3 \times 10^9$ L mol ⁻¹ s ⁻¹	761122
35.	2-Aminoanthracene, conjugate acid Water/Alcohol	FP	424 400 ^a		Solvent was 1:1 water to alcohol; pK_a 3.3; pH 1.25	68B008
36.	1-Aminoanthraquinone Benzene	PR/ET	550 ^b , 2000		†Triplet ET from triplet donors and triplet ET to triplet acceptors; ϵ relative to naphthalene in benzene ($\epsilon_{425} = 13200$ L mol ⁻¹ cm ⁻¹); $\tau_T = 5$ μ s	720392
37.	2-Aminoanthraquinone Benzene	PR/ET	599 ^a , 7140 ^a		†Triplet ET from triplet donors and triplet ET to triplet acceptors; ϵ relative to naphthalene in benzene ($\epsilon_{425} = 13200$ L mol ⁻¹ cm ⁻¹); $\tau_T = 5$ μ s	720392
38.	6-Amino-7H-benz[de]anthracen-7-one 2-PrOH	I.P	610 ^a 570 ^a 515 ^a		Shoulder at 540 nm; half-life = 3.7 ± 0.2 μ s	757427
	Benzene	LP	617 ^a 579 ^a 540 ^a		†Oxygen quenching (1.2×10^9 L mol ⁻¹ s ⁻¹); half-life = 2.3 ± 0.1 μ s	757427
39.	2-Aminobenzoic acid EtOH (93 K)	PS	472 ^a 395 ^a		Shoulder at 435 ^a nm	66B001
40.	4-Aminobenzoic acid EtOH (93 K)	PS	412 ^a 364 ^a			66B001
41.	3-Aminobenzophenone Propylene glycol (203 K)	FP	561 ^a 455 ^a		†Phosphorescence decay	67E105
42.	4-Aminobenzophenone Propylene glycol (203 K)	FP	638 ^a 466 ^a		†Phosphorescence decay	67E105
43.	4-Aminobiphenyl Toluene/EtOH (77 K)	MOD	438 ^a		Glass was 19:1 toluene to EtOH	719059
44.	2-Aminofluorene EtOH/Et ₂ O (77 K)	MOD/KM	425, 10600 \pm 4200		Glass was 2:1 EtOH to Et ₂ O; temperature was not explicitly stated, but 77 K was inferred from the context	719059
	Toluene/EtOH (77 K)	MOD	430 ^a		Glass was 19:1 toluene to EtOH; shoulder at 466 ^a	719059

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
45.	1-Amino-4-hydroxyanthraquinone Benzene	PR-ET/ET	500, 28000		{Triplet ET from biphenyl; ϵ relative to biphenyl in benzene assuming ground state dimerization; $\tau_T = 6.7$ μ s; $E_T = 123 - 176$ kJ mol ⁻¹ ; $k_{et} = 3.0 \times 10^9$ L mol ⁻¹ s ⁻¹	761122
46.	4-Amino-4'-hydroxybiphenyl EtOH/Et ₂ O (77 K)	MOD/KM	425, 24000 \pm 9600		Glass was 2:1 EtOH to Et ₂ O; temperature was not explicitly stated, but 77 K was inferred from the context	719059
	Toluene/EtOH (77 K)	MOD	428 ^a		Glass was 19:1 toluene to EtOH	719059
47.	2-Amino-3-(4-methoxy-6-benzothiazolyl)propionate ion Water	LP/ET	370, 8000		{Oxygen quenching and triplet ET to crocetin; ϵ relative to crocetin in water ($\epsilon_{470} = 75000$ L mol ⁻¹ cm ⁻¹); ϵ considered an upper limit; $\tau_T = 64$ μ s; pH 13; $k_{et} = 1.7 \times 10^{10}$ L mol ⁻¹ s ⁻¹	84A024
	Water	LP/TD	370, 5000		{Oxygen quenching and triplet ET to crocetin; ϵ considered a lower limit; pH 13	84A024
48.	1-Amino-4-(N-methylamino)anthraquinone Benzene	PR-ET/ET	575, 27000		{Triplet ET from biphenyl; ϵ relative to biphenyl in benzene assuming ground state dimerization; $\tau_T = 3.6$ μ s; $E_T = 97 - 123$ kJ mol ⁻¹ ; $k_{et} = 4.2 \times 10^9$ L mol ⁻¹ s ⁻¹	761122
49.	7-Amino-4-methylcarbostyryl EtOH	FP/ET	600 ^b , 46200 \pm 4500 430 ^b , 7050 \pm 700 350 ^b , 28800 \pm 2900		ϵ relative to anthracene in EtOH ($\epsilon_{420} = 75000$ L mol ⁻¹ cm ⁻¹); there were no distinct maxima between 350 and 600 nm; $k_{et} = (1.4 \pm 0.5) \times 10^{10}$ L mol ⁻¹ s ⁻¹	747049
50.	3-Amino-N-methylphthalimide Benzene	FP	450		Lifetime limited by aeration; $\tau_T = 0.260$ μ s	757522
	EtOH	FP/TD	495, 1100		Lifetime limited by aeration; $\tau_T = 0.920$ μ s	757522
	Water	FP	540		Lifetime limited by aeration; $\tau_T = > 1$ μ s	757522
51.	4'-Aminomethyl-4,5',8-trimethylpsoralen EtOH	LP/ET	460 ^a , 24200		{Triplet ET to retinol; ϵ relative to retinol (ϵ_{ref} unspecified); quenched by ground state (10^9 L mol ⁻¹ s ⁻¹); $\tau_T = 100$ μ s	80E156
52.	1-Amino-7-nitrofluorene EPA (77 K)	PS	770 ^a , 570 ^a			78E057
53.	1-Amino-4-nitronaphthalene Benzene	LP	470		{Oxygen quenching (2.0×10^9 L mol ⁻¹ s ⁻¹); $\tau_T = 1.5$ μ s	776195
	EPA (77 K)	PS/KM	650, 2730 \pm 135 480, 7500 \pm 375 440, 28000 \pm 14000		{Phosphorescence decay; $\tau_T = (1.88 \pm 0.11) \times 10^5$ μ s	78E057
	PMMA	FP	665 470		Relative intensities (2:5)	776195
	PMMA (123 K)	FP	665		{Phosphorescence decay; $\tau_T = 1.5 \times 10^5$ μ s	776195
	PVA (123 K)	FP	695 ^a			776195
54.	trans-4-Amino-4'-nitrostilbene Glycerol triacetate	LP	772			78B088
	Glycerol triacetate (198 K)	LP	760			78B088
55.	N-[2-(3-Amino-1-oxopropyl)phenyl]formamide EtOH	LP	460		{Oxygen quenching (3×10^8 L mol ⁻¹ s ⁻¹); $\tau_T = 1$ μ s	78E308
	Water	LP	435		{Oxygen quenching (2×10^9 L mol ⁻¹ s ⁻¹); ionic strength 0.01 mol L ⁻¹ ; $\tau_T = 3.4$ μ s; pH 7.5	78E308

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
56.	9-Aminophenanthrene Toluene/EtOH (77 K)	MOD	473 ^a		Glass was 19:1 toluene to EtOH; shoulder at 431 ^a nm	719059
57.	2-Amino-4-[3H]pteridinone Water	LP/HAT	550, 2000 385 ^b , 2770 360 ^a , 4700 ^a 325 ^b , 3780		†Oxygen quenching (1.3×10^9 L mol ⁻¹ s ⁻¹); hydrogen abstraction from guanosine; ϵ_{550} relative to semireduced radical in water ($\epsilon_{480} = 2300$ L mol ⁻¹ cm ⁻¹); ϵ_{385} and ϵ_{325} are at isosbestic points in difference spectrum; 400 ns delay; $\tau_T = 2.3 \pm 0.2$ μ s; pH 9.2	81E151
	Water	LP	460		400 ns delay, $pK_a = 9.5$ -10; $\tau_T = 0.5$ μ s; pH 12	81E151
	Water	LP	600 415		100 ns delay; $\tau_T = 0.3$ μ s; pH 9.2	81E151
58.	2-Aminopurine Acetonitrile (300 K)	FP	430		†Oxygen quenching and quenching by 2,4-hexadienol; $\tau_T = 83 \pm 10$ μ s	756270
	Water (300 K)	FP	405		†Oxygen quenching and quenching by 2,4-hexadienol; $\tau_T = 55 \pm 4$ μ s	756270
59.	2-Aminopyridine 2-PrOH (163 K)	FP	390 ^a		†Phosphorescence decay; $\tau_T = \sim 1.8 \times 10^3$ μ s	78E062
	EPA (107 K)	FP	390		†Phosphorescence decay; $\tau_T = \sim 2.6 \times 10^3$ μ s	78E062
60.	4-Aminopyridine EPA (103 K)	FP	395		†Phosphorescence decay; $\tau_T = \sim 4.0 \times 10^4$ μ s	78E062
61.	Aniline EPA (77 K)	FP	420 ^a 320 ^a		†Phosphorescence decay; 320 nm was the more intense peak	69E215
62.	1-Anilinonaphthalene Cyclohexane	LP	550			77E543
	MeOH/Water	LP	580		†Quenching by oxygen and <i>trans</i> -stilbene	82E359
63.	Anisole Water	LP	252 ^a		Lifetime was measured at pH 7.5; there was a broad maximum between ~ 350 and ~ 430 nm; shoulder at 279 ^a nm; spectrum is the difference between spectra at 20 ns and 15 μ s; $\tau_T = 3.3 \pm 0.2$ μ s; pH 8.5	757161
64.	Anthracene Acetonitrile	FP	438 ^a 400 ^a		Relative intensities (2:1); solvent contains dimethylmercury	78A324
	Anthracene	PR	630 ^a 610 ^a 570 ^a 540 ^a		Triplet exciton	80E230
	Benzene	FP	430			58E002
	Benzene	FP	430		A value was given for an "integrated extinction coefficient" of 35000, but no details were provided; $\tau_T = 2900$ μ s	60E005
	Benzene	PR/ET	427.5, 71500		ϵ relative to benzophenone in benzene ($\epsilon_{532.5} = 10300$ L mol ⁻¹ cm ⁻¹)	690087
	Benzene	PR/ET	427.5, 61300		ϵ relative to naphthalene in benzene ($\epsilon_{\max} = 17500$ L mol ⁻¹ cm ⁻¹)	690087
	Benzene	PR/ET	430, 45500 408 ^a , 15600 ^a 391 ^a , 3700 ^a		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700$ L mol ⁻¹ cm ⁻¹); ϵ obtained from a simultaneous least squares fit of data from several compounds making use of cyclohexane to benzene ϵ_{\max} ratios of 1.83 for naphthalene and 1.45 for anthracene; λ_{\max} originally quoted as 432.5 nm but corrected in [757282]	71E360
	Benzene	PR	427.5			720206
	Benzene	FP/SD	428.5, 53000 \pm 1900		ϵ method assumes linear variation of triplet spectrum in SD region	767147

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
Benzene		LP/SD	431, 69000		ϵ is upper limit (authors)	80B090
Benzene		LP/ET	431, 42000 \pm 4000		\dagger Triplet ET from benzophenone; ϵ relative to benzophenone in benzene ($\epsilon_{533} = 7200$ L mol ⁻¹ cm ⁻¹); $k_{et} = 0.49 \times 10^9$ L mol ⁻¹ s ⁻¹	80B090
Benzonitrile		LP/RF	429 ^a , 24000 \pm 1500 405 ^a , 10200 ^a		ϵ relative to anthracene in toluene ($\epsilon_{428.5} = 42000$ L mol ⁻¹ cm ⁻¹)	83F075
Benzophenone (77 K)		PS-ET	441		Relative intensities (1:1); sample was a single crystal	64B005
Benzophenone (77 K)		PS	415 442.5		Single crystal; from polarization measurements, the 442.5 nm band was assigned $^3B_{3g} \leftarrow ^3B_{1u}$; the $T_1 \leftarrow S_0$ absorption of benzophenone starting at 413.4 nm obscured the shape of the second T-T band of anthracene	68B004
Benzophenone (303 K)		PR	435		Solution; triplet ET from benzophenone; $k_{et} = (1.6 \pm 0.1) \times 10^9$ L mol ⁻¹ s ⁻¹	700114
Bromobenzene		FP	436		Half-life = 430 \pm 50 μ s	58E002
Bromobenzene		FP	439		A value was given for an "integrated extinction coefficient" of 32500, but no details were given; $\tau_T = 1590$ μ s	60E005
CTAB		LP	420		Aqueous micelles	83N082
Chlorobenzene		FP	431		Half-life = 790 \pm 150 μ s	58E002
Chloroform		FP	430			58E002
Cyclohexane		FP	425		$\tau_T = 160$ μ s	62E009
Cyclohexane		PR/ET	420, 57200 \pm 14300		ϵ relative to benzophenone ketyl radical in water ($\epsilon_{537.5} = 3220$ L mol ⁻¹ cm ⁻¹), assuming this value for $\epsilon_{542.5}$ in cyclohexane	680727
Cyclohexane (77 K)		PS	468.2 431.0 425.2 418.1 402.0 381.0 316.5			69E211
Cyclohexane		PR	427.5			710617
Cyclohexane		PR/ET	422.5, 64700 402 ^a , 19800 ^a		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700$ L mol ⁻¹ cm ⁻¹); reference ϵ obtained by starting from $\epsilon_{\max} = 3220$ L mol ⁻¹ cm ⁻¹ for this ketyl radical in water and assuming the f of the ketyl radical is independent of solvent; final ϵ obtained from a simultaneous least squares fit to data from several compounds; shoulder at 388 ^a nm; λ_{\max} originally quoted as 425 nm but corrected in [757282]	71E360
Cyclohexane		FP/SD	422.5, 85700 \pm 3200		ϵ method assumes linear variation of triplet spectrum in SD region	767147
Cyclohexane		LP/RF	423, 52500		ϵ relative to anthracene in benzene ($\epsilon_{431} = 42000$ L mol ⁻¹ cm ⁻¹) assuming oscillator independent of solvent	80B090
Cyclohexane		LP/SD	423, 69500		ϵ is upper limit (authors)	80B090
Decalin		FP	430		$\tau_T = 2630$ μ s	60E005
Dioxane		PR	425 410			650006
EPA (77 K)		PS/IV	427, 2800 ^c \pm 130		\dagger Phosphorescence decay; ϵ is too low; 1st transition assigned as $^1L_a \rightarrow ^1C_0$	51E001
EPA (77 K)		PS/SD	427.0, 45000 404.0, 12000		ϵ 's are lower limits; $\tau_T = < 1 \times 10^3$ μ s	54B001
EPA (77 K)		PS/KM	516 ^a , 1600 ^a 497 ^a , 800 ^a 481 ^a , 2100 ^a 467 ^a , 1300 ^a 428 ^a , 52800 ^a 402 ^a , 14100 ^a		Solvent, temperature and extinction method assumed from earlier work; polarization also measured	69E213
EPA (77 K)		PS/IV	427, 58000		λ_{\max} assumed from previous work; ϵ estimated by extrapolation to infinite excitation rate	69E212
EPA (77 K)		FP/SD	427.3, ≥ 90000			69F389

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Epoxy plastic (296 K)	LP	420			686058
			390			
	EtOH	FP	420.0		$\tau_T = 150 \pm 45 \mu s$	54E001
			398.0			
	EtOH (93 K)	PS	427		Relative intensities (10:3)	66B001
			403			
	EtOH	FP/ET	440 ^b , 2300		Triplet ET from eosin and proflavine; ϵ relative to eosin in EtOH ($\epsilon_{580} = 9400$ L mol ⁻¹ cm ⁻¹) and proflavine in EtOH ($\epsilon_{550} = 11000$ L mol ⁻¹ cm ⁻¹)	716235
			426 ^b , 21000			
			424 ^b , 36000			
			421, 52000			
			420 ^b , 49000			
	EtOH	FP/SD	420, 75000 \pm 5000		$\tau_T = 3000 \pm 2000 \mu s$	747049
			400, 28000 \pm 2000			
			376, 5750 \pm 400			
			356, 3350 \pm 250			
			335, 2250 \pm 150			
	EtOH	FP/TD	419, 50000 \pm 5000			756293
	EtOH (77 K)	MOD	424		Halfwidth 380 cm ⁻¹	777538
	EtOH	FP/TD	420, 60000			776258
	EtOH	FP-ET/RA	694 ^b , 52		ϵ relative to anthracene in EtOH ($\epsilon_{421} = 52000$ L mol ⁻¹ cm ⁻¹)	78E019
	EtOH	LP/ET	420, 63500		ϵ relative to Methylene Blue cation ($\epsilon_{600} = 27000$ L mol ⁻¹ cm ⁻¹); solvent and temperature assumed; $\tau_T = 25 \mu s$	78E394
	EtOH	LP/SD	421, 54000		ϵ is upper limit (authors)	80B090
	EtOH	LP/RF	421, 48500		ϵ relative to anthracene in benzene ($\epsilon_{431} = 42000$ L mol ⁻¹ cm ⁻¹) assuming oscillator independent of solvent	80B090
	EtOH	LP/COM	418, 72000		Triplet concentration calculated by kinetic simulation and corrected for optical geometry; ϵ value assumes $\Phi_T = 0.7$; oxygen quenching (3.8×10^9 L mol ⁻¹ s ⁻¹)	82B121
	EtOH/MeOH (113 K)	FP/TD	425, 90000		Solvent was 3:1 EtOH to MeOH	67B004
	EtOH/MeOH (104 K)	PS	426		Solvent was 3:1 EtOH to MeOH; polarization also measured; shoulder at 382 nm	68E102
			403			
	EtOH/MeOH/2-MTHF (113 K)	FP/SD	885 ^a , 250 ^a		Glass was 3:1:4 EtOH to MeOH to 2-MTHF	69B006
			840 ^a , 260 ^a			
			800 ^a , 160 ^a			
			787 ^a , 150 ^a			
			763 ^a , 180 ^a			
			752 ^a , 110 ^a			
			730 ^a , 130 ^a			
			719 ^a , 150 ^a			
			709 ^a , 130 ^a			
			699 ^a , 130 ^a			
			690 ^a , 120 ^a			
			645 ^a , 130 ^a			
	EtOH/MeOH (113 K)	FP/TD	888, 260 ^a		1Phosphorescence decay; shoulders at 657 ^a , 551 ^a , 526 ^a , 486 ^a , 453 ^a , 271 ^a , and 268 ^a nm; 6 electronic transitions were assigned with 0-0's at 888, 516, 425.3, 328.7, 271.3, and 262.4 nm; oscillator strength = 0.002, 0.04, 0.25, 0.03, ~0.01, 0.8	72B001
			844 ^a , 300			
			801 ^a , 190 ^a			
			788 ^a , 190 ^a			
			764 ^a , 210 ^a			
			751 ^a , 140 ^a			
			729 ^a , 160 ^a			
			719 ^a , 190 ^a			
			709 ^a , 150 ^a			
			697 ^a , 180 ^a			
			687 ^a , 170 ^a			
			639 ^a , 170 ^a			
			538 ^a , 540 ^a			
			516, 3000			
			504 ^a , 2000 ^a			
			497 ^a , 2300 ^a			
			479 ^a , 3600 ^a			

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	EtOH/MeOH (Continued)		463 ^a , 2900 ^a 444 ^a , 5000 ^a 425.3, 90000 402 ^a , 25000 ^a 381 ^a , 10000 ^a 328.7, 1400 ^a 314 ^a , 2200 ^a 302 ^a , 2300 ^a 262.4, 90000			
	Ethyl acetate	PR	420		0.8 μ s delay; G (triplets) estimated assuming ϵ independent of solvent	761080
	Ethylene glycol	FP	469 425 401		Relative intensities (—:10:3); $\tau_T = 3800 \pm 150 \mu$ s	61E005
	Glycerol	FP	426.6 402.4		$\tau_T = 5500 \pm 600 \mu$ s	54E001
	Glycerol	FP	428		Half-life = $9400 \pm 1500 \mu$ s	58E002
	Glycerol (283 K)	FP/SD	425, 66800 ± 13400 404, 20700 ± 4140		Lifetime measured at 298 K; $\tau_T = 18000 \mu$ s	65E032
	Heptane	FP	421			58E002
	Heptane/Carbon disulfide	FP	421		Solvent was 9:1 heptane to carbon disulfide	58E002
	Heptane (77 K)	PS	429.5 405			67B007
	Hexane	FP	420.3 398		Half-life = 100μ s	53E001
	Hexane	FP/SD	420.3, $118000^e \pm 23600$ 398.0, $27400^e \pm 5480$		†Oxygen quenching; $\tau_T = 84.0 \pm 11 \mu$ s	54E001
	Hexane	FP	420		A value was given for an "integrated extinction coefficient" of 46000, but no details were provided; $\tau_T = 2900 \mu$ s	60E005
	Hexane	FP	465 423 400		Relative intensities (—:10:3); $\tau_T = 910 \pm 83 \mu$ s	61E005
	Hexane	LP	420 400		250 ns delay; relative intensities (4:1); formed from diethylaniline exciplex; rise time of ~ 110 ns	776081
	Liquid paraffin	FP	420.0 401.0		$\tau_T = 370 \pm 110 \mu$ s	54E001
	Liquid paraffin	FP/SD	468, 2000 424, 71500 401, 24300		Solvent had viscosity of 0.03 N-s/m ² ; bands at 424 and 401 nm were assigned to the same electronic transition; oscillator strength = 0.01, 0.4	58E001
	Liquid petrolatum	FP	425		Half-life = $1100 \pm 200 \mu$ s	58E002
	Liquid paraffin	FP	424		Viscosity of solvent was 0.167 N-s/m ² ; $\tau_T = 9100 \mu$ s	62E009
	Liquid paraffin	FP/TD	424 ^b , 63000 ± 5000			67E031
	MCH	LP	421 ^a 402 ^a		421 nm peak was the more intense	82F367
	PMMA (300 K)	FP	423.7 401.1		Relative intensities (100:26); half-life = $1.22 \times 10^4 \mu$ s	62E010
	PMMA (298 K)	FP/RA	895, 250 ± 50 852 ^a 807 ^a 796 ^a 771 ^a		†Decay matches the triplet decay at 424 nm; ϵ relative to anthracene in liquid paraffin ($\epsilon_{424} = 71500$ L mol ⁻¹ cm ⁻¹); $\tau_T = 3.4 \times 10^4 \mu$ s	66E085
	PMMA	FP	894 ^a 851 ^a 812 ^a 801 ^a 776 ^a 717 ^a		Shoulders at $\sim 860^a$ and 640^a nm	71B006
	PMMA (293 K)	LP	426 ^a 402 ^a 386 ^a		426 nm peak was the most intense	747056
	Polystyrene	FP	433		$\tau_T = 2.6 \times 10^4 \mu$ s	64E016

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Polystyrene	PR	435		$G\epsilon$ varied with the concentration of anthracene; 72% of the anthracene copolymerizes with polystyrene	672086
	Polystyrene	PR	436 ^a		$G\epsilon_{436} = 10000^a$ L mol ⁻¹ cm ⁻¹ / (100 eV absorbed)	701073
	Propane	FP/TD	422.5, 52000		ϵ assumes OD saturation is equivalent to total depletion	79A346
	THF	FP	428		$\tau_T = 3770$ μ s	60E005
	Toluene	LP	430 ^a 409 ^a		430 nm was the more intense peak	717126
	Toluene (77 K)	MOD	434		Halfwidth 490 cm ⁻¹	777538
	Toluene	LP/RA	428.5, 42000		ϵ relative to anthracene in benzene ($\Phi_T = 0.72$, $\epsilon_{428.5} = 42000$ L mol ⁻¹ cm ⁻¹) assuming $\Phi_T = 0.72$ for anthracene in toluene	82E451
	Triton X-100	LP	420		Aqueous micelles	83N082
	Water/ <i>tert</i> -BuOH	FP	420 ^a		Solvent mixture contains "1-5%" <i>tert</i> -BuOH for solubility; pH \sim 6	767189
65.	Anthracene- <i>d</i> ₁₀ Cyclohexane (77 K)	PS	467.1 445.0 427.5 421.5 413.6 407.0 398.0		Relative intensities (1:1:2:10:6:4:4)	69E211
	EPA (77 K)	PS/SD	426, 115000			68E105
	Et ₂ O/EtOH/Toluene (77 K)	PS/ESR	425, 75000		‡ESR; solvent was 2:1:1 Et ₂ O to EtOH to toluene; only most intense visible peak reported; assignment $^3B_{3g} \leftarrow ^3B_{2u}$; oscillator strength = 0.27	696115
	EtOH/MeOH (104 K)	PS	424 401		Solvent was 3:1 EtOH to MeOH; polarization also measured; shoulder at 382 nm	68E102
	PMMA (298 K)	FP	894 ^a 851 ^a 811 ^a 796 ^a 760 ^a		‡Decay matches the triplet decay at 424 nm; $\tau_T = 8.4 \times 10^4$ μ s	66E085
	PMMA (293 K)	CWL	428 ^a 404 ^a		‡Phosphorescence decay; maxima shift to the red with increase in applied pressure; spectrum measured at 5.4 kbar and assigned to the $^3B_{1g} \leftarrow ^3B_{2g}$ transition; relative intensities (8:1); at atmospheric pressure principal maximum is at 424 nm; $\tau_T = 85 \times 10^3$ μ s	76B012
	PMMA (293 K)	CWL	888 ^a 840 ^a 785 ^a 715 ^a		‡Phosphorescence decay; spectrum measured at 2.0 kbar and assigned to the $^3B_{1g} \leftarrow ^3B_{2g}$ transition; only principal maxima reported here; at atmospheric pressure lowest energy maximum is at 895 nm; $\tau_T = 85 \times 10^3$ μ s	76B012
66.	Anthracene/Chloranil Liquid paraffin	MOD	420		Mull	71E361
67.	9-Anthraceneacetic acid, methyl ester Water/Acetonitrile	FP	427 ^a 404 ^a		‡Oxygen quenching; solvent 4:1 water to acetonitrile; 427 nm peak was the more intense; $\tau_T = 59$ μ s	747389
68.	Anthracene-9-carboxaldehyde EPA (77 K)	FP	455		$\tau_T = 1700$ μ s	82E338
69.	Anthracene-9-carboxamide EPA (77 K)	FP	433		$\tau_T = 3.96 \times 10^4$ μ s	82E338
70.	Anthracene-9-carboxylic acid EPA (77 K)	FP	431		$\tau_T = 3.45 \times 10^4$ μ s	82E338

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
71.	Anthracene photodimer 2-MTHF (77 K)	LP	433 ^a 408 ^a		Delay 1.5 μ s; species formed by the photolysis of di-anthracene	84B019
	MTHF (100 K)	LP	355 ^a 322 ^a		†Oxygen quenching; $\tau_T = 2 \mu$ s	82A006
72.	Anthracene-tetracene photodimer MCH	LP	442 ^a		Air-saturated solution	82F367
73.	[2.2](1,4)(9,10)-Anthracenophane 2-MTHF (77 K)	LP-ET	510 ^a 480 ^a 440 ^a		†Triplet ET from biacetyl	79B050
74.	anti-[2.2](1,4)-Anthracenophane 2-MTHF (77 K)	LP-ET	510 ^a 475 ^a		†Triplet ET from biacetyl	79B050
75.	syn-[2.2](1,4)-Anthracenophane 2-MTHF (77 K)	LP-ET	680 ^a 650 ^a 500 ^a 460 ^a		†Triplet ET from biacetyl	79B050
76.	Anthraquinone Benzene	PR/ET	390, 10200 370		†Triplet ET from benzophenone and triplet ET to anthracene; ϵ relative to benzophenone in benzene ($\epsilon_{532.5} = 7630$ L mol ⁻¹ cm ⁻¹); $\tau_T = 0.2 \mu$ s	720392
	Benzene	FP	370		†Oxygen quenching, triplet ET to β -carotene; $\tau_T = 0.11 \mu$ s	81F130
	EPA (77 K)	FP	365 ^a		†Phosphorescence decay; $\tau_T = 3400 \mu$ s	83E016
	EtOH/MeOH (113 K)	FP/TD	370, >2000		Solvent was 3:1 EtOH to MeOH; lower limit assumes total ground state depletion	67B004
	Toluene	LP	371 ^a		†Phosphorescence decay in EPA at 77 K; delay 300 ns	83E016
77.	Anthraquinone-2,6-disulfonate ion Water	FP	390		†Oxygen quenching; $\tau_T = < 1 \mu$ s; pH 7	81F130
78.	Anthraquinone-2-sulfonate ion Acetonitrile	LP	579 ^a 473 ^a 451 ^a 378 ^a		Delay 40 ns	83B054
	Water	LP	456 ^a 411 ^a 380 ^a		Spectral data extrapolated back to 15 ns; pH ~ 6	83B054
79.	2-Anthroate ion Water/Alcohol	FP	434 ^a 409 ^a		Solvent was 1:1 water to alcohol; pK_a (triplet 2-anthraic acid) 6.0; pH 13	68B008
80.	1-Anthrol Cyclohexane	FP	440 ^a		Triplet ET from perylene; ϵ relative to perylene in cyclohexane ($\epsilon_{480} = 13000$ L mol ⁻¹ cm ⁻¹); $\tau_T = 295 \mu$ s	767626
	Cyclohexane	LP/ET	440, 28300		†Similarity to 1-anthryl diisopropyl borate spectrum; 510 nm was the more intense peak	777391
	Isooctane	FP	510 439			717266
81.	2-Anthrol Isooctane	FP	442 298		†Similarity to 2-anthryl diisopropyl borate spectrum	717266
82.	9-Anthrol MeOH (300 K)	LP	430			82B102

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
83.	Anthrone ? (81 K)	FP/SD	341, 74000		{Phosphorescence decay; glass was either 1:3:3 1-methylPrOH to Et ₂ O to isooctane or 1:9 MCH to mixture of EtOH and MeOH; $\tau_T = 1800 \pm 200 \mu s$	77B022
	Benzene	LP	530 ^a 385		{Triplet ET, oxygen quenching; decay times limited in oxygen saturated solution; shoulder around 420 nm; relative intensities (1:10); $\tau_T = 0.170 \mu s$; rise time of 0.070 ± 0.020 ns	766464
	Benzene	LP	~580		{Quenching by oxygen, di- <i>tert</i> -butylnitroxide, and dienes	82B102
	Hexane	LP	560		Rise time of 0.050 ± 0.020 ns	84E077
	Isooctane	FP	423 395 380		Some bands at 635 and 545 nm bands were unassigned	717266
84.	1-Anthryl diisopropyl borate Triisopropyl borate	FP	830 745 688 505 428		428 nm was the most intense peak	717266
85.	2-Anthryl diisopropyl borate Triisopropyl borate	FP	520 427.5 405 330			717266
86.	9-Anthryl diisopropyl borate Triisopropyl borate	FP	425 403 340 305 282			717266
87.	9-Anthryl ethenyl ketone EPA (77 K)	FP	430		$\tau_T = 3.13 \times 10^4 \mu s$	82E338
88.	1-(2-Anthryl)-2-(2-naphthyl)ethylene Benzene	PR/ET	635 \pm 3, 62000 \pm 9300 460 \pm 3, 50000 \pm 7500 640 \pm 3 590 ^a 465 \pm 3 439 ^a		{Oxygen quenching; ϵ relative to biphenyl in benzene ($\epsilon_{359} = 27100$ L mol ⁻¹ cm ⁻¹); $\tau_T = 88 \mu s$	84E111
	Bromobenzene	LP	621 ^a 567 ^a 451 ^a 425 ^a		{Azulene quenching; $\tau_T = 58 \mu s$	84E111
	MCH	LP				84E111
89.	1-(2-Anthryl)-2-phenylethylene Benzene	PR/ET	620 \pm 3, 41000 \pm 6000 455 \pm 3, 60000 \pm 9000 630 \pm 3 579 ^a 460 \pm 3 438 ^a		{Oxygen quenching; ϵ relative to biphenyl in benzene ($\epsilon_{359} = 27100$ L mol ⁻¹ cm ⁻¹); $\tau_T = 117 \mu s$	84E111
	Bromobenzene	LP	611 ^a 559 ^a 449 ^a 421 ^a		{Azulene quenching; $\tau_T = 40 \mu s$	84E111
	MCH	LP			Shoulder at 520 ^a nm	84E111
90.	1-(2-Anthryl)-2-(2-thienyl)ethylene Benzene	PR/ET	630 \pm 3, 47000 \pm 7000 455 \pm 3, 38000 \pm 5700		{Oxygen quenching; ϵ relative to biphenyl in benzene ($\epsilon_{359} = 27100$ L mol ⁻¹ cm ⁻¹); $\tau_T = 19 \mu s$	84E111

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Bromobenzene	LP	635 ± 3 581 ^a		†Azulene quenching; $\tau_T = 22 \mu\text{s}$	84E111
	MCH	LP	460 ± 3 619 ^a 566 ^a 448 ^a 420 ^a		Shoulder at 527 ^a nm	84E111
91.	Astacene Benzene	PR-ET	580		†Triplet ET from naphthalene; $\tau_T = 6.2 \mu\text{s}$	80A143
92.	Astaxanthin Benzene	PR-ET	580		†Triplet ET from naphthalene; method not explicit; $\tau_T = 6.2 \mu\text{s}$	81B097
93.	Auramine O PMMA (77 K)	PS	1300 1200 1090 700			65B004
94.	Aurophosphine, conjugate monoacid Glycerol (293 K)	FP	690 ^a 610 ^a 550 ^a		Solvent has added glucose	776171
95.	Azulene Acetonitrile	LP	360		†Triplet ET from benzophenone, oxygen quenching ($8 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); $\tau_T = 3 \mu\text{s}$	81F275
	Benzene	LP/ET	360, 4000		†Triplet ET from benzophenone and other donors, oxygen quenching ($6 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$; ϵ relative to benzophenone in benzene ($\epsilon_{\max} = 7600 \text{ L mol}^{-1} \text{ cm}^{-1}$) assuming unit transfer efficiency; [84E491] sees no maximum down to 345 nm; $\tau_T = 4 \mu\text{s}$	81F275
	Benzene	PR	<345		†Triplet ET to perylene and from 1,3-cyclohexadiene; spectrum corrected for SD by monitoring singlet at 620 nm; $\tau_T = 11 \pm 1 \mu\text{s}$	84E491
96.	Azure A Water	FP	790		$\text{p}K_a = 7.0 \pm 0.1$	67C001
97.	Azure A cation Water	FP/SD	840 ^b , 10000		†Oxygen and I^- quenching; $\text{p}K_b = 6.3$; structure of the spectrum could not be discerned because it was not corrected for SD; pH 9	70E293
98.	Azure B Water	FP	800		$\text{p}K_a = 7.2 \pm 0.1$	67C001
99.	Azure C Water	FP	810		$\text{p}K_a = 7.4 \pm 0.1$	67C001
100.	Bacteriochlorophyll EtOH/Pyridine	LP/SD	631 ^a , 18500 ^a 588 ^a , 19600 ^a 515 ^a , 21500 ^a 400 ^a , 46700 ^a 323 ^a , 38300 ^a		†Triplet ET; solvent was 7:3 EtOH to pyridine; shoulder at 691 ^a nm; $\tau_T = 74 \pm 4 \mu\text{s}$	733184
	Pyridine	FP/SD	655, 9300 620, 11500 410, 22200		$\tau_T = 85 \pm 13 \mu\text{s}$	60E006
	Pyridine	LP/SD	616 ^a , 20800 ^a 512 ^a , 19500 ^a 405 ^a , 41000 ^a		†Triplet ET; shoulder at 690 ^a nm; $\tau_T = 89 \pm 5 \mu\text{s}$	733184

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
101.	Benz[a]acridine EPA (77 K)	PS	497		Relative intensities (10:9); "1,2-benzacridine"; $\tau_T = 3 \times 10^5 \mu s$	54B001
	Isopentane/MCH (77 K)	PS	497		$\tau_T = < 5 \times 10^5 \mu s$	54B001
102.	Benz[c]acridine EPA (77 K)	PS	510 480		Relative intensities (10:9); "3,4-benzacridine"; $\tau_T = 4 \times 10^5 \mu s$	54B001
103.	Benz[b]acridin-12(5H)-one EtOH (300 K)	FP/ET	590, 45000 380		ϵ relative to anthracene in EtOH ($\epsilon_{421} = 52000$ L mol ⁻¹ cm ⁻¹); "6,7-benzacridinone"	81E649
104.	Benzaldehyde Liquid paraffin	FP	470 364 320 307		Solvent viscosity was 0.03 N·s/m ² ; 470 and 364 nm bands were assigned to one electronic transition, and the other two bands were assigned to another electronic transition; relative intensities (11:18:100:86)	58E001
105.	Benzamide Liquid paraffin	FP	505 300		Solvent viscosity was 0.19 N·s/m ² ; bands were assigned to 2 different electronic transitions	58E001
106.	Benz[a]anthracene 2-MTHF (77 K)	PS/ESR	493, 43600 ^c 469 ^a , 22700 ^{a,c} 441 ^a , 19200 ^{a,c}		†ESR; shoulder at 410 ^a nm; "1,2-benzanthracene"; oscillator strength = 0.43	68D211
	Benzene	FP	538 488 458 435 402		Relative intensities (10:100:65:50:30); $\tau_T = 160 \pm 13 \mu s$	61E005
	Benzene	PR/ET	490, 20500		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700$ L mol ⁻¹ cm ⁻¹); ϵ obtained from a simultaneous least squares fit of data from several compounds making use of cyclohexane to benzene ϵ_{\max} ratios of 1.83 for naphthalene and 1.45 for anthracene	71E360
	Benzonitrile	LP/RF	490 ^a , 12000 \pm 800 466 ^a , 9800 ^a		ϵ relative to benz[a]anthracene in toluene ($\epsilon_{490} = 15000$ L mol ⁻¹ cm ⁻¹)	83F075
	Cyclohexane	PR/ET	480, 25100 \pm 6270		ϵ relative to benzophenone ketyl radical in water ($\epsilon_{537.5} = 3220$ L mol ⁻¹ cm ⁻¹), assuming this value for $\epsilon_{542.5}$ in cyclohexane	68O727
	Cyclohexane	LP	480		†Rise time of transient was the same as the decay time of the singlet	68B006
	Cyclohexane	LP	540 490 461			70E288
	Cyclohexane	PR/ET	480, 28800		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700$ L mol ⁻¹ cm ⁻¹); reference ϵ obtained by starting from $\epsilon_{\max} = 3220$ L mol ⁻¹ cm ⁻¹ for this ketyl radical in water and assuming the f of the ketyl radical is independent of solvent; final ϵ obtained from a simultaneous least squares fit to data from several compounds	71E360
	Cyclohexane	LP	543 483 457 429		350 ns delay	737463
	EPA (77 K)	PS	489.0 473.3 459.3		†Phosphorescence decay; 1st transition assigned as ³ L _a → ³ C ₀	51E001
	EPA (77 K)	PS	489.0 460.8 434		†Phosphorescence decay; relative intensities (100:58:54); unresolved shoulder at 474 nm; $\tau_T = (3 \pm 1) \times 10^5 \mu s$	54B001

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
EPA (77 K)	PS		710 645 590		Weak bands in the red only studied	67B005
EPA (77 K)	PS/SD		487.5, 29000 465 ^a , 18200 ^a 433 ^a , 13600 ^a		EPA in this work was 4:4:1 isopentane to Et ₂ O to absolute EtOH	68E104
EPA (77 K)	PS		709.2 645.2 589.9 543.5 489.0 471.9 460.8 434.8 411.0 384.0		Relative intensities (7:5:6:4:100:42:70:44:27:20)	69E211
EPA (77 K)	PS/IV		488, 28800		λ_{\max} assumed from previous work; ϵ estimated by extrapolation to infinite excitation rate	69E212
EtOH (293 K)	FP		540 482			68E098
EtOH (293 K)	LP		442 ^a		Halfwidth 440 cm ⁻¹	697226
EtOH (77 K)	MOD		486			777538
EtOH	LP		480 ^a 450 ^a 430 ^a		Relative intensities (6:4.2:4); 1.8 μ s delay; solution contains Ag ions	78E554
Heptane (77 K)	PS		487 460 437.4 413 402			67B007
Hexane	FP		485.0 450.1 429.9			54E001
Hexane	MOD/SD		480 ^a , 24500 ^a 455 ^a , 18200 ^a 430 ^a , 13800 ^a 360 ^a , 2800 ^a 295 ^a , 43700 ^a 279 ^a , 31600 ^a		Shoulders at 410 nm and 387 nm; $\tau_T = 100 \mu$ s	64E015
Hexane (300 K)	MOD/SD		531, 2500 481, 26000 455, 21000 427, 17000 407, 9000 360, 2800 345, 3600 302, 40000 274, 30000 247, 24000			69E208
Isopentane/MCH (77 K)	PS		486.6 459 430		Relative intensities (100:62:55); unresolved shoulder at 472 nm	54B001
Liquid paraffin	FP/SD		540, 3000 485, 23400 461, 19700 434.5, 12400 415, 8900		Solvent had viscosity of 0.03 N·s/m ² ; 3 electronic transitions were assigned; all bands belong to the 2nd electronic transition except the bands at 540 and 315 nm; oscillator strength = 0.02, 0.34, —	58E001
Naphthalene (273 K)	PR		315 495 475 435		Solution; relative intensities (100:83:66); delay 300 ns; $k_{et} = 1.1 \times 10^{10}$ L mol ⁻¹ s ⁻¹	700114

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Nonane (77 K)	PS	490.0 463.6 460.0 432.0 410.0		Relative intensities (100:42:70:44:27:20)	69E211
	PMMA (296 K)	LP	490 460 430			686058
	PMMA (296 K)	PS	488 ^a 461 ^a 435 ^a			68E104
	PMMA	LP	540 490 461			70E288
	Toluene (77 K)	MOD	495		Halfwidth 460 cm ⁻¹	777538
	Toluene	LP/ET	490 ^a , 15000 ± 1000 466 ^a , 10600 ^a		ϵ relative to anthracene in toluene ($\epsilon_{428.5} = 42000$ L mol ⁻¹ cm ⁻¹)	83F075
107.	7H-Benz[de]anthracen-7-one					
	1,1,2-Trichloro- trifluoroethane	LP	484 ^a		{Oxygen quenching (1.2×10^9 L mol ⁻¹ s ⁻¹)	757427
	Benzene	LP	490		Half-life = 1.0 ± 0.1 μ s	757427
	Benzene (298 K)	LP	500 460			82E204
	MeOH	LP	490 ^a		Half-life = 1.8 ± 0.1 μ s	757427
108.	Benz[a]azulene					
	Benzene	LP	430			81F275
109.	Benzene					
	Cyclohexane	LP/IV	235, 11000		{Oxygen quenching; 106 ns delay; shoulder ~310 nm ($f' \sim 0.12$; ϵ assumes $\Phi_T = 0.23$; assignment $2^3E_{2g} \leftarrow ^3B_{1u}$; oscillator strength = ~0.35	80B023
	EtOH/MeOH (113 K)	FP	429.6 ± 0.5 ~412.5 ± 1.0		{Phosphorescence decay; glass was 3:1 EtOH to MeOH; band assigned to $^3E_{2g} \leftarrow ^3B_{1u}$	69E207
110.	Benzene-<i>d</i>₆					
	3-MP/MCH (103 K)	FP	426.5 ± 0.5 ~412.5 ± 1.0		{Phosphorescence decay; glass was 1:1 3-MP to MCH; band assigned to $^3E_{2g} \leftarrow ^3B_{1u}$	69E207
	EtOH/MeOH (113 K)	FP	425.0 ± 0.5 ~407.5 ± 1.0		{Phosphorescence decay; glass was 3:1 EtOH to MeOH; band assigned to $^3E_{2g} \leftarrow ^3B_{1u}$	69E207
111.	Benzidine					
	EtOH (77 K)	MOD/KM	460, 22000 ± 5500		"4,4'-Diaminobiphenyl"	737055
	EtOH/Et ₂ O (77 K)	MOD/KM	460, 49000 ± 20000		Glass was 2:1 EtOH to Et ₂ O; temperature was not explicitly stated, but 77 K was inferred from the context	719059
	Toluene/EtOH (77 K)	MOD	461 ^a		Glass was 19:1 toluene to EtOH	719059
112.	Benzil					
	Acetonitrile/Water	LP	480 ^a		Solvent was 88:12 acetonitrile to water by volume	79E690
	Benzil (295 K)	LP	510		Triplet exciton in crystal; halfwidth 2860 cm ⁻¹	78E538
	Benzil (77 K)	LP	510		Triplet exciton in crystal; halfwidth 2320 cm ⁻¹	78E538
	Liquid paraffin	FP	486.5 380		Solvent viscosity was 0.03 N·s/m ² ; bands were assigned to different electronic transitions	58E001
	Polystyrene	PR	469 ^a		$G\epsilon_{469} = 4200^a$ L mol ⁻¹ cm ⁻¹ / (100 eV absorbed); $\tau_T = 36$ μ s	701073
113.	Benzil/Triethylamine					
	Acetonitrile/Water	LP	600 ^a		Solvent was 88:12 acetonitrile to water; possible triplet exciplex, but resembles benzil radical cation; $\tau_T = 0.055$ μ s	79E690

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
114.	Benzo[a]carbazole EtOH	FP	680 ^a 569 438 390 365		Most intense peak at 569 nm	766421
115.	Benzo[b]carbazole EtOH	FP	620 ^a 500 420 345		Most intense peak at 500 nm; shoulder around 475 nm	766421
116.	Benzo[c]carbazole EtOH	FP	528 380 352		Most intense peak at 528 nm; shoulder around 600 nm	766421
117.	Benzo[b]chrysene Benzene	MOD	570 530 450		Relative intensities (100:64:39)	71E361
	Liquid paraffin	MOD	570 450		Mull; relative intensities (100:78); shoulder at 530 nm with relative intensity 71; $\tau_T = 53 \mu s$	71E361
118.	Benzo[a]coronene 2-MTHF (77 K)	PS/ESR	588, 25500 423, 38800		\uparrow ESR; assignments $^3B_2 \leftarrow ^3B_2$, $^3A_1 \leftarrow ^3B_2$; oscillator strength = 0.28, 0.18	69E115
	EPA (77 K)	PS/SD	571 ^a , 25800 ^a 420 ^a , 37600 ^a		EPA in this work was 4:4:1 isopentane to Et ₂ O to absolute EtOH; "1,2-benzcoronene"	68E104
	PMMA (296 K)	PS	568 ^a 418 ^a			68E104
	PMMA (77 K)	PS/SD	568 ^a , 21900 ^a		Shoulders at 588 ^a , 532 ^a , and 488 ^a nm; $\tau_T = 4.3 \times 10^6 \mu s$	69E216
	PMMA (296 K)	PS/SD	568 ^a , 20300 ^a		Shoulders at 585 ^a and 538 ^a nm; $\tau_T = 2.9 \times 10^6 \mu s$	69E216
	PMMA (293 K)	CWL/KM	1090, ~200 943 826 588, 18000 568 417, 36200		Three transitions were identified with the 0-0 bands identified by their extinction coefficients; the lowest triplet state was assigned as $^3B_2^+$, and the excited states were assigned as $^3B_2^+$, $^3B_2^-$, and $^3A_1^-$ in increasing energy; the ϵ_{417} is for the compound in EPA at 77 K; oscillator strength = 0.06, 0.24, 0.19	707230
119.	Benzo[flavine PMMA (77 K)	PS	1300 1210 1100 930 800 750			65B004
120.	Benzo[a]fluorene Heptane	FP	620 ^a 600 ^a 440 ^a 405 ^a		Most intense peak at 405 nm	766421
121.	Benzo[b]fluorene Heptane	FP	620 ^a 441 419 392 377 365		Most intense peak at 441 nm; shoulder around 465 nm	766421
122.	Benzo[c]fluorene Heptane	FP	490 ^a 350 ^a		More intense peak at 490 nm; shoulder around 460 nm	766421

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
123.	Benzoic acid Isopentane/MCH	FP/COM	320, ~1000 315 ^a		†Phosphorescence decay; solvent was 5:1 isopentane to MCH; ϵ_{\max} was estimated based on Φ_T being 0.99; shoulders at 344 ^a and 304 ^a nm; delay 10 μ s; $\tau_T = (2.80 \pm 0.10) \times 10^6 \mu$ s	756077
	Liquid paraffin	FP	310		Solvent viscosity was 0.19 N·s/m ²	58E001
124.	Benzoin Benzoin (295 K)	LP	455 ^a 430 ^a 390 ^a		Triplet exciton in crystal	78E538
	Liquid paraffin	FP	476 370		Solvent viscosity was 0.19 N·s/m ² ; relative intensities (95:100)	58E001
125.	Benzo[c][1,5]naphthyridine EtOH/Et ₂ O (77 K)	PS	758 676 610 570 532 505 474 439		Solvent was 3:2 EtOH to Et ₂ O; relative intensities (14:34:43:77:94:100:77:70); $E_T = 270$ kJ mol ⁻¹	80B130
126.	Benzo[c][1,6]naphthyridine EtOH/Et ₂ O (77 K)	PS	557 525		Solvent was 3:2 EtOH to Et ₂ O; relative intensities (100:85); $E_T = 276$ kJ mol ⁻¹	80B130
127.	Benzo[c][1,7]naphthyridine EtOH/Et ₂ O (77 K)	PS	571 531 495 435		Solvent was 3:2 EtOH to Et ₂ O; relative intensities (77:100:97:65); $E_T = 269$ kJ mol ⁻¹	80B130
128.	Benzonitrile EPA (77 K)	FP	490 430		†Phosphorescence decay, molecular orbital calculations; ϵ roughly 200 L mol ⁻¹ cm ⁻¹ (method unspecified); $\tau_T = 3.5 \mu$ s	776213
129.	Benzo[<i>rst</i>]pentaphene Benzene	MOD	490 460		Relative intensities (100:44); "3,4,9,10-dibenzpyrene"; $\tau_T = 170 \mu$ s	71E361
	Cyclohexane	LP	495			70E288
	Cyclohexane	LP	714 694 676 654 637 621 613 581		370 ns delay	737463
	Liquid paraffin	MOD	490 460		Mull; relative intensities (100:45); $\tau_T = 150 \mu$ s	71E361
130.	Benzo[<i>ghi</i>]perylene Benzene	MOD	570 520 470		Relative intensities (11:23:100); "1,12-benzoperylene"; $\tau_T = 150 \mu$ s	71E361
	Cyclohexane	LP	690 667 637 575 529 483 461		370 ns delay; authors reassign "490" nm band to $T_n \leftarrow T_1$, [686058] suggested that it was $S_n \leftarrow S_1$	737463

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	EPA (77 K)	PS/SD	467, 40000 ± 4000			68E105
	PMMA (296 K)	LP	465			686058
			425			
	PMMA (77 K)	PS/SD	465 ^a , 39900 ^a			696020
	PMMA (296 K)	PS/SD	463 ^a , 38900 ^a			696020
	PMMA (293 K)	CWL/KM	1000, ~150		Four transitions were identified with the 0-0 bands identified by their extinction coefficients; the lowest triplet was ³ B ₂ ⁺ , and the other states were (³ A ₁ ⁺), ³ A ₁ ⁻ , ³ B ₁ ⁻ , and ³ A ₁ ⁻ in increasing energy; the ϵ_{568} is for the compound in EPA at 77 K; oscillator strength = ~10 ⁻³ , 0.02, >0.07, 0.39	707230
			781, ~500			
			699			
			637			
			568, ~7000			
			529			
			493			
			465, 38300			
			439			
131.	Benzo[c]phenanthrene Acetonitrile	FP	520		"3,4-Benzphenanthrene"; radical cation also observed at 450 nm; solvent contains dimethylmercury; τ_T = ~200 μ s	78A324
	Cyclohexane	LP	517			70E288
	Hexane	FP	517.1			54E001
	Hexane	FP/SD	517, 4800		Bands were assigned to 2 different electronic transitions; oscillator strength = 0.05, 0.03	58E001
			400, 1580			
	PMMA	LP	517			70E288
132.	Benzophenazine Toluene (293 K)	FP	530 ^a 500 ^a		†Phosphorescence decay in EPA at 77 K, oxygen quenching; isomer unspecified	80E778
133.	Benzophenone 2-PrOH	LP	535		τ_T = 0.046 ± 0.006 μ s	70E288
	Acetonitrile	LP	520 330 ^a		†Triplet ET to <i>cis</i> -3-methylpent-2-ene; relative intensities (1:2); τ_T = 0.71 μ s; k_{et} = 2.7 × 10 ⁸ L mol ⁻¹ s ⁻¹	80B087
	Acetonitrile	LP/HAT	520, 6500 ± 400 320, ~11500		ϵ relative to benzophenone ketyl radical in water (ϵ_{540} = 3220 L mol ⁻¹ cm ⁻¹) assuming ϵ_{ref} unchanged in acetonitrile; hydrogen atom transfer from 2-piperidone (k_{HAT} = 9.7 × 10 ⁶ L mol ⁻¹ s ⁻¹); τ_T = 14 μ s	80E416
	Acetonitrile	LP	525		20 ps delay	80A206
	Acetonitrile/Water	LP/RF	520, 6500 315, 11800		Solvent was 9:1 acetonitrile to water; ϵ relative to benzophenone in acetonitrile (ϵ_{520} = 6500 L mol ⁻¹ cm ⁻¹)	84B033
	Benzene	FP/HAT	525 ^b , 6000 ± 3000		†Specific quenching by heavy metal chelates; ϵ relative to benzophenone ketyl radical in benzene (ϵ_{525} = 7000 L mol ⁻¹ cm ⁻¹); ϵ calculated by estimating that 10% of the triplets form ketyl radicals; τ_T = 10 ± 5 μ s	63E007
	Benzene	PR	530		$G\epsilon_{530}$ = 11700 L mol ⁻¹ cm ⁻¹ / (100 eV absorbed); half-life = 2.11 μ s	64B006
	Benzene	PR/ET	532.5, 9100		ϵ relative to anthracene in cyclohexane (ϵ_{420} = 57200 L mol ⁻¹ cm ⁻¹); author reported mean of 4 measurements (this one and 3 others with different ref. cmpds.) as 10300 ± 2570 L mol ⁻¹ cm ⁻¹	680727
	Benzene	PR/ET	532.5, 12000		ϵ relative to naphthalene in cyclohexane ($\epsilon_{412.5}$ = 22600 L mol ⁻¹ cm ⁻¹); author reported mean of 4 measurements (this one and 3 others with different ref. cmpds.) as 10300 ± 2570 L mol ⁻¹ cm ⁻¹	680727
	Benzene	PR/ET	532.5, 9700		ϵ relative to benz[a]anthracene in cyclohexane (ϵ_{480} = 25100 L mol ⁻¹ cm ⁻¹); author reported mean of 4 measurements (this one and 3 others with different ref. cmpds.) as 10300 ± 2570 L mol ⁻¹ cm ⁻¹	680727
	Benzene	PR/ET	532.5, 10600		ϵ relative to phenanthrene in cyclohexane ($\epsilon_{482.5}$ = 21000 L mol ⁻¹ cm ⁻¹); author reported mean of 4 measurements (this one and 3 others with different ref. cmpds.) as 10300 ± 2570 L mol ⁻¹ cm ⁻¹	680727

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
Benzene	LP		535	$\tau_T = 2.5 \pm 0.3 \mu s$		70E288
Benzene- <i>d</i> ₆	LP		535	$\tau_T = 3.45 \pm 0.30 \mu s$		70E288
Benzene	PR/ET		532.5, 7630	ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700$ L mol ⁻¹ cm ⁻¹); ϵ obtained from a simultaneous least squares fit of data from several compounds making use of cyclohexane to benzene ϵ_{\max} ratios of 1.83 for naphthalene and 1.45 for anthracene		71E360
Benzene	FP/RA		525, 7000	ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 4160$ L mol ⁻¹ cm ⁻¹), taking $\Phi_T = 1$ in benzene and $\Phi_{\text{ketyl}} = 1$ in cyclohexane; reference ϵ of ketyl radical was taken as the average of 3220 in water and 5100 L mol ⁻¹ cm ⁻¹ in 2-PrOH/water; $\tau_T = 12 \pm 2 \mu s$		717009
Benzene	PR		528 ^a	† Triplet ET from benzene; $k_{et} = (1.3 \pm 0.2) \times 10^{11}$ L mol ⁻¹ s ⁻¹		751124
Benzene	LP		528 ^a			751124
Benzene	LP		535	$\tau_T = 3.57 \mu s$		79A171
Benzene	LP		525			80B021
			325			
Benzene	LP/RA		533, 7200	ϵ relative to potassium ferrioxalate actinometer assuming $\Phi_T = 1.0$ for benzophenone in benzene		80B090
Benzene	LP/ET		533 ^a , 7100 \pm 800	† Triplet ET to naphthalene; ϵ relative to naphthalene in benzene ($\epsilon_{425} = 13000$ L mol ⁻¹ cm ⁻¹)		81A174
Benzene	LP/PIV		530, 7220 \pm 320			83B067
Benzophenone (303 K)	PR		540	Liquid; if 90% of the absorption at 540 nm is due to triplets and if $\epsilon = 10300$ L mol ⁻¹ cm ⁻¹ , then $G(\text{triplets}) = 2.2 \pm 0.1$ triplets/(100 eV absorbed); half-life = 0.46 μs		700114
Benzophenone (77 K)	LP		543	Triplet exciton in crystal; halfwidth of 543 nm band		78E538
			450	2450 cm ⁻¹ ; $\tau_T = \sim 1000 \mu s$		
Benzophenone (295 K)	LP		543	Triplet exciton in crystal; halfwidth 3000 cm ⁻¹		78E538
Benzophenone	LP		540 ^a	† Phosphorescence decay; absorption monitored by diffuse-reflectance spectroscopy		84E044
Cyclohexane	LP		535	$\tau_T = 0.30 \pm 0.02 \mu s$		70E288
Cyclohexane	PR		535 ^a			751124
Dioxane	LP		535	$\tau_T = 0.20 \pm 0.02 \mu s$		70E288
Diphenyl ether	LP		~ 540	$\tau_T = (160 \pm 20) \times 10^{-3} \mu s$		78B071
EPA (77 K)	FP		535			55E002
EtOH	LP		535	$\tau_T = 0.104 \pm 0.015 \mu s$		70E288
EtOH	LP/COM		545, 7300	ϵ estimated by scaling by halfwidth of band (3220 cm ⁻¹) relative to that in hexane; $\tau_T = 0.08 \mu s$		79B141
			510 ^a , 6500 ^a			
Heptane	LP		530	Delay 320 ps		84B154
Hexafluorobenzene	LP		535	$\tau_T = 0.435 \pm 0.40 \mu s$		70E288
Hexane	LP/IV		525, 10300	ϵ estimated assuming $\Phi_T = 1.0$ for benzophenone in hexane, corrected for triplet decay and ketyl radical formation during the pulse, and extrapolated to zero intensity; halfwidth 2650 cm ⁻¹ ; $\tau_T = 0.29 \mu s$		79B141
			500 ^a , 9600 ^a			
			445 ^a , 5000 ^a			
			310 ^a , 15000 ^a			
Isopentane (77 K)	FP		543 ^a	† Phosphorescence decay		67E105
			326 ^a			
MCH	LP		529 ^a			757112
PFMCH	FP		530 ^a	† Phosphorescence decay; most intense peak was at 319 nm; $\tau_T = 400 \mu s$		747390
			520 ^a			
			319 ^a			
PMMA (293 K)	FP/RA		606	† Phosphorescence decay; ϵ relative to benzophenone ketyl radical in 2-PrOH/water ($\epsilon_{346} = 5100$ L mol ⁻¹ cm ⁻¹), assuming Φ_T was the same in PMMA and ethylene glycol and assuming ϵ for ketyl radical same in 2-PrOH/water and ethylene glycol; delay 100 μs		66E090
			532, 4200 \pm 100			
PMMA (80 K)	FP		606	† Decay of absorption was equal to decay of phosphorescence at 293, 220, and 100 K; delay 100 μs		66E090
			532			

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	PPFO	LP/SM	520, 2460 ^a ± 350		Aqueous micelle, potassium perfluorooctylsulfonate; ϵ method needs the assumption of $\Phi_T = 1$ for benzophenone in the micelle since only linear intensity dependence observed; $\tau_T = 7.87 \mu\text{s}$	81N156
	SPFO	LP/SM	520, 2660 ^a ± 380		†Phosphorescence decay; aqueous micelle, sodium perfluorooctanoate; ϵ method needs the assumption of $\Phi_T = 1$ for benzophenone in the micelle since only linear intensity dependence observed; $\tau_T = 7.87 \mu\text{s}$	81N156
	Trichlorotrifluoroethane	FP	526 317		†Oxygen quenching and no hydrogens available in the solvent to form ketyl radicals; $\tau_T = 20 \mu\text{s}$	67B006
	Water	FP	525 315		pH 12.7	727098
	Water	FP	525		Delay 20 μs ; $\tau_T = 65 \pm 6 \mu\text{s}$	72A022
	Water	FP	525 315		$\tau_T = \sim 100 \mu\text{s}$; pH Neutral	727098
	Water	LP/HAT	520, 5750 ± 400 320, ~8350		ϵ relative to benzophenone ketyl radical in water ($\epsilon_{540} = 3220 \text{ L mol}^{-1} \text{ cm}^{-1}$); hydrogen atom transfer from 2-piperidone ($k_{\text{HAT}} = 14.8 \times 10^6 \text{ L mol}^{-1} \text{ s}^{-1}$); $\tau_T = 40 \mu\text{s}$	80E416
	Water	LP/RA	520, 5800 ± 400		ϵ relative to naphthalene in cyclohexane ($\epsilon_{414} = 24500 \text{ L mol}^{-1} \text{ cm}^{-1}$, $\Phi_T = 0.75$) assuming $\Phi_T = 1$ for benzophenone in water	80E416
	Water/Acetonitrile	LP	525		Solvent was 4:1 water to acetonitrile; 200 ns delay; $\tau_T = 31 \mu\text{s}$; $E_T = 288 \text{ kJ mol}^{-1}$	82A082
134.	Benzophenone, conjugate acid Sulfuric acid	LP	505		5 mol L ⁻¹ H ₂ SO ₄ ; solvent was 4:1 water to acetonitrile; pK_a 0.20 ± 0.02; $\tau_T = 0.017 \mu\text{s}$	84E456
135.	Benzo[a]pyrene/DNA Water	LP	480		†Oxygen quenching ($2.0 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$); deproteinated calf thymus DNA; $\tau_T = 3.5 \times 10^4 \mu\text{s}$	766403
136.	Benzo[e]pyrene/DNA Water	LP	560 ^a		†Oxygen quenching ($1.6 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$); deproteinated calf thymus DNA; $\tau_T = 1.55 \times 10^4 \mu\text{s}$	766403
137.	Benzo[a]pyrene Benzene	MOD	475		Called "3,4-benzpyrene" in paper; $\tau_T = 84 \mu\text{s}$	71E361
	Cyclohexane	LP	500 475		†Rise time of transient was the same as the decay time of the singlet; "3,4-benzpyrene"	68B006
	Cyclohexane (296 K)	FP	465		Shoulder at 440 nm; $\tau_T = 8700 \mu\text{s}$	78A345
	EPA (77 K)	PS	504 477.3 467.9 444		Relative intensities (31.95:100:76); "3,4-benzopyrene"	54D001
	PMMA (77 K)	PS	470 442 419		Named "3,4-benzopyrene" in paper	70E291
	Toluene/EtOH (77 K)	MOD	557 ^a		Glass was 19:1 toluene to EtOH; shoulders at 606 ^a , 427 ^a , and 380 ^a nm; called "1,2-benzopyrene" in paper	719059
	Water (296 K)	FP	465		Solution contains 2% SDS; shoulder at 445 nm; radical cation also observed ($\lambda_{\max} = 550 \text{ nm}$); $\tau_T = 2.38 \times 10^4 \mu\text{s}$	78A345
	Water (296 K)	FP	470		Caffeine (1%) added as solubilizing agent; radical cation also observed ($\lambda_{\max} = 550 \text{ nm}$); $\tau_T = 2040 \mu\text{s}$	78A345
138.	Benzo[e]pyrene Benzene	MOD	560		Called "1,2-benzpyrene" in paper; $\tau_T = 120 \mu\text{s}$	71E361
	EPA (77 K)	PS/IV	555, 17800		λ_{\max} assumed from previous work; ϵ estimated by extrapolation to infinite excitation rate	69E212
	EPA (77 K)	PS/KM	588 ^a , 7500 ^a 555 ^a , 16100 ^a 519 ^a , 9300 ^a		Solvent, temperature and extinction method assumed from earlier work; polarization also measured	69E213

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	EtOH/MeOH/THF (assumed 77 K)	PS	587.5 558.5 456		Glass was 3:1:4 EtOH to MeOH to THF; the origin of another band was assigned to 392 nm; the 4 bands were assigned to 4 different electronic transitions; T ₁ was assigned as ³ B ₂ ⁺ , and the other 4 states were assigned as ³ A ₁ ⁻ for 587.5 nm and 456 nm and ³ B ₂ ⁻ for the other 2; "1,2-benzopyrene"	70B003
139.	Benzo[f]quinoline EPA (77 K)	PS	517 482 ~452		Relative intensities (10:8:5); "5,6-benzoquinoline"; $\tau_T = 2.6 \times 10^6 \mu s$	54B001
	EtOH/Et ₂ O (77 K)	PS	833 735 662 515 448		Solvent was 3:2 EtOH to Et ₂ O; relative intensities (<5:<5:<5:100:74:49); $E_T = 262 \text{ kJ mol}^{-1}$	80B130
	Isopentane/MCH (77 K)	PS	510		$\tau_T = 1.8 \times 10^6 \mu s$	54B001
	PMMA (77 K)	PS	500 470 440		†Phosphorescence decay (?); $\tau_T = 3.3 \times 10^6 \mu s$	70E291
140.	Benzo[h]quinoline EPA (77 K)	PS	500 465 ~440		Relative intensities (10:8:5); "7,8-benzoquinoline"; $\tau_T = 2.1 \times 10^6 \mu s$	54B001
	EtOH/Et ₂ O (77 K)	PS	500 465 437		Solvent was 3:2 EtOH to Et ₂ O; relative intensities (100:84:55); $E_T = 262 \text{ kJ mol}^{-1}$	80B130
	Isopentane/MCH (77 K)	PS	505 ~465		Relative intensities (10:9); $\tau_T = 1.4 \times 10^6 \mu s$	54B001
	PMMA (293 K)	FP	490 460		†Phosphorescence decay (?); decay was non-exponential, and lifetime was calculated from a terminal 1st-order rate constant; $\tau_T = 1.1 \times 10^6 \mu s$	70E291
	PMMA (77 K)	PS	490 460 433		†Phosphorescence decay (?); $\tau_T = 1.9 \times 10^6 \mu s$	70E291
141.	Benzo[f]quinolizinium Acetonitrile	FP	575*		$\tau_T = 125 \pm 5 \mu s$	81A345
142.	Benzo[h]quinolizinium Acetonitrile	FP	575*		$\tau_T = 120 \pm 5 \mu s$	81A345
143.	1,4-Benzquinone EtOH Water	LP LP	450 410		100 ps delay †Oxygen quenching ($2.4 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); $\tau_T = 0.53 \mu s$	79B007 80B112
144.	Benzo[b]triphenylene/Chloranil PMMA	PS	620 540 450* 420* 410* 400*		Charge transfer complex; most intense peak at 450 nm	766652
145.	Benzo[b]triphenylene 2-MTHF (77 K)	PS/ESR	454, 31000		†ESR; only most intense visible peak reported; "1,2,3,4-dibenzanthracene"; oscillator strength = 0.20	696115
	Benzene	MOD	450		$\tau_T = 90 \mu s$	71E361
	Cyclohexane	LP	440		†Rise time of transient was the same as the decay time of the singlet	68B006
	Epoxy plastic (296 K)	LP	425 412			686058

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	PMMA	LP	445			70E288
	PMMA (293 K)	CWL/KM	752, 1300		Two transitions were assigned with the 0-0 bands identified by their extinction coefficients; both transitions are assigned as $^3B_2^- \leftarrow ^3B_2^+$; oscillator strength = 0.10, 0.27	707230
			680			
			613			
			568			
			446, 28800			
			422			
			400			
	PMMA	PS	612 ^a		Most intense peak at 455 nm	766652
			596 ^a			
			578 ^a			
			540 ^a			
			498 ^a			
			455 ^a			
			425 ^a			
			412 ^a			
			396 ^a			
146.	1-Benzoylacetone					
	EtOH/MeOH (118 K)	FP	438		Solvent was 3:1 EtOH to MeOH	68B005
147.	Benzoylamino-2- Δ^2 -thiazoline					
	Cyclohexane (293 K)	FP/RA	551, 7200		\uparrow Triplet ET to biacetyl, oxygen quenching; ϵ relative to naphthalene in cyclohexane ($\epsilon_{414} = 24500$ L mol ⁻¹ cm ⁻¹); $\tau_T = 1000$ μ s	82E373
148.	9-Benzoylanthracene					
	Benzene	LP	434			84B110
			408			
	EPA (77 K)	FP	430		430 nm peak was the more intense; $\tau_T = 3.31 \times 10^4$ μ s	82E338
			406 ^a			
	Toluene	LP	420		\uparrow Oxygen quenching; rise time of 0.02 ± 0.004 ns	777635
149.	9-Benzoyl-10-bromoanthracene					
	EPA (77 K)	FP	430		$\tau_T = 170$ μ s	82E338
150.	9-Benzoyl-10-chloroanthracene					
	EPA (77 K)	FP	432		$\tau_T = 3800$ μ s	82E338
	Toluene	LP	415		\uparrow Oxygen quenching; rise time of 0.052 ± 0.002 ns	777635
151.	9-Benzoyl-10-cyanoanthracene					
	Toluene	LP	430		\uparrow Oxygen quenching	777635
152.	2-Benzoyl-N-methyl- β -naphthiazoline					
	EtOH	FP	630		\uparrow Oxygen quenching (6.6×10^{10} L mol ⁻¹ s ⁻¹); 50 μ s delay; $\tau_T = 480$ μ s	78E534
	THF	FP/SD	640, 16000		\uparrow Oxygen quenching; 50 μ s delay; $\tau_T = 380$ μ s	78E534
			620 ^b , 15000			
			440, 8000 ^a			
	THF	FP/SD	640, 16000		\uparrow Oxygen quenching (6.6×10^{10} L mol ⁻¹ s ⁻¹); 50 μ s delay; $\tau_T = 390$ μ s	79E304
			440, 13000			
153.	8-Benzoylnaphtho[de-2,3,4]bicyclo[3.2.2]nona-2,6,8-triene					
	EPA (103 K)	LP	430		\uparrow Phosphorescence decay, oxygen quenching	81F390
			380			
	MCH (173 K)	LP	430		\uparrow Phosphorescence decay, oxygen quenching; $\tau_T = 1.0$ μ s	81F390
			380			
154.	9-Benzoyl-10-nitroanthracene					
	Benzene	LP	431 ^a		Delay 400 ns	80E884
	Benzene (293 K)	LP	~450		170 ps delay; growth followed at 452 nm; rise time of $(82 \pm 6) \times 10^{-3}$ ns	80B084
	EPA (77 K)	FP	~450		$\tau_T = 1.6 \times 10^4$ μ s	80B084
	EPA (77 K)	FP	436 ^a		$\tau_T = 1.6 \times 10^4$ μ s	80E884
	EPA (77 K)	LP	435 ^a			84F385

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
155.	2-Benzoylpyridine PFMCH	FP	526 ^a 331 ^a		Shoulder at 539 ^a nm; 331 nm peak was the more intense; $\tau_T = 480 \mu\text{s}$	747390
156.	3-Benzoylpyridine PFMCH	FP	523 ^a 305 ^a		\dagger Phosphorescence decay; 305 nm peak was the more intense; $\tau_T = 110 \mu\text{s}$	747390
157.	4-Benzoylpyridine PFMCH	FP	510 ^a 490 ^a 319 ^a		\dagger Phosphorescence decay; most intense peak at 319 nm; "phenyl-4-pyridyl ketone"; $\tau_T = 240 \mu\text{s}$	747390
158.	1-Benzoyl-3,3,3-trifluoroacetone EtOH/MeOH (118 K)	FP	633		Solvent was 3:1 EtOH to MeOH	68B005
159.	Benzyl 9-anthroate Acetonitrile	FP	425		Radical anion ($\lambda_{\max} = 550 \text{ nm}$) also observed; $\tau_T = 1100 \mu\text{s}$	80A030
	Benzene	FP/?	430, 63000 410, 28000 ^a		ϵ method unspecified; $\tau_T = 6100 \mu\text{s}$	80A030
160.	1-Benzylisoquinoline- <i>N</i> -oxide EtOH	FP	392		$\tau_T = 2.9 \mu\text{s}$	727105
161.	Benzyl phenyl ketone, conjugate acid Sulfuric acid	LP	359 ^a		5 mol L ⁻¹ H ₂ SO ₄ ; solvent was 4:1 water to acetonitrile; pK _a 0.35 \pm 0.05	84E456
162.	3-Benzyl-3-phenylphenanthro[9,10- <i>b</i>]furan-2(3 <i>H</i>)-one Benzene	LP	490 ^a		\dagger Oxygen quenching and triplet ET to 1-methylnaphthalene	84E520
163.	3-Benzyl-3,4,5-triphenyl-2(3 <i>H</i>)-furanone Benzene	LP/ET	375 \pm 5, 14000 \pm 2800		\dagger Oxygen quenching; ϵ relative to benzophenone in benzene ($\epsilon_{332} = 7600 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 8.2 \pm 1.2 \mu\text{s}$; $k_{et} = 2.7 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	84E520
164.	Biacetyl 2-PrOH	FP	1060 910 800 725 330 304 210		\dagger Oxygen quenching; delay 0 μs	697092
	Benzene	PR	317		\dagger Phosphorescence half-life; half-life = 120 μs	680386
	Benzene	PR/ET	315, 6200		\dagger Phosphorescence decay; ϵ relative to benz[a]anthracene in cyclohexane ($\epsilon_{480} = 25100 \text{ L mol}^{-1} \text{ cm}^{-1}$); author reported the mean of 2 measurements with this and another ref. compd. as 6400 \pm 1600 L mol ⁻¹ cm ⁻¹	680727
	Benzene	PR/ET	315, 6600		\dagger Phosphorescence decay; ϵ relative to anthracene in cyclohexane ($\epsilon_{420} = 57200 \text{ L mol}^{-1} \text{ cm}^{-1}$); author reported the mean of 2 measurements with this and another ref. compd. as 6400 \pm 1600 L mol ⁻¹ cm ⁻¹	680727
	Benzene	PR/ET	315, 5160		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700 \text{ L mol}^{-1} \text{ cm}^{-1}$); ϵ obtained from a simultaneous least squares fit of data from several compounds making use of cyclohexane to benzene ϵ_{\max} ratios of 1.83 for naphthalene and 1.45 for anthracene	71E360

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	CCl ₄	FP	1056 916 796 720 330 310 220		†Oxygen quenching	697092
	EtOH	LP/RF	740, 1760		ϵ relative to biacetyl in 2-PrOH ($\epsilon_{725} = 1710$ L mol ⁻¹ cm ⁻¹); $\tau_T = 3.4 \pm 1.0$ μ s	84E162
	Liquid paraffin	FP	317		Solvent viscosity was 0.03 N·s/m ²	58E001
165.	9,9'-Bianthryl MeOH	LP	425			776431
166.	9,9'-Bicarbazole Cyclohexane	FP	410 ^a 370 ^a		†Triplet ET to anthracene; carbazyl radicals ($\lambda_{\max} = 600$ nm) also observed; $\tau_T = 120$ μ s	78A368
	Hexane	FP	420		†Oxygen and piperylene quenching; "dicarbazyl"	757573
167.	Bifluorenylidene Liquid paraffin	FP/TD	440, ≥ 110000		†Phosphorescence decay in MCH/Isopentane glass at 77K; solvent was "viscous paraffin", viscosity 1.75 P; $\tau_T = (2.2 \pm 0.2) \times 10^3$ μ s	767488
	MCH/Isopentane (77 K)	FP	440		†Phosphorescence decay; glass was 1:1 MCH to isopentane by volume; $\tau_T = (0.94 \pm 0.03) \times 10^6$ μ s	767488
168.	Bilirubin Benzene	PR/ET	500, 8800		†Triplet ET from biphenyl and to β -carotene, oxygen quenching (8.2×10^8 L mol ⁻¹ s ⁻¹); ϵ relative to biphenyl in benzene ($\epsilon_{365} = 27100$ L mol ⁻¹ cm ⁻¹); half-life = 9 μ s; $E_T = \sim 150$ kJ mol ⁻¹ ; $k_{et} = 4.4 \times 10^9$ L mol ⁻¹ s ⁻¹	761168
169.	Biliverdin Acetone	PR-ET/ET	700 ^a , 16000 ^a		†Triplet ET from biphenyl; ϵ relative to biphenyl in benzene ($\epsilon_{360} = 27100$ L mol ⁻¹ cm ⁻¹), assuming ϵ_{ref} independent of solvent; more intense maximum below 400 nm not observed; half-life = 11.7 μ s; $E_T = \sim 90$ kJ mol ⁻¹ ; $k_{et} = 7.4 \times 10^9$ L mol ⁻¹ s ⁻¹	79E136
170.	Biliverdin, dimethyl ester Benzene	PR-ET/ET	800 ^a , 14000 ^a 400 ^a , 55000 ^a		†Triplet ET from biphenyl; ϵ relative to biphenyl in benzene ($\epsilon_{360} = 27100$ L mol ⁻¹ cm ⁻¹); half-life = 6.7 μ s; $E_T = \sim 90$ kJ mol ⁻¹ ; $k_{et} = 1.05 \times 10^{10}$ L mol ⁻¹ s ⁻¹	79E136
171.	1,1'-Binaphthyl 2-MTHF	LP-ET	610 ^a		†Triplet ET from benzoquinone, oxygen quenching, triplet quenching; 2 μ s delay; shoulder at 550 nm	777241
	2-MTHF (123 K)	LP-ET	610 ^a 530 ^a 415 ^a 395 ^a		†Oxygen quenching, triplet quenching; 10 μ s delay	777241
	2-MTHF (83 K)	LP-ET	530 ^a 415 ^a 395 ^a		†Oxygen quenching, triplet quenching; 14 ms delay	777241
	Benzene (299 K)	PR/ET	615 ^b , 12000 608 ^a		Shoulder at 407 ^a nm; ϵ relative to benzophenone in benzene ($\epsilon_{332.5} = 7630$ L mol ⁻¹ cm ⁻¹)	741006
	Toluene	PR	605		†Triplet ET to fluorenone, oxygen quenching; 2 μ s delay; $\tau_T = 14$ μ s	771048
172.	2,2'-Binaphthyl Benzene (299 K)	PR/ET	637 ^a 601 ^a 450 ^b , 24000 440 ^a		Shoulder at 554 ^a nm; ϵ relative to benzophenone in benzene ($\epsilon_{332.5} = 7630$ L mol ⁻¹ cm ⁻¹); 440 nm was the most intense peak	741006

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
173.	Biphenyl					
	2-PrOH/Isooctane/Et ₂ O (81 K)	FP	371 ^a		‡Phosphorescence decay; glass was 1:3:3 2-PrOH to isooctane to Et ₂ O	81B119
	3-MH (77 K)	PR	375		‡Phosphorescence decay; $\tau_T = 3.7 \times 10^6 \mu\text{s}$	771059
	3-MP	PR	360		$G\epsilon_{360} = 16600 \text{ L mol}^{-1} \text{ cm}^{-1} / (100 \text{ eV absorbed})$; half-life = 3.4 μs	64B006
	3-MP (77 K)	PS/ESR	375, 28200		‡ESR; oscillator strength = ~ 0.27	69B002
	Acetone	PR	360			710186
	Alcohol/Ether (77 K)	MOD	370		Glass was 2:1 alcohol to ether	76E682
	Benzene	PR/ET	359, 27100		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700 \text{ L mol}^{-1} \text{ cm}^{-1}$); ϵ obtained from a simultaneous least squares fit of data from several compounds making use of cyclohexane to benzene ϵ_{\max} ratios of 1.83 for naphthalene and 1.45 for anthracene; corrected λ_{\max} quoted from reference [78Z194]	71E360
	Benzene	PR	370			761024
	Boric acid	PS	360		Glass	777388
	BuOH (~ 80 K)	FP/COM	363, 300 ^d		ϵ estimated from numerical simulation of triplet state kinetics; $\tau_T = (3.93 \pm 0.1) \times 10^6 \mu\text{s}$	67B010
	BuOH (~ 80 K)	FP/TV	363, 250 ^d		ϵ estimated by extrapolation to infinite excitation rate	67B010
	Cyclohexane	PR/ET	361.3, 33000		ϵ relative to anthracene in cyclohexane ($\epsilon_{420} = 57200 \text{ L mol}^{-1} \text{ cm}^{-1}$); author reported mean of 2 measurements (this one and another with a different ref. compd.) as $35400 \pm 8850 \text{ L mol}^{-1} \text{ cm}^{-1}$	680727
	Cyclohexane	PR/ET	361.3, 37700		ϵ relative to benz[a]anthracene in cyclohexane ($\epsilon_{480} = 25100 \text{ L mol}^{-1} \text{ cm}^{-1}$); author reported mean of 2 measurements (this one and another with a different ref. compd.) as $35400 \pm 8850 \text{ L mol}^{-1} \text{ cm}^{-1}$	680727
	Cyclohexane	PR/ET	361.3, 42800		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700 \text{ L mol}^{-1} \text{ cm}^{-1}$); reference ϵ obtained by starting from $\epsilon_{\max} = 3220 \text{ L mol}^{-1} \text{ cm}^{-1}$ for this ketyl radical in water and assuming the f of the ketyl radical is independent of solvent; final ϵ obtained from a simultaneous least squares fit to data from several compounds	71E360
	EPA (77 K)	PS/KM	630 ^a , 1600 ^a 577 ^a , 2000 ^a 529 ^a , 1800 ^a 488 ^a , 2100 ^a 458 ^a , 1900 ^a 428 ^a , 1500 ^a 370, 35800 357 ^a , 27000 ^a		Two electronic bands, the 2nd starting at 370 nm; ϵ -method assumed on basis of earlier work by authors; oscillator strength = 0.07, 0.42	747347
	EtOH/Et ₂ O (77 K)	MOD/KM	365, 7000 ^c \pm 2100		Glass was 2:1 EtOH to Et ₂ O; temperature was not explicitly stated, but 77 K was inferred from the context	719059
	EtOH/MeOH (113 K)	FP/SD	625 ^a , 500 ^a 571 ^a , 800 ^a 526 ^a , 750 ^a 484, 1200 ^a 452 ^a , 900 ^a 424 ^a , 600 ^a 370, 50000 \pm 10000 352 ^a , 30000 ^a		‡Phosphorescence decays; shoulders at 685 and 334 nm; glass was 3:1 EtOH to MeOH; three electronic states were tentatively assigned as $^3A_g^+$, $^3B_{3g}$, and $^3A_g^-$ for the 0-0 bands of 685, 484, and 370 nm, respectively; T^1 was $^3B_{1u}^+$	70E290
	Ethyl acetate	PR	360		0.8 μs delay; $G(\text{triplets})$ estimated assuming ϵ independent of solvent	761080
	Hexane (300 K)	MOD/SD	360, 38500		$\tau_T = 130 \pm 5 \mu\text{s}$	69E208
	Liquid paraffin	FP	368.5		Solvent viscosity was 0.03 N·s/m ² ; relative intensities (100:69)	58E001
	Liquid paraffin (290 K)	PR	363		Delay 20 μs ; $\tau_T = 85 \mu\text{s}$	700277
	PMMA	PR	370			692001
	PVA	FP	365		Stretched polymer film; polarized spectrum obtained with some structure	776279

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	PVA (81 K)	FP	373 ^a		†Phosphorescence decay	81B119
	Polystyrene	PR	375		$G\epsilon = 11000 \text{ L mol}^{-1} \text{ cm}^{-1} / (100 \text{ eV absorbed})$; $\tau_T = 2 \times 10^4 \mu\text{s}$	672086
	Polystyrene (77 K)	COM	370		Method was "photolysis"	692001
	Polystyrene	PR	385 ^a		$G\epsilon_{385} = 6000^b \text{ L mol}^{-1} \text{ cm}^{-1} / (100 \text{ eV absorbed})$	701073
	Toluene/EtOH (77 K)	MOD	366 ^a		Glass was 19:1 toluene to EtOH	719059
174.	Biphenyl- <i>d</i> ₁₀ EtOH/Et ₂ O (77 K)	MOD/KM	365, 7000 ± 2100		Glass was 2:1 EtOH to Et ₂ O; temperature was not explicitly stated, but 77 K was inferred from the context	719059
175.	4-Biphenylcarboxaldehyde Alcohol/Ether (77 K)	MOD	440		Glass was 2:1 alcohol to ether	76E682
176.	Biphenylene Benzene	PR/ET	350, 10000		†Triplet ET from naphthalene and triplet ET to anthracene; ϵ relative to naphthalene in benzene ($\epsilon_{\max} = 13200 \text{ L mol}^{-1} \text{ cm}^{-1}$); half-life = 80 μs	720464
	Cyclohexane	FP-ET/SD	339, 10000		†Triplet ET from naphthalene and triplet ET to anthracene; $\tau_1 = 100 \mu\text{s}$; oscillator strength = 0.13	720464
177.	2-Biphenylphenylbenzoxazole Benzene	LP/SD	560, 45000		†Oxygen quenching ($1.5 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); 100 ns delay; "PBBO"; $\tau_T = 0.325 \mu\text{s}$	777265
178.	2-(4-Biphenyl)benzoxazole Pentane	LP/TD	460, 2500 ± 250		Shoulders at 536 ^a , 495 ^a , and 405 ^a nm; $\tau_T = 0.70 \pm 0.07 \mu\text{s}$	82E632
179.	1-(4-Biphenyl)-3-chloro-1-propanone Benzene	FP	425		†Physical quenching by free radical; $\tau_T = 140 \mu\text{s}$	83P212
	CCl ₄	FP	425		†Physical quenching by free radical; $\tau_T = 90 \mu\text{s}$	83P212
180.	1-(2-Biphenyl)-1-phenylethylene Cyclohexane	LP/?	370, 17000 ± 3000		†Oxygen quenching; $\tau_T = 1.75 \mu\text{s}$	78E448
181.	2-(4-Biphenyl)-5-phenyl-1,3,4-oxadiazole Benzene	LP	510		†Oxygen quenching ($1.0 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); 100 ns delay; $\tau_T = 0.460 \mu\text{s}$	777265
182.	2-(4-Biphenyl)-5-phenyloxazole Dioxane	LP/TD	530, 61000 ± 250 498 ^a 455 ^a 426 ^a		$\tau_T = 0.62 \pm 0.06 \mu\text{s}$	82E632
	Pentane	LP/TD	540, 71500 ± 250 499 ^a 417 ^a		$\tau_T = 0.35 \pm 0.04 \mu\text{s}$	82E632
183.	5',1'-[1,1'-Biphenyl]-4-yl-1,1':4',1'':3'',1'':4''',1''':5''''-quinquephenyl 2-PrOH/Isooctane/Et ₂ O (81 K)	FP	576 ^a 533 ^a 498 ^a 455 ^a 349 ^a		†Phosphorescence decay; glass was 1:3:3 2-PrOH to isooctane to ether; most intense peak at 455 nm; "1,3,5-tris(biphenyl-4-yl)benzene"; $\tau_T = 2.2 \times 10^6 \mu\text{s}$	81B119
	Alcohol/Ether (77 K)	MOD	572 446		Glass was 2:1 alcohol to ether	76E682
184.	2,2'-Biquinoline Water/EtOH	FP	600 ^a 460 ^a 420 ^a		Solvent was 3:2 water to EtOH by volume; 20 μs delay; pH 7.5	79E799

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
185.	6,6'-Biquinoline, conjugate diacid Water/EtOH	FP	590 ^a 450 ^a		Relative intensities (1:2); solvent was 2:1 water to EtOH by volume; 20 μ s delay; $pK_a = 3.8 \pm 0.5$; pH 1.2	79E799
186.	2,2'-Biquinoline, conjugate monoacid Water/EtOH	FP	610 ^a 415 ^a		Solvent was 3:2 water to EtOH by volume; 20 μ s delay; $pK_a = 5.8 \pm 0.5$; pH 0.4	79E799
187.	6,6'-Biquinoline, conjugate monoacid Water/EtOH	FP	580 ^a 410 ^a		Relative intensities (1:2); solvent was 3:2 water to EtOH; 20 μ s delay; $pK_a = 5.6 \pm 0.5$; pH 5.0	79E799
188.	Bis(2,2'-bipyridine)dichloroiridium(III) ion DMF/Water	LP	577 ^a 554 ^a 490 ^a 470 ^a 458 ^a 442 ^a 431 ^a		Shoulders at 528 ^a and 509 ^a nm; solvent was 9:11 dimethylformamide to water	79B098
	DMF/Water	LP	600 ^a 565 ^a 505 ^a 469 ^a 454 ^a		Shoulders at 534 ^a , 439 ^a , and 419 ^a nm; solvent was 19:1 dimethylformamide to water	79B098
	Water	LP	488 ^a 468 ^a 459 ^a 439 ^a		Shoulder at 419 ^a nm	79B098
189.	2,5-Bis(5- <i>tert</i> -butyl-2-benzoxazolyl)thiophene Benzene	LP/SD	520, 160000		\uparrow Oxygen quenching (2.1×10^9 L mol ⁻¹ s ⁻¹); 100 ns delay; "BBOT"; $\tau_T = 0.240$ μ s	777265
190.	4,4'-Bis(dimethylamino)benzophenone EtOH	FP	500 ^a 405 ^a		\uparrow Triplet ET to naphthalene, oxygen quenching; decay measured at 500 nm, lifetime = 25 μ s in cyclohexane; "Michler's ketone"; $\tau_T = 20$ μ s; $k_{et} = (9.9 \pm 0.1) \times 10^9$ L mol ⁻¹ s ⁻¹	777603
191.	6,6'-Bis(dimethylamino)-3,3'-diethyl-2,2'-thiacarbocyanine iodide BuOH	FP	700 ^a			777036
192.	1,7-Bis(dimethylamino)-1,4,6-heptatrien-3-one PrOH	FP	800		\uparrow Oxygen quenching and triplet ET from anthracene and benzo[<i>rsi</i>]pentaphene and to azulene, perylene, and tetracene	83E737
193.	2,5-Bis[7-(dimethylamino)-2,4,6-heptatrienylidene]cyclopentanone Toluene	FP	850 730		\uparrow Oxygen quenching and triplet ET from anthracene and benzo[<i>rsi</i>]pentaphene and to azulene, perylene, and tetracene	83E737
194.	1,9-Bis(dimethylamino)-1,3,6,8-nonatetraen-5-one PrOH	FP	750		\uparrow Oxygen quenching and triplet ET from anthracene and benzo[<i>rsi</i>]pentaphene and to azulene, perylene, and tetracene	83E737
	Toluene	FP	820 610		\uparrow Oxygen quenching and triplet ET from anthracene and benzo[<i>rsi</i>]pentaphene and to azulene, perylene, and tetracene	83E737
195.	1,9-Bis(dimethylamino)-1,4,6,8-nonatetraen-3-one PrOH	FP	550		\uparrow Oxygen quenching and triplet ET from anthracene and benzo[<i>rsi</i>]pentaphene and to azulene, perylene, and tetracene	83E737

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{max} , /nm	ϵ_{max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
196.	<i>all-trans</i> -1,15-Bis(dimethylamino)-1,3,6,8,10,12,14-pentadecaheptaen-5-one Toluene	FP	630		†Oxygen quenching and triplet ET from anthracene and benzo[<i>rst</i>]pentaphene and to azulene, perylene, and tetracene	83E737
197.	2,6-Bis[5-(dimethylamino)-2,4-pentadienylidene]cyclohexanone Heptane	FP	886 ^a 782 ^a 619 ^a		†Oxygen quenching and triplet ET from anthracene and benzo[<i>rst</i>]pentaphene and to azulene, perylene, and tetracene	83E737
	PrOH	FP	950 680		†Oxygen quenching and triplet ET from anthracene and benzo[<i>rst</i>]pentaphene and to azulene, perylene, and tetracene	83E737
	Toluene	FP	910 640		†Oxygen quenching and triplet ET from anthracene and benzo[<i>rst</i>]pentaphene and to azulene, perylene, and tetracene	83E737
198.	2,5-Bis[5-(dimethylamino)-2,4-pentadienylidene]cyclopentanone Toluene	FP	900 635		†Oxygen quenching and triplet ET from anthracene and benzo[<i>rst</i>]pentaphene and to azulene, perylene, and tetracene	83E737
199.	2,6-Bis[3-(dimethylamino)-2-propenylidene]cyclohexanone PrOH	FP	800		†Oxygen quenching and triplet ET from anthracene and benzo[<i>rst</i>]pentaphene and to azulene, perylene, and tetracene	83E737
	Toluene	FP	750 600		†Oxygen quenching and triplet ET from anthracene and benzo[<i>rst</i>]pentaphene and to azulene, perylene, and tetracene	83E737
200.	4,4'-Bis(dimethylamino)thiobenzophenone Benzene	LP/RA	490 ± 5, 7200 ± 1400 335 ± 5, 14400 ± 2900		ϵ relative to benzophenone in benzene ($\epsilon_{332.5} = 7630$ L mol ⁻¹ cm ⁻¹), taking $\Phi_T = 0.4$ at 337 nm excitation and taking $\Phi_T = 1$ for benzophenone; $\tau_T = 1.3 \pm 0.1$ μ s	84A221
201.	<i>all-trans</i> -1,13-Bis(dimethylamino)-1,3,5,8,10,12-tridecahexaen-7-one Toluene	FP	900 670		†Oxygen quenching and triplet ET from anthracene and benzo[<i>rst</i>]pentaphene and to azulene, perylene, and tetracene	83E737
202.	<i>all-trans</i> -1,13-Bis(dimethylamino)-1,3,6,8,10,12-tridecahexaen-5-one PrOH	FP	740		†Oxygen quenching and triplet ET from anthracene and benzo[<i>rst</i>]pentaphene and to azulene, perylene, and tetracene	83E737
	Toluene	FP	600		†Oxygen quenching and triplet ET from anthracene and benzo[<i>rst</i>]pentaphene and to azulene, perylene, and tetracene	83E737
203.	<i>all-trans</i> -1,11-Bis(dimethylamino)-1,3,6,8,10-undecapentaen-5-one PrOH	FP	870		†Oxygen quenching and triplet ET from anthracene and benzo[<i>rst</i>]pentaphene and to azulene, perylene, and tetracene	83E737
	Toluene	FP	680		†Oxygen quenching and triplet ET from anthracene and benzo[<i>rst</i>]pentaphene and to azulene, perylene, and tetracene	83E737
204.	Bis(dimethylformamide)phthalocyaninoruthenium(II) Acetonitrile	LP	527 ^a		Another maximum at 359 ^a was near a SD region; $\tau_T = 0.165$ μ s	83E262
205.	Bis(dimethylsulfoxide)phthalocyaninoruthenium(II) Acetonitrile	LP	485 ^a		Another maximum at 352 ^a nm was near a SD region	83E262

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
206.	Bis(dimethylsulfoxide)phthalocyanine/ruthenium(II)/1,4-Dinitrobenzene Acetonitrile	LP	525 ^a		Triplet exciplex; another maximum at 409 ^a nm was near a SD region	83E262
207.	1,4-Bis(methylamino)anthraquinone Benzene	PR-ET/ET	550, 20000		†Triplet ET from biphenyl; ϵ relative to biphenyl in benzene assuming ground state dimerization; $\tau_T = 5.9$ μ s; $E_T = 97 - 151$ kJ mol ⁻¹ ; $k_{et} = 8.1 \times 10^9$ L mol ⁻¹ s ⁻¹	761122
208.	1,1-Bis(1-naphthyl)ethylene Benzene	LP-ET	444 ^a		†Triplet ET from xanthone and oxygen quenching; $\tau_T = 0.22$ μ s	84B007
209.	trans-1,2-Bis(1-naphthyl)ethylene Benzene	LP/ET	530 \pm 3, 30000 \pm 4500		†Quenching by oxygen, a nitroxide free radical, and azulene; ϵ relative to fluorenone in benzene ($\epsilon_{435} = 6000$ L mol ⁻¹ cm ⁻¹); $\tau_T = 0.91 \pm 0.14$ μ s; $k_{et} = (7.4 \pm 1.1) \times 10^9$ L mol ⁻¹ s ⁻¹	84E237
	Bromobenzene	LP	535 \pm 3		$\tau_T = 0.67 \pm 0.10$ μ s	84E237
210.	trans-1,2-Bis(2-naphthyl)ethylene Benzene	LP/ET	430 \pm 3, 20000 \pm 3000		†Quenching by oxygen, a nitroxide free radical, and azulene; ϵ relative to fluorenone in benzene ($\epsilon_{435} = 6000$ L mol ⁻¹ cm ⁻¹); $\tau_T = 0.21 \pm 0.03$ μ s; $k_{et} = (7.7 \pm 1.2) \times 10^9$ L mol ⁻¹ s ⁻¹	84E237
	Bromobenzene	LP	435 \pm 3		$\tau_T = 0.17 \pm 0.03$ μ s	84E237
211.	9,10-Bis(phenylethynyl)anthracene Acetonitrile	LP-ET	$\sim 480^a$		†Triplet ET from benzophenone; delay 5 μ s	82E509
212.	rac-Bis[1-(1-pyrenyl)ethyl]ether Hexane	LP	513 ^a 479 ^a 415 390		Most intense peak at 415 nm	83B137
213.	Bonellin EtOH (298 K)	LP/TD	625 ^a , 40000 ^a 443 ^b , 9500 \pm 500 390 ^a , 10000 ^a		†Triplet ET from anthracene, oxygen quenching (1.7×10^8 L mol ⁻¹ s ⁻¹); $\tau_T = 430 \pm 50$ μ s; $E_T = 180 \pm 20$ kJ mol ⁻¹ ; $k_{et} = (2.0 \pm 1.0) \times 10^8$ L mol ⁻¹ s ⁻¹	80E593
214.	Brilliant Sulfaflavine anion EPA	CWL/SD	575, 1800		Glass was 1:1 mixture of standard EPA and EtOH	73E346
215.	9-Bromoanthracene Acetonitrile	LP	424 ^a 400 ^a		†Oxygen quenching; 424 nm peak was the more intense; $\tau_T = 19.5$ μ s	84B110
	Benzene	PR/ET	430, 48000		ϵ relative to naphthalene in benzene ($\epsilon_{\max} = 17500$ L mol ⁻¹ cm ⁻¹)	690087
	Benzene	PR/ET	430, 47700		ϵ relative to benzophenone in benzene ($\epsilon_{532.5} = 10300$ L mol ⁻¹ cm ⁻¹)	690087
	Cyclohexane	FP	425		$\tau_T = 43$ μ s	62E009
	Cyclohexane	PR	425			690087
	Cyclohexane	FP/SD	423.5, 66500 \pm 3250		ϵ method assumes linear variation of triplet spectrum in SD region	767147
	Dioxane	PR	425			690087
	EPA (77 K)	LP	430 405		$\tau_T = 200$ μ s	84B110
	Liquid paraffin	FP	419		Viscosity of solvent was 0.167 N-s/m ² ; $\tau_T = 110$ μ s	62E009
216.	4-Bromobiphenyl Toluene	PR	390		†Oxygen quenching; $\tau_T = \sim 4$ μ s	80A235

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
217.	4-Bromo-1-cyanoisquinoline-N-oxide					
	Cyclohexane	FP	427		$\tau_T = 1.7 \mu s$	727105
	EtOH	FP	420		$\tau_T = 4.0 \mu s$	727105
218.	N-[2-[(2-Bromo-6-cyano-4-nitrophenyl)azo]-5-[(2-cyanoethyl)(2-hydroxyethyl)amino]-4-methoxyphenyl]acetamide					
	2-MTHF (103 K)	LP	680		Shoulder at 780 nm; triplet not observable above 211 K; below 116 K lifetime constant; $\tau_T = 130 \mu s$	80B101
	Dibutyl terephthalate	LP	685		†Possibilities other than triplet character excluded; $\tau_T = 0.023 \mu s$	80B101
	Glycerol triacetate	LP	685		Shoulder at 780 nm; $\tau_T = 0.014 \mu s$	80B101
	Glycerol triacetate (203 K)	LP	685		Shoulder at 790 nm; below 221 K lifetime constant; $\tau_T = 83 \mu s$	80B101
			400			
219.	N-[2-[(2-Bromo-4,6-dinitrophenyl)azo]-5-[(2-cyanoethyl)(2-hydroxyethyl)amino]-4-methoxyphenyl]acetamide					
	1,3-Dibromobenzene	LP	700		Shoulder at 780 nm; $\tau_T = 0.010 \mu s$	80B101
			390			
	1-Phenylethanol	LP	700		Shoulder at 780 nm; $\tau_T = 0.012 \mu s$	80B101
			400			
	2-MTHF (103 K)	LP	700		†Oxygen quenching; triplet virtually not observable above ~273 K; below 111 K lifetime constant; $\tau_T = 100 \mu s$	80B101
			400			
	2-MTHF	LP	800-700		$\tau_T = < 0.008 \mu s$	80B101
	2-PrOH	LP	800-700		$\tau_T = 0.012 \mu s$	80B101
	Biphenyl/Diphenyl ether	LP	700		Shoulder at 800 nm; solvent was 1:3 biphenyl to diphenyl ether; $\tau_T = 0.008 \mu s$	80B101
			420			
	Dibutyl terephthalate	LP	700		†Possibilities other than triplet character of the transient excluded; shoulder at 750 nm; $\tau_T = 0.026 \mu s$	80B101
			400			
	Glycerol triacetate	LP	695		Shoulder at 770 nm; $\tau_T = 0.012 \mu s$	80B101
			400			
	Glycerol triacetate (203 K)	LP	720		Below 220 K lifetime constant; $\tau_T = 100 \mu s$	80B101
	PMMA	LP	680		Shoulder at 740 nm	80B101
	<i>o</i> -Terphenyl/Diphenyl ether	LP	695		Shoulder at 780 nm; solvent was 1:1 <i>o</i> -terphenyl to diphenyl ether; $\tau_T = 0.010 \mu s$	80B101
			400			
220.	4-[(2-Bromo-4,6-dinitrophenyl)azo]-N,N-dimethylbenzenamine					
	2-MTHF (103 K)	LP	700		Lifetime measured at 138 K; triplet not observable above 138 K; below 100 K lifetime (500 μs) constant; $\tau_T = 0.014 \mu s$	80B101
			410			
	Glycerol triacetate (203 K)	LP	700		Lifetime measured at 247 K; triplet not observable above 247 K; $\tau_T = 0.020 \mu s$	80B101
221.	4-Bromoisoquinoline-N-oxide					
	Water	FP	382		$\tau_T = 0.96 \mu s$	727105
222.	Bromo(methanol)(phthalocyaninato)rhodium(III)					
	Acetonitrile/2-PrOH	LP/SD	640 ^b , 900		Solvent was 2.6 mol L ⁻¹ 2-PrOH	83F405
			590 ^b , 22000			
223.	1-(Bromomethyl)naphthalene					
	?	LP	426		Solvent was either MeOH or cyclohexane	84A222
224.	1-Bromonaphthalene					
	CTAB	LP/TD	425, 11500		Aqueous micelle; ϵ method corrects for TT-annihilation in micellar environment and gives upper limit; shoulder at 370 nm; $\tau_T = 50 \mu s$	81N070
			400 ^a , 7000 ^a			
	Cyclohexane	FP	424		$\tau_T = 270 \mu s$	62E009
	Cyclohexane (77 K)	PS	414.5		Relative intensities (2:1)	69E211
			392.0			
	Et ₂ O/Pentane (77 K)	PS	415.5		Relative intensities (2:1)	69E211
			393			
	Ethylene glycol	FP	420		Relative intensities (100:60:25); $\tau_T = 830 \pm 140 \mu s$	61E005
			398			
			378			

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Hexane	FP	419 398 375		Relative intensities (100:55:20); $\tau_T = 83 \pm 14 \mu\text{s}$	61E005
	Hexane	LP	425		Delay 500 ps	82E303
	Toluene	PR	420		†Oxygen quenching; $\tau_T = \sim 9 \mu\text{s}$	80A235
225.	2-Bromonaphthalene					
	Cyclohexane	FP	425		$\tau_T = 150 \mu\text{s}$	62E009
	Cyclohexane (77 K)	PS	417 390 370		Relative intensities (100:55:10)	69E211
	EPA (77 K)	PS	421.8		†Phosphorescence decay	51E001
	EPA (77 K)	FP	423 398			55E002
	Hexane	FP	417.5 392.9			54E001
	Hexane	FP	417.5 394 372		Relative intensities (100:61:29)	58E001
	Liquid paraffin	FP	423		Viscosity of solvent was 0.167 N·s/m ² ; $\tau_T = 2300 \mu\text{s}$	62E009
226.	4-[(2-Bromo-4-nitrophenyl)azo]-N,N-dimethylbenzenamine					
	2-MTHF (77 K)	LP	700 400		Triplet not observable above 125 K; below 106 K lifetime constant; $\tau_T = 100 \mu\text{s}$	80B101
	EPA (89 K)	LP	710 390			80B101
	EtOH/MeOH (113 K)	LP	700 390		Solvent was 4:1 EtOH to MeOH	80B101
	Glycerol triacetate (203 K)	LP	680 400		Lifetime measured at 247 K; triplet not observable above 247 K; $\tau_T = 0.014 \mu\text{s}$	80B101
	MCH/Toluene (113 K)	LP	700 390		Solvent was 1:1 MCH to toluene	80B101
	PMMA	LP	690			80B101
227.	9-Bromophenanthrene					
	Cyclohexane	FP	482.5		$\tau_T = 115 \mu\text{s}$	62E009
	Liquid paraffin	FP	482.5		Viscosity of solvent was 0.167 N·s/m ² ; $\tau_T = 710 \mu\text{s}$	62E009
228.	1-Bromopyrene					
	EtOH (293 K)	LP	505 ^a 476 ^a		More intense absorption below 425 nm	697226
229.	trans-4-Bromostilbene					
	2-MTHF (135 K)	LP	392 373 352		First peak most intense; $\tau_T = 0.02 \mu\text{s}$	79E378
	2-MTHF (135 K)	LP	392 373 352		Triplet absorption not observable above 135 K; reported lifetime constant below 104 K; $\tau_T = 480 \mu\text{s}$	79E640
	3-MP (88 K)	LP	393 376 355			79E640
	EPA (130 K)	LP	393 372 355		First peak most intense; $\tau_T = 0.02 \mu\text{s}$	79E378
	EPA (108 K)	LP	393 372 355		Triplet absorption not observable above 130 K; reported lifetime constant below 97 K; $\tau_T = 500 \mu\text{s}$	79E640
	EtOH (138 K)	LP	391 372 352		Relative intensities (4:3:1); triplet absorption not observable above 143 K; reported lifetime constant below 115 K; $\tau_T = 480 \mu\text{s}$	79E640
	EtOH (143 K)	LP	391 372 352		First peak most intense; $\tau_T = 0.1 \mu\text{s}$	79E378

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	EtOH/MeOH (103 K)	LP	392 372 352		Solvent was 4:1 EtOH to MeOH	79E640
	Glycerol (286 K)	LP	395 375 354		First peak most intense; $\tau_T = 0.03 \mu s$	79E378
	Glycerol (193 K)	LP	395 375 354		Relative intensities (4:3:1); triplet absorption not observable above 286 K; reported lifetime constant below 209 K; $\tau_T = 450 \mu s$	79E640
	Glycerol triacetate (233 K)	LP	390 370 352		Triplet absorption not observable above 243 K; reported lifetime constant below 207 K; $\tau_T = 440 \mu s$	79E640
	Glycerol triacetate (243 K)	LP	390 370 352		First peak most intense; $\tau_T = 0.03 \mu s$	79E378
	MCH/Isohexane (114 K)	LP	392 372 352		First peak most intense; $\tau_T = 0.3 \mu s$	79E378
	MCH/Isohexane (77 K)	LP	392 372 352		Solvent was 1:1 MCH to isohexane by volume; triplet absorption not observable above 114 K; reported lifetime constant below 100 K; $\tau_T = 450 \mu s$	79E640
230.	2-Bromotriphenylene EtOH/MeOH (113 K)	FP/TD	430, >8000		Solvent was 3:1 EtOH to MeOH; lower limit assumes total ground state depletion	67B004
231.	2-Butanone Methylene chloride	LP	245		†Phosphorescence decay and oxygen quenching; $\tau_T = 1.4 \mu s$	84B051
232.	Butyl benzoylformate Benzene (298 K)	LP	451 ^a 430 ^a 400 ^a		$\tau_T = 0.5 \mu s$	84A030
233.	4-tert-Butyl-3,5-dinitroanisole CF ₃ CH ₂ OH	LP	441 ^a		†Quenching by oxygen and tetramethyldiazetidine; $\tau_T = 3.5 \mu s$	84E033
234.	2-[1-(Butylimino)ethyl]-5-methylphenol Benzene (295 K)	LP	430 ^a		†Oxygen quenching and triplet ET to compound by triplet donors	727374
235.	2-tert-Butyl-4-methylindazole EtOH (103 K)	MOD/SD	409 ^a , 8400 ^a		Shoulders at 388 ^a and 294 ^a nm; another maximum < 250 nm; $\tau_T = (9.1 \pm 0.8) \times 10^3 \mu s$	716244
	EtOH (178 K)	MOD/SD	409 ^a , 7600 ^a		Shoulders at 393 ^a and 280 ^a nm; another maximum < 250 nm; $\tau_T = (1.9 \pm 0.3) \times 10^4 \mu s$	716244
236.	N-Butyl-5-nitro-2-furamide 2-PrOH (298 K)	LP	505 ± 5		$\tau_T = 0.037 \mu s$	84A263
	Acetonitrile (298 K)	LP	498 ± 5		†Quenching by oxygen and azulene; $\tau_T = 0.26 \mu s$	84A263
	CCl ₄ (298 K)	LP	500 ± 5		$\tau_T = 0.052 \mu s$	84A263
	MeOH (298 K)	LP	515 ± 5		$\tau_T = 0.022 \mu s$	84A263
	Water (298 K)	LP	515 ± 5		$\tau_T = 0.019 \mu s$	84A263
237.	1-Butynylbenzene Isopentane/2-PrOH (77 K)	PS	255 ^a		†ESR; glass was 7:3 isopentane to 2-propanol by volume; "1-ethyl-2-phenylacetylene"; by assuming that radical formation occurs only from T [*] and that it is wavelength-independent, the author devise an ESR method that determines the shape of TTA even in the SD region	68B007

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
238.	Butyraldehyde Pentane	LP	355 320		†Triplet ET to 2,5-dimethyl-2,4-hexadiene; both peaks have the same decay and quenching behavior and are attributed to a single species; $\tau_T = 0.210 \mu\text{s}$; $k_{et} = (1.5 \pm 0.2) \times 10^{10} \text{ L mol}^{-1} \text{ s}^{-1}$	81E442
239.	9-Butyrylanthracene EPA (77 K)	FP	430		$\tau_T = 3.42 \times 10^4 \mu\text{s}$	82E338
240.	Cadmium(II) tetrabenzoporphyrin Pyridine	FP	500 415			73E345
241.	Cadmium(II) tetraphenylporphyrin MCH	FP/TD	490, 57000 415, 22000		†Oxygen quenching; $\tau_T = 265 \mu\text{s}$	81E271
242.	Caffeine Water	LP-ET	351		†Triplet ET from acetone; $\tau_T = 4.5 \mu\text{s}$; pH 7.1; $k_{et} = 5.6 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	82B045
243.	Camphoroquinone 2-PrOH	FP	1060 908 790 702 630 ~500 320 276		†Triplet ET from benzophenone, triplet ET to anthracene, and oxygen quenching; another maximum <200 nm; delay 0 μs	697092
	Benzene	FP	1070 916 796 706 638 404 314		†Triplet ET from benzophenone, triplet ET to anthracene, and oxygen quenching	697092
	CCl ₄	FP	1068 916 796 705 638 590 500 316 280		†Triplet ET from benzophenone, triplet ET to anthracene, and oxygen quenching; another maximum <220 nm	697092
244.	Canthaxanthin ?	FP-ET	541 ^a		†Triplet ET from anthracene; it was not reported whether the solvent was hexane or benzene; lifetime was measured in hexane; $\tau_T = 7.1 \mu\text{s}$	733001
	Benzene	PR-ET	555		†Triplet ET from naphthalene; $\tau_T = 3.8 \mu\text{s}$	80A143
245.	Carbazole 2-MTHF (77 K)	PS/ESR	433, 12200		†ESR; only most intense visible peak reported; oscillator strength encompasses two electronic transitions; oscillator strength = 0.18	696115
	Benzene	FP	418 ^a			84F248
	Cyclohexane	FP/?	435 ^a , 14000 ^a 400 ^a , 12500 ^a		ϵ determination unspecified; $\tau_T = 167 \mu\text{s}$	77A178
	Cyclohexane	FP/ET	420, 14500		ϵ relative to naphthalene in cyclohexane ($\epsilon_{415} = 24500 \text{ L mol}^{-1} \text{ cm}^{-1}$); carbazyl radicals ($\epsilon_{600} = 7000 \text{ L mol}^{-1} \text{ cm}^{-1}$) also observed; $\tau_T = 170 \mu\text{s}$	78A368

TABLE 6 Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{max} , /nm	ϵ_{max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Cyclohexane	LP	~620 ^a 420		†Rises with fluorescence decay; relative intensities (1:8); shoulder at 400 nm; $S_0 \leftarrow S_1$ absorption also recorded ($\lambda_{620} = 20000 \text{ L mol}^{-1} \text{ cm}^{-1}$)	80E023
	EPA (77 K)	PS	428 406		†Phosphorescence decay; $\tau_T = (8.0 \pm 0.2) \times 10^6 \mu\text{s}$	67E106
	EPA (77 K)	PS/KM	431 ^a , 14300 407 ^a , 11200 ^a		Shoulders at 506 ^a and 375 ^a nm; two electronic bands were assigned: the 1st was the shoulder at 506 nm with $f = 0.06$, and the 2nd started at 431 nm; ϵ -method assumed on basis of earlier work by authors; oscillator strength = 0.24	747347
	EtOH (77 K)	MOD/KM	425, 19000 \pm 4700 406, 14000 \pm 3500 318, 7000 \pm 1700 305, 5700 \pm 1400			737055
	EtOH	FP	639 ~480 425 ^a 405 ^a		Most intense peak at 425 nm	766421
	EtOH/Et ₂ O (77 K)	MOD/KM	425, 10100 \pm 4000		Glass was 2:1 EtOH to Et ₂ O; temperature was not explicitly stated, but 77 K was inferred from the context	719059
	Hexane	FP	420		†Oxygen and piperylene quenching	757573
	PMMA (77 K)	CWL	510 ^a 430 ^a 415 ^a 378 ^a		Most intense peak 430 nm	837042
246.	3-Carboethoxypsoralen					
	Benzene	PR/ET	450 ^b , 7500 400 ^a		†Triplet ET from biphenyl, oxygen quenching ($1.9 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); ϵ relative to biphenyl in benzene ($\epsilon_{367} = 27100 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 20 \mu\text{s}$	82E133
	Benzene	LP	407 ^a		†Oxygen quenching and triplet ET from biphenyl	84B025
	EtOH	LP/ET	590 ^a 500 ^a 450 ^b , 6800 370 ^a		†Triplet ET to retinol, oxygen quenching ($2.7 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$); ϵ relative to retinol in hexane ($\epsilon_{405} = 80000 \text{ L mol}^{-1} \text{ cm}^{-1}$) assuming ϵ independent of solvent; $\tau_T = 5.5 \mu\text{s}$; $k_{et} = 9.6 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$	82E133
	EtOH	LP/ET	450 ^b , 5457		ϵ relative to retinol, but actual reference ϵ used was not specified; $\tau_T = 5.6 \mu\text{s}$	83E324
	Water	LP/ELT	605 ^a 590 ^a 450 ^b , 3700 \pm 900 400 ^a		†Oxygen quenching ($3.0 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$); ϵ relative to radical anion ($\epsilon_{600} = 2500 \text{ L mol}^{-1} \text{ cm}^{-1}$) assuming electron transfer from tyrosine 100% efficient; $\tau_T = 17 \mu\text{s}$	82E133
	Water	PR/ELT	605 ^a 590 ^a 450 ^b , 3150 400 ^a		†Oxygen quenching ($3.0 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$); ϵ relative to radical anion ($\epsilon_{600} = 2500 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 17 \mu\text{s}$	82E133
	Water	LP/ELT	605 ^a 590 ^a 450 ^b , 3700 \pm 900 400 ^a		†Oxygen quenching ($3.0 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$); ϵ relative to radical anion ($\epsilon_{600} = 2000 \text{ L mol}^{-1} \text{ cm}^{-1}$) assuming electron transfer from tryptophan 100% efficient; $\tau_T = 17 \mu\text{s}$	82E133
	Water	LP/TD	450 ^b , 2000		$\tau_T = 5 \mu\text{s}$	83E324
	Water	LP	612 ^a 578 ^a 397 ^a		†Oxygen quenching	84B025
247.	Carbostyryl					
	EPA (77 K)	PS	427 ^a		Phosphorescence decay; shoulder at 465 ^a nm; $\tau_T = 9.6 \times 10^5 \mu\text{s}$	717171
	EtOH/Water	FP	451 ^a 430 ^a		Solvent was 7:3 EtOH to water; peaks were of almost equal intensity	737046
	Glycerol	FP	444 ^a		Shoulder at 488 ^a nm	717171
	Liquid paraffin (268 K)	FP	488 ^a 446 ^a		446 nm was the more intense peak	717171

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Water	FP	431 ^a 431 ^a		Shoulder at 476 ^a nm; 451 nm was the more intense peak; $\tau_T = 17 \mu\text{s}$	737046
248.	Carbostyryl, negative ion Water	FP	460 ^a		Shoulders at 478 ^a and 440 ^a nm; $\tau_T = 30 \mu\text{s}$; pH 14	737046
249.	4-Carboxybenzophenone Water	LP/RA	535, 6600 525 ^b , 6150		†Phosphorescence decay; carboxylate anion form; ϵ relative to benzophenone in benzene ($\epsilon_{525} = 7630 \text{ L mol}^{-1} \text{ cm}^{-1}$) assuming triplet yields of unity for both species; corresponding ketyl radical ($\epsilon_{570} = 5500 \text{ L mol}^{-1} \text{ cm}^{-1}$) and radical anion ($\epsilon_{660} = 8000 \text{ L mol}^{-1} \text{ cm}^{-1}$) also measured; lifetime limited by ground state quenching; $\tau_T = 5.0\text{--}6.7 \mu\text{s}$; pH 7	81A314
250.	11- <i>cis</i> - β -apo-14'-Carotenal Hexane (298 K) MeOH	LP/SD LP-ET	470 \pm 5, 90000 \pm 10000 490		"11- <i>cis</i> -C ₂₂ aldehyde" †Triplet ET from anthracene	84B026 84B026
251.	β -apo-14'-Carotenal Acetonitrile	PR/RF	490, 112000		"all- <i>trans</i> -C ₂₂ aldehyde"; ϵ relative to compound in cyclohexane ($\epsilon_{470} = 121000 \text{ L mol}^{-1} \text{ cm}^{-1}$) assuming oscillator strength independent of solvent; $\tau_T = 11.9 \mu\text{s}$	79E546
	Benzene	PR/RF	490, 119000		ϵ relative to compound in cyclohexane ($\epsilon_{470} = 121000 \text{ L mol}^{-1} \text{ cm}^{-1}$) assuming oscillator strength independent of solvent; $\tau_T = 8.3 \mu\text{s}$	79E546
	Benzene	LP	485			82F477
	CF ₃ CH ₂ OH	LP-ET	510		†Triplet ET from anthracene; $\tau_T = \sim 17 \mu\text{s}$	83E026
	Cyclohexane	PR/ET	470, 121000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361.3} = 42800 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 7.1 \mu\text{s}$	79E546
	Cyclohexane	LP	473			82F477
	Cyclohexane	LP-ET	475		†Triplet ET from anthracene; solution 0.08 mol L ⁻¹ in (CF ₃) ₃ CHOH; H-bonded species shows as a shoulder around 510 nm; $\tau_T = 8 \mu\text{s}$	83E026
	DFMeOH	LP-ET	560		†Triplet ET from anthracene; $\tau_T = > 40 \mu\text{s}$	83E026
	Hexane	LP/SD	470, 127000 \pm 19100		ϵ assumes triplet does not absorb where singlet depletion is followed; $\tau_T = 5 \mu\text{s}$; $E_T = \sim 148 \text{ kJ mol}^{-1}$	78E721
	MeOH	PR/RF	475, 149000 ^c		ϵ relative to compound in cyclohexane ($\epsilon_{470} = 121000 \text{ L mol}^{-1} \text{ cm}^{-1}$) assuming oscillator strength independent of solvent; $\tau_T = 10.3 \mu\text{s}$	79E546
252.	β -apo-8'-Carotenal ?	FP-ET	517 ^a		†Triplet ET from anthracene; it was not reported whether the solvent was hexane or benzene; lifetime was measured in hexane; $\tau_T = 10 \mu\text{s}$	733001
	Hexane	LP-ET/SD	520, 223000 \pm 33500		†Triplet ET from biphenyl; ϵ assumes triplet does not absorb where singlet depletion is followed; $\tau_T = 2.5 \mu\text{s}$; $E_T = \sim 122 \text{ kJ mol}^{-1}$	78E721
253.	15,15'- <i>cis</i> - β -Carotene ?	FP-ET	507 ^a		†Triplet ET from anthracene; it was not reported whether the solvent was hexane or benzene; lifetime was measured in hexane; shoulder at 499 ^a nm; $\tau_T = 7.1 \mu\text{s}$	733001
	Hexane	PR/ET	515, 250000 \pm 37500		†Triplet ET from biphenyl, ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800 \text{ L mol}^{-1} \text{ cm}^{-1}$), assuming authors used standard for cyclohexane; $\tau_T = 5.9 \pm 0.6 \mu\text{s}$; $k_{et} = (2.0 \pm 0.20) \times 10^{10} \text{ L mol}^{-1} \text{ s}^{-1}$	776412
254.	β -Carotene Acetic acid	LP-ET	512		†Triplet ET from hematoporphyrin; $\tau_T = 9 \mu\text{s}$	81B115
	Benzene (296 K)	LP-ET	526		Triplet ET from chlorophyll <i>a</i> ; oxygen quenching ($3.6 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); $\tau_T = 5.9 \mu\text{s}$; $k_{et} = 1.2 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	73E347

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Benzene	LP-ET	520		†Triplet ET from anthracene; $\tau_T = 8.0 \pm 0.5 \mu\text{s}$	767094
	Benzene	PR	540			761024
	CTAB/Triton X-100	LP	525		Surfactant ratio 1:2 CTAB to Triton X-100; mixed aqueous micelle; $\tau_T = 4 \mu\text{s}$	80N033
			475			
	Carbon disulfide (293 K)	LP	550		†Triplet ET from chlorophyll <i>a</i> ; $\tau_T = 15 \mu\text{s}$; $k_{\text{et}} = 4 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	69N001
	Hexane (188 K)	FP-ET	522		†Triplet ET from benz[<i>a</i>]anthracene; $\tau_T = 300 \mu\text{s}$	66E089
	Hexane (298 K)	FP-ET	514		†Triplet ET from chlorophyll <i>a</i> , tetracene, benz[<i>a</i>]anthracene, and anthracene; the data presented is for anthracene; $\tau_T = 70 \mu\text{s}$; $k_{\text{et}} = 1.1 \times 10^{10} \text{ L mol}^{-1} \text{ s}^{-1}$	66E089
	Hexane (293 K)	LP	515		†Triplet ET from chlorophyll <i>a</i>	69N001
	Hexane	PR/ET	515, 170000 \pm 40000		ϵ relative to naphthalene in cyclohexane ($\epsilon_{412.5} = 22600 \text{ L mol}^{-1} \text{ cm}^{-1}$), exact ref. used by author not stated but surmised; $\tau_T = 9 \mu\text{s}$	703001
	Hexane	PR/SD	515, 230000		$\tau_T = 9 \mu\text{s}$	703001
	Hexane	PR/ET	515, 130000 \pm 10000 482 ^a		†Triplet ET from naphthalene and oxygen quenching; shoulder at 453 ^a nm; ϵ relative to naphthalene in cyclohexane ($\epsilon_{415} = 24500 \text{ L mol}^{-1} \text{ cm}^{-1}$), except authors used 412.5 nm as reference λ ; $\tau_T = 9 \mu\text{s}$; $k_{\text{et}} = 1.5 \times 10^{10} \text{ L mol}^{-1} \text{ s}^{-1}$	713035
	Hexane	PR/ET	515, 242000 \pm 36300		†Triplet ET from biphenyl; ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800 \text{ L mol}^{-1} \text{ cm}^{-1}$), assuming authors used standard for cyclohexane; $\tau_T = 5.9 \pm 0.6 \mu\text{s}$; $k_{\text{et}} = (1.8 \pm 0.18) \times 10^{10} \text{ L mol}^{-1} \text{ s}^{-1}$	776412
255.	ζ-Carotene Cyclohexane	PR/ET	445, 81000		†Triplet energy transfer from biphenyl; ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 23 \mu\text{s}$; $E_T = < 123 \text{ kJ mol}^{-1}$; $k_{\text{et}} = 1.4 \times 10^{10} \text{ L mol}^{-1} \text{ s}^{-1}$	761035
256.	β- <i>apo</i> -8'-Carotenoic acid, methyl ester ?	FP-ET	507 ^a		†Triplet ET from anthracene; it was not reported whether the solvent was hexane or benzene; lifetime was measured in hexane; shoulder at 497 ^a nm; $\tau_T = 8.3 \mu\text{s}$	733001
257.	4-(β- <i>apo</i> -7'-Carotenyl)benzyl pyropheophorbide Toluene	LP	~540		$\tau_T = 4.63 \mu\text{s}$	82B094
258.	Chalcone Heptane	LP	430 \pm 15		"1,3-Diphenyl-2-propen-1-one"; $\tau_T = 0.013 \mu\text{s}$	83E347
259.	Chloranil 1,2-Dichloroethane	LP/RA	510, 7200 \pm 1300 370, 4700 ^a		†Oxygen quenching; lifetime measured for the 510 nm band; ϵ relative to benzophenone in benzene ($\epsilon_{532.5} = 7630 \text{ L mol}^{-1} \text{ cm}^{-1}$) and assuming Φ_T for chloranil is unity; $\tau_T = 5.6 \mu\text{s}$	776335
	1,4-Dioxane	LP/RA	515, 5800 \pm 100		ϵ relative to benzophenone in benzene ($\epsilon_{532.5} = 7600 \text{ L mol}^{-1} \text{ cm}^{-1}$) assuming $\Phi_T = 1.0$ for chloranil in 1,4-dioxane; subsidiary peak at 740 nm due to complexation with solvent; $\tau_T = 3.4 \mu\text{s}$	79A178
	Acetone	FP	520 503 471			727069
	Acetonitrile	FP	510 487 479 372			727069
	Acetonitrile	LP/RA	510, 7700 \pm 300		ϵ relative to benzophenone in benzene ($\epsilon_{532.5} = 7630 \text{ L mol}^{-1} \text{ cm}^{-1}$) assuming $\Phi_T = 1$ for chloranil in acetonitrile; ϵ_{ref} assumed from earlier work	78A195
	Benzene	FP	516			727069

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Butyronitrile	LP/RA	510, 7570 480 380		†Oxygen quenching; ϵ relative to the chloranil radical anion in acetonitrile ($\epsilon_{\max} = 9700$ L mol ⁻¹ cm ⁻¹); other peaks of roughly equal intensity at 480 and 380 nm; $\tau_T = 6.3$ μ s	79B061
	Cyclohexane	LP	500		$\tau_T = 2.0 \pm 0.3$ μ s	697272
	Cyclohexane	FP	539 507 495 474			727069
	Dioxane	LP/RA	515, 5800 \pm 300		ϵ relative to benzophenone in benzene ($\epsilon_{532.5} = 7630$ L mol ⁻¹ cm ⁻¹) assuming $\Phi_T = 1$ for chloranil in dioxane; ϵ_{ref} assumed from earlier work	78A195
	EtOH	LP	500		†Oxygen and anthracene quenching; $\tau_T = 1.2$ μ s	697272
	MeOH	FP	\sim 500			727069
	THF	FP	\sim 500			727069
	Trichlorotrifluoroethane	FP	522 507 497 473 410 379		†Oxygen quenching; $\tau_T = 30$ μ s	727069
260.	Chloranil/Triphenylamine					
	Benzene/MeOH (293 K)	LP	650 550		650 nm band was the more intense; solvent was 19:1 benzene to MeOH; $\tau_T = 0.04$ μ s	81E715
	Benzene (293 K)	LP	650 550		650 nm band was the more intense; $\tau_T = 0.067$ μ s	81E715
261.	Chloroaluminum phthalocyanine					
	Dimethylacetamide-/Water	LP/SD	490 ^b , 26600		Solvent was 7:3 dimethylacetamide to water	83E088
	Dimethylsulfoxide/Water	LP/SD	490 ^b , 26600		Solvent was 8:2 dimethylsulfoxide to water	83E088
262.	1-Chloroanthracene					
	Benzene	PR/ET	437.5, 60900		ϵ relative to benzophenone in benzene ($\epsilon_{532.5} = 10300$ L mol ⁻¹ cm ⁻¹)	690087
	Benzene	PR/ET	437.5, 66500		ϵ relative to naphthalene in benzene ($\epsilon_{\max} = 17500$ L mol ⁻¹ cm ⁻¹)	690087
	Cyclohexane	FP	425		$\tau_T = 670$ μ s	62E009
	Cyclohexane	PR	430			690087
	Dioxane	PR	432.5			690087
	Hexane	FP	430.7 407.5			54E001
	Hexane	FP	431 407.5		Relative intensities (100:25)	58E001
	Liquid paraffin	FP	419		Viscosity of solvent was 0.167 N·s/m ² ; $\tau_T = 4200$ μ s	62E009
263.	1-Chloroanthraquinone					
	EPA (77 K)	LP	480 ^a 375 ^a		†Phosphorescence decay; delay 500 ns; 375 nm was the more intense peak; $\tau_T = 170 \pm 10$ μ s	83E016
	EtOH	LP	380 ^a		Delay 50 ns	84A253
	Toluene	LP	495 ^a 385 ^a		†Phosphorescence decay in EPA at 77 K; delay 300 ns; 385 nm was more intense peak	83E016
264.	2-Chloroanthraquinone					
	EPA (77 K)	FP	376 ^a		†Phosphorescence decay; $\tau_T = 3400$ μ s	83E016
	Toluene	LP	381 ^a		†Phosphorescence decay in EPA at 77 K; delay 300 ns	83E016
265.	Chlorobenzene					
	Cyclohexane	LP	295 ^a		†Triplet ET to piperylene, oxygen quenching (4.5×10^9 L mol ⁻¹ s ⁻¹); 400 ns delay; $\tau_T = 0.48$ μ s	707561
	Cyclohexane	LP/RA	300, 6250 \pm 300		ϵ relative to naphthalene in cyclohexane ($\epsilon_{414} = 24500$ L mol ⁻¹ cm ⁻¹), taking 0.64 for Φ_T of chlorobenzene and 0.75 for naphthalene; $\tau_T = 1.6 \pm 0.1$ μ s	84E529

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
266.	4-Chlorobenzophenone Acetonitrile/Water	LP/ET	535, 7000 320, 12800		Solvent was 9:1 acetonitrile to water; ϵ relative to 1-methylnaphthalene in acetonitrile/water ($\epsilon_{415} = 11200$ L mol ⁻¹ cm ⁻¹)	84B033
267.	(2'-Chlorobenzoyl)amino-2- Δ^2 -thiazoline Cyclohexane (293 K)	FP/RA	557, 11600		{Triplet ET to biacetyl, oxygen quenching; ϵ relative to naphthalene in cyclohexane ($\epsilon_{414} = 24500$ L mol ⁻¹ cm ⁻¹); $\tau_T = 2000$ μ s	82E373
268.	(4'-Chlorobenzoyl)amino-2- Δ^2 -thiazoline Cyclohexane (293 K)	FP/RA	563, 16900		{Triplet ET to biacetyl, oxygen quenching; ϵ relative to naphthalene in cyclohexane ($\epsilon_{414} = 24500$ L mol ⁻¹ cm ⁻¹); $\tau_T = 2000$ μ s	82E373
269.	4-Chloro-6-[4-(diethylamino)phenyl]- <i>N,N</i> -dimethyl-1,3,5-triazin-2-amine EPA (77 K) PMMA (293 K)	PS PS	485 490		{Phosphorescence decay; $\tau_T = 4.2 \times 10^6$ μ s	83E427 83E427
270.	<i>N</i> -[2-(2-Chloro-4,6-dinitrophenyl)azo]-5-[(2-cyanoethyl)(2-hydroxyethyl)amino]-4-methoxyphenylacetamide 2-MTHF (108 K) Dibutyl terephthalate Glycerol triacetate (203 K) Glycerol triacetate	LP LP LP LP	690 410 690 400 690 695 400		Shoulder at 780 nm; lifetime measured at 189 K; triplet not observable above 189 K; below 100 K lifetime (1000 μ s) constant; $\tau_T = 0.091$ μ s Shoulder at 770 nm; $\tau_T = 0.028$ μ s Shoulder at 780 nm; below 216 K lifetime constant; $\tau_T = 500$ μ s Shoulder at 780 nm; $\tau_T = 0.020$ μ s	80B101 80B101 80B101 80B101
271.	3-Chloro-2,6-diphenyl-1,4-benzoquinone/4-Methoxy- <i>N,N</i> -dimethylaniline Benzene (293 K) Benzene/MeOH (293 K)	LP LP	560 460 460		560 nm peak was the more intense; $\tau_T = 0.083$ μ s Solvent was 19:1 benzene to MeOH; $\tau_T = 0.056$ μ s	81E715 81E715
272.	3-Chloro-2,6-diphenyl-1,4-benzoquinone/Naphthalene Benzene (293 K) Benzene/MeOH (293 K)	LP LP	600 600		$\tau_T = 0.11$ μ s Solvent was 19:1 benzene to MeOH; $\tau_T = 0.091$ μ s	81E715 81E715
273.	3-Chloro-2,6-diphenyl-1,4-benzoquinone/ <i>N,N,N',N'</i> -Tetramethyl- <i>p</i> -phenylenediamine Benzene (293 K) Benzene/MeOH (293 K)	LP LP	620 550 620 570		550 nm peak was the more intense 570 nm peak was the more intense; solvent was 19:1 benzene to MeOH	81E715 81E715
274.	3-Chloro-2,6-diphenyl-1,4-benzoquinone/1,2,3-Trimethoxybenzene Benzene/MeOH (293 K) Benzene (293 K)	LP LP	460 560 460		Solvent was 19:1 benzene to MeOH; $\tau_T = 0.37$ μ s 560 nm peak was the more intense; $\tau_T = 0.63$ μ s	81E715 81E715
275.	3-Chloro-2,6-diphenyl-1,4-benzoquinone/Triphenylamine Benzene (293 K) Benzene/MeOH (293 K)	LP LP	650 550 650 550		550 nm band was the more intense; $\tau_T = 0.16$ μ s 650 nm band was the more intense; solvent was 19:1 benzene to MeOH; $\tau_T = 0.077$ μ s	81E715 81E715
276.	Chloroindium(III) phthalocyanine Dimethylacetamide- /Water	LP/SD	490, 23000		Solvent was 7:3 dimethylacetamide to water	84A122
277.	Chloro(methanol)(phthalocyaninato)rhodium(III) Acetonitrile/2-PrOH	LP/SD	640 ^b , 900 590 ^b , 22000		Solvent was 2.6 mol L ⁻¹ 2-PrOH	83F405

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
278.	1-(Chloromethyl)naphthalene					
	Cyclohexane	LP	422		†Oxygen quenching	84A222
	MeOH	LP	422		†Triplet ET from benzophenone and thioxanthane	84A222
279.	1-Chloronaphthalene					
	1-Chloronaphthalene	FP-ET	427 403		†Triplet ET from benzophenone; 30 μ s delay; evidence for triplet excimer at long wavelengths presented; relative intensities (3:2); $\tau_T = 20 \mu$ s	767159
	Cyclohexane	FP	420		$\tau_T = 280 \mu$ s	62E009
	Cyclohexane (77 K)	PS	413.5 319		Relative intensities (10:6)	69E211
	Cyclohexane	LP/ET	420, 20700 \pm 3000		†Triplet ET from TMPD; ϵ relative to TMPD in cyclohexane ($\epsilon_{570} = 11900$ L mol ⁻¹ cm ⁻¹)	757282
	Durene (77 K)	PS	424.5 400		Single crystal	67B008
	Et ₂ O/Pentane (77 K)	PS	419 395.5 374.5		Relative intensities (10:6:1)	69E211
	Hexane	FP-ET	424 397 380		†Triplet ET from benzophenone; 30 μ s delay; relative intensities (6:3:1)	767159
	Hexane (293 K)	LP	423 ^a		Delay 500 ps	82E303
	PMMA (294 K)	FP	524 ^a		Shoulder at 552 nm ^a ; $\tau_T = (1.4 \pm 0.1) \times 10^5 \mu$ s	69B004
280.	2-Chloronaphthalene					
	Cyclohexane	FP	420		$\tau_T = 180 \mu$ s	62E009
	Cyclohexane (77 K)	PS	414 411 392.5 390 370		Relative intensities (10:-6:-2)	69E211
	EPA (77 K)	PS	420.7 397		†Phosphorescence decay	51E001
	PMMA (294 K)	FP	578 ^a 532 ^a		Shoulder at 637 ^a nm; $\tau_T = (2.5 \pm 0.2) \times 10^5 \mu$ s	69B004
281.	3-Chloro-1-(2-naphthyl)-1-propanone					
	Benzene	FP	425		$\tau_T = 220 \mu$ s	83P212
	CCl ₄	FP	400		$\tau_T = 251 \mu$ s	83P212
282.	1-(2-Chlorophenyl)naphthalene					
	3-MP (77 K)	PS/ESR	495, 9500 465 ^a , 7100 ^a 433 ^a , 3900 ^a		†ESR; oscillator strength = 0.12	69B002
283.	1-(4-Chlorophenyl)naphthalene					
	3-MP (77 K)	PS/ESR	505, 32500 472 ^a , 15800 ^a 435 ^a , 7400 ^a		†ESR; oscillator strength = 0.29	69B002
284.	Chlorophyll <i>a</i>					
	Acetone	LP/SD	460, 56000			84R057
	Benzene	FP	460		Half-life = $150 \pm 40 \mu$ s	55E003
	Cyclohexanol	FP	465 360		Half-life = $600 \pm 180 \mu$ s	55E003
	Cyclohexanol (294 K)	LP	1150 750		Half-life = 500μ s	81B111
	Digitonin (293 K)	LP	670 ^a 510 ^a		†Oxygen quenching (2.4×10^8 L mol ⁻¹ s ⁻¹); aqueous micellar suspension; $\tau_T = 830 \mu$ s	69N001
	Et ₂ O	FP/SD	430 ^b , 40000			67E027
	Heptane	FP/TD	480 ^b , 17700 \pm 3000 470 ^b , 23000 \pm 3000 460 ^b , 24000 \pm 4000		Solution contains pyridine; ϵ method assumed	77C001
	Heptane	FP/TD	480 ^b , 17100 \pm 3000		Solution contains ether; ϵ method assumed	77C001

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	MeOH	FP	462 425(?) 390(?)		Half-life = 200 ± 50 μ s	55E003
	MeOH	FP	550 450		550 nm maximum was judged to be of "doubtful significance" by the experimenters	58R002
	Polyvinylbutyral (77 K)	LP	472 ^a			80E867
	Polyvinylbutyral (77 K)	FP	460 ^a 425 ^a		Polymer film; $\tau_T = 2 \times 10^3$ μ s	82E129
	Pyridine (298 K)	FP/SD	530, 17800 462, 32000		$\tau_T = 1500$ μ s	58R001
	Pyridine	LP/COM	540, 35000 460, 48000		Triplet concentration calculated by kinetic simulation; $S_n \leftarrow S_1$ spectrum also resolved ($\epsilon_{440} = 68000$ L mol ⁻¹ cm ⁻¹); ϵ_{460} corrected from text	79B037
	Pyridine	LP	450 ^a		$\tau_T = 70$ μ s	81F121
	SHS	LP	740 ^a 465		Aqueous microemulsion, droplet radius ~ 100 Å; difference spectrum only; $\tau_T = 800$ μ s	80N057
285.	Chlorophyll <i>b</i>					
	3-MP	FP/SD	450 ^b , 26000			67E027
	Benzene	FP	482 440 420(?)		Half-life = 1300 ± 300 μ s	55E003
	Benzene	FP	580 475 440		440 and 580 nm maxima were judged to be of "doubtful significance" by the experimenters	58R002
	Benzene	FP	480 430		430 nm maximum was judged to be of "doubtful significance" by the experimenters; benzene was wet	58R002
	Benzene	FP/SD	316 ^a , 36500 ^a		$\tau_T = 2500$ μ s	59B002
	Et ₂ O	FP/SD	450 ^b , 21000			67E027
	MeOH	FP	496 454 420(?) 390		Half-life = 100 ± 40 μ s	55E003
	MeOH/Water	FP	~700 ~525 ~405		Solvent was 19:1 MeOH to water	55R001
	Pyridine (298 K)	FP/SD	605, 12100 550, 21500 485, 34700 445, 26000 380, 19700		$\tau_T = 3200$ μ s	58R001
286.	Chlorophyll <i>a</i> dimer					
	Heptane	FP/TD	480 ^b , 13500 ± 4000		Solution contains ether; ϵ method assumed	77C001
	Heptane	FP/TD	480 ^b , 17700 ± 3000 470 ^b , 21500 ± 4000 460 ^b , 25000 ± 4000		Solution contains pyridine; ϵ method assumed	77C001
287.	Chlorophylline					
	Water	FP	520 470		Sodium salt	65R023
288.	<i>cis</i> -4-Chlorostilbene					
	EtOH (123 K)	LP	390 370 347		Triplet state absorption observed only after repetitive (~30) flashes	79E640
289.	<i>trans</i> -4-Chlorostilbene					
	EPA (108 K)	LP	391 370 346		Triplet absorption not observable above 122 K; reported lifetime constant below 93 K; $\tau_T = 10^4$ μ s	79E640
	EtOH (103 K)	LP	390 369 348		Triplet absorption not observable above 135 K; reported lifetime constant below 111 K; $\tau_T = 10^4$ μ s	79E640

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Glycerol (197 K)	LP	395 376 355		Relative intensities (3:2:1); triplet absorption not observable above 275 K; reported lifetime constant below 202 K; $\tau_T = 5000 \mu\text{s}$	79E640
290.	2-Chlorothioxanthen-9-one Benzene	FP	650-670		\dagger Oxygen quenching ($5.5 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); $\tau_T = 61 \mu\text{s}$	81A294
291.	6-Chloro-1,3,3-trimethyl-6'-nitrospiro-(2,2'-indolin[2H-1]benzopyran) Toluene (298 K)	FP	450		\dagger Oxygen quenching, activationless decay; assigned to $\pi \rightarrow \pi^*$ transition in uncolored form	67B003
292.	Chlorpromazine 2-PrOH	LP	460		\dagger Triplet ET to β -carotene, oxygen quenching; peak at 510 nm assigned to dechlorinated radical; $\tau_T = 3.2 \mu\text{s}$	767023
293.	4-Chromanone Benzene	LP	$\sim 450^a$		110 ps delay; $\tau_T = 0.3 \mu\text{s}$; rise time of $(50 \pm 10) \times 10^{-3} \text{ ns}$	81E222
	EPA (77 K)	FP	427 ^a		\dagger Phosphorescence decay	80E884
	EPA (77 K)	FP	430 ^a		\dagger Phosphorescence decay; $\tau_T = 93 \times 10^3 \mu\text{s}$	81E222
	EtOH	LP	557 ^a		\dagger Oxygen quenching; shoulder at 511 ^a nm; 442 nm peak more intense; delay 150 ns; $\tau_T = 0.406 \mu\text{s}$	80E884
			442 ^a			
	EtOH	LP	560 ^a 510 ^a 450 ^a		150 ns delay; relative intensities (1:2:3); $\tau_T = 0.4 \mu\text{s}$; rise time of $(50 \pm 10) \times 10^{-3} \text{ ns}$	81E222
294.	Chromone EPA (77 K)	FP	650 ^a		\dagger Phosphorescence decay; lifetime lengthens in polar glasses; $\tau_T = (35 \pm 10) \times 10^3 \mu\text{s}$	81E222
295.	Chrysene 1-BuOH/Isopentane (77 K)	PS	583.0 540.5 406.0		A distinction between a 1:7 and a 3:7 glass was not made	63B001
	2-MTHF (77 K)	PS/ESR	585, 48600 461 ^a , 3900 ^a 418 ^a , 6600 ^a		\dagger ESR; shoulders at 571 ^a , 543 ^a , and 495 ^a nm; the band was assigned as $^3A_g \leftarrow ^3B_u$; oscillator strength = 0.40	68D211
	2-MTHF (77 K)	PS	586 ^a 460 ^a 419 ^a		Shoulders at 574 ^a , 545 ^a , and 496 ^a nm	69E210
	Benzene	PR	575			761024
	Cyclohexane	LP/ET	565, 21600 \pm 2000		\dagger Triplet ET to anthracene; ϵ relative to anthracene in cyclohexane ($\epsilon_{422.5} = 64700 \text{ L mol}^{-1} \text{ cm}^{-1}$)	757282
	EPA (77 K)	PS	584.1		\dagger Phosphorescence decay; 1st transition assigned as $^3L_a \rightarrow ^3C_b$	51E001
	EPA (77 K)	PS	629 583.1 406 385		\dagger Phosphorescence decay; relative intensities (11:100:18:15); unresolved shoulders at 563 nm and 541 nm; $\tau_T = (2.6 \pm 0.2) \times 10^6 \mu\text{s}$	54B001
	EPA (77 K)	PS	730 625		Weak bands in the red only studied	67B005
	EPA (77 K)	PS/SD	585, 48000 \pm 4800			68E105
	EPA (77 K)	PS/IV	585, 36200		λ_{\max} assumed from previous work; ϵ estimated by extrapolation to infinite excitation rate	69E212
	EPA (77 K)	PS	729.9 625.0 580.0 559.9 535.0 400.0 379.9		Relative intensities (6:3:100:60:30:12:10)	69E211
	EtOH (293 K)	FP	562.5			68E098
	EtOH	LP	560 ^a		1.8 μs delay; solution contains Ag ions	78E554

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	EtOH/Et ₂ O (77 K)	MOD/KM	580, 23100 ± 6900		Glass was 2:1 EtOH to Et ₂ O; temperature was not explicitly stated, but 77 K was inferred from the context	719059
	Hexane	FP	554.9			54E001
	Hexane	FP/SD	570, 8800 401, 1800 378, 1500		Bands were assigned to 2 electronic transitions; 570 nm band was assigned to the 1st electronic transition; oscillator strength = 0.15, 0.025	58E001
	Hexane (77 K)	PS	580.0 534.8 400.0		Relative intensities (100:30:12)	69E211
	Hexane (300 K)	MOD/SD	564 ^a , 23500 ^a 400 ^a , 1700 ^a 285, 26200 ^a 254, 25800 ^a		Shoulder at 517 ^a nm; in their later compilation [73E284], the authors normalize to $\epsilon_{585} = 48000$ L mol ⁻¹ cm ⁻¹ which means all of their values reported here should be multiplied by a factor of about 2; $\tau_T = 710 \pm 25$ μ s	69E208
	Polystyrene	PR	569 ^a		$G\epsilon_{569} = 6500^a$ L mol ⁻¹ cm ⁻¹ / (100 eV absorbed)	701073
	Water/ <i>tert</i> -BuOH	FP	560 ^a		Solvent mixture contains "46%" <i>tert</i> -BuOH for solubility; pH ~6	767189
296.	9-Cinnamoylanthracene EPA (77 K)	FP	430		"9-Anthryl styryl ketone"; $\tau_T = 3.43 \times 10^4$ μ s	82E338
297.	Copper(II) phthalocyanine 1-Chloronaphthalene	LP/TD	480 470 ^b , 3600		†Triplet ET to β -carotene; $\tau_T = 0.035 \pm 0.005$ μ s; $E_T = 150 \pm 10$ kJ mol ⁻¹	78A378
298.	Coproporphyrin III MeOH/Water	LP/TD	401 ^a , 144000 ^a		Solvent 9:1 MeOH to 7.4 phosphate buffer; shoulders at 373 ^a and 326 ^a nm	83E667
	Water	LP/TD	401 ^a , 86100 ^a 372 ^a , 95800 ^a		Phosphate buffer; pH 7.4	83E667
299.	Coproporphyrin III, tetramethyl ester Benzene	LP/TD	440 ^b , 26000 401 ^a		†Oxygen quenching (1.6×10^9 L mol ⁻¹ s ⁻¹); $\tau_T = 240$ μ s	80E200
	Benzene	PR/ET	440 ^b , 27000 401 ^a		†Oxygen quenching (1.6×10^9 L mol ⁻¹ s ⁻¹); ϵ relative to biphenyl in benzene ($\epsilon_{360} = 27100$ L mol ⁻¹ cm ⁻¹); $\tau_T = 240$ μ s	80E200
300.	Coproporphyrin I, tetramethyl ester Benzene	PR/ET	440 ^b , 26500 398 ^a 330 ^a		†Oxygen quenching (1.8×10^9 L mol ⁻¹ s ⁻¹); ϵ relative to biphenyl in benzene ($\epsilon_{360} = 27100$ L mol ⁻¹ cm ⁻¹); $\tau_T = 250$ μ s	80E200
	Benzene	LP/TD	440 ^b , 32000 398 ^a 330 ^a		†Oxygen quenching (1.8×10^9 L mol ⁻¹ s ⁻¹); $\tau_T = 250$ μ s	80E200
301.	Coronene 2-MTHF (77 K)	PS/ESR	484, 15800		†ESR; only most intense visible peak reported; assignment $^3A_1 \leftarrow ^3B_2$; oscillator strength = 0.23	696115
	?	FP/?	700 ^b , 900 460, 15000		Method and solvent unspecified	78A163
	Benzene	PR	480			761024
	Dioxane	LP	490		†Rise time of transient was the same as the decay time of the singlet	68B006
	Dioxane	LP	525 480			70E288
	EtOH (293 K)	FP	478			68E098
	EtOH	LP	480 ^a		1.8 μ s delay; solution contains Ag ions	78E554
	EtOH/Et ₂ O (77 K)	MOD/KM	460, 14300 ± 4300		Glass was 2:1 EtOH to Et ₂ O; temperature was not explicitly stated, but 77 K was inferred from the context	719059
	Hexane	FP	630.1 490.2 461.9		Band at 461.9 was later assigned as $S_n \leftarrow S_1$ by authors in [737463]	54E001

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Hexane	FP/SD	524.5, 330 ^d 480, 680 ^d 461, 1000 ^d		Band at 461 nm was assigned to a second electronic transition; band at 461 nm was later reassigned as S _n ← S ₁ by authors in [737463]; oscillator strength = —, 0.02 ‡Decay of singlet was roughly the same as the rise of the transient	58E001 686058
	PMMA (296 K)	LP	480 390			70E288
	PMMA	LP	525 480			
	Polystyrene	PR	459 ^a		$G\epsilon_{459} = 7900^a$ L mol ⁻¹ cm ⁻¹ / (100 eV absorbed)	701073
	Toluene	LP	595 565 530 495 465 ^a		Delay 880 ns	737463
	Water/2-PrOH	FP			Solvent mixture contains "40%" 2-propanol for solubility; pH ~6	767189
302.	Coronene- <i>d</i> ₁₂					
	PMMA (77 K)	PS/SD	637 ^a , 10300 ^a 588 ^a , 5100 ^a 549 ^a , 5400 ^a 483 ^a , 17500 ^a 398 ^a , 13400 ^a		$\tau_T = 3.5 \times 10^5$ μ s	69E216
	PMMA (296 K)	PS/SD	633 ^a , 8600 ^a 588 ^a , 5200 ^a 549 ^a , 5400 ^a 483 ^a , 16800 ^a 398 ^a , 14600 ^a		$\tau_T = 2.0 \times 10^5$ μ s	69E216
303.	Coumarin					
	Benzene	PR/ET	400, 11000		‡Triplet ET from biphenyl; ϵ relative to biphenyl in benzene ($\epsilon_{367} = 27100$ L mol ⁻¹ cm ⁻¹); 20 ns delay; $\tau_T = 3.8$ μ s; $k_{et} = 3.4 \times 10^9$ L mol ⁻¹ s ⁻¹	79E282
	EPA (77 K)	PS	460.8 451 432.1 423 406.5 389 ^a		‡Phosphorescence decay; shoulders at 441 ^a and 413 ^a nm; most intense peaks were at 460.8, 432.1, and 406.5 nm; $\tau_T = (7 \pm 2) \times 10^5$ μ s	71B001
	EPA	CWL	460 ^a 433 ^a 417 ^a 392 ^a		Shoulder at 449 ^a nm; glass, temperature unspecified	73E344
	Water	LP	400		0.1 μ s delay; $\tau_T = 1.3$ μ s	79E282
304.	<i>p</i> -Cresol					
	Water	LP	255 ^a		‡Triplet ET to tryptophan; lifetime was measured at pH 7.5; there was a broad maximum between ~350 and ~420 nm; spectrum is the difference between spectra at 20 ns and 15 μ s; $\tau_T = 3.4 \pm 0.2$ μ s; pH 7.7	757161
305.	α -Crocin					
	Water	LP/ET	470 ^b , 73000 445 ^a , 94800 ^a		‡Oxygen quenching; ϵ relative to psoralen in water ($\epsilon_{460} = 10000$ L mol ⁻¹ cm ⁻¹); $\tau_T = 8$ μ s; pH 8	83B068
306.	Croconate Blue dianion					
	EtOH	LP-ET/SD	590 ^b , 25000		‡Triplet ET from 9,10-dibromoanthracene and oxygen quenching; $\tau_T = 69$ μ s; $k_{et} = 4.5 \times 10^9$ L mol ⁻¹ s ⁻¹	84N017
	Water	LP-ET/SD	585 ^b , 20000		‡Triplet ET from 9,10-dibromoanthracene and oxygen quenching; $\tau_T = 32$ μ s	84N017
307.	Crystal Violet					
	PMMA (193 K)	PS	640		Shoulder at 620 nm	59B001
	PMMA (77 K)	PS	1280 1210 1100 750 640 620			65B004

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	PMMA (77 K)	PS	1241 ^a 613 ^a		Relative intensities (3:20); shoulder at 532 ^a nm; solvent contains 2-chloroethanol	69B003
308.	9-Cyanoanthracene					
	Acetonitrile	FP	433		$\tau_T = 560 \mu s$	737140
	Acetonitrile	LP	425		†Oxygen quenching; intersystem crossing promoted by thioanisole	83E025
	Benzene	PR/ET	435, 10300		ϵ relative to naphthalene in benzene ($\epsilon_{\max} = 17500$ L mol ⁻¹ cm ⁻¹)	690087
	Benzene	FP	440		$\tau_T = 530 \mu s$	737140
	Cyclohexane	PR	425			690087
	Cyclohexane	FP	433		$\tau_T = 1700 \mu s$	737140
	Dioxane	PR	432.5			690087
309.	4-Cyanobiphenyl					
	Toluene/EtOH (77 K)	MOD	404 ^a		Glass was 19:1 toluene to EtOH	719059
310.	4'-Cyanochalcone					
	Heptane	LP	360 ± 15		"4-(1-Oxo-3-phenyl-2-propenyl)benzonitrile"; $\tau_T = 0.020 \mu s$	83E347
311.	<i>trans</i> -4-Cyano-4'-dimethylaminostilbene					
	2-MTHF (133 K)	LP	~700 460			80F299
	Glycerol triacetate (198 K)	LP	~700 460			80F299
312.	<i>N</i> -[5-(2-Cyanoethyl)(2-hydroxyethyl)amino]-2-[(2,4-dinitrophenyl)azo]-4-methoxyphenylacetamide					
	2-MTHF (77 K)	LP	700 410		†Oxygen quenching; triplet not observable above 153 K; below 117 K lifetime constant; $\tau_T = 1000 \mu s$	80B101
	Dibutyl terephthalate	LP	~700		$\tau_T = <0.012 \mu s$	80B101
	Glycerol triacetate	LP	~700			80B101
	Glycerol triacetate (198 K)	LP	680 420		Lifetime measured at 268 K; triplet virtually not observable above 268 K; below 213 K lifetime constant; $\tau_T = 0.025 \mu s$	80B101
	PMMA	LP	660			80B101
313.	<i>trans</i> -4-Cyano-4'-methoxystilbene					
	2-MTHF (143 K)	LP	490 410		Triplet absorption not observable above 143 K; below 106 K lifetime ($1 \times 10^4 \mu s$) constant; $\tau_T = 3.3 \mu s$	80F299
	EtOH (132 K)	LP	480 420		Triplet absorption not observable above 132 K; below 113 K lifetime ($1.3 \times 10^4 \mu s$) constant; lifetime measured at 132 K; $\tau_T = 5.0 \mu s$	80F299
	Glycerol triacetate (198 K)	LP	450		No triplet observed at room temperature; $\tau_T = \sim 7000 \mu s$	78B088
	Glycerol (198 K)	LP	490 420			80F299
	Glycerol triacetate (223 K)	LP	450		Triplet absorption not observable above 238 K; below 206 K lifetime ($1 \times 10^4 \mu s$) constant; lifetime measured at 238 K; $\tau_T = 2.0 \mu s$	80F299
	PMMA (298 K)	LP	430			80F299
314.	1-Cyanonaphthalene					
	Cyclohexane (77 K)	PS	413 390			69E211
	Et ₂ O/Pentane (77 K)	PS	445 423.5 412 387			69E211
	MeOH	FP	440 ^a			736218
315.	9-Cyano-10-nitroanthracene					
	Benzene	LP	~460 ^a		Delay 3.0 μs	80E884
	EPA (77 K)	LP	430 ^a			84F385

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
316.	<i>trans</i> -4-Cyanostilbene					
	2-MTHF (98 K)	LP	405		Triplet absorption not observable above 127 K; below 104 K lifetime constant; $\tau_T = 1 \times 10^4 \mu s$	80F299
	EtOH (118 K)	LP	400		Triplet absorption not observable above 138 K; below 110 K lifetime ($1.7 \times 10^4 \mu s$) constant; lifetime measured at 138 K; $\tau_T = 1.7 \mu s$	80F299
	Glycerol triacetate (198 K)	LP	405		Triplet absorption not observable above 239 K; below 210 K lifetime constant; $\tau_T = 1 \times 10^4 \mu s$	80F299
	PMMA (298 K)	LP	395			80F299
317.	1,3-Cycloheptadiene					
	Benzene	PR	310		$\tau_T = 0.39 \mu s$	82B057
318.	2-Cyclohepten-1-one					
	Cyclohexane	LP	270 ^a		†Oxygen quenching; solvent uncertain; $\tau_T = 0.011 \mu s$	80B055
319.	1-(1-Cyclohepten-1-yl)naphthalene					
	Benzene	LP-ET	443 ^a 381 ^a		†Triplet ET from xanthone and oxygen quenching; $\tau_T = 0.053 \mu s$	84B007
320.	2-(1-Cyclohepten-1-yl)naphthalene					
	Benzene	LP-ET	390 ^a		†Triplet ET from xanthone and oxygen quenching ($7.0 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); $\tau_T = 0.04 \mu s$	84B090
321.	1,3-Cyclohexadiene					
	Benzene	LP-ET/TD	302.5, 2300 \pm 300		†Triplet ET from 2-fluorenyl phenyl ketone and other triplet sensitizers; ϵ assumes ET proceeds with unit efficiency; $\tau_T = 30 \mu s$; $E_T = \sim 220 \text{ kJ mol}^{-1}$	80B021
	Benzene	PR	310		$\tau_T = 5.2 \mu s$	82B057
322.	2-Cyclohexen-1-one					
	Cyclohexane	LP/RA	270 ^b , ≥ 2100 265 ^a		†Oxygen quenching ($5 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); solvent uncertain; ϵ relative to benzophenone in benzene ($\epsilon_{535} = 7600 \text{ L mol}^{-1} \text{ cm}^{-1}$, $\Phi_T = 1$); ϵ is lower limit since $\Phi_T < 1$; $\tau_T = 0.023 \mu s$	80B055
323.	1-(1-Cyclohexen-1-yl)naphthalene					
	Benzene	LP-ET	442 ^a		†Triplet ET from xanthone and oxygen quenching; $\tau_T = 2 \mu s$	84B007
324.	1,3-Cyclooctadiene					
	Benzene	PR	315		$\tau_T = 0.14 \mu s$	82B057
325.	1-(1-Cycloocten-1-yl)naphthalene					
	Benzene	LP-ET	446 ^a 381 ^a		†Triplet ET from xanthone and oxygen quenching; $\tau_T = 0.12 \mu s$	84B007
326.	Cyclopentadiene					
	Benzene	PR	300		$\tau_T = 1.6 \mu s$	82B057
327.	2-Cyclopentenone					
	Cyclohexane	LP/RA	270 ^b , ≥ 1900 260 ^a		†Oxygen quenching ($5 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); solvent uncertain; ϵ relative to benzophenone in benzene ($\epsilon_{535} = 7600 \text{ L mol}^{-1} \text{ cm}^{-1}$, $\Phi_T = 1$); ϵ is lower limit since $\Phi_T < 1$; $\tau_T = 0.030 \mu s$	80B055
328.	1-(1-Cyclopenten-1-yl)naphthalene					
	Benzene	LP-ET	441 ^a		†Triplet ET from xanthone and oxygen quenching; $\tau_T = 3.6 \mu s$	84B007
329.	2-(1-Cyclopenten-1-yl)naphthalene					
	Benzene	LP-ET	581 ^a 394 ^a		†Triplet ET from xanthone and oxygen quenching ($3.3 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); 394 nm peak was the more intense	84B090

TABLE 6 Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
330.	1,2,3,4,5,6,7,8,9,10-Decahydro-1,1,5,5,6,6,10,10-octamethylcyclobuta[1,2,3,4]dicycloheptene Benzene	LP-ET	< 300		†Triplet ET from 2,3-dimethyl-1,4-naphthoquinone; $\tau_T = 0.24 \mu\text{s}$; $E_T = > 120 \text{ kJ mol}^{-1}$; $k_{et} = 7 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	81E098
331.	Decapreno- β -carotene Benzene (296 K)	LP-ET	595		Triplet ET from chlorophyll <i>a</i> ; oxygen quenching ($7.0 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); $\tau_T = 3.2 \mu\text{s}$; $k_{et} = 1.8 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	73E347
	Benzene	PR-ET	590		†Triplet ET from naphthalene; $\tau_T = 6.5 \mu\text{s}$	80A143
332.	Deoxybenzoin Benzene	LP	500 < 390 ^a		†Oxygen quenching ($\sim 2 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); $\tau_T = > 0.24 \mu\text{s}$	79A289
	Benzene	LP	450 315		†Oxygen quenching ($> 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); relative intensities (1.4); $\tau_T = 0.77 \mu\text{s}$	79A028
	MeOH	LP	495 ^a		Oxygen quenching ($2.4 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); $\tau_T = 0.14 \mu\text{s}$	77E797
333.	Deuteroporphyrin, dimethyl ester Benzene	LP/TD	440 ^b , 22000 410 ^a		†Oxygen quenching ($2.3 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); $\tau_T = 220 \mu\text{s}$	80E200
	Benzene	PR/ET	440 ^b , 28000 410 ^a		†Oxygen quenching ($2.3 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); ϵ relative to biphenyl in benzene ($\epsilon_{360} = 27100 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 210 \mu\text{s}$	80E200
334.	<i>trans</i> - <i>N,N'</i> -Diacetylidindigo EPA (83 K)	LP	575 ^a 365 ^a		Relative intensities (2:1); data corrected from paper (H. Goerner, private communication, 1985); $\tau_T = 80 \mu\text{s}$	79E543
	EPA (283 K)	LP	580 ^a 375 ^a		Relative intensities (2:1); no transient spectrum observed at RT; data corrected from paper (H. Goerner, private communication, 1985); $\tau_T = 0.4 \mu\text{s}$	79E543
	Glycerol triacetate (203 K)	LP	580 ^a 390 ^a		Data corrected from paper (H. Goerner, private communication, 1985)	79E543
335.	1,4-Diaminoanthraquinone Benzene	PR-ET/ET	575, 22000		†Triplet ET from biphenyl; ϵ relative to biphenyl in benzene assuming ground state dimerization; $\tau_T = 4.5 \mu\text{s}$; $E_T = 123 - 151 \text{ kJ mol}^{-1}$; $k_{et} = 2.2 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	761122
336.	3,6-Diaminophthalimide EtOH	FP	545		Lifetime limited by aeration; $\tau_T = 0.560 \mu\text{s}$	757522
337.	5,5'-Di- <i>tert</i> -amylthioindigo 2-MTHF (93 K)	LP	595 390		$\tau_T = 120 \mu\text{s}$	79E543
	2-MTHF	LP	590 385		$\tau_T = 0.25 \mu\text{s}$	79E543
	EPA (77 K)	FP	590		$\tau_T = 154 \mu\text{s}$	767582
	EPA (93 K)	LP	580 370		$\tau_T = 53 \mu\text{s}$	79E543
	EPA (283 K)	LP	590 370		Φ_T strongly temperature dependent in this solvent; $\tau_T = 0.40 \mu\text{s}$	79E543
	Glycerol triacetate (203 K)	LP	605 390		Relative intensities (5:4); $\tau_T = 33 \mu\text{s}$	79E543
	Glycerol triacetate	LP	600 385		Relative intensities (5:4); $\tau_T = 0.36 \mu\text{s}$	79E543
	MCH	LP	595 380		$\tau_T = 0.25 \mu\text{s}$	79E543
	PMMA	LP	590 390			79E543

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
338.	Dibenz[a,j]acridine/PrOH 3-MP/PrOH (77 K)	PS	550		Glass was 99:1 3-MP to PrOH; which isomer of PrOH was not specified; no T-T absorption in 3-MP; "1,2,7,8-dibenzacridine"	706135
339.	Dibenz[a,h]acridine 3-MP (77 K)	PS	585 545 520		"1,2,5,6-Dibenzacridine"	706135
340.	Dibenz[a,h]anthracene/Chloranil PMMA	PS	668 ^a 626 ^a 580 ^a 540 ^a 508 ^a 495 ^a 470 ^a 455 ^a 440 ^a 424 ^a		Most intense peak at 580 nm	766652
341.	Dibenz[a,h]anthracene 2-MTHF (77 K)	PS/ESR	587, 25000		†ESR; only most intense visible peak reported; assignment ³ A _g ← ³ B _u ; oscillator strength encompasses two electronic transitions; "1,2,5,6-dibenzanthracene"; oscillator strength = 0.25	696115
	Benzonitrile	LP/RF	583 ^a , 10500 ± 700 548 ^a , 9400 ^a 509 ^a , 7900 ^a 476 ^a , 6000 ^a 445 ^a , 5000 ^a		ε relative to compound in toluene (ε ₅₈₄ = 13000 L mol ⁻¹ cm ⁻¹)	83F075
	EPA (77 K)	PS	585 546 506 495		†Phosphorescence decay	51E001
	EPA (77 K)	PS	584.8 547.1 508 500 472 428		†Phosphorescence decay; relative intensities (100:67:39:40:30:24); τ _T = (1.4 ± 0.1) × 10 ⁶ μs	54B001
	EPA (77 K)	PS/SD	585, 35000 ± 3500			68E105
	EPA (77 K)	PS/IV	585, 28700		λ _{max} assumed from previous work; ε estimated by extrapolation to infinite excitation rate	69E212
	EtOH/Et ₂ O (77 K)	MOD/KM	560, 17600 ± 5300		Glass was 2:1 EtOH to Et ₂ O; temperature was not explicitly stated, but 77 K was inferred from the context	719059
	Hexane	FP	535.1 498.0 452.9 421.1			54E001
	Hexane	FP/SD	532.5, 67000 499, 57600 469.5, 46900 424, 33400 398, 18100 376.5, 9400 327		The bands between 532.5 and 469.5 nm were assigned to the 1st electronic transition, the next three bands to the 2nd electronic transition, and the last band to the 3rd electronic transition; oscillator strength = 2.2, 0.4, —	58E001
	Liquid paraffin	FP/TD	535 ^b , 22000 ± 3000			67E031
	Liquid paraffin	MOD	573 538 510		Mull; relative intensities (100:91:59); τ _T = 90.3 μs	71E361

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	PMMA	LP	532 480			70E288
	Toluene	LP/ET	584 ^a , 13000 ± 1000 544 ^a , 12100 ^a 509 ^a , 8300 ^a 478 ^a , 7000 ^a 444 ^a , 3300 ^a	ϵ relative to anthracene in toluene ($\epsilon_{428.5} = 42000$ L mol ⁻¹ cm ⁻¹)		83F075
342.	7H-Dibenzo[c,g]carbazole Cyclohexane	LP	~600			82E497
343.	Dibenzo[b,def]chrysene Hexane	MOD	511 ^a 480 ^a		"3,4,8,9-Dibenzpyrene"; 511 nm was the more intense peak; $\tau_T = 130$ μ s	70E295
	Hexane	MOD	510 470		Relative intensities (100:28); $\tau_T = 130$ μ s	71E361
	Liquid paraffin	MOD	520 490		Mull; relative intensities (100:37); $\tau_T = 110$ μ s	71E361
344.	Dibenzo[def,mno]chrysene Heptane	LP	662 633 581 535 524 508		"Anthanthrene"; delay 370 ns	737463
	Liquid paraffin	MOD	585 420 395		Mull; relative intensities (—:100:19); shoulder at 560 nm with relative intensity 34; $\tau_T = 103$ μ s	71E361
345.	Dibenzo[def,p]chrysene Hexane	MOD	573 477		Relative intensities (23:100); "1,2,3,4-dibenzpyrene"; $\tau_T = 120$ μ s	71E361
346.	Dibenzo[def,mno]chrysene-6,12-dione Benzene	FP	688 ^a 520 ^a		"Anthanthrone"; 688 nm peak was the more intense	84F379
347.	Dibenzo[g,p]chrysene Heptane	LP	633 613 575 532 503 483 442		"1,2,3,4,5,6,7,8-Tetrabenzonaphthalene"; delay 380 ns	737463
348.	5H-Dibenzo[a,d]cycloheptene Cyclohexane	FP	425		Decay followed in acetonitrile and mixed hydrocarbons as a function of temperature	80E375
	Isopentane/3-MP (77 K)	FP/IV	426, 20000 402 ^a		Glass was 6:1 isopentane to 3-MP; shoulder at 373 ^a nm; $\tau_T = 7100$ μ s	687111
349.	Dibenzophenazine Toluene (293 K)	FP	530 ^a 510 ^a 480 ^a		‡Phosphorescence decay in EPA at 77 K, oxygen quenching; isomer unspecified	80E778
350.	6H-Dibenzo[b,d]pyran-6-one EtOH (300 K)	FP	500 390			81E650
351.	Dibenzo[f,h]quinoxaline PMMA (77 K)	PS	460 429 387		‡Phosphorescence decay (?); "5,6,7,8-dibenzoquinoxaline"; $\tau_T = 9.4 \times 10^3$ μ s	70E291

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
352.	Dibenzo[<i>c,g</i>]triphenylene MCH	LP	605 420		†Oxygen quenching; relative intensities (2:1); $\tau_T = 120$ μ s	79A237
353.	2,3-Dibenzoylbicyclo[2.2.1]hepta-2,5-diene MeOH	LP	445 ^a		Assignment uncertain; shoulder around 400 nm; $\tau_T = 0.054$ μ s	80F372
354.	2,3-Dibenzoylbicyclo[2.2.2]octa-2,5-diene MeOH	LP	450 ^a		Assignment uncertain; shoulder around 520 nm; $\tau_T = 0.040$ μ s	80F372
355.	Dibenzoylmethane EtOH/MeOH (118 K)	FP	630		Solvent was 3:1 EtOH to MeOH	68B005
356.	<i>N</i> -(Dibenzylamino)methylphthalimide EtOH	FP	590 350		†Oxygen quenching, diene quenching; maxima assumed from text; $\tau_T = 12$ μ s	79A147
357.	2,5-Di(4-biphenyl)oxazole 2-MTHF (77 K) Benzene	CWL LP/SD	590 560, 110000		"BBO" †Oxygen quenching (1.7×10^9 L mol ⁻¹ s ⁻¹); 100 ns delay; $\tau_T = 0.285$ μ s	74B003 777265
358.	9,10-Dibromoanthracene Acetonitrile Cyclohexane EPA (77 K) EtOH Liquid paraffin Toluene	LP FP LP I.P./ET FP LP/ET	421 ^a 419 427 420, 82000 424 427.5, 48000 \pm 4000		†Oxygen quenching; $\tau_T = 19.5$ μ s $\tau_T = 36$ μ s $\tau_T = 22$ μ s ϵ relative to Methylene Blue cation ($\epsilon_{660} = 27000$ L mol ⁻¹ cm ⁻¹); solvent and temperature assumed; $\tau_T = 11$ μ s Viscosity of solvent was 0.167 N·s/m ² ; $\tau_T = 44$ μ s ϵ relative to rubrene in toluene ($\epsilon_{500} = 26000$ L mol ⁻¹ cm ⁻¹)	84B110 62E009 84B110 78E394 62E009 84E393
359.	4,4'-Dibromobiphenyl Hexane (293 K) Toluene	LP PR	~400 390		Delay 500 ps †Oxygen quenching; $\tau_T = \sim 3$ μ s	82E303 80A235
360.	Dibromofluorescein dianion Water	FP/SD	506 ^b , 18000 \pm 6000		pH 9	67E031
361.	1,4-Dibromonaphthalene Cyclohexane Liquid paraffin	FP FP	425 423		$\tau_T = 150$ μ s Viscosity of solvent was 0.167 N·s/m ² ; $\tau_T = 2000$ μ s	62E009 62E009
362.	(<i>E</i>)- <i>N,N</i> -Dibutyl-2,3-dihydro-3-oxo-2-(3-oxonaphtho[1,2- <i>b</i>]thien-2(3 <i>H</i>)-ylidene)naphtho[1,8- <i>bc</i>]thiopyran-6-sulfonamide Chloroform	LP	730		A substituted "trans-perinaphthioindigo"	83A355
363.	(<i>Z</i>)- <i>N,N</i> -Dibutyl-2,3-dihydro-3-oxo-2-(3-oxonaphtho[1,2- <i>b</i>]thien-2(3 <i>H</i>)-ylidene)naphtho[1,8- <i>bc</i>]thiopyran-6-sulfonamide Chloroform	LP	670		A substituted "cis-perinaphthioindigo"	83A355
364.	5,5'-Dibutylthioindigo EPA (77 K)	FP	585		$\tau_T = 152$ μ s	767582
365.	5,5'-Di- <i>tert</i> -butylthioindigo EPA (77 K)	FP	595		$\tau_T = 133$ μ s	767582
366.	<i>trans</i> -1,2-Di(<i>N</i> -carbazolyl)cyclobutane DMF THF	LP LP	420 424 ^a		†Oxygen quenching; 500 ns delay Delay 800 ns; shoulders at 504 ^a and 404 ^a nm	81P245 83E662
367.	<i>meso</i> (DL)-2,4-Di(<i>N</i> -carbazolyl)pentane THF	LP	424 ^a		Delay 700 ns; shoulders at 501 ^a and 399 ^a nm	83E662

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
368.	<i>rac</i> (DD,LL)-2,4-Di(<i>N</i> -carbazolyl)pentane THF	LP	425 ^a		Delay 1 μ s; shoulders at 498 ^a and 402 ^a nm	83E662
369.	1,5-Dichloroanthracene					
	Benzene	PR/ET	445, 63400		ϵ relative to naphthalene in benzene ($\epsilon_{\max} = 17500$ L mol ⁻¹ cm ⁻¹)	690087
	Benzene	PR/ET	445, 64300		ϵ relative to benzophenone in benzene ($\epsilon_{532.5} = 10300$ L mol ⁻¹ cm ⁻¹)	690087
	Cyclohexane	FP	425		$\tau_T = 60$ μ s	62E009
	Cyclohexane	PR	440			690087
	Cyclohexane	FP/RA	955, 1000 850 760 442.5 420		†Oxygen quenching; ϵ relative to 1,5-dichloroanthracene in benzene ($\epsilon_{445} = 64300$ L mol ⁻¹ cm ⁻¹)	71B005
	Dioxane	PR	442.5			690087
	Liquid paraffin	FP	419		Viscosity of solvent was 0.167 N·s/m ² ; $\tau_T = 2000$ μ s	62E009
	PFMCH	FP	951 ^a 840 ^a 755 ^a 430 ^a 410 ^a		†Oxygen quenching	71B005
	PMMA	FP	855			66E086
370.	9,10-Dichloroanthracene					
	Acetonitrile	FP	418 ^a			82F494
	Benzene	PR/ET	422.5, 41700		ϵ relative to naphthalene in benzene ($\epsilon_{\max} = 17500$ L mol ⁻¹ cm ⁻¹)	690087
	Benzene	PR/ET	422.5, 51100		ϵ relative to benzophenone in benzene ($\epsilon_{532.5} = 10300$ L mol ⁻¹ cm ⁻¹)	690087
	Cyclohexane	FP	419		$\tau_T = 100$ μ s	62E009
	Cyclohexane	PR	417.5			690087
	Dioxane	PR	420			690087
	EtOH (298 K)	FP	419 ^a			66E086
	EtOH (201 K)	FP	421 ^a			66E086
	EtOH	FP/ET	694 ^b , 64 417, 46000		Triplet ET from eosin; ϵ relative to eosin in EtOH ($\epsilon_{580} = 9400$ L mol ⁻¹ cm ⁻¹)	78E019
	Liquid paraffin	FP	423		Solvent viscosity was 0.03 N·s/m ²	58E001
	Liquid paraffin	FP	419		Viscosity of solvent was 0.167 N·s/m ² ; $\tau_T = 500$ μ s	62E009
	Toluene	LP/ET	426, 46000 \pm 4000		ϵ relative to rubrene in toluene ($\epsilon_{500} = 26000$ L mol ⁻¹ cm ⁻¹)	84E393
371.	1,8-Dichloroanthraquinone					
	EPA (77 K)	LP	500 ^a 370 ^a		†Phosphorescence decay; delay 100 ns; 370 nm was the more intense peak; $\tau_T = 40 \pm 10$ μ s	83E016
	EPA (77 K)	LP	375 ^a		Delay 100 ns	84A253
	EtOH	LP	380 ^a		Delay 50 ns	84A253
	Toluene	LP	500 ^a 379 ^a		†Phosphorescence decay in EPA at 77 K; delay 300 ns; 379 nm was more intense peak	83E016
372.	4,4'-Dichlorobenzophenone					
	Acetonitrile/Water	LP/ET	545, 9600 320, 12900		Solvent was 9:1 acetonitrile to water; ϵ relative to 1-methylnaphthalene in acetonitrile/water ($\epsilon_{415} = 11200$ L mol ⁻¹ cm ⁻¹)	84B033
373.	Dichlorobis(4,7-dimethyl-1,10-phenanthroline)iridium(III) ion					
	DMF/Water	LP	608 ^a 573 ^a 543 ^a 489 ^a 464 ^a 447 ^a 425 ^a		Shoulders at 530 ^a and 511 ^a nm; solvent was 9:11 dimethylformamide to water	79B098

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	DMF/Water	LP	626 ^a 566 ^a 547 ^a 535 ^a 516 ^a 497 ^a 468 ^a		Shoulders at 588 ^a , 458 ^a , and 439 ^a nm; solvent was 19:1 dimethylformamide to water	79B098
	Water	LP	588 ^a 552 ^a 531 ^a 490 ^a 461 ^a 444 ^a 424 ^a		Shoulders at 509 ^a and 469 ^a nm	79B098
374.	Dichlorobis(5,6-dimethyl-1,10-phenanthroline)iridium(III) ion					
	DMF/Water	LP	650 ^a 566 ^a 512 ^a 440 ^a		Solvent was 19:1 dimethylformamide to water	81E786
	DMF/Water	LP	572 ^a 510 ^a 438 ^a		Solvent was 9:11 dimethylformamide to water; shoulder at 479 ^a nm	81E786
	EtOH/MeOH	LP	435 ^a		Solvent was 4:1 EtOH to MeOH; shoulders at 559 ^a , 487 ^a , and 410 ^a nm	81E786
	Water	LP	606 ^a 568 ^a 530 ^a			81E786
375.	Dichlorobis(1,10-phenanthroline)iridium(III) ion					
	DMF/Water	LP	566 ^a 502 ^a 474 ^a		Solvent was 9:11 dimethylformamide to water; shoulders at 590 ^a , 538 ^a , 460 ^a , 445 ^a and 423 ^a nm	79B098
	DMF/Water	LP	516 ^a 570 ^a 531 ^a 499 ^a		Solvent was 19:1 dimethylformamide to water; shoulders at 473 ^a and 450 ^a nm	79B098
	DMF/Water	LP	613 ^a 477 ^a		Solvent was 19:1 dimethylformamide to water; shoulders at 559 ^a and 507 ^a nm	81E786
	DMF/Water	LP	483 ^a 434 ^a		Solvent was 9:11 dimethylformamide to water; shoulder at 537 ^a nm	81E786
	EtOH/MeOH	LP	451 ^a		Solvent was 4:1 EtOH/MeOH; shoulders at 522 ^a and 482 ^a nm	81E786
	EtOH/MeOH (77 K)	LP	508 ^a 481 ^a 450 ^a		Glass was 4:1 EtOH to MeOH	81E786
	Water	LP	598 ^a 561 ^a 530 ^a 509 ^a 476 ^a 451 ^a 418 ^a			79B098
	Water	LP	532 ^a 440 ^a			81E786
376.	3,5-Dichloro-2,6-diphenyl-1,4-benzoquinone/Triphenylamine					
	Benzene (293 K)	LP	650 550		550 nm band was the more intense; $\tau_T = 0.1 \mu s$	81E715
	Benzene/MeOH (293 K)	LP	650 550		650 nm band was the more intense; solvent was 19:1 benzene to MeOH; $\tau_T = 0.048 \mu s$	81E715
377.	1,4-Dichloronaphthalene					
	Cyclohexane	FP	422.5		$\tau_T = 360 \mu s$	62E009

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
378.	4,4'-Dichlorostilbene Isopentane/3-MP (77 K)	FP/IV	380 ^a 359, 28000		Glass was 6:1 isopentane to 3-MP; the ϵ at 359 was listed as 280000 in the paper, but we assume that this was a misprint; shoulder at 344 ^a nm and some unresolved feature centered at 394 ^a nm; $\tau_T = 4800 \mu s$	687111
379.	4-(4,6-Dichloro-1,3,5-triazin-2-yl)-N,N-diethylaniline EPA (77 K) PMMA (293 K)	PS/KM PS/KM	490, 31000 \pm 5000 490, 31000 \pm 5000		\dagger Phosphorescence decay; $\tau_T = (4.3 \pm 0.4) \times 10^6 \mu s$ $\tau_T = (1.7 \pm 0.1) \times 10^6 \mu s$	83E427 83E427
380.	Dicumarol EPA (77 K)	PS	499.5 465 437 413 ^a 380 ^a		\dagger Phosphorescence decay; most intense peaks were at 499.5, 465, and 437 nm; $\tau_T = (1.3 \pm 0.2) \times 10^6 \mu s$	71B001
381.	9,10-Dicyanoanthracene Acetonitrile Heptane Toluene	LP FP LP/ET	440 426 440, 9000 \pm 2000		\dagger Oxygen quenching; intersystem crossing promoted by dimethyliodobenzene; shoulder at 425 nm \dagger Oxygen quenching and enhanced triplet yield in presence of dibromoethane; $\tau_T = \sim 100 \mu s$ ϵ relative to rubrene in toluene ($\epsilon_{300} = 26000$ L mol ⁻¹ cm ⁻¹)	83E025 78E414 84E393
382.	1,2-Dicyanobenzene EPA (77 K)	FP	510 459		\dagger Phosphorescence decay, molecular orbital calculations; ϵ roughly 200 L mol ⁻¹ cm ⁻¹ (method unspecified); $\tau_T = 2.5 \mu s$	776213
383.	1,3-Dicyanobenzene EPA (77 K) Et ₂ O/Isopentane (77 K)	FP PS	530 440 290 ^a 300		\dagger Phosphorescence decay, molecular orbital calculations; ϵ roughly 200 L mol ⁻¹ cm ⁻¹ (method unspecified); $\tau_T = 5.0 \mu s$ Glass was 1:1 Et ₂ O to isopentane	776213 75B004
384.	1,4-Dicyanobenzene EPA (77 K) Et ₂ O/Isopentane (77 K)	FP PS	500 297		\dagger Phosphorescence decay, molecular orbital calculations; ϵ roughly 200 L mol ⁻¹ cm ⁻¹ (method unspecified); shoulder at 450 nm; $\tau_T = 1.9 \mu s$ Glass was 1:1 Et ₂ O to isopentane	776213 75B004
385.	1,4-Dicyanonaphthalene Acetonitrile Acetonitrile Heptane	LP LP/ET LP	455 275 455, 7000 \pm 1000 275 455 275		\dagger Oxygen quenching (2.1×10^9 L mol ⁻¹ s ⁻¹); ϵ relative to benzophenone in acetonitrile ($\epsilon_{320} = 6500$ L mol ⁻¹ cm ⁻¹); $\tau_T = 40 \mu s$	84B017 84B066 84E236
386.	trans-4,4'-Dicyanostilbene 2-MTHF (133 K) EtOH (135 K) Glycerol triacetate (198 K) PMMA (298 K)	LP LP LP LP	415 410 410 405		Triplet absorption not observable above 134 K; below 99 K lifetime ($1.3 \times 10^4 \mu s$) constant; lifetime measured at 134 K; $\tau_T = 1.0 \mu s$ Triplet absorption not observable above 138 K; below 110 K lifetime ($1.3 \times 10^4 \mu s$) constant; lifetime measured at 138 K; $\tau_T = 2.5 \mu s$ Triplet absorption not observable above 243 K; below 214 K lifetime constant; $\tau_T = 6700 \mu s$	80F299 80F299 80F299 80F299

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
387.	<i>all-trans</i> -3',4'-Didehydro- β,ψ -16'-carotenal Hexane	LP-ET/SD	580, 363000 \pm 54500		{Triplet ET from biphenyl; ϵ assumes triplet does not absorb where singlet depletion is followed; "torularhodinaldehyde"; $\tau_T = 2.5 \mu\text{s}$; $E_T = \sim 94 \text{ kJ mol}^{-1}$	78E721
388.	6,6'-Diethoxythioindigo Benzene	LP	560 ^a 420 ^a		{Oxygen quenching ($2.9 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); spectrum independent of amount of <i>cis</i> or <i>trans</i> isomers used; $\tau_T = 0.143 \mu\text{s}$	78F030
	Dichloromethane	LP	560 ^a 420 ^a		{Oxygen quenching; spectrum independent of amount of <i>cis</i> or <i>trans</i> isomers used; $\tau_T = 0.065 \mu\text{s}$	78F030
389.	<i>N</i> -[2-(Diethylamino)ethyl]phthalimide EtOH	FP	590 350		{Oxygen quenching, diene quenching; maxima assumed from text; $\tau_T = 42 \mu\text{s}$	79A147
390.	7-Diethylamino-4-methylcoumarin EPA EtOH	CWL FP/ET	596 ^a 620, 21600 \pm 2200		Glass, temperature unspecified ϵ relative to anthracene in EtOH ($\epsilon_{420} = 75000 \text{ L mol}^{-1} \text{ cm}^{-1}$); there was another maximum at wavelength shorter than 400 nm; $\tau_T = 3000 \pm 1000 \mu\text{s}$; $k_{et} = (6.9 \pm 1.0) \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	73E344 747049
391.	1,1'-Diethyl-6-bromo-2,2'-cyanine iodide PVA	FP/SD	651 ^a , 33800 ^a			67F505
392.	1,1'-Diethyl-2,2'-carbocyanine chloride Ethylene glycol Glycerol MeOH PVA	FP FP FP-ET/SD FP	635 635 635, 58000 \pm 3000 625 ^b , 52000 741 ^a 673 ^a		"Pinacyanol chloride"; $\tau_T = 256 \pm 20 \mu\text{s}$ $\tau_T = 417 \pm 50 \mu\text{s}$ {Triplet ET from naphthalene, oxygen quenching; $\tau_T = 190 \pm 18 \mu\text{s}$; $k_{et} = (1.0 \pm 0.1) \times 10^{10} \text{ L mol}^{-1} \text{ s}^{-1}$ $\tau_T = 2.0 \times 10^4 \mu\text{s}$	79E243 79E243 79E243 69B005
393.	1,1'-Diethyl-4,4'-carbocyanine iodide EtOH	FP-ET/ ET&SD	778 ^a , 59000 ^a 728 ^b , 32000 593 ^b , 6000 504 ^a , 25100 ^a 425 ^a , 36400 ^a		ϵ relative to naphthalene in EtOH ($\epsilon_{415} = 40000 \text{ L mol}^{-1} \text{ cm}^{-1}$); in ET measurement, compensation was made for the simultaneous decay and buildup of triplet; shoulder at 580 ^a nm; $\tau_T = 1100 \mu\text{s}$; $k_{et} = 1.0 \times 10^{10} \text{ L mol}^{-1} \text{ s}^{-1}$	736051
394.	1,1'-Diethyl-2,2'-cyanine iodide MeOH PVA PVA	LP FP/SD FP	600 635 ^a , 31000 ^a 642 ^a		{Triplet ET from naphthalene; "pseudoisocyanine"; $\tau_T = 49 \pm 1 \mu\text{s}$; $k_{et} = (1.2 \pm 0.1) \times 10^{10} \text{ L mol}^{-1} \text{ s}^{-1}$ $\tau_T = 1.2 \times 10^4 \mu\text{s}$	79E828 67F505 69B005
395.	1,1'-Diethyl-2,2'-dicarbocyanine iodide EtOH	FP-ET/ ET&SD	791 ^a , 111000 ^a 780 ^b , 115000 \pm 11000 760 ^b , 96800 \pm 9700 740 ^b , 40000 627.5 ^b , 40000 611 ^a , 48000 ^a		ϵ relative to naphthalene in EtOH ($\epsilon_{415} = 40000 \text{ L mol}^{-1} \text{ cm}^{-1}$); in ET measurement, compensation was made for the simultaneous decay and buildup of triplet; shoulder at 504 ^a nm; $\tau_T = 480 \mu\text{s}$; $k_{et} = 8.9 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	736051
396.	1,1'-Diethyl-6-iodo-2,2'-cyanine iodide PVA	FP/SD	660 ^a , 28500 ^a		Shoulder at 501 ^a nm	67F505
397.	3,3'-Diethyl-9-methoxy-2,2'-thiacarbocyanine iodide 2-PrOH BuOH PrOH	FP FP FP	640 ^a 650 648 ^a		Isomer of PrOH unspecified	777036 777036 757441

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
398.	3,3'-Diethyl-2,2'-oxacarbocyanine iodide BuOH	FP-ET	605		†Triplet ET from naphthalene	777036
399.	3,3'-Diethyl-2,2'-oxadibocarbocyanine iodide EtOH	FP-ET/ET	670 ^b , 95500 ± 9500 660 ^b , 110000 ± 10000 650, 135000 ^a		†Triplet ET from naphthalene; ϵ relative to naphthalene in EtOH (ϵ_{415} = 40000 L mol ⁻¹ cm ⁻¹); τ_T = ~5000 μ s; k_{et} = 6.7×10^9 L mol ⁻¹ s ⁻¹	726156
400.	N,N'-Diethylrhodamine 1-ProH	FP/SD	920 615, 12000 410		†Triplet ET to anthracene sulfonic acid and from naphthalene, oxygen quenching; 250 μ s delay	78A304
	Water	FP/SD	920 625 590 ^b , 8000 410		250 μ s delay; blue band decays with same lifetime as others	777041
	Water	FP	920 615 410		†Triplet ET to anthracene sulfonic acid and from naphthalene, oxygen quenching; 250 μ s delay; pH 7.0	78A304
401.	5,5'-Diethylselenoindigo EPA (77 K)	FP	630		τ_T = 303 μ s	767582
402.	9,9-Diethyl-9H-9-stannafuorene MCH/Isopentane (77 K)	FP	382 ^a		†Phosphorescence lifetime; glass was 3:1 MCH to isopentane; shoulder at 366 ^a nm; τ_T = 4.3×10^4 μ s	81E648
403.	3,3'-Diethyl-2,2'-thiacarbocyanine iodide BuOH PVA	FP-ET FP	605 ^a 930 ^a 911 ^a		†Triplet ET from naphthalene Shoulder at 869 ^a nm; τ_T = 6500 μ s	777036 69B005
404.	3,3'-Diethyl-2,2'-thiacyanine iodide 2-ProH PVA	FP-ET FP	660 711 ^a 629 ^a		†Triplet ET from naphthalene Shoulder at 566 ^a nm; τ_T = 1.6×10^4 μ s	777036 69B005
405.	3,3'-Diethyl-2,2'-thiadibocarbocyanine iodide 2-ProH EtOH	FP-ET FP-ET/ET	705 770 ^b , 130000 ± 9000 760 ^b , 156000 ± 10000 ~690 ^a , 267000 ^a		†Triplet ET from naphthalene †Triplet ET from naphthalene; ϵ relative to naphthalene in EtOH (ϵ_{415} = 40000 L mol ⁻¹ cm ⁻¹); shoulder at ~780 ^a nm; τ_T = 1100 μ s; k_{et} = $(5.6 \pm 0.4) \times 10^9$ L mol ⁻¹ s ⁻¹	777036 726156
	PVA	FP	1020 ^a		Shoulder at 982 ^a nm; τ_T = 4000 μ s	69B005
406.	3,3'-Diethyl-2,2'-thiatricarbocyanine iodide Propane	FP/ET	900, 145000		†Triplet ET from anthracene, oxygen quenching; ϵ relative to anthracene in propane ($\epsilon_{422.5}$ = 52000 L mol ⁻¹ cm ⁻¹) assuming unit transfer probability; radical cations and anions also observed (λ_{\max} = 520 nm); τ_T = 67 μ s	79A346
407.	6,6'-Dihexyloxythioindigo EPA (77 K)	FP	560		τ_T = 323 μ s	767582
408.	12a,12b-Dihydrobenzo[ghi]perylene MCH/Isohexane	LP	428 360		†First order production from well characterized triplet state, oxygen quenching (1.1×10^9 L mol ⁻¹ s ⁻¹); relative intensities (1:1); solvent was 2:1 MCH to isohexane; τ_T = 0.81 ± 0.03 μ s	79A237
409.	3,4-Dihydro-1,1'-binaphthyl Benzene	LP-ET	567 ^a 401 ^a		†Triplet ET from xanthone and oxygen quenching; τ_T = 1.8 μ s	84B007

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
410.	4a,5a-Dihydrocarbazole					
	MCH	LP-ET	430		†Triplet ET from triphenylamine and oxygen quenching; $\tau_T = 0.5 \mu\text{s}$	72B002
	MCH	LP	435		†Kinetic precursor of ground state ($\lambda_{\max} = 610 \text{ nm}$); $\tau_T = 0.67 \mu\text{s}$	78B072
411.	4',5'-Dihydro-3-carbethoxypsoralen					
	EtOH	LP	535 ^a		†Oxygen quenching ($1.8 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); $\tau_T = 26 \mu\text{s}$	82E133
	Water	LP/ELT	550 ^a , 10000 \pm 1500		†Oxygen quenching ($3 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); ϵ relative to radical anion ($\epsilon_{600} = 1500 \text{ L mol}^{-1} \text{ cm}^{-1}$) assuming electron transfer from tyrosine 100% efficient; $\tau_T = 25 \mu\text{s}$	82E133
412.	7,7'-Dihydro- β -carotene					
	?	FP-ET	478 ^a 445 ^a		†Triplet ET from anthracene; it was not reported whether the solvent was hexane or benzene; lifetime was measured in hexane; 478 nm peak was the more intense; $\tau_T = 14 \mu\text{s}$	733001
	Hexane	PR/ET	447.5, 420000 \pm 63000		†Triplet ET from biphenyl; ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800 \text{ L mol}^{-1} \text{ cm}^{-1}$), assuming authors used standard for cyclohexane; $\tau_T = 14 \pm 1.4 \mu\text{s}$; $k_{et} = (2.1 \pm 0.21) \times 10^{10} \text{ L mol}^{-1} \text{ s}^{-1}$	776412
413.	8b,9a-Dihydro-9H-cyclopropa[e]pyrene					
	Hexane	LP	521 ^a 485 ^a		$\tau_T = 4 \pm 1 \mu\text{s}$	84F068
414.	10,11-Dihydro-5,7:14,16-dietheno-8H,13H-diindeno[2,1-h:1',2'-i][1,4]dioxacyclotridecin					
	2-MTHF (77 K)	PS/KM	385 ^a , 21200 ^a		Compound "XI" in paper	83E383
415.	12a,12b-Dihydro-5,8-dimethyl(indolino[2,3-c]carbazole)					
	MCH (278 K)	LP	500		Delay 0.5 μs	83A225
416.	5,10-Dihydro-5,10-dimethylphenazine					
	3-MP	LP/RA	450, 62000 \pm 20000 337 ^a , 16000		ϵ relative to anthracene in cyclohexane ($\epsilon_{422} = 61000 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 8 \mu\text{s}$	777088
417.	(R)-4,5-Dihydrodinaphtho[2,1-e:1',2'-g][1,4]dioxocin					
	2-MTHF (96 K)	PS	417 ^a		Compound "VI" in paper; shoulder at 407 ^a nm	83E383
418.	(R)-5,6-Dihydro-4H-dinaphtho[2,1-f:1',2'-h][1,5]dioxonin					
	2-MTHF (96 K)	PS	427 ^a 408 ^a		Compound "VII" in paper	83E383
419.	5,10-Dihydro-5,10-diphenylphenazine					
	3-MP	LP/RA	444, 98000 \pm 30000 400 ^a , 88000 ^a 337 ^a , 10000 280 ^a , 91000 ^a		ϵ relative to anthracene in cyclohexane ($\epsilon_{422} = 61000 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 1 \mu\text{s}$	777088
420.	5,10-Dihydroindeno[2,1-a]indene					
	Benzene (298 K)	LP-ET	392 ~375		†Triplet ET from benzophenone, oxygen quenching ($3.4 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); $\tau_T = > 3 \mu\text{s}$; $k_{et} = 5 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	81E214
	Benzene (298 K)	LP-ET	392 ~375		†Triplet ET from xanthone, oxygen quenching; $\tau_T = > 5 \mu\text{s}$	81E214
	EPA (77 K)	FP	395 375 355		Most intense peak at 395 nm; $\tau_T = 88 \mu\text{s}$	737069
	EPA (298 K)	LP	384 365			81E214

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
421.	3,4-Dihydro-2'-methyl-1,1'-binaphthyl Benzene	LP-ET	481 ^a		†Triplet ET from xanthone	84B007
422.	5,10-Dihydro-5-methyl-10-phenylphenazine 3-MP	LP/RA	444, 94000 ± 30000 337 ^b , 12000		ϵ relative to anthracene in cyclohexane ($\epsilon_{422} = 61000$ L mol ⁻¹ cm ⁻¹); $\tau_T = 6$ μ s	777088
423.	5,12-Dihydro-5-methylquino[2,3- <i>b</i>]acridine-7,14-dione EtOH (300 K)	FP	710 350		†Triplet ET to anthracene and oxygen quenching; $\tau_T = 70$ μ s	79E965
	EtOH (300 K)	FP/SD	710, 15300 ± 1500		"N-Methylquinacridinone"	81E649
424.	9,10-Dihydro-9-oxo-2-acridinesulfonate ion EtOH	FP	580		†Oxygen quenching; $\tau_T = 140$ μ s	81A390
425.	9,10-Dihydro-9-oxo-2-acridinesulfonic acid PVA	FP	585		Lifetime quoted from [79E964]; $\tau_T = 2 \times 10^5$ μ s	81A390
426.	2,3-Dihydrophenalene EPA	FP	430 400		†Oxygen quenching (2×10^9 L mol ⁻¹ s ⁻¹) at 260 K; $\tau_T = 2$ μ s	81F390
427.	3,4-Dihydrophenanthrene Benzene	LP-ET	580 430		†Oxygen and azulene quenching and triplet ET from benzophenone; $\tau_T = 1.2$ μ s	83E278
428.	9,10-Dihydrophenanthrene Hexane (300 K)	MOD/SD	480, 7800 365, 26000		$\tau_T = 120 \pm 9$ μ s	69E208
429.	1,2-Dihydro-3-phenylnaphthalene Isopentane/3-MP (77 K)	FP/IV	384, 40000 364 ^a		Glass was 6:1 isopentane : 3-MP; $\tau_T = 3400$ μ s	687111
430.	4',5'-Dihydropsoralen Benzene	PR/ET	500, 15700		†Triplet ET from biphenyl; ϵ relative to biphenyl in benzene ($\epsilon_{367} = 27100$ L mol ⁻¹ cm ⁻¹); 5 μ s delay; $k_{et} = 5.6 \times 10^9$ L mol ⁻¹ s ⁻¹	79E282
	Water	LP	510		4 μ s delay	79E282
431.	5,12-Dihydroquino[2,3- <i>b</i>]acridine-7,14-dione EtOH (300 K)	FP	710 350		"Linear quinacridinone"; [79E965] quotes a $\lambda_{\max} = 690$ nm for the same system	81E649
432.	1,2-Dihydro-2,2,4,6-tetramethylquinoline Benzene	FP	647 ^a			84F248
433.	1,4-Dihydro-5,6,11,12-tetraphenyl-1,4-epidioxytetracene Cyclohexane	LP	432 ^a		A rubreneperoxide, called "II" in paper	84E056
434.	5,12-Dihydro-5,6,11,12-tetraphenyl-5,12-epidioxytetracene Cyclohexane	LP	455 ^a		A rubreneperoxide, called "I" in paper	84E056
435.	(<i>E</i>)-5[(3,4-Dihydro[1,4]thiazino[3,4- <i>b</i>]benzothiazol-1-yl)methylene]-3-ethyl-2-thioxo-4-thiazolidinone Toluene	FP-FT	650 ^a		†Triplet ET from fluorenone; "merocyanine dye photoisomer"	767764
436.	10 <i>H</i> ,10' <i>H</i> -10,10'-Dihydroxybianthrylidene 1-PrOH/2-PrOH (83 K)	FP	465 ^a 405 ^a		Solvent was 2:3 1-PrOH to 2-PrOH	777238
437.	4,4'-Dihydroxybiphenyl PVA (77 K)	FP	550 400 390		Stretched polymer film; most intense peak at 400 nm polarization also measured	78B129

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Toluene/EtOH (77 K)	MOD	411 ^a		Glass was 19:1 toluene to EtOH	719059
438.	Dihydroxylicopene Benzene	PR-ET	540		†Triplet ET from naphthalene; $\tau_T = 7.9 \mu\text{s}$	80A143
439.	1,5-Dihydroxynaphthalene 3-MP/Et ₂ O (77 K)	PS	447 420 394.5		†Phosphorescence decay; glass 19:1 3-MP to Et ₂ O; 447 nm peak was the most intense; $\tau_T = 1.03 \times 10^6 \mu\text{s}$	74B004
440.	1,8-Dihydroxynaphthalene 3-MP/Et ₂ O (77 K)	PS	448 423		†Phosphorescence decay; glass 19:1 3-MP to Et ₂ O; 448 nm peak was the more intense; $\tau_T = 1.11 \times 10^6 \mu\text{s}$	74B004
441.	2,3-Dihydroxynaphthalene 3-MP/Et ₂ O (77 K)	PS	443 418 394		†Phosphorescence decay; glass 19:1 3-MP to Et ₂ O; 443 nm peak was the most intense; $\tau_T = 1.95 \times 10^6 \mu\text{s}$	74B004
	Toluene/EtOH (77 K)	MOD	445 ^a		Glass was 19:1 toluene to EtOH	719059
	Water	FP	430		†Oxygen quenching; $pK_a = 10.0 \pm 0.2$; 5 μs delay; pH 1.8	77A196
442.	2,7-Dihydroxynaphthalene 3-MP/Et ₂ O (77 K)	PS	444 419		†Phosphorescence decay; glass 19:1 3-MP to Et ₂ O; 444 nm peak was the more intense; $\tau_T = 1.01 \times 10^6 \mu\text{s}$	74B004
443.	2,3-Dihydroxynaphthalene, conjugate base Water	FP	545 ^a 500 ^a 480 ^a 455 ^a 440 ^a		†Oxygen quenching; 5 μs delay; pH 13.5	77A196
444.	3,6-Dihydroxyphthalimide Dioxane	FP/TD	440, 2900		Lifetime limited by aeration; $\tau_T = 0.250 \mu\text{s}$	7575
445.	4,4'-Dimethoxybenzophenone Acetonitrile/Water	LP/ET	675, 3600 545, 5200 440, 7100 350, 7800		Solvent was 9:1 acetonitrile to water; ϵ relative to 1-methylnaphthalene in acetonitrile/water ($\epsilon_{415} = 11200 \text{ L mol}^{-1} \text{ cm}^{-1}$)	84B000
446.	2,3-Dimethoxy-1,4-benzoquinone Benzene	PR	434 ^a		$\tau_T = 0.37 \pm 0.05 \mu\text{s}$	743062
	Cyclohexane	PR/ET	430 \pm 5, 5300 \pm 1200		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800 \text{ L mol}^{-1} \text{ cm}^{-1}$)	743062
	Cyclohexane	LP/ET	430 \pm 5, 5700 \pm 1200		†Triplet ET to β -carotene; ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 0.59 \pm 0.10 \mu\text{s}$	743062
447.	2,5-Dimethoxy-1,4-benzoquinone Benzene	PR	452 ^a		Shoulder at 481 ^a nm; $\tau_T = 5.0 \pm 0.7 \mu\text{s}$	743062
	Cyclohexane	PR/ET	489 ^a 440 \pm 5, 5400 \pm 1100		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800 \text{ L mol}^{-1} \text{ cm}^{-1}$)	743062
	Cyclohexane	LP/ET	489 ^a 440 \pm 5, 5300 \pm 1100		†Triplet ET to β -carotene; ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 5.9 \pm 1.0 \mu\text{s}$	743062
448.	4,4'-Dimethoxybiphenyl Boric acid	PS	415		Glass	777388
449.	5,7-Dimethoxycoumarin Benzene	PR/ET	450, 11400		†Triplet ET from biphenyl; ϵ relative to biphenyl in benzene ($\epsilon_{367} = 27100 \text{ L mol}^{-1} \text{ cm}^{-1}$); 20 ns delay; $\tau_T = 10 \mu\text{s}$; $k_{\text{et}} = 5.6 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	79E282

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
450.	2,3-Dimethoxy-5-methyl-1,4-benzoquinone					
	Benzene	PR	423 ^a		$\tau_T = 0.59 \pm 0.10 \mu\text{s}$	743062
	Cyclohexane	PR/ET	430 \pm 5, 6300 \pm 1400		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800 \text{ L mol}^{-1} \text{ cm}^{-1}$)	743062
	Cyclohexane	LP/ET	430 \pm 5, 6800 \pm 1400		\uparrow Triplet ET to β -carotene; ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 0.55 \pm 0.09 \mu\text{s}$	743062
	Trichlorotrifluoroethane	LP	413 ^a		$\tau_T = 0.43 \pm 0.08 \mu\text{s}$	743062
451.	1,4-Dimethoxynaphthalene					
	3-MP (77 K)	PS	469 445		\uparrow Phosphorescence decay; 469 nm peak was the more intense; $\tau_T = 1.08 \times 10^6 \mu\text{s}$	74B004
452.	1,2-Dimethoxy-4-nitrobenzene					
	Acetonitrile	LP	490 ^a			82B014
453.	trans-2,5-Dimethoxy-4'-nitrostilbene					
	Glycerol triacetate (198 K)	LP	\sim 780 540 467		Most intense peak at 467 nm	78B088
	Glycerol triacetate	LP	\sim 830 554 465		Most intense peak at 465 nm	78B088
454.	4,4'-Dimethoxythiobenzophenone					
	Benzene	LP/RA	520 \pm 5, 3500 \pm 700 400 \pm 5, 5500 \pm 1100 295 \pm 5, 19900 \pm 4000		ϵ relative to benzophenone in benzene ($\epsilon_{322.5} = 7630 \text{ L mol}^{-1} \text{ cm}^{-1}$), taking $\Phi_T = 0.5$ at 337 nm excitation and taking $\Phi_T = 1$ for benzophenone; SD method gave similar ϵ ; $\tau_T = 1.4 \pm 0.1 \mu\text{s}$	84A221
455.	4-(Dimethylamino)chalcone					
	Heptane	LP	550 \pm 15		"3-[4-(Dimethylamino)phenyl]-1-phenyl-2-propen-1-one"; $\tau_T = 0.5 \mu\text{s}$	83E347
456.	10-(Dimethylamino)-3,5,7,9-decatetraen-2-one					
	Toluene	FP-ET	520		\uparrow Triplet ET from anthracene	83E737
457.	2-(Dimethylamino)ethyl benzoate					
	Cyclohexane (290 K)	FP	352 ^a 321 ^a		10 μs delay	767556
458.	2-[7-(Dimethylamino)-2,4,6-heptatrienylidene]-6-[5-(dimethylamino)-2,4-pentadienylidene]cyclohexanone					
	Toluene	FP	680		\uparrow Oxygen quenching and triplet ET from anthracene and benzo[<i>rst</i>]pentaphene and to azulene, perylene, and tetracene	83E737
459.	N-[(Dimethylamino)methyl]phthalimide					
	EtOH	FP	590 350		\uparrow Oxygen quenching, diene quenching; maxima assumed from text; $\tau_T = 10 \mu\text{s}$	79A147
460.	1-Dimethylamino-4-nitronaphthalene					
	2-PrOH	LP	505		Lifetime extrapolated from low temperature; $\tau_T = 0.002 \mu\text{s}$	77A216
	Acetone	LP	505		\uparrow Oxygen quenching ($0.7 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); $\tau_T = 1.0 \mu\text{s}$	77A216
	Acetonitrile	LP	510		$\tau_T = 0.2 \mu\text{s}$	77A216
	Benzene	LP	510		\uparrow Oxygen quenching ($1.6 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); $\tau_T = 4.5 \mu\text{s}$	77A216
	Cyclohexane	LP	500		\uparrow Phosphorescence decay in PMMA at 77 K; $\tau_T = 9.1 \mu\text{s}$	77A216
	Hexane	LP	500		$\tau_T = 8.4 \mu\text{s}$	77A216

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
461.	<i>trans</i> -4-Dimethylamino-4'-nitrostilbene					
	2,2-Dimethylbutane- /Pentane (77 K)	FP	845 ^a		Glass 3:8 pentane to 2,2-dimethylbutane; shoulder at 750 ^a nm	747022
	2-PrOH	LP	680			78B088
	Benzene	LP	800		†Oxygen quenching and triplet ET from triphenylene; shoulder at 745 nm; $\tau_T = 2.1 \mu s$	747022
	Benzene	LP	790			78B088
	Cyclohexane	LP	735		Shoulder at 650 nm; $\tau_T = 0.43 \mu s$	747022
	DMF	LP-ET	910		†Triplet ET from triphenylene; $\tau_T = 21 \mu s$; $k_{et} = 8 \times 10^9$ L mol ⁻¹ s ⁻¹	747022
	DMF	LP	820			
	DMF	LP	805			78B088
	EtOH	LP	630			78B088
	Glycerol triacetate (198 K)	LP	830			78D088
	Glycerol triacetate	LP	835			78B088
	MeOH	LP	600			78B088
	Toluene	LP	800		$\tau_T = 1.8 \mu s$	82F498
	<i>tert</i> -BuOH	LP	760			78B088
462.	8-(Dimethylamino)-3,5,7-octatrien-2-one					
	Toluene	FP-ET	480		†Triplet ET from anthracene	83E737
463.	9-[4-(Dimethylamino)phenyl]anthracene					
	Acetonitrile	LP	442 ^a		†Oxygen quenching	776464
	Acetone	LP	440 ^a		†Oxygen quenching	776464
	Et ₂ O	LP	444 ^a		†Oxygen quenching	776464
464.	4-[2-[4-(Dimethylamino)phenyl]ethyl]benzophenone					
	Acetonitrile	LP	490 ^a		Possible triplet exciplex, structure unspecified; "benzophenone (CH ₂) ₂ (N,N'-dimethylaniline)"	78B112
465.	<i>trans</i> -1-(4-Dimethylaminophenyl)-2-nitroethylene					
	Benzene/Acetonitrile	LP	660 ^a		Solvent was 8:2 benzene to acetonitrile by volume	78E057
	Benzene	LP	650 ^a		$\tau_T = 0.56 \mu s$	78E057
	Cyclohexane	LP	570 ^a		$\tau_T = 0.379 \mu s$	78E057
	EPA (77 K)	PS/SD	600, 19100 ± 3820 440, 35000 ± 7000		†Phosphorescence decay; $\tau_T = (7.9 \pm 1.5) \times 10^4 \mu s$	78E057
466.	2-(Dimethylamino)purine					
	Acetonitrile (300 K)	FP	485		†Oxygen quenching and quenching by 2,4-hexadienol; $\tau_T = 59 \pm 4 \mu s$	756270
467.	2-(Dimethylamino)pyridine					
	2-PrOH (179 K)	FP	~470		†Phosphorescence decay in EPA; $\tau_T = (1.4 \pm 0.3) \times 10^3 \mu s$	767526
	3-MP (179 K)	FP	~465 ^a		†Phosphorescence decay in EPA	767526
	EPA (100 K)	FP	~470		†Phosphorescence decay in EPA; $\tau_T = (4.73 \pm 0.46) \times 10^3 \mu s$	767526
	EPA (103 K)	FP	470 ^a		Lifetime measured at 93 K; $\tau_T = 0.4 \times 10^6 \mu s$	80E318
468.	4-(Dimethylamino)pyridine					
	2-PrOH (188 K)	FP	500		$\tau_T = 200 \pm 12 \mu s$	737127
	EPA (93 K)	FP	500		†Phosphorescence decay	72B004
	EPA (77 K)	FP	500		†Phosphorescence decay	737127
	Et ₂ O (183 K)	FP	500			737127
469.	2-(Dimethylamino)pyridine/4-Nitroaniline					
	EPA (100 K)	FP	590 460		At 93 K lifetimes are 4.5 ms (590 nm) and 0.4 s (460 nm); relative intensities (2:1)	80E318
470.	N,N-Dimethylaniline					
	3-MP (77 K)	FP	465 ^a 456 ^a 440 ^a 341 ^a		†Phosphorescence decay; another maximum was estimated to be at ~392 nm which was in the SD region of 390-430 nm; most intense peak at 341 nm	69E215

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Cyclohexane	PR	460 ± 10		†Oxygen quenching (10 ¹⁰ L mol ⁻¹ s ⁻¹); further peak at 370 nm attributed to the cyclohexadienyl radical; half-life = 0.95 µs	761069
	Liquid paraffin	FP	480		†Phosphorescence decay in low temperature glass; delay 5 µs; half-life = 100 µs	68B002
	<i>N,N</i> -Dimethylaniline	LP/ET	465, 5000		†Triplet ET to anthracene and to naphthalene; ϵ relative to naphthalene in dimethylaniline ($\epsilon_{427.5} = 14400$ L mol ⁻¹ cm ⁻¹); half-life = 0.25 µs; $k_{et} = 4.8 \times 10^9$ L mol ⁻¹ s ⁻¹	720440
471.	<i>N,N'</i> -Dimethyl-3-anilinocarbazole MCH (278 K)	LP	610		†Oxygen quenching and rise time same as fluorescence decay; delay 30 ns; rise time of 13 ns	83A225
472.	1,3-Dimethylantracene EtOH (93 K)	PS	442 417			66B001
473.	9,10-Dimethylantracene Benzene	PR/ET	435, 47000		ϵ relative to benzophenone in benzene ($\epsilon_{532.5} = 10300$ L mol ⁻¹ cm ⁻¹)	690087
	Benzene	PR/ET	435, 42200		ϵ relative to naphthalene in benzene ($\epsilon_{\max} = 17500$ L mol ⁻¹ cm ⁻¹)	690087
	Cyclohexane	PR	427.5			690087
	Dioxane	PR	430			690087
474.	2,5-Dimethyl-1,4-benzoquinone SDS	LP	470		Aqueous micelles; $\tau_T = 0.035$ µs	83N133
	Water	LP	480 440		$\tau_T = 0.83$ µs	80B112
475.	2,6-Dimethyl-1,4-benzoquinone SDS	LP	455		Aqueous micelles; $\tau_T = 0.020$ µs	83N133
	Water	LP	445		$\tau_T = 0.060$ µs	83N133
476.	3,3'-Dimethylbiphenyl 3-MH (77 K)	PR	390		1 µs delay; radical ions also contribute to observed peak at 390 nm; $\tau_T = 3 \times 10^6$ µs	771059
477.	(<i>all-E</i>)-3,8-Dimethyl-1,10-bis(2,6,6-trimethyl-1-cyclohexen-1-yl)-1,3,5,7,9-decapentaene Benzene (296 K)	LP-ET	457		Triplet ET from chlorophyll <i>a</i> ; oxygen quenching (2.8×10^9 L mol ⁻¹ s ⁻¹); $\tau_T = 20$ µs; $k_{et} = 0.9 \times 10^9$ L mol ⁻¹ s ⁻¹	73E347
478.	1-(3,3-Dimethyl-1-buten-2-yl)naphthalene Benzene	LP-ET	426 ^a		†Triplet ET from xanthone and oxygen quenching; $\tau_T = 0.44$ µs	84B007
479.	<i>N,N'</i> -Dimethyl-5,11-dihydroindolo[3,2- <i>b</i>]carbazole SDS	LP	658 ^a 395 ^a		Aqueous micelle	79N005
480.	<i>cis</i> -2,3-Dimethyl-2,3-di-(2-naphthyl)oxirane Benzene	LP/ET	425, 10000 ± 2000		†Oxygen and azulene quenching; ϵ relative to benzophenone in benzene ($\epsilon_{533.5} = 7630$ L mol ⁻¹ cm ⁻¹); $\tau_T = 1.2 \pm 0.2$ µs	84A344
481.	<i>trans</i> -2,3-Dimethyl-2,3-di-(2-naphthyl)oxirane Benzene	LP/ET	425, 12000 ± 2400		†Oxygen and azulene quenching; ϵ relative to benzophenone in benzene ($\epsilon_{533.5} = 7630$ L mol ⁻¹ cm ⁻¹); $\tau_T = 1.4 \pm 0.2$ µs	84A344

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
482.	2,4-Dimethyl-1,5-diphenyl-1,5-pentanedione ?	LP	455 ^a		Solvent benzene (?); conformation such that carbonyl groups are in the plane formed by the three central carbons of the pentane chain; more intense maximum below 350 nm; $\tau_T = 0.060 \mu\text{s}$	81A404
483.	<i>N,N'</i> -Dimethyl- <i>N,N'</i> -diphenyl-1,4-phenylenediamine MCH (278 K)	LP	635		\dagger Oxygen quenching and risetime same as fluorescence decay; delay 50 ns; $\tau_T = 3 \mu\text{s}$; rise time of 5 ns	83A225
484.	3,9-Dimethyl- <i>trans</i> -fluorenone 3-MP (77 K)	MOD/ SD&KM	481, 93500 \pm 4300 457, 55600 \pm 3500			757141
485.	2,5-Dimethyl-2,4-hexadiene Benzene	PR	310		$\tau_T = 0.08 \mu\text{s}$	82B057
486.	1,3-Dimethylindazole EtOH (178 K)	MOD/SD	420 ^a , 8100 ^a		Shoulders at 439 ^a and 286 ^a nm; another maximum < 250 nm; $\tau_T = (1.5 \pm 0.2) \times 10^4 \mu\text{s}$	716244
	EtOH (103 K)	MOD/SD	420 ^a , 8600 ^a 287 ^a , 4700 ^a		Shoulder at 448 ^a nm; another maximum < 250 nm; $\tau_T = (3.6 \pm 0.4) \times 10^5 \mu\text{s}$	716244
487.	<i>N,N'</i> -Dimethylindigo EPA (98 K)	LP	740 470		No triplet spectrum observed at RT; data corrected from paper (H. Goerner, private communication, 1985)	79E543
488.	1,4-Dimethyl-7-isopropylazulene Benzene	LP-ET	380-420		\dagger Triplet ET from benzophenone, oxygen quenching; $\tau_T = 3 \mu\text{s}$; $k_{et} = 7 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	81F275
489.	1,2-Dimethylnaphthalene Cyclohexane (77 K)	PS	420 400			67B007
490.	2,3-Dimethylnaphthalene Cyclohexane (77 K)	PS	426.5 420.5 404 398 392.5			67B007
	EtOH (77 K)	MOD	422		Halfwidth 540 cm ⁻¹	777538
	Toluene (77 K)	MOI	433		Halfwidth 920 cm ⁻¹	777538
491.	2,7-Dimethylnaphthalene EtOH (77 K)	MOD	419		Halfwidth 460 cm ⁻¹	777538
	Toluene (77 K)	MOD	431		Halfwidth 880 cm ⁻¹	777538
492.	<i>N,N</i> -Dimethyl-2-naphthylamine Cyclohexane	FP	476 ^a		Delay 30 μs	686045
493.	<i>N,N</i> -Dimethyl-2-naphthylamine, conjugate acid Water	FP	409 388		$\text{p}K_a = 2.7$; pH Acidic	61E008
494.	<i>cis</i> -3,3-Dimethyl-1-(2-naphthyl)-1-butene Benzene	LP-ET	430 400		\dagger Oxygen and azulene quenching and triplet ET from benzophenone and xanthone; $\tau_T = 0.14 \mu\text{s}$	83E210
495.	<i>trans</i> -3,3-Dimethyl-1-(2-naphthyl)-1-butene Benzene	LP-ET	430 400		\dagger Oxygen and azulene quenching and triplet ET from benzophenone and xanthone; $\tau_T = 0.13 \mu\text{s}$	83E278

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
496.	2,2-Dimethyl-1-(1-naphthyl)-1-phenylethylene Benzene	LP-ET	410 ^a		†Triplet ET from xanthone and oxygen quenching; τ_T = 0.10 μ s	84B007
497.	<i>N,N</i> -Dimethyl-4-nitroaniline EPA (77 K)	PS/KM	900, 3370 \pm 170 760, 2780 \pm 140 550, 2700 \pm 135 390, 20600 \pm 1030		†Phosphorescence decay; τ_T = (4.37 \pm 0.10) \times 10 ⁵ μ s	78E057
	EPA (77 K)	PS	550 450		†Phosphorescence decay; τ_T = 8.3 \times 10 ⁵ μ s	83E427
498.	<i>N,N</i> -Dimethyl-4-[(4-nitrophenyl)azo]benzenamine 2-MTHF (103 K)	LP	700 390		Triplet not observable above 125 K; below 103 K lifetime constant; τ_T = 500 μ s	80B101
	Glycerol triacetate (203 K)	LP	690		Lifetime measured at 233 K; triplet not observable above 233 K; τ_T = 0.050 μ s	80B101
499.	3,8-Dimethyl-4,7-phenanthroline EtOH/Et ₂ O (77 K)	PS	495 465 435		Solvent was 3:2 EtOH to Et ₂ O; relative intensities (100:60:25); E_T = 264 kJ mol ⁻¹	80B130
500.	<i>N,N</i> -Dimethyl- <i>p</i> -phenylenediamine EtOH (77 K)	MOD/KM	530, 23000 \pm 5700			737055
501.	(<i>E,E,E,E,E,E</i>)-7,11-Dimethyl-7-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2,4,6,8,10,12-tridecahexaenal Cyclohexane	COM	500, 201000		" <i>all-trans</i> -C ₂₄ aldehyde"; ϵ average of that obtained by PR/ET, LP/ET and LP/SD; ϵ relative to C ₁₇ aldehyde in cyclohexane (ϵ_{410} = 63000 L mol ⁻¹ cm ⁻¹) in LP/ET; ϵ relative to biphenyl in cyclohexane ($\epsilon_{361.3}$ = 42800 L mol ⁻¹ cm ⁻¹) in PR/ET; τ_T = 7.1 μ s	79E546
	MeOH	LP/SD	510, 136000		ϵ is upper limit; τ_T = 8.3 μ s	79E546
502.	1,3-Dimethyluracil Acetonitrile	LP/?	380 ^a , 8000 ^a		ϵ method unspecified	81E042
503.	(<i>S</i>)-Dinaphtho[2,1- <i>d</i> :1',2'- <i>f</i>][1,3]dioxepin 2-MTHF (96 K)	PS	420 ^a 400 ^a		Compound "V" in paper	83E383
504.	Di-2-naphthylamine Toluene/EtOH (77 K)	MOD	508 ^a 439 ^a		Glass was 19:1 toluene to EtOH; 508 nm peak was the more intense	719059
505.	1,4-Di-1-naphthylbutane Isooctane	FP	430 ^a 400 ^a 380 ^a 362 ^a		Isomer assumed; additional absorption in the red assigned to intramolecular triplet excimer	78B089
	Isooctane	LP	405 ^a 390 ^a		Shoulder at 370 nm; rise time of \sim 100 ns	81E015
506.	<i>t</i> -3, <i>t</i> -4-Di-(1-naphthyl)- <i>r</i> -1, <i>c</i> -2-dimethoxycarbonylcyclobutane Acetonitrile	LP-ET	430		†Triplet ET from benzophenone, oxygen quenching (1.5 \times 10 ¹⁰ L mol ⁻¹ s ⁻¹); 100 ns delay; τ_T = 0.13 μ s; k_{et} = (7 \pm 1) \times 10 ⁹ L mol ⁻¹ s ⁻¹	80B010
	Acetonitrile	LP-ET	470		†Triplet ET from benzophenone; 400 ns delay; intramolecular triplet exciplex; τ_T = 4.0 μ s; k_{et} = (7 \pm 1) \times 10 ⁹ L mol ⁻¹ s ⁻¹	80B010
507.	1,2-Di-1-naphthylethane Isooctane	FP	430 ^a 396 ^a 355 ^a		Isomer assumed; shoulder \sim 380 nm; additional absorption in the red assigned to intramolecular triplet excimer	78B089

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Isooctane	LP	410 ^a 390 ^a		Shoulder at 370 nm; additional absorption around 470 nm (rise time < 17 ns) ascribed to triplet excimer; rise time of ~100 ns	81E015
508.	Di-1-naphthylmethane					
	Isooctane	FP	430 ^a 396 ^a 350 ^a		Shoulder ~370 nm; additional absorption in the red assigned to intramolecular triplet excimer; $\tau_T = \sim 50 \mu s$	78B089
	Isooctane	LP	410 ^a 390 ^a		Shoulder at 370 nm; rise time of ~100 ns	81E015
509.	2,5-Di(1-naphthyl)-1,3,4-oxadiazole					
	Cyclohexane	FP/TD	602, 7150 ^a 569, 5200 ^a 454, 5500 ^a		Only peaks reported by authors given - extensive vibronic structure observed; oscillator strength = 0.04, 0.04, 0.06	78B081
	Cyclohexane	FP/TD	600 ^a , 7100 ^a 570 ^a , 5200 ^a 460 ^a , 5700 ^a		0-0 bands only reported here though vibrational structure seen; oscillator strength = 0.04, 0.04, 0.06	79E297
510.	cis-2,3-Di-(2-naphthyl)oxirane					
	Benzene	LP/ET	430, 10000 \pm 2000		\uparrow Oxygen and azulene quenching; ϵ relative to benzophenone in benzene ($\epsilon_{533.5} = 7630 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 0.36 \pm 0.05 \mu s$	84A344
511.	trans-2,3-Di-(2-naphthyl)oxirane					
	Benzene	LP/ET	430, >8000		\uparrow Oxygen and azulene quenching; ϵ relative to benzophenone in benzene ($\epsilon_{533.5} = 7630 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 0.16 \pm 0.02 \mu s$	84A344
512.	1,3-Di-1-naphthylpropane					
	Isooctane	FP	430 ^a 400 ^a 375 ^a		Isomer assumed; additional absorption in the red assigned to intramolecular triplet excimer	78B089
	Isooctane	LP	415 ^a 390 ^a		Shoulder at 370 nm; rise time of ~100 ns	81E015
513.	1,3-Di-2-naphthylpropane					
	Methylene chloride	LP	430 ^a		Shoulder at 405 ^a	84P257
514.	5,5'-Dineopentylthioindigo					
	EPA (77 K)	FP	592		$\tau_T = 130 \mu s$	767582
	EPA (93 K)	LP	580 375		$\tau_T = 53 \mu s$	79E543
	EPA (283 K)	LP	590 370		Φ_T strongly temperature dependent in this solvent; $\tau_T = 0.46 \mu s$	79E543
	Glycerol triacetate (203 K)	LP	600 390		$\tau_T = 50 \mu s$	79E543
	Glycerol triacetate	LP	595 385		$\tau_T = 0.40 \mu s$	79E543
515.	3,5-Dinitroanisole					
	Acetonitrile/Water	LP	475		Solvent 1:1 acetonitrile to water; $\tau_T = 0.055 \mu s$	737466
	Water	LP	~435			82F150
516.	3,5-Dinitroanisole/Water					
	Water	LP	435		\uparrow Oxygen quenching and triplet ET to 3,3',4,4'-tetramethyldiazetidine dioxide; transient was assigned to an H-bonding exciplex with water; $\tau_T = 1.4 \mu s$	777345
517.	1,2-Dinitronaphthalene					
	EtOH	LP	550		\uparrow Phosphorescence decay in EPA glass at 77 K, triplet ET to tetracene, oxygen quenching ($3.3 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); $\tau_T = 4.3 \mu s$	776194

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Hexane	LP	490		†Phosphorescence decay in EPA glass at 77 K, triplet ET to tetracene, oxygen quenching (1.9×10^9 L mol ⁻¹ s ⁻¹); $\tau_T = 1.0$ μ s	776194
518.	1,4-Dinitronaphthalene					
	EtOH	LP	550		Lifetime measured in neutral EtOH; spectrum shifts to the red (30 nm) with increasing sulfuric acid concentration ($\{10$ mol L ⁻¹ }); $\tau_T = 4.76$ μ s	767270
	Formamide	LP	570			767270
	Hexane	LP	545		†Phosphorescence decay in EPA at 77K, triplet ET to tetracene; $\tau_T = 4.55$ μ s; $k_{et} = 1.0 \times 10^{10}$ L mol ⁻¹ s ⁻¹	767270
519.	1,8-Dinitronaphthalene					
	EtOH	LP	590		†Phosphorescence decay in EPA glass at 77 K, triplet ET to tetracene, oxygen quenching (2.1×10^9 L mol ⁻¹ s ⁻¹); $\tau_T = 5.3$ μ s	776194
	Hexane	LP	550		†Phosphorescence decay in EPA glass at 77 K, triplet ET to tetracene, oxygen quenching (8.1×10^8 L mol ⁻¹ s ⁻¹); $\tau_T = 2.5$ μ s	776194
520.	4-[(2,4-Dinitrophenyl)azo]-N,N-dimethylbenzenamine					
	2-MTHF (103 K)	LP	700		Triplet not observable above 126 K; below 103 K lifetime constant; $\tau_T = 1300$ μ s	80B101
			420			
	Glycerol triacetate (198 K)	LP	690		Lifetime measured at 233 K; triplet not observable above 233 K; $\tau_T = 0.033$ μ s	80B101
521.	3-[[4-[(2,4-Dinitrophenyl)azo]phenyl](2-hydroxyethyl)amino]-4-propanenitrile					
	2-MTHF (103 K)	LP	700		Triplet not observable above 123 K; below 103 K lifetime constant; $\tau_T = 670$ μ s	80B101
			400			
	Glycerol triacetate (203 K)	LP	680		Lifetime measured at 248 K; triplet not observable above 248 K; $\tau_T = 0.025$ μ s	80B101
522.	trans-2,4-Dinitrostilbene					
	Glycerol triacetate	LP	542		542 nm peak was the more intense	78B088
			428			
	Glycerol triacetate (198 K)	LP	560		560 nm peak was the more intense	78B088
			434			
523.	cis-4,4'-Dinitrostilbene					
	Cyclohexane	LP	500		$\tau_T = 0.083$ μ s	747022
	MeOH	LP	500			747022
524.	trans-4,4'-Dinitrostilbene					
	Cyclohexane	LP	500		†Oxygen quenching in benzene; $\tau_T = 0.080$ μ s	747022
	EPA (77 K)	FP	550 ^a			747022
	EPA	LP	472			78B088
	EPA (88 K)	LP	486			78B088
	Glycerol triacetate (198 K)	LP	~735		496 nm peak was the more intense	78B088
			496			
	Glycerol triacetate	LP	500			78B088
	MeOH	LP	500		†Oxygen quenching in benzene	747022
525.	N,N'-Dipentylpyromellitic diimide					
	Acetonitrile	LP	555		Triplet detected in the presence of N-benzyl-1,4-dihydronicotinamide	83B041
			510			
			420			
526.	1,3-Di(9-phenanthryl)propane					
	Decane (293 K)	LP	481 ^a		Shoulder at 402 ^a nm; The shortest wavelength band and shoulder are mainly due to intramolecular triplet excimers	82E287
			447 ^a			
			427 ^a			
	THF	LP	480 ^a		Delay 1 μ s; shoulder at 423 ^a nm	83E554
			450 ^a			

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
527.	Diphenylacetylene Glycerol	FP	418 380		418 nm was the stronger band; $\tau_T = 267 \mu\text{s}$	747417
528.	Diphenylamine 3-MP (77 K) Cyclohexane	PS PR/ET	545 530, 10400		$\tau_T = (2.0 \times 0.1) \times 10^6 \mu\text{s}$ ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700 \text{ L mol}^{-1} \text{ cm}^{-1}$); reference ϵ obtained by starting from $\epsilon_{\max} = 3220 \text{ L mol}^{-1} \text{ cm}^{-1}$ for this ketyl radical in water and assuming the f of the ketyl radical is independent of solvent; final ϵ obtained from a simultaneous least squares fit to data from several compounds	67E106 71E360
	EPA (~90 K)	PS	558 ^a			42B002
	EPA (77 K)	PS	549		\dagger Phosphorescence decay; $\tau_T = (1.9 \pm 0.2) \times 10^6 \mu\text{s}$	67E106
	EtOH (77 K)	MOD/KM	550, 31000 \pm 7700			737055
	EtOH/Et ₂ O (77 K)	MOD/KM	550, 28600 \pm 11000		Glass was 2:1 EtOH to Et ₂ O; temperature was not explicitly stated, but 77 K was inferred from the context	719059
	Hexane	LP	520			84E043
	Toluene/EtOH (77 K)	MOD	555 ^a		Glass was 19:1 toluene to EtOH	719059
529.	9,10-Diphenylanthracene Benzene	PR/ET	445, 17700		ϵ relative to naphthalene in benzene ($\epsilon_{\max} = 17500 \text{ L mol}^{-1} \text{ cm}^{-1}$)	690087
	Benzene	PR/ET	445, 13800		ϵ relative to benzophenone in benzene ($\epsilon_{532.5} = 10300 \text{ L mol}^{-1} \text{ cm}^{-1}$)	690087
	Benzene (298 K)	FP	~450		$\tau_T = 5000 \pm 2500 \mu\text{s}$	746270
	Benzene	PR/ET	450, 20000 \pm 1000		ϵ relative to biphenyl in benzene ($\epsilon_{359} = 27100 \text{ L mol}^{-1} \text{ cm}^{-1}$)	83E281
	Bromobenzene	LP	452			83E281
	Cyclohexane	PR	445			690087
	Cyclohexane	PR/ET	440, 16000 \pm 1000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800 \text{ L mol}^{-1} \text{ cm}^{-1}$)	83E281
	Dioxane	PR	425			690087
	EtOH (298 K)	FP	~440		$\tau_T = 3000 \pm 1500 \mu\text{s}$	746270
	EtOH	FP/ET	421, 26000 ^c		Triplet ET from eosin; ϵ relative to eosin in EtOH ($\epsilon_{580} = 9400 \text{ L mol}^{-1} \text{ cm}^{-1}$)	78E019
	Heptane (298 K)	FP	~450		$\tau_T = 2500 \pm 1300 \mu\text{s}$	746270
	Liquid paraffin	FP	440			65F031
	MeOH	LP	440			83E281
	Toluene	LP/ET	452, 15500 ^a		Triplet ET from anthracene; ϵ relative to anthracene in benzene ($\Phi_T = 0.72$, $\epsilon_{428.5} = 42000 \text{ L mol}^{-1} \text{ cm}^{-1}$) assuming $\Phi_T = 0.72$ for anthracene in toluene; oxygen quenching ($3.4 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); paper shows a spectrum with a peak at 428 nm with an ϵ of 18500 $\text{L mol}^{-1} \text{ cm}^{-1}$ which we take to be the spectrum of the sensitizer anthracene; $\tau_T = 10^3 \mu\text{s}$	82E451
	Toluene	PR/ET	448, 16000 \pm 1000		ϵ relative to biphenyl in benzene ($\epsilon_{359} = 27100 \text{ L mol}^{-1} \text{ cm}^{-1}$), assuming no solvent effect	83E281
530.	4,4'-Diphenylbenzophenone Alcohol/Ether (77 K)	MOD	415		Glass was 2:1 alcohol to ether	76E682
531.	2,6-Diphenyl-1,4-benzoquinone Dibutyl phthalate	LP	600 ^a		$\tau_T = 0.56 \mu\text{s}$; $E_T = 218 - 238 \text{ kJ mol}^{-1}$	79B044
532.	2,6-Diphenyl-1,4-benzoquinone/Diphenylamine Toluene (261 K)	LP	700 ^a		Claimed triplet exciplex; $\tau_T = 0.083 \pm 0.012 \mu\text{s}$	79B044
533.	2,6-Diphenyl-1,4-benzoquinone/TMPD Dibutyl phthalate	LP	610 ^a 560 ^a		Claimed triplet exciplex; $\tau_T = 0.039 \pm 0.003 \mu\text{s}$; $E_T = < 175 \text{ kJ mol}^{-1}$	79B044

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
534.	2,6-Diphenyl-1,4-benzoquinone/Triphenylamine					
	Benzene (293 K)	LP	650		550 nm band was the more intense; $\tau_T = 0.2 \mu\text{s}$	81E715
			550			
	Benzene/MeOH (293 K)	LP	650		650 nm band was the more intense; solvent was 19:1 benzene to MeOH; $\tau_T = 0.083 \mu\text{s}$	81E715
			550			
	Dibutyl phthalate	LP	670 ^a		Claimed triplet exciplex; $\tau_T = 0.14 \pm 0.01 \mu\text{s}$; $E_T = < 175 \text{ kJ mol}^{-1}$	79B044
			550 ^a			
535.	1,4-Diphenyl-1,3-butadiene					
	2-MTHF (123 K)	LP	~395			82E429
	Acetonitrile (298 K)	LP-ET	384		†Triplet ET from benzophenone	82E429
	Benzene (295 K)	LP/SD	395 ± 2, 45000 ± 6800		Triplet yields enhanced by presence of O ₂ ; $\tau_T = 2.6 \pm 0.3 \mu\text{s}$	82E365
	Benzene (298 K)	LP-ET	394		†Triplet ET from biacetyl	82E429
			372			
	Benzene	LP	405		†Oxygen and azulene quenching; $\tau_T = 2.6 \pm 0.2 \mu\text{s}$	84E319
	Bromobenzene (295 K)	LP/SD	416 ± 2, 45000 ± 6800		$\tau_T = 1.5 \pm 0.2 \mu\text{s}$	82E365
	Cyclohexane (295 K)	LP/SD	390 ± 2, 60000 ± 9000		Triplet yields enhanced by presence of O ₂ ; shoulder at 372 nm; $\tau_T = 1.6 \pm 0.2 \mu\text{s}$	82E365
	EPA (77 K)	FP	435		Main peak at 398 nm; $\tau_T = 2450 \mu\text{s}$	707199
			413			
			398			
			357			
	Ethyl bromide (295 K)	LP/SD	392 ± 2, 55000 ± 8300		$\tau_T = 2.5 \pm 0.3 \mu\text{s}$	82E365
	Ethylene glycol	LP	393		†Oxygen and azulene quenching; shoulders at 375 and 365 nm; $\tau_T = 5.0 \mu\text{s}$	84E319
	Glycerol triacetate (298 K)	LP	~390		$\tau_T = 4 \mu\text{s}$	82E429
	Glycerol triacetate (198 K)	LP	396		$\tau_T = 1500 \mu\text{s}$	82E429
	MeOH (295 K)	LP/SD	385 ± 2, 58000 ± 8700		Triplet yields enhanced by presence of O ₂	82E365
536.	1,4-Diphenylbutadiyne					
	PMMA	PS	450		Polarization also measured; phase sensitive detection; $\tau_T = 8.0 \times 10^3 \mu\text{s}$	78E152
537.	1,2-Diphenylcyclobutene					
	Benzene (298 K)	FP	~390		$\tau_T = \geq 200 \mu\text{s}$	81E214
	Benzene (298 K)	LP-ET	~385		†Triplet ET from benzophenone, oxygen quenching ($4 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); bandwidth ~3000 cm ⁻¹ ; $\tau_T = > 3 \mu\text{s}$; $k_{et} = 6.5 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	81E214
	Benzene (298 K)	LP-ET	~385		†Triplet ET from xanthone, oxygen quenching ($3.4 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); bandwidth ~3000 cm ⁻¹ ; $\tau_T = > 3 \mu\text{s}$; $k_{et} = 6 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	81E214
	EPA (77 K)	LP	392			81E214
			373			
	EPA (298 K)	LP	~370			81E214
538.	1,4-Diphenyl-2,3-dibenzoyl-1,4-epoxy-1,4-dihydronaphthalene					
	Benzene	LP/KA	490 ± 5, 5270 ± 1300		†Oxygen and azulene quenching; ϵ relative to benzophenone in benzene ($\epsilon_{532} = 7600 \text{ L mol}^{-1} \text{ cm}^{-1}$), assuming $\Phi_T = 1$ for benzophenone and taking $\Phi_T = 0.66$ for the compound in benzene; $\tau_T = 1.7 \pm 0.3 \mu\text{s}$	84A355
	MeOH	LP	480 ± 5		†Oxygen quenching; $\tau_T = 1.6 \pm 0.2 \mu\text{s}$	84A355
539.	1,1-Diphenylethylene					
	<i>tert</i> -BuOH	LP-ET	~330		†Triplet ET from acetone	82E204
540.	9,9-Diphenyl-9H-9-germafluorene					
	MCH/Isopentane (77 K)	FP	380 ^a		†Phosphorescence lifetime; glass was 3:1 MCH to isopentane; shoulder at 363 ^a nm; $\tau_T = 4.5 \times 10^5 \mu\text{s}$	81E648

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
541.	1,6-Diphenyl-1,3,5-hexatriene					
	1,2-Dichloroethane	LP	425		Radical cation ($\lambda_{\max} \sim 615$ nm) also formed by a bi-	80N033
			400		photonic process	
	1,2-Dichloroethane	PR	425		Radical cation ($\lambda_{\max} \sim 615$ nm) also formed	80N033
			400			
	2-MTHF (123 K)	LP	428		$\tau_T = 400 \mu\text{s}$	82E429
	2-MTHF (298 K)	LP	420			82E429
			396			
	Acetonitrile (298 K)	FP	413		$\tau_T = 70 \mu\text{s}$	82E429
			390			
	Benzene	PR/ET	422.5, 113000		ϵ relative to biphenyl in benzene ($\epsilon_{359} = 27100$ L mol ⁻¹ cm ⁻¹); ϵ_{ref} unstated, but assumed from authors' earlier work	761088
			397 ^a , 70500 ^a			
	Benzene (298 K)	LP-ET	425		†Triplet ET from benzophenone and biacetyl	82E429
			400			
	Benzene (295 K)	LP/SD	426 ± 2 , 104000 ± 16000		Triplet yields enhanced by presence of O ₂ ; $\tau_T = 10 \pm 1 \mu\text{s}$	82E365
	Bromobenzene (295 K)	LP/SD	430 ± 2 , 105000 ± 16000		$\tau_T = 13 \pm 1 \mu\text{s}$	82E365
	CTAB/Triton X-100	LP	415		Surfactant ratio 1:2 CTAB to Triton X-100; mixed aqueous micelle; $\tau_T = \sim 30 \mu\text{s}$	80N033
	Cyclohexane	PR	425		Radical anion ($\lambda_{\max} \sim 660$ nm) and cation ($\lambda_{\max} \sim 630$ nm) also formed; half-life = 20 μs	80N033
			400			
	Cyclohexane (295 K)	LP/SD	416 ± 2 , 114000 ± 17000		Triplet yields enhanced by presence of O ₂ ; $\tau_T = 20 \pm 2 \mu\text{s}$	82E365
	EPA (77 K)	FP	429		Main peak at 429 nm; $\tau_T = 400 \mu\text{s}$	707199
			405			
	EtOH	PR/RF	415, 113000		ϵ relative to 1,6-diphenyl-1,3,5-hexatriene in benzene ($\epsilon_{422.5} = 113000$ L mol ⁻¹ cm ⁻¹) assuming oscillator strength independent of solvent; $E_T = \sim 150$ kJ mol ⁻¹	761088
			388 ^a , 70300 ^a			
542.	1,4-Diphenylnaphthalene					
	3-MP (77 K)	PS/ESR	505, ~ 10000		†ESR; oscillator strength = $\sim 0.02, 0.52$	69B002
			444, 32500			
543.	1,5-Diphenylnaphthalene					
	3-MP (77 K)	PS/ESR	538, 22800		†ESR; oscillator strength = 0.25	69B002
544.	1,8-Diphenyl-1,3,5,7-octatetraene					
	2-MTHF (298 K)	LP	435			82E429
	2-MTHF (123 K)	LP	440			82E429
	Acetonitrile (298 K)	FP	432		$\tau_T = 70 \mu\text{s}$	82E429
			408			
	Benzene	PR/ET	440, 162000		ϵ relative to biphenyl in benzene ($\epsilon_{359} = 27100$ L mol ⁻¹ cm ⁻¹); ϵ_{ref} unstated, but assumed from authors' earlier work	761088
					†Triplet ET from benzophenone and biphenyl; $\tau_T = 100 \mu\text{s}$	82E429
	Benzene (298 K)	FP	444			
			419			
	Benzene (295 K)	LP/SD	445 ± 2 , 188000 ± 28000		Triplet yields enhanced by presence of O ₂ ; $\tau_T = 21 \pm 2 \mu\text{s}$	82E365
	Bromobenzene (295 K)	LP/SD	450 ± 2 , 179000 ± 27000		$\tau_T = 14 \pm 1 \mu\text{s}$	82E365

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Cyclohexane (295 K)	LP/SD	437 ± 2, 32000	210000 ±	Triplet yields enhanced by presence of O ₂ ; $\tau_T = 40 \pm 4 \mu s$	82E365
	EtOH	PR/RF	427, 162000		ϵ relative to 1,8-diphenyl-1,3,5,7-octatetraene in benzene ($\epsilon_{440} = 162000$ L mol ⁻¹ cm ⁻¹) assuming oscillator strength independent of solvent; $E_T = \sim 120$ kJ mol ⁻¹	761088
	Ethyl bromide (295 K)	LP/SD	440 ± 2, 35000	230000 ±	$\tau_T = 11 \pm 1 \mu s$	82E365
	Glycerol triacetate (298 K)	LP	440		$\tau_T = 60 \mu s$	82E429
	Glycerol triacetate (198 K)	LP	444		$\tau_T = 100 \mu s$	82E429
	MeOH (295 K)	LP/SD	430 ± 2, 29000	191000 ±	Triplet yields enhanced by presence of O ₂ ; $\tau_T = 34 \pm 3 \mu s$	82E365
545.	2,5-Diphenyl-1,3,4-oxadiazole					
	Benzene	LP	550		†Oxygen quenching (1.6×10^9 L mol ⁻¹ s ⁻¹); 100 ns delay; "PPD"; $\tau_T = 0.300 \mu s$	777265
	Cyclohexane	FP/TD	449, 670 ^a 440, 880 ^a 432, 900 ^a 425, 980 ^a		Authors claim that the spectrum reported in [777265] is erroneous; oscillator strength = 0.01	83E216
546.	2,5-Diphenyloxazole					
	Benzene	PR	500		†Triplet ET from naphthalene and triplet ET to anthracene; "PPO"	720206
	Benzene	LP	570		†Oxygen quenching (2.5×10^9 L mol ⁻¹ s ⁻¹); 100 ns delay; $\tau_T = 0.200 \mu s$	777265
	Cyclohexane	PR	500			720206
	Cyclohexane	FP/ET	500, 13000		†Triplet ET to anthracene; ϵ relative to anthracene in cyclohexane ($\epsilon_{425} = 64700$ L mol ⁻¹ cm ⁻¹); $\tau_T = 1.7 \times 10^3 \mu s$	80E439
	EPA (77 K)	CWL	515			74B003
	EtOH	FP/ET	500, 28400 ± 1500		ϵ relative to naphthalene in EtOH ($\epsilon_{415} = 40000$ L mol ⁻¹ cm ⁻¹); $\tau_T = 2500 \pm 1300 \mu s$; $k_{et} = (9.2 \pm 1.0) \times 10^9$ L mol ⁻¹ s ⁻¹	747049
547.	N,N'-Diphenyl-p-phenylenediamine					
	Benzene	PR/ET	615, 26800		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700$ L mol ⁻¹ cm ⁻¹); ϵ obtained from a simultaneous least squares fit of data from several compounds making use of cyclohexane to benzene ϵ_{\max} ratios of 1.83 for naphthalene and 1.45 for anthracene	71E360
548.	N,N'-Diphenyl-p-phenylenediamine, conjugate acid					
	EPA/Acetic acid (137.4 K)	FP/ELT	610 600 ^b , 67000 ± 15000		†Phosphorescence decay; ϵ relative to the radical cation of compound ($\epsilon_{390} = 22000$ L mol ⁻¹ cm ⁻¹); solvent was 97:3 EPA to acetic acid; EPA was 8:3:5 Et ₂ O to isopentane to EtOH in this work	67A001
549.	9,9-Diphenyl-9H-9-silafluorene					
	MCH/Isopentane (77 K)	FP	373 ^a		†Phosphorescence lifetime; glass was 3:1 MCH to isopentane; $\tau_T = 3.0 \times 10^6 \mu s$	81E648
550.	N-[(Dipropylamino)methyl]phthalimide					
	EtOH	FP	590 350		†Oxygen quenching, diene quenching; maxima assumed from text; $\tau_T = 10 \mu s$	79A147
551.	Durene/Tetracyanobenzene					
	Et ₂ O/Isopentane (77 K)	MOD	541		Glass was 1:1 Et ₂ O to isopentane; another maximum > 1430 nm	72E276

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
552.	Duroquinone					
	2-PrOH	LP	457 ^a		†Triplet ET to anthracene; shoulder at 484 ^a	717520
	Acetonitrile	LP	485 ^a		50 ns delay; quenched by amines	777602
	Acetonitrile (293 K)	FP	480			82E614
	Benzene	PR/ET	490, 9450		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361.3} = 35400$ L mol ⁻¹ cm ⁻¹); author reports a mean of 9800 L mol ⁻¹ cm ⁻¹ from 3 separate standards	690520
	Benzene	PR/ET	490, 8600		ϵ relative to anthracene in cyclohexane ($\epsilon_{420} = 57200$ L mol ⁻¹ cm ⁻¹); author reports a mean of 9800 L mol ⁻¹ cm ⁻¹ from 3 separate standards	690520
	Benzene	PR/ET	490, 11400 ^c		ϵ relative to naphthalene in cyclohexane ($\epsilon_{412.5} = 22600$ L mol ⁻¹ cm ⁻¹); author reports a mean of 9800 L mol ⁻¹ cm ⁻¹ from 3 separate standards	690520
	Benzene	LP	490		$\tau_T = 3.1$ μ s	70E288
	Benzene	LP	477 ^a		†Triplet ET to anthracene and dibenz[<i>a,h</i>]anthracene; $\tau_T = 10$ μ s	717520
	Benzene	PR/ET	490, 6950		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700$ L mol ⁻¹ cm ⁻¹); ϵ obtained from a simultaneous least squares fit of data from several compounds making use of cyclohexane to benzene ϵ_{\max} ratios of 1.83 for naphthalene and 1.45 for anthracene	71E360
	Benzene (293 K)	FP	490			82E614
	Cyclohexane	PR/ET	490 ^b , 4250		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361.3} = 35400$ L mol ⁻¹ cm ⁻¹); author reports a mean of 4700 L mol ⁻¹ cm ⁻¹ from 3 separate standards	690520
	Cyclohexane	PR/ET	490 ^b , 4950		ϵ relative to anthracene in cyclohexane ($\epsilon_{420} = 57200$ L mol ⁻¹ cm ⁻¹); author reports a mean of 4700 L mol ⁻¹ cm ⁻¹ from 3 separate standards	690520
	Cyclohexane	PR/ET	490 ^b , 4900		ϵ relative to naphthalene in cyclohexane ($\epsilon_{412.5} = 22600$ L mol ⁻¹ cm ⁻¹); author reports a mean of 4700 L mol ⁻¹ cm ⁻¹ from 3 separate standards	690520
	Cyclohexane	PR/ET	490, 5330		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700$ L mol ⁻¹ cm ⁻¹); reference ϵ obtained by starting from $\epsilon_{\max} = 3220$ L mol ⁻¹ cm ⁻¹ for this ketyl radical in water and assuming the f of the ketyl radical is independent of solvent; final ϵ obtained from a simultaneous least squares fit to data from several compounds	71E360
	Cyclohexane	LP	490 450 ^a 310 ^a		Relative intensities (5.4:11); $\tau_T = 21$ μ s; oscillator strength = 0.121 ± 0.005	767144
	Dioxane	FP	485		Half-life = ~ 30 μ s	587005
	Et ₂ O	FP	~ 485		Half-life = ~ 30 μ s	587005
	EtOH	FP	~ 485		Half-life = ~ 30 μ s	587005
	EtOH	LP/RF	490, 5580 \pm 550 455 ^a 310 ^a		ϵ relative to duroquinone in cyclohexane ($\epsilon_{490} = 5330$ L mol ⁻¹ cm ⁻¹) assuming oscillator strength independent of solvent; relative intensities (1:1:2); $\tau_T = 15$ μ s	767144
	EtOH/Sulfuric acid	FP	~ 490		Solvent was 50% 0.1 mol L ⁻¹ H ₂ SO ₄ and 50% alcohol; half-life = 30 μ s	587005
	EtOH/Water	FP	~ 485		Solvent was 1:1 EtOH to water; half-life = ~ 30 μ s	587005
	Liquid paraffin	FP	490 460		Half-life = 100 μ s	587005
	Liquid paraffin	LP	458 ^a		†Quenching by anthracene; shoulder at 493 ^a nm; $\tau_T = 29$ μ s	717520
	MeOH (293 K)	FP	460			82E614
	PMMA	FP	470		$\tau_T = 180 \pm 10$ μ s	707305
	SDS	LP	~ 450		Aqueous micelle	83N056
	Toluene	LP/ET	490, 7600 \pm 500		ϵ relative to anthracene in toluene ($\epsilon_{428.5} = 42000$ L mol ⁻¹ cm ⁻¹)	83F075
	Water	LP/RF	500, 4200 \pm 450 440 420 395 ^a		ϵ relative to duroquinone in cyclohexane ($\epsilon_{490} = 5330$ L mol ⁻¹ cm ⁻¹) assuming oscillator strength independent of solvent; relative intensities (4:5:4:12); $\tau_T = 5.9$ μ s	767144

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Water/EtOH	LP/ELT	490 ^a 450 ^a 420 ^b , 6600 ± 800		ϵ relative to durosemiquinone ($\epsilon_{420} = 4700$ L mol ⁻¹ cm ⁻¹) assuming complete conversion in the presence of Fe ²⁺ ions; solvent was 2:1 water to EtOH by volume; $\tau_T = 6.3 \pm 0.8$ μ s	777045
	Water/MeOH	FP	510 ^a 440 ^a		Solvent was 4:1 water to MeOH by volume; lifetime measured at pH = 0.7; $\tau_T = 1.3$ μ s; pH > 1	80C005
553.	Duroquinone/Triphenylamine					
	Benzene (293 K)	LP	490		$\tau_T = 2$ μ s	81E715
	Benzene/MeOH (293 K)	LP	650 550		650 nm band was the more intense; solvent was 19:1 benzene to MeOH; $\tau_T = 0.27$ μ s	81E715
554.	Duroquinone, conjugate monoacid					
	Water/MeOH	FP	470 ^a ~380 ^a		$pK_a = -0.1$; solvent was 4:1 water to MeOH by volume; lifetime measured at pH = -0.8; $\tau_T = \sim 0.3$ μ s; pH < 1	80C005
555.	Echinenone					
	Benzene	PR-ET	550		†Triplet ET from naphthalene; $\tau_T = 5.2$ μ s	80A143
556.	Eosin					
	Acetonitrile/Water	LP	580		Solvent was 3.5:1.5 acetonitrile to water; $\tau_T = 30$ μ s	84E216
	EtOH	FP/SD	580, 9400			716235
	Water	FP	550		Maximum was near the SD region; pH 9.1	746168
557.	Eosin/Aniline					
	Water	FP	555		Maximum was near SD region; pH 5.5	746168
558.	Eosin/4-Bromophenol					
	Water	FP	660		pH 9.2	746168
559.	Eosin/2-Naphthol					
	Water	FP	584		pH 9.0	746168
560.	Eosin/Resorcinol					
	Water	FP	580		pH 7.3	746168
561.	Eosin/Sulfanilic acid					
	Water	FP	597		pH 4.0	746168
562.	Eosin dianion					
	Water	FP/SD	540, 26000 ^a		Decay time was measured at pH 9; solution was oxygen free; $pK_b > 10$; paper reported a band at 459 nm that was later suggested to be due to semi-oxidized eosin [65A002]; $\tau_T = 2400$ μ s; pH 5.4	63A001
	Water	FP	565 ^a 325		565 nm maximum was near the SD region in an apparent difference spectrum and may not be the true maximum; paper reported 580 nm (instead of 565) as a maximum, but displayed spectrum and other reports were inconsistent with this; solution was a 0.1 mol L ⁻¹ phosphate buffer solution; pH 7.2	66A002
	Water	FP/SD	518 ^b , 28000 ± 4000		pH 9	67E031
	Water	FP/TD	630, 4800		pH Basic	727073
563.	5-(1,4-Epidioxyphenyl)-6,11,12-triphenyltetracene					
	Cyclohexane	LP	390 ^a		A rubreneperoxide, called "III" in paper	84E056
564.	Ergosterol					
	Benzene	LP-ET/TD	305, 2800 ± 500		†Triplet ET from 2-fluorenyl phenyl ketone and other triplet sensitizers; ϵ assumes ET proceeds with unit efficiency; $\tau_T = > 150$ μ s; $E_T = \sim 220$ kJ mol ⁻¹	80B021

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
565.	Erythrosin ?	FP	600 390		Solvent was not specified, probably water at an unknown pH	747229
	Acetonitrile/Water	LP	780		Solvent was 3.5:1.5 acetonitrile to water	84E216
566.	Erythrosin dianion Gelatin	FP/SD	525 ^a , 12000 ^a		Extinction coefficient was calculated on the assumption that 44% of the ground state was converted to triplets in a 10 J flash; $\tau_T = 630 \mu\text{s}$	64E016
	Water	FP/SD	526 ^b , 26000 \pm 7000		pH 9	67E031
567.	2-Ethoxynaphthalene Cyclohexane	FP	440 ^a 411 ^a		440 nm peak was the more intense; delay 30 μs	686045
	Liquid paraffin	FP	436 413 394		Solvent viscosity was 0.19 N·s/m ² ; relative intensities (10:8:5)	58E001
568.	Ethyl Violet PMMA (77 K)	PS	1232 ^a 625 ^a		Relative intensities (3:20); shoulder at 535 ^a nm; solvent contains 2-chloroethanol	69B003
569.	10-Ethyl-9(10H)-acridinone EtOH (300 K)	FP/SD	580, 14600 \pm 1500 310		"N-Ethylacridinone"; 310 nm quoted from [79E964]; $\tau_T = \sim 43 \mu\text{s}$	81E649
570.	N-Ethylcarbazole 2-MTHF	LP	420		Shoulder at 500 nm; $\tau_T = 5 \mu\text{s}$	81E016
	Acetonitrile	LP/ET	420, 27000		ϵ relative to phenanthrene ($\epsilon_{480} = 27000 \text{ L mol}^{-1} \text{ cm}^{-1}$)	77B012
	Cyclohexane	FP	500 ^a 425 ^a		\uparrow Triplet ET to anthracene; $\tau_T = 200 \mu\text{s}$	78A368
	Cyclohexane	FP/ET	420, 13500		ϵ relative to naphthalene in cyclohexane ($\epsilon_{415} = 24500 \text{ L mol}^{-1} \text{ cm}^{-1}$)	84F039
	EtOH	LP/ET	420, 25000		ϵ relative to phenanthrene ($\epsilon_{480} = 27000 \text{ L mol}^{-1} \text{ cm}^{-1}$); solvent and temperature assumed	78E394
	Hexane	FP	420		\uparrow Oxygen and perylene quenching; "ethylcarbazole"	757573
	MeOH	LP	420			77B012
	THF	LP	423 ^a		Delay 1 μs ; shoulders at 498 ^a and 398 ^a nm	83E662
	Toluene	LP	425		\uparrow Rises with fluorescence decay; 100 ns delay	80E441
571.	(E,Z)-3-Ethyl-5[(3-ethyl-2(3H)-benzothiazolylidene)ethylidene]-2-thioxo-4-thiazolidinone Toluene	FP-ET	600 ^a		\uparrow Triplet ET from fluorenone; "merocyanine dye"	767764
572.	1-Ethyl-2-[5-(1-ethyl-1,3-dihydro-3,3-dimethyl-2H-indol-2-ylidene)-3-fluoro-1,3-pentadienyl]-3,3-dimethyl-3H-indolium 1-PrOH	LP-ET	710		\uparrow Triplet ET from anthracene, oxygen quenching; $\tau_T = 109 \mu\text{s}$	78A386
573.	1-Ethyl-2-[5-(1-ethyl-1,3-dihydro-3,3-dimethyl-2H-indol-2-ylidene)-1,3-pentadienyl]-3,3-dimethyl-3H-indolium 1-PrOH	LP-ET	690		\uparrow Triplet ET from anthracene, oxygen quenching; $\tau_T = 455 \mu\text{s}$	78A386
574.	1-Ethynaphthalene Isooctane	LP	415 ^a 390 ^a			81E015
575.	2-Ethynaphthalene Benzene	PR	430 ^a		\uparrow Triplet ET to perylene; $\tau_T = 7.1 \mu\text{s}$; $k_{et} = 1.29 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	81E508
576.	9-Ethylphenanthrene Liquid paraffin (293 K)	LP	482 ^a 452 ^a		Shoulder at 428 ^a nm	82E287
	THF	LP	480 ^a 453 ^a		\uparrow Triplet ET to naphthalene; delay 500 ns; shoulder at 427 ^a nm	83E554

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
577.	<i>N</i> -Ethylphthalimide Acetonitrile	LP	335		†Oxygen quenching; $\tau_T = 31 \mu\text{s}$	80B066
578.	β -Ethylstyrene Cyclohexane	LP/RA-ET	325 ^b , 3500		ϵ relative to naphthalene in cyclohexane ($\epsilon_{413} = 24000$ L mol ⁻¹ cm ⁻¹), but in ref. cited ($\epsilon_{415} = 24500$ L mol ⁻¹ cm ⁻¹); triplet ET from thioxanthone; $\tau_T = \sim 0.030 \mu\text{s}$	82E181
579.	Etiochlorin II Dimethyl phthalate	FP/SD	416 ^a		Shoulders at 466 nm and 424 nm	74B007
580.	Euchrysine, conjugate monoacid Glycerol (293 K)	FP	620 ^a 560 ^a		Solvent has added glucose	776171
581.	Flavine mononucleotide Water	LP	680		$\tau_T = 11 \mu\text{s}$; pH 7.0	81A024
582.	Flavine mononucleotide, conjugate monoacid Water	LP	680		$pK_a = 5.2$; $\tau_T = 10 \mu\text{s}$; pH 4.1	81A024
583.	Flavone Benzene	LP	661 ^a 366 ^a		†Oxygen quenching; 661 nm peak was the more intense; delay 140 ns	80E884
	EPA (77 K)	FP	640 ^a 370 ^a		Lifetime measured at 370 nm; 370 nm peak was the more intense; $\tau_T = 4.1 \times 10^5 \mu\text{s}$	80E884
584.	Fluoranthene PMMA (293 K)	FP	~ 450		$\tau_T = 4 \times 10^5 \mu\text{s}$	70E291
	Water/ <i>tert</i> -BuOH	FP	405 ^a		Solvent mixture contains "1-5%" <i>tert</i> -BuOH for solubility; shoulder at 380 nm; pH ~ 6	767189
585.	Fluorene 2-MTHF (77 K)	PS/ESR	384, 20700		†ESR; only most intense visible peak reported; oscillator strength = ~ 0.25	696115
	2-MTHF (77 K)	PS/KM	386 ^a , 18000 ^a			83E383
	EtOH (293 K)	FP	376			68E098
	EtOH/Et ₂ O (77 K)	MOD/KM	382, 12000 \pm 3600		Glass was 2:1 EtOH to Et ₂ O; temperature was not explicitly stated, but 77 K was inferred from the context	719059
	Heptane (77 K)	PS	421 386 366			67B007
	Heptane	FP	380 ^a 365 ^a		380 nm peak was the more intense	766421
	Hexane	FP	376			54E001
	Hexane (300 K)	MOD/SD	499 ^a , 3200 ^a 376, 40000		$\tau_T = 150 \pm 2 \mu\text{s}$	69E208
	Isopentane/MCH (77 K)	PS	421.1 388		Relative intensities (23:100); unresolved shoulder at 395 nm; $\tau_T = (5.7 \pm 0.3) \times 10^6 \mu\text{s}$	54B001
	Liquid paraffin	FP	503 382.5		Solvent viscosity was 0.03 N·s/m ² ; relative intensities (22:100); bands were assigned to 2 different electronic transitions	58E001
	PVA	FP	385		Stretched polymer film; polarized light indicates underlying band at 400 nm	776279
	Polystyrene	PR	394 ^a		$G \epsilon_{394} = 9900^a$ L mol ⁻¹ cm ⁻¹ / (100 eV absorbed)	701073
	Toluene/EtOH (77 K)	MOD	387 ^a		Glass was 19:1 toluene to EtOH; shoulder at 509 ^a nm	719059

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
586.	Fluoren-9-one					
	Acetonitrile	LP/?	420, 6000		ϵ method unspecified; $\tau_T = 23 \mu s$	767201
	Acetonitrile	LP	630 600 425 387 317.5 285		†Intensity of delayed fluorescence proportional to square of transient absorbance, decay kinetics; most intense peak at 285 nm; $\tau_T = 100 \mu s$	78E495
	Benzene	FP	600 430		†Oxygen quenching; $\tau_T = 60 \mu s$	69E218
	Benzene	LP/ET	430, 5900 \pm 500		†Triplet ET from benzophenone; ϵ relative to benzophenone in benzene ($\epsilon_{332.5} = 7630$ L mol ⁻¹ cm ⁻¹)	757282
	Benzene	LP	610 425		$\tau_T = 100 \mu s$	767201
	Benzene	LP	650 610 435 387 322.5 288		†Intensity of delayed fluorescence proportional to square of transient absorbance, decay kinetics; most intense peak at 288 nm with approximate $\epsilon = 30000$ L mol ⁻¹ cm ⁻¹ ; $\tau_T = 1000 \mu s$	78E495
	Cyclohexane	LP	590 ^a 425		†Triplet ET, oxygen quenching; decay times limited in oxygen saturated solution; shoulder around 445 nm; relative intensities (1:10); $\tau_T = 0.220 \mu s$; rise time of 0.140 \pm 0.030 ns	766464
	Cyclohexane	LP	620 460 420 360		†Oxygen quenching; $\tau_T = 2.5 \mu s$; rise time of < 50 ns	767201
	Hexane	LP	600 430		325 ps delay; relative intensities (1:3)	79B007
	MCH	LP	650 600 425 387 325 285		†Intensity of delayed fluorescence proportional to square of transient absorbance, decay kinetics; most intense peak at 285 nm; $\tau_T = 500 \mu s$	78E495
	Water/ <i>tert</i> -BuOH	FP	440 ^a 403 ^a 370 ^a		Solvent mixture contains "1-5%" <i>tert</i> -BuOH for solubility; pH \sim 6	767189
587.	1-Fluorenyl diisopropyl borate					
	Triisopropyl borate	FP	380			717266
588.	2-Fluorenyl diisopropyl borate					
	Triisopropyl borate	FP	380			717266
589.	Fluoren-9-ylidene					
	Acetonitrile	LP	500 ^a 400 ^a		Relative intensities (8:1); 200 ns delay	80A222
590.	2-Fluorenyl phenyl ketone					
	Benzene	LP/TD	525, 18600 \pm 900		$\tau_T = 360 \mu s$	80B021
591.	Fluorescein					
	Boric acid (178 K)	PS/SD	650, 21000 ^a 505, 17000 ^a		ϵ relative to ground state $\epsilon_{436} = 67000^a$ L mol ⁻¹ cm ⁻¹ .	41E001
	Boric acid (77 K)	PS/SD	1060, 17000 920 650			65B004
	EtOH/Et ₂ O (90 K)	PS/SD	1140, 13500 ^a 640, 7000 ^a 470, 9500 ^a 370, 8300 ^a 320, 4000 ^a		Solvent was 2:1 EtOH to Et ₂ O	57B001

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	PMMA (193 K)	PS	500		Shoulders at 640 and 550 nm	59B001
			465			
	PMMA (77 K)	PS	1100			65B004
			960			
			660			
	PMMA (77 K)	PS	1660			65B004
			1080			
			940			
			790			
			740			
			635			
			550			
			510			
	PMMA (77 K)	PS	1056 ^a		Relative intensities (5:2:2:2); shoulders at 926 ^a and 461 ^a nm; solvent contains 2-chloroethanol; pH 4.0	69B007
			641 ^a			
			552 ^a			
			498 ^a			
592.	Fluorescein, conjugate monoacid					
	Sulfuric acid	FP/SD	640, 7700 ^a		Cation of fluorescein; $pK_a = -2.1$; solution was 9 mol L ⁻¹ H ₂ SO ₄ ; a spectrum was given also for the neutral triplet, at pH 0.7, which showed absorption between 450 and 850 nm, no distinct maximum was evident, but the extinction coefficient of the neutral triplet ranged between 1000 and 10,000 L mol ⁻¹ cm ⁻¹ in this region; $\tau_T = 4500$ μ s; pH Acidic	63F019
593.	Fluorescein dianion					
	EtOH	FP/TD	570, 6800		pH Basic	727073
	Water	FP/SD	535 ^a , 8700 ^a		Another maximum > 950 nm; pK_b 7.0; 535 nm peak was very broad, with 8700 L mol ⁻¹ cm ⁻¹ reading extending from ~532 to ~540 nm; $\tau_T = 2.0 \times 10^4$ μ s; pH 12	60A001
	Water	FP/SD	489 ^b , 15000 \pm 4000		pH 9	67E031
	Water	FP/TD	570, 8400		pH Basic	727073
594.	Fluorobenzene					
	Cyclohexane	LP	290 ^a		400 ns delay; $\tau_T = 0.67$ μ s	707561
595.	4-Fluorobenzophenone					
	Acetonitrile/Water	LP/ET	520, 5900		Solvent was 9:1 acetonitrile to water; ϵ relative to 1-methylnaphthalene in acetonitrile/water ($\epsilon_{415} = 11200$ L mol ⁻¹ cm ⁻¹)	84B033
			315, 12100			
596.	trans-4-Fluorostilbene					
	EPA (98 K)	LP	383		Triplet absorption not observable above 110 K; reported lifetime constant below 98 K; $\tau_T = 10^4$ μ s	79E640
			365			
			349			
	Glycerol (223 K)	LP	385		Relative intensities (3:2:1); triplet absorption not observable above 244 K; reported lifetime constant below 198 K; $\tau_T = 10^4$ μ s	79E640
			365			
			345			
597.	γ -[2-(Formylamino)phenyl]- γ -oxo- α -aminobutyryl-L-lysine					
	EtOH	LP	460		$\tau_T = 0.3$ μ s	78E308
	Water	LP	450		†Oxygen quenching (1.3×10^9 L mol ⁻¹ s ⁻¹); ionic strength 0.01 mol L ⁻¹ ; $\tau_T = 0.85$ μ s; pH 7.5	78E308
598.	3-Formylfurochromene					
	EtOH	LP/ET	440, 28000		†Triplet ET to retinol; ϵ relative to retinol in hexane ($\epsilon_{405} = 80000$ L mol ⁻¹ cm ⁻¹) assuming ϵ independent of solvent; $\tau_T = 50$ μ s; $k_{et} = 4.2 \times 10^9$ L mol ⁻¹ s ⁻¹	79E678

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
599.	<i>N</i> -Formylkynurenine					
	EtOH	LP	450		†Oxygen quenching (4.5×10^8 L mol ⁻¹ s ⁻¹); $\tau_T = 10$ μ s	78E308
	Water	FP	435		†Oxygen and KI quenching; $\tau_T = 2$ μ s; pH 7.6	757361
	Water	LP	440		†Oxygen quenching (2×10^9 L mol ⁻¹ s ⁻¹); ionic strength 0.01 mol L ⁻¹ ; $\tau_T = 6.7$ μ s; pH 7.5	78E308
600.	2-Formyl-5'-methyldifurobenzene					
	EtOH	LP/ET	540, 14000		†Triplet ET to retinol; ϵ relative to retinol in hexane ($\epsilon_{405} = 80000$ L mol ⁻¹ cm ⁻¹) assuming ϵ independent of solvent; $\tau_T = 13$ μ s; $k_{et} = 4.7 \times 10^9$ L mol ⁻¹ s ⁻¹	79E678
601.	3-Formyl-2-methylfurochromene					
	EtOH	LP/ET	450, 49500		†Triplet ET to retinol; ϵ relative to retinol in hexane ($\epsilon_{405} = 80000$ L mol ⁻¹ cm ⁻¹) assuming ϵ independent of solvent; $\tau_T = 29$ μ s; $k_{et} = 3 \times 10^9$ L mol ⁻¹ s ⁻¹	79E678
602.	3-Formyl-8-methylfurochromene					
	EtOH	LP/ET	450, 22000		†Triplet ET to retinol; ϵ relative to retinol in hexane ($\epsilon_{405} = 80000$ L mol ⁻¹ cm ⁻¹) assuming ϵ independent of solvent; $\tau_T = 18$ μ s; $k_{et} = 6 \times 10^9$ L mol ⁻¹ s ⁻¹	79E678
603.	<i>n</i> -Glucose phenyllosazone					
	EtOH	LP/ET	460 ^b , 61200		N-chelate; no ref. compd. given for ϵ -measurement; $\tau_T = 0.46 \pm 0.01$ μ s	83E018
	EtOH	LP/ET	530		N-chelate; ϵ relative to biacetyl in EtOH ($\epsilon_{740} = 1760$ L mol ⁻¹ cm ⁻¹); $k_{et} = 2 \times 10^9$ L mol ⁻¹ s ⁻¹	84E162
			460, 61000 \pm 18000			
604.	Hematoporphyrin IX					
	Acetic acid	FP	450		†Oxygen quenching	776190
	Acetic acid	LP	440		†Triplet ET to β -carotene; $\tau_T = 42$ μ s	81B115
	CTAB	LP	888		Aqueous micelles	82N210
			435			
			320			
	EtOH	FP	437		†Oxygen quenching; $\tau_T = 910$ μ s	82B118
	MeOH/Water	LP/TD	399 ^a , 117000 ^a		Solvent 9:1 MeOH to pH 7.4 phosphate buffer; shoulders at 370 ^a and 320 ^a nm	83E667
	SDS	LP	896		Aqueous micelles	82N210
			438			
			323			
	Triton X-100	LP	895		Aqueous micelles	82N210
			439			
			320			
	Water	FP	440		†Oxygen quenching; $\tau_T = 710$ μ s; pH 10.3	82B118
	Water	LP	890		pH Neutral	82N210
			440			
			320			
	Water	FP	440		†Oxygen quenching; phosphate buffer; $\tau_T = 770$ μ s; pH 7.1	82B118
	Water	LP/TD	390 ^a , 79400 ^a 371 ^a , 85800 ^a		Phosphate buffer; pH 7.4	83E667
605.	<i>all-trans</i> -1,3,5-Heptatriene					
	Toluene	PR	315		†Oxygen quenching; $G\Delta\epsilon_{\max} = (1.8 \pm 0.2) \times 10^4$ L mol ⁻¹ cm ⁻¹ / (100 eV absorbed); $\tau_T = 0.29 \pm 0.03$ μ s	84B139
606.	Hexacene					
	1,2,4-Trichlorobenzene	FP	550		†Triplet ET from pentacene; shoulder at 481 ^a ; 550 nm peak was the more intense; $\tau_T = 10$ μ s	82E280
			513 ^a			
607.	1,1,1,5,5,5-Hexafluoroacetylacetone					
	EtOH/MeOH (118 K)	FP/SD	390, 1000 \pm 200		Solvent was 3:1 EtOH to MeOH	68B005

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
608.	Hexahelicene Et ₂ O/Isopentane (77 K)	PS	670		†Phosphorescence decay; glass 1:1 Et ₂ O to isopentane; $\tau_T = 1.9 \times 10^6 \mu s$	62E008
609.	10,11,13,14,16,17-Hexahydro-5,7:20,22-dietheno-8 <i>H</i> ,19 <i>H</i> -diindeno[2,1- <i>n</i> :1',2'- <i>o</i>][1,4,7,10]tetraoxacyclononadecin 2-MTHF (77 K)	PS/KM	383 ^a , 17100 ^a		Compound "XII" in paper	83E383
610.	9,10,11,12,13,14-Hexahydro-5,7:15,17-dietheno-2,20-heptano- <i>H</i> -cyclotetradeca[1,2- <i>a</i> :1,14- <i>a'</i>]diindene 2-MTHF (77 K)	PS/KM	465 ^a , 35000 ^a		Compound "IX" in paper	83E383
611.	Hexamethylbenzene/Tetracyanobenzene Et ₂ O/Isopentane (77 K)	MOD	1400 532		Glass was 1:1 Et ₂ O to isopentane	72E276
612.	6-Hydroxy-7 <i>H</i> -benz[<i>de</i>]anthracen-7-one 2-PrOH Benzene	LP LP	530 ^a 535 ^a		Half-life = $2.4 \pm 0.4 \mu s$ †Oxygen quenching (1.2×10^9 L mol ⁻¹ s ⁻¹); half-life = $1.0 \pm 0.2 \mu s$	757427 757427
613.	2-Hydroxybenzophenone EtOH MeOH (175 K)	LP LP	450 ~500		Possible tautomeric triplet state; 480 ps delay; $S_n \leftarrow S_1$ absorption ($\lambda_{\max} = 435$ nm) also observed at 7 ps delay $\tau_T = 0.024 \mu s$	80B001 82E475
614.	4-Hydroxybenzophenone Isopentane (77 K)	FP	512 ^a 360 ^a		†Phosphorescence decay	81E105
615.	4-Hydroxybiphenyl PVA (77 K)	FP	520 390 348		Stretched polymer film; most intense peak at 390 nm; polarization also measured	78B129
616.	3-Hydroxychromone Benzene	LP	397 ^a		Delay 200 ns; maximum uncertain due to absorption by a tautomer in the same spectral region	83B113
617.	4-Hydroxycoumarin EPA (77 K)	PS	497.5 465.1 435 413 ^a 382 ^a		†Phosphorescence decay; the 2 smallest wavelength maxima were somewhat in doubt because of photo- chem. in this region of the spectrum; most intense peaks were at 497.5, 465.1, and 435 nm; $\tau_T = (1.4 \pm 0.2) \times 10^6 \mu s$	71B001
618.	1-(2-Hydroxyethyl)-3,3-dimethyl-6'-nitrospiro-(2,2'-indolin[2 <i>H</i> -1]benzopyran) Benzene	PR	<430		†Oxygen quenching; cis-cisoid isomer	78A411
619.	3-Hydroxyflavone 3-MP	LP	400		†Oxygen quenching	83B030
620.	7-Hydroxyflavone MeOH	LP	370		†Oxygen quenching; $\tau_T = 5 \mu s$	84A218
621.	1-Hydroxyfluorene Isooctane PMMA	FP FP	385 385			717266 717266
622.	2-Hydroxyfluorene Isooctane PMMA Toluene/EtOH (77 K)	FP FP MOD	385 385 422 ^a		Glass was 19:1 toluene to EtOH; shoulders at 530 ^a and 387 ^a nm	717266 717266 719059

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
623.	4-Hydroxy-6-methylcoumarin EPA	CWL	501 ^a 464 ^a 434 ^a 383 ^a		Glass, temperature unspecified	73E344
24.	7-Hydroxy-4-methylcoumarin EPA	CWL/SD	505 ^a , 14500 ^a ± 3600		Glass, temperature unspecified	73E344
625.	2-Hydroxy-2-methyl-1-phenylpropan-1-one Benzene	LP	310		†Triplet ET to naphthalene; $\tau_T = 0.030 \mu\text{s}$; $k_{et} = 5 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	80B030
626.	2-Hydroxy-2-methyl-1-[4-(2-propyl)phenyl]propan-1-one Benzene Cyclohexane	LP LP	310 ^a 340		†Triplet ET to naphthalene; $\tau_T = 0.050 \mu\text{s}$	80B030 80B030
627.	9-Hydroxyphenanthrene 1,2-Dichloroethane	LP	455 ^a		†Phosphorescence decay in MTHF at 77K; 400 ns delay	79A177
628.	4'-Hydroxypropiophenone Acetonitrile/Water	LP	387 ^a		Solvent isotope insensitive quenching; solvent was 9:1 acetonitrile to water; lifetime limited by self-quenching; $\tau_T = 0.230 \mu\text{s}$	81A174
629.	4-Hydroxypyrene 1,2-Dichloroethane	LP	410		†Oxygen quenching	79A177
630.	1-(1H-Inden-3-yl)naphthalene Benzene	LP-ET	559 ^a 397 ^a		†Triplet ET from xanthone and oxygen quenching; $\tau_T = 4.5 \mu\text{s}$	84B007
631.	Indole Benzene (297 K) Cyclohexane Cyclohexane Water (298 K) Water	PR LP/RA LP/ET LP LP/RA	430 430, 4000 430 ^b , 4700 ± 500 ~440 430, 3640		†Triplet ET to naphthalene and anthracene, oxygen quenching †Oxygen quenching, triplet quenching; ϵ relative to naphthalene in cyclohexane ($\epsilon_{414} = 24000 \text{ L mol}^{-1} \text{ cm}^{-1}$) and assuming $\Phi_T = 0.77$ for naphthalene and $\Phi_T = 0.51$ for indole; shoulder at 350 nm; $\tau_T = 16 \mu\text{s}$ †Triplet ET to anthracene; ϵ relative to anthracene in cyclohexane ($\epsilon_{422.5} = 64700 \text{ L mol}^{-1} \text{ cm}^{-1}$); ϵ corrected for 4% radical absorption at maximum; and for energy transfer via excited singlet states; $\tau_T = 16 \mu\text{s}$; $k_{et} = 9 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$ Spectrum is the difference between spectra at 50 ns and 4 μs ; $\tau_T = 11.6 \pm 0.5 \mu\text{s}$; pH 7.5 ϵ relative to indole in cyclohexane ($\epsilon_{430} = 4700 \text{ L mol}^{-1} \text{ cm}^{-1}$) scaled by halfwidths; assumes oscillator strength is independent of solvent	771021 777037 81E082 757163 81E082
632.	2-Iodoanthracene Cyclohexane Hexane (293 K)	LP LP	432 435 ~413		122 ps delay; rise time of $\sim 30 \times 10^{-3} \text{ ns}$ Delay 100 ps	81E309 82E303
633.	9-Iodoanthracene Cyclohexane	LP	432		122 ps delay; rise time of $\sim 50 \times 10^{-3} \text{ ns}$	81E309
634.	4-Iodobiphenyl Toluene	PR	400		†Oxygen quenching	80A235
635.	Iodo(methanol)(phthalocyaninato)rhodium(III) Acetonitrile/2-PrOH	LP/SD	640 ^b , 900 590 ^b , 22000		Solvent was 2.6 mol L ⁻¹ 2-PrOH	83F405

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
636.	1-Iodonaphthalene Toluene	PR	430		‡Triplet ET to benz[a]anthracene, oxygen quenching; $\tau_T = \sim 0.010 \mu\text{s}$	80A235
637.	2-Iodonaphthalene EPA (77 K)	FP	424 400			55E002
	Toluene	PR	430		‡Oxygen quenching	80A235
638.	β -Ionone Hexane	LP/SD	330, 85300 \pm 12800		ϵ assumes triplet does not absorb where singlet depletion is followed; $\tau_T = 0.16 \mu\text{s}$; $E_T = 207 \pm 12 \text{ kJ mol}^{-1}$	78E721
639.	2-Isopropylthioxanthene-9-one Benzene	FP	650-670		$\tau_T = 95 \mu\text{s}$	81A294
640.	Isopsoralen Benzene	PR/ET	450 ^b , 4700		‡Triplet ET to β -carotene; ϵ relative to biphenyl in benzene ($\epsilon_{367} = 27100 \text{ L mol}^{-1} \text{ cm}^{-1}$); "angelicin"; $k_{et} = 5.2 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	78E157
	Benzene	LP	405 ^a			78E157
	Water	LP	440 ^a		$\tau_T = 5.9 \mu\text{s}$	78E157
641.	Isoquinoline 1-BuOH/Isopentane (77 K)	FP/SD	419.0, 14900 \pm 1000 395 ^a , 7400 ^a		Glass was 3:7 1-BuOH to isopentane; oscillator strength = 0.11	707240
	3-MP (77 K)	MOD/SD	417, 8800 395 ^a 375 ^a 362 ^a 346 ^a		Shoulder at 250 nm	73B001
	EPA (77 K)	PS	233, 14000 488 459 421.6 400		Relative intensities (3:9:100:66)	54B001
	Isopentane/MCH (77 K)	PS	414.1 392		Relative intensities (2:1)	54B001
642.	Isoquinoline-N-oxide Cyclohexane	FP	418		$\tau_T = 3.6 \mu\text{s}$	727105
	EtOH	FP	396		$\tau_T = 7.7 \mu\text{s}$	727105
	Water	FP	389		$\tau_T = 2.0 \mu\text{s}$	727105
643.	Kynurenic acid EtOH	LP/ET	570, 29000		‡Triplet ET to naphthalene, oxygen quenching ($2 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$); ϵ relative to naphthalene in cyclohexane ($\epsilon_{415} = 24500 \text{ L mol}^{-1} \text{ cm}^{-1}$), assuming unit transfer efficiency and assuming ϵ independent of solvent (authors see naphthalene peak at 417.5 nm in EtOH); $k_{et} = 4 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$	79B086
	Water	LP	590		‡Triplet ET to naphthalene, oxygen quenching; $\tau_T = 10 \mu\text{s}$; pH 7; $E_T = 278 \text{ kJ mol}^{-1}$	79B086
644.	Lumichrome EtOH	LP/RA	560 ^a , 5200 ^a 450 ^a , 4000 ^a 370 ^a , 22600 ^a		ϵ relative to anthracene in cyclohexane ($\Phi_T = 0.71$, $\epsilon_{423} = 64700 \text{ L mol}^{-1} \text{ cm}^{-1}$) and using $\Phi_T = 0.61$ for lumichrome in ethanol; shoulder around 650 nm	777617
	Water	LP/RA	640, 5600 ^a 570, 4800 ^a 400 ^a , 10300 ^a 350 ^a , 9000 ^a 300 ^a , 15400 ^a		Lifetime measured at 640 nm; ϵ relative to anthracene in cyclohexane ($\Phi_T = 0.71$, $\epsilon_{423} = 64700 \text{ L mol}^{-1} \text{ cm}^{-1}$) and using $\Phi_T = 0.44$ for lumichrome in water; $\tau_T = 12 \mu\text{s}$; pH 2.2	777617

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
645.	Lumiflavine					
	Chloroform	FP/SD	396 ^a , 8130 ^a		$\tau_T = 30 \mu s$	68E100
	EtOH	LP/RA	640 ^a , 10100 ^a		ϵ relative to anthracene in cyclohexane ($\Phi_T = 0.71$, $\epsilon_{423} = 64700$ L mol ⁻¹ cm ⁻¹) and using $\Phi_T = 0.30$ for lumiflavine in ethanol; shoulder around 300 nm	777617
	Water	FP/TD	510 ^a , 5100 ^a 380 ^a , 12900 ^a 650, 5000		†Transient sensitive to oxygen and iodide ions; TD assumed, value for ϵ is a lower limit, upper limit is double this value; pH 6.5	66E087
	Water	FP/SD	640, 5300 ^a 376 ^a , 13100 ^a		†Triplet ET to acridine; $\tau_T = 90 \mu s$	68E100
	Water	MOD	542 ^a		Another transition with a maximum at shorter wavelengths than 500 nm was also present; $\tau_T = 320 \mu s$	70E295
	Water	FP/TD	690, 4400 370, 10700		Shoulder at $\sim 440^a$ nm; pK_a 9.8; delay 8 μs ; $\tau_T = 67 \mu s$; pH 7	757078
646.	Lumiflavine, conjugate monoacid					
	Water	FP/TD	670, 4800 430, 7000 366 ^a , 6400 ^a		pK_a 4.4; delay 8 μs ; $\tau_T = 29 \mu s$; pH 2	757078
	Water	LP/RA	650, 6900 420 ^a , 7000 ^a 350 ^a , 6500 ^a		lifetime measured at 670 nm; ϵ relative to anthracene in cyclohexane ($\Phi_T = 0.71$, $\epsilon_{423} = 64700$ L mol ⁻¹ cm ⁻¹) and using $\Phi_T = 0.42$ for lumiflavine in water; shoulder around 300 nm; $\tau_T = 20 \mu s$; pH 2.2	777617
647.	Lumiflavine, negative ion					
	Water	FP/TD	650, ~ 2000 350, ~ 11000		Shoulder at $\sim 440^a$ nm; delay 8 μs ; $\tau_T = 63 \mu s$; pH 13	757078
648.	<i>all-trans</i> -Lutein					
	Digitonin (293 K)	LP/TD	518, 40000		†Triplet ET from chlorophyll <i>a</i> , oxygen quenching (7×10^8 L mol ⁻¹ s ⁻¹), temperature independent half-life; aqueous micellar suspension; half-life = 8 ns	69N001
649.	<i>all-trans</i> -Lycopene					
	Hexane	FP-ET	534 ^a 527 520 ^a		†Triplet ET from anthracene; 520 nm was the most intense peak; $\tau_T = 43 \mu s$; $k_{et} = 1.7 \times 10^{10}$ L mol ⁻¹ s ⁻¹	69E209
	Hexane	PR/ET	525, 390000 \pm 20000 488 ^a ~ 460		†Triplet ET from naphthalene; ϵ relative to naphthalene in cyclohexane ($\epsilon_{415} = 24500$ L mol ⁻¹ cm ⁻¹), except authors used 412.5 nm as reference λ ; $\tau_T = 8 \mu s$; $k_{et} = 1.4 \times 10^{10}$ L mol ⁻¹ s ⁻¹	713035
	Hexane	PR/ET	525, 495000 \pm 74300		†Triplet ET from biphenyl; ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹), assuming authors used standard for cyclohexane; $\tau_T = 5.9 \pm 0.6 \mu s$; $k_{et} = (2.0 \pm 0.20) \times 10^{10}$ L mol ⁻¹ s ⁻¹	776412
650.	Lysyltryptophanyllysine					
	Water (293 K)	LP	450		Buffered; $\tau_T = 20 \mu s$; pH 7.0	81A232
651.	Magnesium(II) mesoporphyrin					
	EtOH (77 K)	FP	550 ^a 440 ^a		Relative intensities (1:7)	82E129
652.	Magnesium(II) octaethylporphyrin					
	MeOH	LP/ELT	440 ^b , 100000 430 ^a		†Oxygen and ethyl iodide quenching; ϵ relative to methyl viologen radical cation in MeOH ($\epsilon_{440} = 580$ L mol ⁻¹ cm ⁻¹)	83A102
653.	Magnesium(II) phthalocyanine					
	Dimethylacetamide-Water	LP/SD	490, 31000		Solvent was 7:3 dimethylacetamide to water	84A122
	Dioxane	FP	500 400			63B003
	Dioxane	FP	470		$\tau_T = 100 \mu s$	65A001

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Polyvinylbutyral (293 K)	LP	470			80E867
	Polyvinylbutyral (293 K)	LP	400			
	Polyvinylbutyral (293 K)	LP	676		Relative intensities (5:1); polymer film; $\tau_T = 10 \mu\text{s}$	82E129
	PrOH	FP/SD	610			
	Pyridine	FP	470, 33600			73E345
	Pyridine	FP	470			58R002
	Pyridine	FP	400			
	Pyridine	FP	500			63B003
	Pyridine	FP	400			
654.	Magnesium(II) tetraphenylporphyrin					
	CTAC	LP	840		†Oxygen quenching; aqueous micelles	83N073
			460			
	MCH	FP/TD	485, 72000		†Oxygen quenching; $\tau_T = 1400 \mu\text{s}$	81E271
			415, 27000			
	SHS/2-Pentanol	LP	840		Aqueous microemulsion, hexadecane in water; shoulder at 720 nm	83N073
			460			
655.	Malachite Green					
	PMMA (77 K)	PS	780			65B004
656.	Mercury(II) tetraphenylporphyrin					
	MCH	FP/TD	495, 86000		†Oxygen quenching; biexponential decay, reported value is shorter component; $\tau_T = 10 \mu\text{s}$	81E271
			420, 24000			
657.	Mesoporphyrin					
	Pyridine	FP	450		Isomer of mesoporphyrin unspecified	58R002
658.	Mesoporphyrin, dimethyl ester					
	Benzene	PR/ET	440 ^b , 32000		†Oxygen quenching ($1.4 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); ϵ relative to biphenyl in benzene ($\epsilon_{360} = 27100 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 220 \mu\text{s}$	80E200
			398 ^a			
	Benzene	LP/TD	440 ^b , 32000		†Oxygen quenching ($1.4 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); $\tau_T = 220 \mu\text{s}$	80E200
			398 ^a			
659.	1-(β -Methacryloxyethyl)-3,3-dimethyl-6'-nitrospiro(indoline-2,2'-[2H]benzopyran)					
	Acetonitrile	LP	430 ^a			83B117
	Dioxane	LP	440		†Oxygen quenching; $\tau_T = 1.0 \mu\text{s}$; rise time of $< 10 \text{ ns}$	82E041
	Toluene	LP	440			83B117
660.	8,11b-Methanocyclodeca[<i>cde</i>]naphthalene					
	Hexane	LP	490		†Oxygen quenching; $\tau_T = 0.20 \pm 0.05 \mu\text{s}$	84F068
			390			
661.	3-Methoxyacetophenone					
	Water/Acetonitrile	LP	390 ^a		Solvent was 4:1 water to acetonitrile; 200 ns delay; $\tau_T = 3.2 \mu\text{s}$; $E_T = 303 \text{ kJ mol}^{-1}$	82A082
662.	4-Methoxyacetophenone					
	Cyclohexane	LP/RA	360 ^a , 10800 ^a		ϵ relative to benzophenone in cyclohexane ($\epsilon_{333} = 7630 \text{ L mol}^{-1} \text{ cm}^{-1}$, assuming no solvent effect from benzene), and taking all the triplet yields to be unity	737198
	Toluene	LP	390			84E144
663.	3-Methoxy-7H-benz[<i>de</i>]anthracen-7-one					
	Benzene	LP	$\sim 505^a$		†Oxygen quenching ($1.3 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); half-life $= 2.4 \pm 0.2 \mu\text{s}$	757427
664.	4-Methoxybenzophenone					
	Acetonitrile/Water	LP/ET	680, 3400		Solvent was 9:1 acetonitrile to water; ϵ relative to 1-methylnaphthalene in acetonitrile/water ($\epsilon_{415} = 11200 \text{ L mol}^{-1} \text{ cm}^{-1}$)	84B033
			520, 5400			
			450, 4700			
			335, 5600			
	Water/Acetonitrile	LP	465 ^a		Solvent was 4:1 water to acetonitrile; 200 ns delay; $\tau_T = 7.2 \mu\text{s}$; $E_T = 288 \text{ kJ mol}^{-1}$	82A082
			450 ^a			

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
665.	9-[2-(4-Methoxycarbonylbenzoxy)ethyl]carbazole 2-MTHF	LP	500 ^a 420		Relative intensities (1:2); $\tau_T = 5 \mu\text{s}$; rise time of 10 ns	81E016
666.	9-[3-(4-Methoxycarbonylbenzoxy)propyl]carbazole 2-MTHF	LP	490 ^a 420		Relative intensities (1:2); $\tau_T = 5 \mu\text{s}$	81E016
667.	4-Methoxychalcone Heptane	LP	480 \pm 15		"3-(4-Methoxyphenyl)-1-phenyl-2-propen-1-one"; $\tau_T = 0.023 \mu\text{s}$	83E347
	MeOH	LP	525 \pm 15		$\tau_T = 0.029 \mu\text{s}$	83E347
668.	(R)-4-Methoxydinaphtho[2,1-d:1',2'-f][1,3,2]dioxaphosphepin 4-oxide 2-MTHF (96 K)	PS/KM	420 ^a , 11100 ^a 400 ^a , 9100 ^a		Compound "IV" in paper	83E383
669.	4-Methoxydinaphtho[2,1-d:1',2'-f][1,3,2]dioxaphosphepin 4-oxide MTHF (102 K)	PS	418 ^a 400 ^a		Relative intensities (5:4); solvent and temperature assumed; $\tau_T = 10^6 \mu\text{s}$	80E361
670.	8'-Methoxy-5-methyl-6'-nitrospiro[(5,6-dihydrophenanthridine-6,2)-(2H)chromene] Toluene	FP	640 ^a 510 ^a		\uparrow Triplet ET from fluorene; $E_T = 217\text{--}288 \text{ kJ mol}^{-1}$	767647
671.	2-Methoxy-2-methyl-1-phenyl-1-propanone Cyclohexane	LP	475 430 400		\uparrow Phosphorescence decay, oxygen quenching ($1.1 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); relative intensities (2:3:6); lifetime limited by reaction with solvent; $\tau_T = 0.12 \mu\text{s}$	80E642
672.	1-Methoxynaphthalene 3-MP (77 K)	PS	49. 462 441 417		441 nm peak was the most intense	74B004
	Cyclohexane	LP/ET	440, 7000 \pm 500		\uparrow Triplet ET from TMPD; ϵ relative to TMPD in cyclohexane ($\epsilon_{570} = 11900 \text{ L mol}^{-1} \text{ cm}^{-1}$)	757282
	EPA (77 K)	FP	490 465 440 420 395 370		440 nm was the most intense peak; $\tau_T = 2.3 \times 10^6 \mu\text{s}$	677485
	EtOH (293 K)	FP	436			68E098
	Isopentane/MCH/Et ₂ O (77 K)	PS	488 ^a 459 ^a 435 ^a 417 ^a		Glass was 6:1:0.7 isopentane to MCH to Et ₂ O; shoulders at 397 ^a nm; 435 nm was the most intense peak	71B003
	MeOH	FP	445 ^a			736218
673.	2-Methoxynaphthalene 3-MP (77 K)	PS	432 408 388 368 351.5		432 nm peak was the most intense	74B004
	Cyclohexane	LP/ET	435, 15000 \pm 800		\uparrow Triplet ET from TMPD; ϵ relative to TMPD in cyclohexane ($\epsilon_{570} = 11900 \text{ L mol}^{-1} \text{ cm}^{-1}$)	757282
	Heptane	FP	440 ^a		Lifetimes suggest solution not fully deaerated; $\tau_T = 50 \pm 5 \mu\text{s}$	77E663
	Isopentane/MCH/Et ₂ O (77 K)	PS	435 ^a 412 ^a		Glass was 6:1:0.7 isopentane to MCH to Et ₂ O; shoulders at 391 ^a and 375 ^a nm; 435 nm was the more intense peak	71B003

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
674.	4-Methoxy-1-naphthonitrile EPA (77 K)	FP	522.5 495 469 450 425		495 was the most intense peak; $\tau_T = 2.2 \times 10^6 \mu s$	677485
675.	<i>cis,cis</i> -1-(3'-Methoxy-5'-nitro-2'-oxo-3',5'-cyclohexadienyl)-3,4-dimethyl-1,3-pentadiene Toluene	LP/SD	520, ~9000 330, ~12000		†Oxygen quenching; ϵ 's are uncertain because of spectral overlaps with another triplet and a photoproduct	83F081
676.	<i>trans</i> -3-Methoxy-4'-nitrostilbene Glycerol triacetate (198 K)	LP	755 552 420		Relative intensities (1:2:2)	78B088
	Glycerol triacetate	LP	755 554 412		Relative intensities (1:2:2)	78B088
677.	<i>cis</i> -4-Methoxy-4'-nitrostilbene Benzene	LP	600 ^a 420 ^a		Relative intensities (3:2); identical spectrum derived from <i>trans</i> isomer	78B088
	Cyclohexane	LP	545 416		Relative intensities (9:10); identical spectrum derived from <i>trans</i> isomer	78B088
	DMF	LP	652 466		Relative intensities (2:1); identical spectrum derived from <i>trans</i> isomer	78B088
	MeOH	LP	680 ^a 440 ^a		Relative intensities (2:1); identical spectrum derived from <i>trans</i> isomer	78B088
678.	<i>trans</i> -4-Methoxy-4'-nitrostilbene 2,2-Dimethylbutane- /Pentane (77 K)	FP	650 ^a 600 ^a		Glass 3:8 pentane to 2,2-dimethylbutane	747022
	Benzene	LP	600 ^a 420 ^a		Relative intensities (3:2); identical spectrum derived from <i>cis</i> isomer	78B088
	Cyclohexane	LP	600		†Oxygen quenching in benzene; $\tau_T = 0.090 \mu s$	747022
	Cyclohexane	LP	580		Identical spectrum obtained from <i>cis</i> -isomer; oxygen quenching (5.3×10^9 L mol ⁻¹ s ⁻¹); $\tau_T = 0.077 \mu s$	777455
	Cyclohexane	LP	545 416		Relative intensities (9:10); identical spectrum derived from <i>cis</i> isomer	78B088
	DMF	LP	652 466		Relative intensities (2:1); identical spectrum derived from <i>cis</i> isomer	78B088
	EPA (77 K)	FP	645 612 450		Shoulders at 540 and 490 nm; 645 nm was the most intense peak; $\tau_T = 1.5 \times 10^6 \mu s$	677485
	EPA (88 K)	LP	610 526		610 nm peak was the more intense	78B088
	EPA	LP	600 526		600 nm peak was the more intense	78B088
	Glycerol triacetate	LP	~780 624 444		Most intense peak at 624 nm	78B088
	Glycerol triacetate (198 K)	LP	~780 625 460		Most intense peak at 625 nm	78B088
	MeOH	LP	~700 600		†Oxygen quenching in benzene; $\tau_T = 0.33 \mu s$	747022
	MeOH	LP	680 ^a 440 ^a		Relative intensities (2:1); identical spectrum derived from <i>cis</i> isomer	78B088
679.	4-Methoxyphenol EtOH (77 K)	MOD	444 ^a 392 ^a 337 ^a		†Phosphorescence decay; $\tau_T = (2.9 \pm 0.2) \times 10^6 \mu s$	707152

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
680.	2-[1-(4-Methoxyphenyl)-hydrazinyl-2-ylidene]-1-(4-nitrophenyl)-1,3-diketobutane					
	Hexane	LP	510		{Triplet ET to β -carotene and from biphenyl, oxygen quenching; half-life = 0.045 μ s; E_T = 190 \pm 10 kJ mol ⁻¹	80A051
681.	trans-1-(4-Methoxyphenyl)-2-nitroethylene					
	Benzene	LP	520 ^a		τ_T = 0.015 μ s	78E057
	Benzene/Acetonitrile	LP	525 ^a		Solvent was 8:2 benzene to acetonitrile by volume; τ_T = 0.019 μ s	78E057
	Cyclohexane	LP	450 ^a		τ_T = 0.0098 μ s	78E057
	EPA (77 K)	PS/KM	740 530			78E057
682.	4'-Methoxypropiophenone					
	Benzene	LP/ET	390, 12000		{Triplet ET to naphthalene; ϵ relative to naphthalene in benzene (ϵ_{425} = 13000 L mol ⁻¹ cm ⁻¹)	81A174
	Benzene	LP	382 ^a			84E018
683.	5-Methoxypsoralen					
	Benzene	LP	455 ^a		"Bergapten"	78E157
	Benzene	PR/ET	450 ^b , 10200		{Triplet ET to β -carotene; ϵ relative to biphenyl in benzene (ϵ_{367} = 27100 L mol ⁻¹ cm ⁻¹); k_{et} = 4.6 \times 10 ⁹ L mol ⁻¹ s ⁻¹	78F157
	EtOH	LP/ET	461 ^a , 11100 ^a 450 ^b , 9500		ϵ relative to retinol, but actual reference ϵ used was not specified; τ_T = 4.2 μ s	83E324
684.	8-Methoxypsoralen					
	Benzene	LP	470 ^a 390 ^a		"Xanthotoxin"; τ_T = 1.1 μ s	78E157
	Benzene	FP-ET	600 480		{Triplet ET from acetophenone; relative intensities (1:2)	79R026
	EtOH	LP/ET	370, 24000		{Triplet ET to retinol; ϵ relative to retinol in hexane (ϵ_{405} = 80000 L mol ⁻¹ cm ⁻¹) assuming ϵ independent of solvent; τ_T = 10 μ s; k_{et} = 5.3 \times 10 ⁹ L mol ⁻¹ s ⁻¹	79E678
	EtOH	LP/ET	370 ^b , 21000		ϵ relative to retinol, but actual reference ϵ used was not specified; τ_T = 2.3 μ s	83E324
	Glycerol	FP	600 480		{Oxygen quenching; there was a long-lived species that was attributed to a secondary product; there was another maximum <340 nm	757485
	Glycerol/Water	FP-ET	630 480 360		Solvent contains 57% (wt) glycerol; relative intensities (5:4:10)	79R026
	MeOH	LP	600 370			79B042
	Water	LP	>580 ^a 350 ^a			78E157
	Water	LP	360		{Oxygen quenching (3.3 \times 10 ⁹ L mol ⁻¹ s ⁻¹); radical contributions subtracted and no infrared peak observed; τ_T = 1.6 μ s	79A114
	Water	LP/SD	370 ^b , 16500		τ_T = 2.4 μ s	83E324
685.	trans-4-Methoxystilbene					
	EPA (77 K)	FP	400 382.5		Shoulder at 365 nm; 400 nm was the more intense peak; τ_T = 2.35 \times 10 ⁴ μ s	677485
686.	8-Methoxy-2,2,3-trimethyl-6-nitro-2H-chromene					
	Toluene	LP/SD	520, ~8000 330, ~10000		{Oxygen quenching; ϵ 's are uncertain because of spectral overlaps with another triplet and a photoproduct	83F081
687.	6-Methoxy-1,3,3-trimethyl-6'-nitrospiro-(2,2'-indolin[2H-1]benzopyran)					
	Toluene (298 K)	FP	470		{Oxygen quenching, activationless decay; assigned to $\pi \rightarrow \pi^*$ transition in uncolored form	67B003

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
688.	7'-Methoxy-1,3,3-trimethyl-6'-nitrospiro-(2,2'-indolin[2H-1]benzopyran) Toluene (298 K)	FP	445		†Oxygen quenching, activationless decay; assigned to $\pi \rightarrow \pi^*$ transition in uncolored form	67B003
689.	Methyl Violet PMMA (193 K)	PS	630			59B001
	PMMA (77 K)	PS	1290 1200 1100 770 665 640 620 1112 ^a 613 ^a			65B004
	PMMA (77 K)	PS			Relative intensities (1:5); shoulder at 524 ^a nm; solvent contains 2-chloroethanol	69B003
690.	2-Methylaceanthren-1-one EPA (77 K)	FP	455		†Phosphorescence decay; $\tau_T = 5900 \mu s$	82E338
691.	3-Methylacetophenone 2-PrOH	LP	339 ^a		†Quenching with oxygen, piperylene, and naphthalene	737473
	Cyclohexane	LP	332 ^a		†Quenching with oxygen, piperylene, and naphthalene	737473
692.	4-Methylacetophenone Cyclohexane	LP/RA	331 ^a , 11400 ^a		ϵ relative to benzophenone in cyclohexane ($\epsilon_{333} = 7630 \text{ L mol}^{-1} \text{ cm}^{-1}$, assuming no solvent effect from benzene), and taking all the triplet yields to be unity	737198
693.	9-Methylacridine EtOH (298 K)	FP/?	521 ^a , 3000 ^a 452 ^a , 30000 ^a 408 ^a , 10000 ^a 330 ^a , 10000 ^a			697225
694.	10-Methyl-9(10H)-acridinethione Acetonitrile (296 K)	LP/SD	520 \pm 3, 9000 \pm 900		†Oxygen ($4.5 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$) and nitroxide radical quenching and triplet ET to anthracene; $\tau_T = 1.6 \pm 0.3 \mu s$	84E342
	Benzene (296 K)	LP/ET&SD	520 \pm 3, 9300 \pm 900		†Oxygen ($5.2 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$) and nitroxide radical quenching and triplet ET to anthracene; ϵ relative to 1,6-diphenyl-1,3,5-hexatriene in benzene ($\epsilon_{426} = 104000 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 2.5 \pm 0.5 \mu s$	84E342
	Cyclohexane (296 K)	LP/ET&SD	520 \pm 3, 8800 \pm 900		†Oxygen ($4.8 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$) and nitroxide radical quenching and triplet ET to anthracene; ϵ relative to 1,6-diphenyl-1,3,5-hexatriene in cyclohexane ($\epsilon_{416} = 114000 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 2.6 \pm 0.5 \mu s$	84E342
	MeOH (296 K)	LP/ET&SD	520 \pm 3, 8500 \pm 900		†Oxygen ($4.5 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$) and nitroxide radical quenching and triplet ET to anthracene; ϵ relative to 1,6-diphenyl-1,3,5-hexatriene in MeOH ($\epsilon_{410} = 121000 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 2.3 \pm 0.5 \mu s$	84E342
695.	4-(Methylamino)benzoic acid EtOH (93 K)	PS	435 ^a 405 ^a 363 ^a			66B001
696.	N-Methyl-1-anilinonaphthalene MCH (296 K)	LP	602 ^a 420 ^a		$\tau_T = 2000 \mu s$	82E527
697.	N-Methyl-2-anilinonaphthalene MCH (153 K)	FP	500		†Phosphorescence decay; lifetime = 120 μs at 294 K	82E527

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
698.	<i>N</i> -Methyl-2-anilino-6-naphthalenesulfonate ion EtOH (293 K)	FP	500		$\tau_T = 200 \mu s$	82E527
699.	2-Methylanthracene EtOH	LP	427 ^a 405 ^a			706157
	EtOH/MeOH (113 K)	FP/TD	901 ^a , 195 ^a 854 ^a , 150 ^a 814 ^a , 100 ^a 799 ^a , 90 ^a 790 ^a , 60 ^a 775 ^a , 110 ^a 762 ^a , 90 ^a 736 ^a , 80 ^a 724 ^a , 90 ^a 714 ^a , 90 ^a 702 ^a , 100 ^a 693 ^a , 80 ^a 658 ^a , 70 ^a 569 ^a , 150 ^a 522 ^a , 1200 ^a 481 ^a , 1600 ^a 431 ^a , 73000 ^a 407 ^a , 22300 ^a 384 ^a , 3900 ^a 366 ^a , 1700 ^a 317 ^a , 2300 ^a 305 ^a , 2200 ^a 263 ^a , 68300 ^a	†Phosphorescence decay; shoulders at 687 ^a , 672 ^a , 648 ^a , 634 ^a , 532 ^a , 328 ^a , and 272 ^a nm	72B001	
	PMMA	FP	901 ^a 855 ^a 803 ^a 770 ^a 430	‡Lifetime same as 430 nm triplet band; if $\epsilon_{430} \sim 50000$ L mol ⁻¹ cm ⁻¹ , then $\epsilon \sim 700$ L mol ⁻¹ cm ⁻¹ for near infrared band; $\tau_T = (2.1 \pm 1) \times 10^4 \mu s$	66E086	
700.	9-Methylanthracene Benzene	PR/ET	430, 64200	ϵ relative to benzophenone in benzene ($\epsilon_{532.5} = 10300$ L mol ⁻¹ cm ⁻¹)	690087	
	Benzene	PR/ET	430, 51300	ϵ relative to naphthalene in benzene ($\epsilon_{\max} = 17500$ L mol ⁻¹ cm ⁻¹)	690087	
	Cyclohexane	PR	425		690087	
	Dioxane	PR	425		690087	
	EtOH	LP	425 ^a 405 ^a		706157	
	EtOH	FP/ET	440 ^b , 2800 426 ^b , 37400 424, 43000 420 ^b , 24000	Triplet ET from eosin and proflavine; ϵ relative to eosin in EtOH ($\epsilon_{380} = 9400$ L mol ⁻¹ cm ⁻¹) and proflavine in EtOH ($\epsilon_{550} = 11000$ L mol ⁻¹ cm ⁻¹)	716235	
	EtOH	FP-ET/RA	694 ^b , 47	ϵ relative to 9-methylanthracene in EtOH ($\epsilon_{424} = 43000$ L mol ⁻¹ cm ⁻¹)	78E019	
	Liquid paraffin	FP	428		65F031	
701.	9-Methylanthracene- <i>d</i> ₁₂ EtOH	LP	424 ^a		706157	
702.	Methyl benzoate Cyclohexane (290 K)	FP	333 ^a	10 μs delay	767556	
703.	2-Methylbenzoic acid Isopentane/MCH (77 K)	FP	325 ^a 310 ^a	†Phosphorescence decay; solvent was 5:1 isopentane to MCH by volume; dimeric species; shoulder at 350 ^a nm; $\tau_T = 1.16 \times 10^6 \mu s$	776222	

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
704.	3-Methylbenzoic acid Isopentane/MCH (77 K)	FP	325 ^a 310 ^a		†Phosphorescence decay; solvent was 5:1 isopentane to MCH by volume; dimeric species; $\tau_T = 3.95 \times 10^6$ μ s	776222
705.	4-Methylbenzoic acid Isopentane/MCH (77 K)	FP	330 ^a 315 ^a		†Phosphorescence decay; solvent was 5:1 isopentane to MCH by volume; dimeric species; $\tau_T = 3.45 \times 10^6$ μ s	776222
706.	2-Methylbenzophenone EtOH	LP	520		$\tau_T = 0.0028$ μ s	84B003
707.	4-Methylbenzophenone Acetonitrile/Water	LP/ET	525, 6600 315, 11700		Solvent was 9:1 acetonitrile to water; ϵ relative to 1-methylnaphthalene in acetonitrile/water ($\epsilon_{415} = 11200$ L mol ⁻¹ cm ⁻¹)	84B033
708.	Methyl-1,4-benzoquinone Water	LP	460 410		$\tau_T = 0.37$ μ s	80B112
709.	(2'-Methylbenzoyl)amino-2- Δ^2 -thiazoline Cyclohexane (293 K)	FP/RA	550, 7400		†Triplet ET to biacetyl, oxygen quenching; ϵ relative to naphthalene in cyclohexane ($\epsilon_{414} = 24500$ L mol ⁻¹ cm ⁻¹); $\tau_T = 450$ μ s	82E373
710.	(4'-Methylbenzoyl)amino-2- Δ^2 -thiazoline Cyclohexane (293 K)	FP/RA	550, 9600		†Triplet energy transfer to biacetyl, oxygen quenching; ϵ relative to naphthalene in cyclohexane ($\epsilon_{414} = 24500$ L mol ⁻¹ cm ⁻¹); $\tau_T = 2000$ μ s	82E373
711.	Methyl benzoylformate Benzene (298 K)	LP	446 ^a		Shoulders at 431 ^a and 400 ^a nm; $\tau_T = 1$ μ s	84A030
712.	Methylchlorophyllide <i>a</i> Acetone	FP	520 470			65R023
	Benzene	FP	520 470			65R023
	EtOH	FP	520 470			65R023
713.	1-Methyl-4-[4-cyanostyryl]pyridinium EtOH (103 K)	LP	415		Lifetime constant below 113 K; $\tau_T = \sim 500$ μ s	83E509
714.	10-Methyl-5-deazaalloxazine-3-propanesulfonate ion Water	LP	660 ^a 516 ^a		Delay 200 ns; half-life = 7.5 μ s	82B043
715.	3-Methyl-5-deazalumiflavine Water	FP/TD	700 ^b , 3600 \pm 200 684 ^a , 3700 ^a 501 ^a , 8100 ^a		pH 9.2	81A434
716.	1-Methyldeoxybenzoin Benzene	LP	325		†Oxygen quenching ($> 10^9$ L mol ⁻¹ s ⁻¹); $\tau_T = 0.09$ μ s	79A028
717.	Methyl 4-dimethylaminobenzoate EtOH (93 K)	PS	435 ^a 412 ^a 391 ^a 357 ^a		357 nm was the most intense peak	66B001

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
718.	<i>N</i> -Methyldiphenylamine EPA (118 K)	FP/COM	540, 26000 ± 3000		‡Phosphorescence decay; ϵ based on quantitative decay of triplet to <i>N</i> -methyl-4a,4b-dihydrocarbazole whose room temperature $\epsilon_{610} = 21000 \pm 2000$ L mol ⁻¹ cm ⁻¹ in MCH	737125
719.	Methylene Blue cation Acetonitrile	FP	870 ^a 750 ^a 520 ^a 422 ^a		$\tau_T = 150$ μ s	77A203
	Acetonitrile	LP	870 420		1 μ s delay; semioxidized cation radical ($\lambda_{\max} = 520$ nm) also observed; half-life = 12.5 μ s	81A347
	Acetonitrile	LP	860 ^a 780 ^a		‡Oxygen quenching (1.7×10^9 L mol ⁻¹ s ⁻¹); relative intensities (7:4); 1 μ s delay; $\tau_T = 14$ μ s	84E216
	Acetonitrile/Water	LP	840 ^a 750 ^a		‡Oxygen quenching; relative intensities (13:6); 1 μ s delay; solvent was 1:1 acetonitrile to water by volume	84E216
	CTAB	LP	420		Aqueous micelles	83N062
	EtOH	LP	847 ^a 775 ^a 418 ^a		847 nm was the most intense peak; $\tau_T = 450$ μ s	756162
	EtOH	LP/SD	860 780 420 410 ^b , 20000			767803
	EtOH	LP/SD	860, 27000 420, 14000		ϵ assumes triplet does not absorb at 658 nm where SD was followed; solvent and temperature assumed	78E394
	EtOH/Water	LP/TD	840, 20000 420, 9000		Solvent was "30% EtOH"; no acid added; $\tau_T = 36$ μ s; pH 8.2	79B104
	MeOH	LP/SD	420, 11000		‡Oxygen quenching; ϵ estimated assuming no triplet absorption at 650 nm; $\tau_T = 5.9$ μ s	767574
	MeOH	LP/SD	410, 22000		‡Oxygen quenching (2.2×10^9 L mol ⁻¹ s ⁻¹); ϵ assumes triplet does not absorb between 560 and 570 nm, where SD is monitored; decay followed at 420 nm; less intense peak at 525 nm assigned to triplet excimer; $\tau_T = 11 \pm 2$ μ s	79A230
	Water	FP	420 280		$pK_b = 7.2$; pH Neutral; half-life = 130 μ s	64B003
	Water	FP	800 415		$pK_b = 7.3$; delay 20 μ s; $\tau_T = \sim 30$ μ s; pH Neutral	64E013
	Water	LP/SD	820 420, 9000		‡Oxygen quenching; extinction coefficient was a lower limit; half-life = 30 μ s	67E107
	Water	FP/SD	825, 18000		$pK_a = 7.5 \pm 0.1$	67C001
	Water	LP	830 420		1 μ s delay; semioxidized cation radical ($\lambda_{\max} = 520$ nm) also observed; pH 8.2; half-life = 25 μ s	81A347
720.	Methylene Blue cation, conjugate monoacid					
	Acetonitrile/Water	LP/TD	470 ^a , 4500 ^a 428 ^a , 6200 ^a 372 ^a , 14000 ^a		Solvent was 1:1 acetonitrile to water by volume; $\tau_T = 4.0$ μ s; pH Acidic	777062
	Acetonitrile/Water	LP/TD	710-670, >16200 370, 15300		Solvent was 1:1 acetonitrile to water by volume; 0.01N HCl added; pH Acidic	79B104
	EtOH/Sulfuric acid	LP/RA	700 ^b , 55000		ϵ relative to unprotonated form ($\epsilon_{410} = 20000$ L mol ⁻¹ cm ⁻¹); 500 ns delay; pH Acidic	767803
	EtOH/Water	LP/TD	703, 18700 680, 19000 370, 13200		Solvent was "30% EtOH"; 0.01N H ₂ SO ₄ added; $pK_a = 7.2$; $\tau_T = 4.5$ μ s; pH 2.0	79B104
	Hydrochloric acid	FP	375		$pK_a = 6.7$; delay 20 μ s; pH 2	64E013
	Sulfuric acid	FP	375		$pK_a = 6.8$; pH 1	64B003
	Water	LP/TD	490 ^a , 3300 ^a 430 ^a , 3300 ^a 372 ^a , 7000 ^a		$\tau_T = 4.5$ μ s; pH Acidic	777062
	Water	LP/TD	690, 14500 370, 12500		0.01N H ₂ SO ₄ added; pH Acidic	79B104

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
721.	Methylene Blue, dimer Water	LP	~550		†Oxygen quenching; half-life = ~2 μ s	67E107
722.	Methylene Green cation Water	FP	794 ^a 761 ^a 518 ^a 412 ^a		SD region between 540 and 720 nm was not measured; pH 9	707364
723.	4,4'-Methylenebis(1,1'-biphenyl) Alcohol/Ether (77 K)	MOD	380		Glass was 2:1 alcohol to ether	76E682
724.	α -Methyl- β -ethylstyrene Cyclohexane	LP/RA-ET	325 ^b , 4000		ϵ relative to naphthalene in cyclohexane ($\epsilon_{413} = 24000$ L mol ⁻¹ cm ⁻¹), but in ref. cited ($\epsilon_{415} = 24500$ L mol ⁻¹ cm ⁻¹); triplet ET from thioxanthone; $\tau_T = \sim 0.038$ μ s	82E181
725.	5-Methyl-endo-cis-fluorenacene MCH/Isopentane (77 K)	PS	458 ^a		Glass was 4:1 MCH to isopentane; shoulder at 430 nm	74B006
726.	7-Methylguanine Water	LP-ET	385		†Triplet ET from acetone; pH 7.1; $k_{et} = 7.5 \times 10^9$ L mol ⁻¹ s ⁻¹	82B045
727.	1-Methylindazole MeOH	FP	420		†Oxygen quenching, piperylene quenching, and ESR; $\tau_T = 53$ μ s	757167
728.	2-Methylindazole MeOH	FP	405		†Oxygen quenching, piperylene quenching, and ESR; $\tau_T = 26$ μ s	757167
729.	1-Methylindole Benzene (293 K)	LP-ET/ET	460, 5000 \pm 1500		†Triplet ET from xanthone, from acetophenone and to naphthalene, oxygen quenching; ϵ relative to naphthalene in benzene ($\epsilon_{425} = 13200$ L mol ⁻¹ cm ⁻¹); $\tau_T = 2$ μ s	771021
	Benzene (297 K)	PR	460		†Triplet ET to naphthalene and anthracene, oxygen quenching; $\tau_T = 1.8$ μ s	771021
	Water	FP	460		†Oxygen quenching; $\tau_T = 29$ μ s; pH basic	83A213
730.	2-Methylindole Water	LP-ET	435		†Triplet ET from acetone; pH 7.1	82B045
731.	3-Methylindole Benzene (297 K)	PR	450		†Triplet ET to naphthalene and anthracene, oxygen quenching	771021
732.	1-Methylisoquinoline-N-oxide Cyclohexane	FP	422		$\tau_T = 5.9$ μ s	727105
	EtOH	FP	418 395		†Oxygen quenching and triplet ET to anthracene; $\tau_T = 7.7$ μ s	727105
	Water	FP	388		$\tau_T = 7.7$ μ s	727105
733.	1-Methylnaphthalene 1-Methylnaphthalene	PR/RF	420, 6350		Liquid; ϵ relative to 1-methylnaphthalene in cyclohexane ($\epsilon_{420} = 14200$ L mol ⁻¹ cm ⁻¹)	720243
	1-Methylnaphthalene	FP-ET	422 ^a 397 ^a		†Triplet ET from benzophenone; 30 μ s delay; evidence for triplet excimer at long wavelengths presented; relative intensities (3:2); 30 μ s delay; $\tau_T = 25$ μ s	767159
	Acetonitrile/Water	LP/ET	415, 11200		Solvent was 9:1 acetonitrile to water; ϵ relative to benzophenone in acetonitrile ($\epsilon_{520} = 6500$ L mol ⁻¹ cm ⁻¹)	84B033

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Benzene	LP-ET	425		\uparrow Triplet ET from benzophenone; $k_{\text{et}} = 9.4 \times 10^5$ L mol ⁻¹ s ⁻¹	84E092
	Cyclohexane (77 K)	PS	415.5		Relative intensities (10:6:1)	69E211
			393			
	Cyclohexane	PR/ET	370			
			420, 14200		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{542.5} = 3220$ L mol ⁻¹ cm ⁻¹), assuming ϵ is the same as $\epsilon_{537.5}$ for ketyl radical in water	720243
	Cyclohexane	LP/ET	420, 22400 \pm 3000		\uparrow Triplet ET from TMPD; ϵ relative to TMPD in cyclohexane ($\epsilon_{570} = 11900$ L mol ⁻¹ cm ⁻¹)	757282
	EtOH/MeOH (113 K)	FP/TD	581 ^a , 100 ^a		Glass was 3:1 EtOH to MeOH; shoulders at 629 ^a and 469 ^a nm	717460
			562 ^a , 70 ^a			
			538 ^a , 150 ^a			
			503 ^a , 125 ^a			
			485 ^a , 230 ^a			
			457 ^a , 480 ^a			
	EtOH/MeOH (113 K)	FP/TD	628 ^a , 10 ^a		\uparrow Phosphorescence decay; shoulders at 561 ^a , 469 ^a , 408 ^a , 367 ^a , 349 ^a , 285 ^a , and 236 ^a nm	72B001
			578 ^a , 100 ^a			
			535 ^a , 160 ^a			
			500 ^a , 120 ^a			
			484 ^a , 240 ^a			
			457 ^a , 480 ^a			
			419 ^a , 21900 ^a			
			396 ^a , 11300 ^a			
			375 ^a , 2500 ^a			
			356 ^a , 730 ^a			
			239 ^a , 49200 ^a			
	Hexane	FP-ET	418 ^a		\uparrow Triplet ET from benzophenone; 30 μ s delay; relative intensities (3:2:1)	767159
			395 ^a			
			383 ^a			
	Isooctane	FP	425 ^a		Shoulder at 360 nm	78B089
			397 ^a			
			375 ^a			
	Liquid paraffin	FP	422		Solvent viscosity was 0.03 N-s/m ² ; relative intensities (100:60:31)	58E001
			399			
			380			
	PMMA (294 K)	FP	625 ^a		Shoulder at 568 ^a nm; $\tau_T = (1.10 \pm 0.05) \times 10^6$ μ s	69B004
734.	2-Methylnaphthalene					
	Cyclohexane	LP/ET	420, 25100 \pm 3000		\uparrow Triplet ET from TMPD; ϵ relative to TMPD in cyclohexane ($\epsilon_{570} = 11900$ L mol ⁻¹ cm ⁻¹)	757282
	EtOH/MeOH (113 K)	FP/TD	575 ^a , 100 ^a		Glass was 3:1 EtOH to MeOH; shoulder at 633 ^a nm	717460
			529 ^a , 170 ^a			
			490 ^a , 200 ^a			
			461 ^a , 280 ^a			
	EtOH/MeOH (113 K)	FP/TD	631 ^a , 20 ^a		\uparrow Phosphorescence decay; shoulders at 402 ^a , 263 ^a , 234 ^a , and 231 ^a nm	72B001
			578 ^a , 100 ^a			
			532 ^a , 160 ^a			
			492 ^a , 200 ^a			
			462 ^a , 290 ^a			
			415 ^a , 25400 ^a			
			393 ^a , 12800 ^a			
			381 ^a , 1800 ^a			
			372 ^a , 3400 ^a			
			360 ^a , 600 ^a			
			353 ^a , 700 ^a			
			344 ^a , 350 ^a			
			238 ^a , 25500 ^a			
	PMMA (294 K)	FP			Shoulder at 575 ^a nm; $\tau_T = (1.30 \pm 0.05) \times 10^6$ μ s	69B004
735.	1-Methylnaphthalene/Pyromellitic dianhydride					
	Et ₂ O/Isopentane (77 K)	FP	600 ^a		\uparrow Phosphorescence decay; glass was 1:1 Et ₂ O to isopentane; shoulder at 645 ^a nm; $\tau_T = 9.8 \times 10^4$ μ s	747587
			420 ^a			

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
736.	Methyl 2-naphthoate Cyclohexane (290 K)	FP	412 ^a 401 ^a		35 μ s delay; $\tau_T = 66 \mu$ s	767556
737.	2-Methyl-1,4-naphthoquinone Water	LP/ELT	400 ^b , 6600		†Oxygen quenching and triplet ET to β -carotene; ϵ relative to 2-methylnaphthosemiquinone ion in water ($\epsilon_{390} = 11000$ L mol ⁻¹ cm ⁻¹); pH 7.0	83E311
738.	1-Methyl-1-(1-naphthyl)ethylene Benzene	LP-ET	439 ^a		†Triplet ET from xanthone and oxygen quenching; $\tau_T = 0.55 \mu$ s	84B007
739.	1-Methyl-1-(2-naphthyl)ethylene Benzene	LP-ET	599 ^a 415 ^a 388 ^a		†Triplet ET from xanthone and oxygen quenching (5×10^9 L mol ⁻¹ s ⁻¹); 415 nm peak was the most intense; $\tau_T = 0.13 \mu$ s	84B090
740.	Methyl-1-naphthylphenylgermane Cyclohexane	LP	430 ^a 410 ^a		430 nm peak was the more intense; delay 0.8 μ s	83A218
741.	5-Methyl-6'-nitrospiro[(5,6-dihydrophenanthridine-6,2)-(2H)chromene] Toluene	FP	670 ^a 470 ^a		†Triplet ET from fluorene; $\tau_T = 300 \mu$ s; $E_T = 217$ -288 kJ mol ⁻¹	767647
742.	1-Methyl-4-[4-nitrostyryl]pyridinium EtOH (298 K)	LP	450		†Oxygen quenching and triplet ET from acetone and biacetyl; $\tau_T = 0.055 \mu$ s	83E509
	EtOH (103 K)	LP	450		Lifetime constant below 108 K; $\tau_T = 3000 \mu$ s	83E509
743.	5-Methylphenazinium methyl sulfate Water	FP	525		†Oxygen quenching; $\tau_T = 50 \pm 5 \mu$ s; pH 3	80A196
744.	N-Methylphenothiazine Hexane	LP/ET	465, 23000 \pm 300		ϵ relative to naphthalene in cyclohexane ($\epsilon_{413} = 24500$ L mol ⁻¹ cm ⁻¹) assuming unit transfer efficiency and assuming ϵ independent of solvent	79N005
	SDS	LP/RA	465, 23000		Aqueous micelle; ϵ relative to N-methylphenothiazine in hexane ($\epsilon_{465} = 23000$ L mol ⁻¹ cm ⁻¹)	79N005
	Water/EtOH	LP	465		Solvent was 2:1 water to EtOH	82A297
745.	N-Methylphthalimide EtOH	FP	590 350		†Oxygen quenching, diene quenching; maxima assumed from text; $\tau_T = 43 \mu$ s	79A147
746.	4-(2-Methylpropionyloxy)benzophenone Benzene	LP	530		†Triplet ET to naphthalene	84P005
747.	Methyl pyropheophorbide <i>a</i> Toluene	LP	\sim 460			82B094
748.	<i>all-trans</i> -Methyl retinoate Benzene	PR/ET	435, 89000		ϵ relative to biphenyl in benzene ($\epsilon_{367.5} = 27100$ L mol ⁻¹ cm ⁻¹); $\tau_T = 17 \mu$ s; $k_{et} = 8.4 \times 10^9$ L mol ⁻¹ s ⁻¹	80E137
749.	2-Methylthioxanthene-9-one 1,2-Dichloroethane	FP	650-670		$\tau_T = 20 \mu$ s	81A294
	Acetonitrile	FP	630		†Oxygen quenching (5.5×10^9 L mol ⁻¹ s ⁻¹); $\tau_T = 40 \mu$ s	81A294
	Benzene	FP	655 ^a		†Oxygen quenching (5.5×10^9 L mol ⁻¹ s ⁻¹); $\tau_T = 95 \mu$ s	81A294
	Cyclohexane	FP	650-670		$\tau_T = 10 \mu$ s	81A294
	EtOH	FP	605		$\tau_T = 60 \mu$ s	81A294
	MeOH	FP	605		$\tau_T = 68 \mu$ s	81A294

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
750.	(E,E,E)-5-Methyl-7-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2,4,6-heptatrienal					
	Acetonitrile	LP/SD	440, 51000		"all-trans-C ₁₇ aldehyde"; ϵ is upper limit; $\tau_T = 10.6 \mu s$	79E546
	Acetonitrile	LP/RF	440, 61000		ϵ relative to compound in cyclohexane ($\epsilon_{410} = 63000$ L mol ⁻¹ cm ⁻¹) assuming oscillator strength independent of solvent; $\tau_T = 10.6 \mu s$	79E546
	Benzene	PR/ET	430, 63000		ϵ relative to biphenyl in benzene ($\epsilon_{367.5} = 27100$ L mol ⁻¹ cm ⁻¹); $\tau_T = 9.1 \mu s$	79E546
	Benzene	PR/RF	430, 71000		ϵ relative to compound in cyclohexane ($\epsilon_{410} = 63000$ L mol ⁻¹ cm ⁻¹) assuming oscillator strength independent of solvent; $\tau_T = 9.1 \mu s$	79E546
	Cyclohexane	PR/ET	410, 63000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361.3} = 42800$ L mol ⁻¹ cm ⁻¹); $\tau_T = 6.2 \mu s$	79E546
	MeOH	LP/SD	440, 51000		ϵ is upper limit; $\tau_T = 10.9 \mu s$	79E546
	MeOH	LP/RF	440, 57000		ϵ relative to compound in cyclohexane ($\epsilon_{410} = 63000$ L mol ⁻¹ cm ⁻¹) assuming oscillator strength independent of solvent; $\tau_T = 10.9 \mu s$	79E546
751.	(E,E,E)-6-Methyl-8-(2,6,6-trimethyl-1-cyclohexen-1-yl)-3,5,7-octatrien-2-one					
	Hexane	LP/SD	405-410, 52000		ϵ assumed to be at λ_{\max} ; "all-trans-C ₁₈ ketone"	84E036
752.	(E,E,E)-6-Methyl-8-(2,6,6-trimethyl-1-cyclohexen-1-yl)-3,5,7-octatrien-2-one					
	Hexane	LP/SD	405-410, 66000		ϵ assumed to be at λ_{\max} ; "7-cis-C ₁₈ ketone"	84E036
	Hexane	PR/ET	405-410, 37000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹), assuming authors used standard for cyclohexane; ϵ assumed to be at λ_{\max}	84E036
753.	(E,Z,E)-6-Methyl-8-(2,6,6-trimethyl-1-cyclohexen-1-yl)-3,5,7-octatrien-2-one					
	Hexane	LP/SD	405-410, 54000		ϵ assumed to be at λ_{\max} ; "9-cis-C ₁₈ ketone"	84E036
	Hexane	PR/ET	405-410, 23000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹), assuming authors used standard for cyclohexane; ϵ assumed to be at λ_{\max}	84E036
754.	(E,Z,Z)-6-Methyl-8-(2,6,6-trimethyl-1-cyclohexen-1-yl)-3,5,7-octatrien-2-one					
	Hexane	LP/SD	405-410, 23000		ϵ assumed to be at λ_{\max} ; "7,9-dicis-C ₁₈ ketone"	84E036
	Hexane	PR/ET	405-410, 42000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹), assuming authors used standard for cyclohexane; ϵ assumed to be at λ_{\max}	84E036
755.	(E,E)-3-Methyl-5-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2,4-pentadienal					
	Hexane	LP/SD	385, 38500 ± 5800		ϵ assumes triplet does not absorb where singlet depletion is followed; "all-trans-C ₁₅ aldehyde"; $\tau_T = 0.1 \mu s$; $E_T = 188 \pm 12$ kJ mol ⁻¹	78E721
	Hexane	PR/ET	380-390, 20000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹), assuming authors used standard for cyclohexane; ϵ assumed to be at λ_{\max}	84E036
	Hexane	LP/SD	380-390, 41000		ϵ assumed to be at λ_{\max}	84E036
	MeOH	LP/RF	400, 34000		ϵ relative to compound in hexane ($\epsilon_{385} = 38500$ L mol ⁻¹ cm ⁻¹) assuming oscillator strength independent of solvent; $\tau_T = 0.19 \mu s$	79E546
756.	(E,Z)-3-Methyl-5-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2,4-pentadienal					
	Hexane	LP/SD	380-390, 17000		ϵ assumed to be at λ_{\max} ; "7-cis-C ₁₅ aldehyde"	84E036
	Hexane	PR/ET	380-390, 14000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹), assuming authors used standard for cyclohexane; ϵ assumed to be at λ_{\max}	84E036
757.	(Z,E)-3-Methyl-5-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2,4-pentadienal					
	Hexane	PR/ET	380-390, 28000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹), assuming authors used standard for cyclohexane; ϵ assumed to be at λ_{\max}	84E036
	Hexane	LP/SD	380-390, 14000		ϵ assumed to be at λ_{\max} ; "9-cis-C ₁₅ aldehyde"	84E036

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
758.	(Z,Z)-3-Methyl-5-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2,4-pentadienal					
	Hexane	PR/ET	380-390, 21000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹), assuming authors used standard for cyclohexane; ϵ assumed to be at λ_{\max}	84E036
	Hexane	LP/SD	380-390, 17000		ϵ assumed to be at λ_{\max} , "7,9-dicis-C ₁₅ aldehyde"	84E036
759.	3-Methyl-3,4,5-triphenyl-2(3H)-furanone					
	Benzene	LP/ET	370 ± 5, 11000 ± 2200		†Oxygen quenching; ϵ relative to benzophenone in benzene ($\epsilon_{332} = 7600$ L mol ⁻¹ cm ⁻¹); $\tau_T = 12 \pm 2$ μ s; $k_{et} = 2.9 \times 10^9$ L mol ⁻¹ s ⁻¹	84E520
760.	1-Methyl-L-tryptophan					
	EtOH	FP	460		†Triplet ET from acetophenone	727041
	Water	FP	460		†Air quenching; $\tau_T = < 5$ μ s; pH 5.4	727041
761.	Methyl vinyl ketone					
	Cyclohexane	LP	260 ^a		†Oxygen quenching; solvent uncertain; $\tau_T = 0.008$ μ s	80B055
762.	1-Methyl-2-vinylnaphthalene					
	Benzene	LP-ET	416 ^a 401 ^a		†Triplet ET from xanthone and oxygen quenching (7.5×10^9 L mol ⁻¹ s ⁻¹); 401 nm peak was the more intense; $\tau_T = 0.06$ μ s	84B090
763.	2-Methyl-3-vinylnaphthalene					
	Benzene	LP-ET	571 ^a 412 ^a 391 ^a		†Triplet ET from xanthone and oxygen quenching (4.9×10^9 L mol ⁻¹ s ⁻¹); 412 nm peak was the most intense; $\tau_T = 0.255$ μ s	84B090
764.	1-Naphthaldehyde					
	Benzene	FP	495		$\tau_T = 140$ μ s	63E007
	Cyclohexane (77 K)	PS	412.5 405 387			69E211
	Cyclohexane	LP	494 475 450		Shoulders at 427 and 405 nm; delay 660 ps; rise time of 0.026 ± 0.003 ns	84B028
	Et ₂ O/Pentane (77 K)	PS	440 425 412 405 387			69E211
	MeOH	LP	478 ^a 456 ^a 410 ^a		Shoulders at 497 ^a , 437 ^a , and 425 ^a nm; delay 660 ps; 478 nm was the most intense peak; rise time of 0.020 ± 0.002 ns	84B028
	Toluene/MCH (50 K)	LP	479 ^a		Glass was 1:1 toluene to MCH; delay 42 ps; spectrum showed evidence of structure between 410 and 460 nm	84B028
	Water/ <i>tert</i> -BuOH	FP	520 ^a		Solvent mixture contains "1-5%" <i>tert</i> -BuOH for solubility; shoulders at 540 and 470 nm; pH ~6	767189
765.	2-Naphthaldehyde					
	Cyclohexane	LP	434 408		Delay 600 ps; rise time of 0.055 ± 0.005 ns	84B028
	MeOH	LP	442 ^a		Delay 600 ps; rise time of 0.025 ± 0.003 ns	84B028
	Toluene/MCH (69 K)	LP	446 ^a		Glass was 1:1 toluene to MCH; delay 42 ps; shoulders at 431 ^a and 405 ^a nm	84B028
766.	2-Naphthalenamine					
	Liquid paraffin	FP	457 377			61E008

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
767.	2-Naphthalenamine, conjugate acid					
	MeOH/Water (290 K)	LP/ET	410, 7800		†Oxygen quenching; ϵ relative to benzophenone ketyl radical in water ($\epsilon_{345} = 3220$ L mol ⁻¹ cm ⁻¹); solvent was 19:1 MeOH to water; $\tau_T = 1.8$ μ s; pH 1.8	83A249
	Water	FP	391 ^a , 4300 ^a		"2-Naphthylamine cation"; $pK_a = 3.3$; pH Acidic	61E008
			377 ^a , 2150 ^a			
			414			
			387			
768.	Naphthalene					
	1,2-Dichloroethane	FP	414		$\tau_T = 125$ μ s	80E720
	2-ProH	FP	414 ^a		Relative intensities (5:3:1); 30 μ s delay; $\tau_T = 58$ μ s	737158
			388 ^a			
			372 ^a			
	3-MP	PR	411 ^a		$G\epsilon$ values were 9870 and 27300 L mol ⁻¹ cm ⁻¹ / (100 eV absorbed)	680553
			388 ^a			
	3-MP (77 K)	PR	413 ^a		$G\epsilon$ values were 17300 and 17900 L mol ⁻¹ cm ⁻¹ / (100 eV absorbed)	680553
			390 ^a			
	3-MP (77 K)	PS/ESR	415, 31900		†ESR; oscillator strength = 0.12	69B002
	3-MP (77 K)	PS	412.4		All 8 bands were assigned to the electronic transition $^3B_{1g} \leftarrow ^3B_{2u}$	70B005
			399.3			
			390.0			
			377.7			
			369.8			
			358.2			
			351.5			
			341.2			
	3-MP (77 K)	MOD	415			73E342
			400			
			391.5			
			380			
			371			
			360			
	3-MP (85 K)	LP	415 ^a		415 nm peak was the most intense	747056
			400 ^a			
			391 ^a			
			379 ^a			
			372 ^a			
	3-MP (293 K)	LP	412		412 nm peak was the most intense; $\tau_T = 6$ μ s	747056
			390			
			370			
	Acetonitrile	FP	410 ^a		Relative intensities (5:3:1); 30 μ s delay; $\tau_T = 38$ μ s	737158
			385 ^a			
			375 ^a			
	Benzene	FP	415		Relative intensities (100:60:15)	61E005
			392			
			372			
	Benzene	PR	420		$G\epsilon_{420} = 15100$ L mol ⁻¹ cm ⁻¹ / (100 eV absorbed); half-life = 1.59 μ s	64B006
	Benzene	PR/ET	415, 17500		ϵ relative to benzophenone in benzene ($\epsilon_{532.5} = 10300$ L mol ⁻¹ cm ⁻¹); λ_{\max} unspecified, assumed 415 nm	690087
	Benzene	PR/ET	425, 13200		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700$ L mol ⁻¹ cm ⁻¹); ϵ obtained from a simultaneous least squares fit of data from several compounds making use of cyclohexane to benzene ϵ_{\max} ratios of 1.83 for naphthalene and 1.45 for anthracene	71E360
			402 ^a , 7300 ^a			
			382 ^a , 4200 ^a			
	Benzene	PR	420			720206
			400			
	Benzene	LP/ET	425, 11300		†Triplet ET from benzophenone; ϵ relative to benzophenone in benzene ($\epsilon_{533} = 7200$ L mol ⁻¹ cm ⁻¹); $k_{et} = 0.74 \times 10^9$ L mol ⁻¹ s ⁻¹	80B090
	Benzophenone (77 K)	PS-ET	425 ^a		Relative intensities (100:58 ^a); sample was a single crystal	64B005
			403 ^a			

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Benzophenone (77 K)	PS/COM	427 ^a , 3500 ^c 413 ^a , 1400 ^{a,c} 403 ^a , 1800 ^{a,c} 400 ^a , 1800 ^{a,c}		Mixed crystals; shoulder at 418 ^a ; ϵ was measured using a technique involving reabsorption of phosphorescence	68E106
	Butane/Isopentane (77 K)	PS/IV	414, 14000 \pm 3000 393 ^a , ~7800		Glass was 3:7 butane to isopentane; ϵ_T was computed from OD = $\epsilon_T \tau \rho_i (1 - \Phi_f)$ after the optical density, OD, was extrapolated to zero intensity of the monitoring beam (namely $i_0 \rightarrow 0$), here τ was lifetime of triplet, and Φ_f was fluorescence quantum yield; oscillator strength = 0.05	65E031
	Cyclohexane	FP	414		$\tau_T = 175 \mu s$	62E009
	Cyclohexane (77 K)	PS	415 ^a 412 ^a 410 ^a		Other shoulders were present in the spectrum; the shoulders appeared in different places depending on whether the samples were frozen rapidly or slowly	63B004
	Cyclohexane	PR	415 390		$G\epsilon_{415} = 10200 \text{ L mol}^{-1} \text{ cm}^{-1} / (100 \text{ eV absorbed})$; half-life = 5.85 μs	64B006
	Cyclohexane	PR	415 391 375		Relative intensities (100:55:22)	676011
	Cyclohexane (77 K)	PS	417.5 415 410.5 393.5 391 386 370 366 361 347.5			67B007
	Cyclohexane	PR/ET	412.5, 22600 \pm 5650		ϵ relative to benzophenone ketyl radical in water ($\epsilon_{537.5} = 3220 \text{ L mol}^{-1} \text{ cm}^{-1}$), assuming this value for $\epsilon_{542.5}$ in cyclohexane	680727
	Cyclohexane	LP	414			68E107
	Cyclohexane (77 K)	PS	415.0 411.5 409.0 393.0 389.5 387.5 373.0		Shoulder at 422 nm; peak positions depend on cooling rate	69E211
	Cyclohexane (77 K)	PS	412.0 409.0 406.8 390.0 386.5 385.5 370.0		Shoulder at 418.5 nm; peak positions depend on cooling rate	69E211
	Cyclohexane	PR/ET	415, 24500 393 ^a , 12000 ^a 373 ^a , 4600 ^a		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700 \text{ L mol}^{-1} \text{ cm}^{-1}$); reference ϵ obtained by starting from $\epsilon_{\max} = 3220 \text{ L mol}^{-1} \text{ cm}^{-1}$ for this ketyl radical in water and assuming the f of the ketyl radical is independent of solvent; final ϵ obtained from a simultaneous least squares fit to data from several compounds	71E360
	Cyclohexane	PR	412.5 390 370			720206
	Cyclohexane	LP	414 391 370		$\tau_T = 10 \mu s$	747056
	Cyclohexane	FP/SD	414, 24000 \pm 3000			756293
	Dipropyl ether (118 K)	FP	417 ^a 405 ^a 394 ^a			696073

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
Durene (77 K)	PS		419 395		Polarization of the bands was found to be along the long molecular axis; transition was assigned as ³ B _{2g} ← ³ B _{1g} ; sample was a mixed crystal	64E014
Durene (4.2 K)	PS		417.4 394.4 383 374		Single crystal; relative intensities (100:46:2:10)	67B008
Durene (1.5 K)	PS		417.4 403 394 382 373		Mixed crystal; maxima were taken from a fit to the spectrum	75B003
EPA (77 K)	PS/IV		417, 480 ^d ± 50 392.8, 215 ^d ± 40		†Phosphorescence decay; ϵ 's are too low; 1st transition assigned as ³ L _a → ³ C _b	51E001
EPA (77 K)	PS/SD		416.7, 10000 392.7, 4300 371, 1000		ϵ 's are lower limits	54B001
EPA (77 K)	FP		416 393 373		416 nm was the most intense peak; $\tau_T = 2.6 \times 10^6 \mu s$	677485
EPA (77 K)	PS/KM		530 ^a , 900 ^a 499 ^a , 800 ^a 420 ^a , 20600 ^a 396 ^a , 11200 ^a		Solvent, temperature and extinction method assumed from earlier work; polarization also measured	69E213
EPA (77 K)	PS/IV		417, 23600		λ_{\max} assumed from previous work; ϵ estimated by extrapolation to infinite excitation rate	69E212
EPA (77 K)	PS/KM		417, 23300		λ_{\max} assumed from previous work; $\tau_T = 2.3 \times 10^6 \mu s$	69E212
EPA (77 K)	FP/SD		414.0, 40000 ± 6000 393 ^a , 17000 ^a 374 ^a , 4000 ^a		$\tau_T = 2.3 \times 10^6 \mu s$; oscillator strength = 0.14	69F389
EtOH (77 K)	PS		414 390 371			67B007
EtOH (293 K)	FP		414			68E098
EtOH	FP/ET		415, 40000 ± 2000 391 ^a		ϵ relative to anthracene in EtOH ($\epsilon_{420} = 75000$ L mol ⁻¹ cm ⁻¹); $\tau_T = 1800 \pm 700 \mu s$; $k_{et} = (1.26 \pm 0.1) \times 10^{10}$ L mol ⁻¹ s ⁻¹	747049
EtOH (77 K)	MOD		411 388 ^a 368 ^a		Relative intensities (6:3:1); halfwidth of principal maximum 480 cm ⁻¹	777538
EtOH/MeOH (113 K)	FP/TD		575 ^a , 140 ^a 559 ^a , 75 ^a 529 ^a , 250 ^a 515 ^a , 150 ^a 490 ^a , 330 ^a 476 ^a , 150 ^a 360 ^a , 459 ^a		Glass was 3:1 EtOH to MeOH; shoulders at 625 ^a , 588 ^a , 541 ^a , 500 ^a , and 467 ^a nm	717460
EtOH/MeOH (113 K)	FP/TD		629.0, 10 ^a 574 ^a , 120 ^a 526 ^a , 250 488.6, 320 ^a 456 ^a , 350 ^a 414.0, 33000 400.2, 3200 ^a 390 ^a , 16700 ^a 376 ^a , 2100 ^a 369 ^a , 3500 ^a 359 ^a , 1000 ^a 351 ^a , 1000 ^a 341 ^a , 700 ^a 335 ^a , 700 ^a 323 ^a , 730 ^a 236.0, 55000		†Phosphorescence decay; shoulders at 589 ^a , 556 ^a , 536 ^a , 511 ^a , 498 ^a , 467 ^a , and 266 ^a nm; 6 electronic transitions were assigned with 0-0's at 629, 488.6, 414, 400.2, 275, and 236 nm; oscillator strength = 0.002, 0.002, 0.12, ~0.01, 0.13, 0.5	72B001

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Ethylene glycol	FP	415 392 378		Relative intensities (100:55:10); $\tau_T = 1000 \pm 100 \mu s$	61E005
	Hexamethylbenzene (77 K)	PS	418 395		Single crystal; relative intensities (2:1)	67B008
	Hexane	FP	413.1 390.0			54E001
	Hexane	FP	410 389 369		Relative intensities (107:55:15); $\tau_T = 91 \pm 8 \mu s$	61E005
	Hexane (77 K)	PS	416 391			67B007
	Isopentane/MCH (77 K)	PS	413.5 389.5 368		Relative intensities (100:41:9)	54B001
	Isopentane/MCH/Et ₂ O (77 K)	PS	414.5 392.0 372.5		‡Phosphorescence decay; glass was 6:1:0.7 isopentane to MCH to Et ₂ O; 414.5 nm was the most intense peak	71B003
	Liquid paraffin	FP/SD	415, 10000 391.5, 6800 372, 2700		Solvent had viscosity of 0.03 N·s/m ² ; all bands were assigned to same electronic transition; oscillator strength = 0.06	58E001
	Liquid paraffin MCH (77 K)	FP MOD	415 414 ^a 390 ^a 370 ^a		Viscosity of solvent was 0.167 N·s/m ² ; $\tau_T = 2850 \mu s$ Relative intensities (6:3:1)	62E009 777538
	MCH (77 K)	MOD/KM	405 ^b , 1600 366 ^b , 740			82E648
	MCH/Isopentane (90 K)	MOD	417 ^a 400 ^a 391 ^a 382 ^a 372 ^a 360 ^a 351 ^a		Glass was 4:1 MCH to isopentane	72B003
	Naphthalene (373 K)	PR/RF	420, 7100 ^c		Liquid; ϵ was found by assuming that the oscillator strength remained 0.12 in all solvents; $\tau_T = 1.2 \mu s$	720243
	Naphthalene- <i>d</i> ₈ (4.2 K)	MOD	415 402.5 392.5 380 372.5 360		Mixed crystals; the bands were embedded in a broad absorption band that was explained as due to mixing of guest triplet states with host conduction band or host charge-transfer states	73E342
	Naphthalene- <i>d</i> ₈ (1.5 K)	PS	414.8 392 370		Mixed crystal; maxima were taken from a fit to the spectrum	75B003
	Naphthalene (78 K)	PR	600 565 535 416 392		Triplet exciton; most intense peak at 416 nm	761030
	Naphthalene	PR	415 ^a 390 ^a		Microcrystals; spectrum obtained by diffuse reflectance spectroscopy; 415 nm peak was the more intense; half-life = 20 μs	84B120
	PMMA (293 K)	PS	415.5 393.5		Relative intensities (100:45)	67B008
	PMMA (294 K)	FP/RF	571, ~220 526, ~310 415 ^a 391 ^a 373 ^a		‡Phosphorescence decay; ϵ relative to naphthalene in cyclohexane ($\epsilon_{412.5} = 22600 \text{ L mol}^{-1} \text{ cm}^{-1}$), assuming $\epsilon_{417} = 22000 \text{ L mol}^{-1} \text{ cm}^{-1}$ in PMMA; $\tau_T = (1.5 \pm 0.05) \times 10^6 \mu s$; oscillator strength = ~0.002	69B004
	PMMA	PR	415 385		Relative intensity (10:6)	692001
	PMMA (86 K)	PR	415 390			710278

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Pentane (77 K)	PS	413.6 401.6 390.6 370.1		Relative intensities (10:1:6:1)	69E211
	Polystyrene	PR	428 400 380		Relative intensities (100:53:33)	672086
	Polystyrene (77 K)	PR	425			692001
	Polystyrene	PR	424 ^a		$G\epsilon_{424} = 12000^a$ L mol ⁻¹ cm ⁻¹ / (100 eV absorbed)	701073
	Toluene (77 K)	MOD	422 402 ^a		Relative intensities (2:1); halfwidth of principal maximum 1180 cm ⁻¹	777538
769.	Naphthalene- <i>d</i> ₈ 1-BuOH/Isopentane (77 K)	PS	413.2 389.7 371.0		A distinction between a 1:7 and a 3:7 glass was not made; the 3 bands were assigned to a single electronic transition $^3B_{3g}^- \leftarrow ^3B_{1g}^+$	63B001
	3-MP (77 K)	PS/ESR	417, 31900 403 ^a , 5200 ^a 391 ^a , 14000 ^a		†ESR; band was assigned as $^3B_{3g}^- \leftarrow ^3B_{2u}^-$; oscillator strength = 0.12	68D211
	3-MP (77 K)	PS	575.0 547.0 531.2 509 ^a 491.6 473 ^a 456.0 412.2 400.5 390.0 379.3 369.3 359.4 351.3 342.2		Bands starting at 412.2 nm were assigned to the $^3B_{3g}^- \leftarrow ^3B_{2u}^+$ transition; other bands may be a mixture of electronic transitions, but one of these was assigned as $^3A_{1g}^- \leftarrow ^3B_{2u}^+$; the bands at <412 nm are about 20 times stronger than the bands >412 nm	70B005
	? (77 K)	PS	419 ^a 406 ^a 393 ^a		†Phosphorescence decay and ESR lifetime; glass used was either 2-MTHF or 3-MP; $\tau_T = 2.30 \times 10^7$ μ s	67E108
	Butane/Isopentane (77 K)	PS/IV	414, 14000 \pm 3000 393 ^a , ~1800		Glass was 3:7 butane to isopentane; ϵ_T was computed from $OD = \epsilon_T \tau_{p10}(1 - \Phi_f)$ after the optical density, OD, was extrapolated to zero intensity of the monitoring beam (namely $i_0 \rightarrow 0$), here τ was lifetime of triplet, and Φ_f was fluorescence quantum yield; oscillator strength = 0.05	65E031
	Cyclohexane (77 K)	PS	413.5 409.4 407.2 402.0 389.8 386.0 383.2 379.5 369.2 367.2		Shoulders at 418.2 nm and 397.4 nm; relative intensities (100:80:60:20:60:50:20:15:20:15)	69E211
	Dipropyl ether (100 K)	FP	415 ^a 394 ^a		Shoulders at 405 ^a and 373 ^a nm; $\tau_T = 1.9 \times 10^7$ μ s	696073
	Dipropyl ether (110 K)	PS	417 ^a 395 ^a		Relative intensities (2:1); polarization also measured	70B002
	Durene (77 K)	PS	417 ^a 395 ^a 372 ^a		Single crystal; the points were taken from the spectrum of the <i>b</i> -axis polarized absorption	67B008
	Durene (4.2 K)	PS	415.9 392.9 382 371 354		Single crystal; relative intensities (100:47:2:14:2)	67B008

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	EPA (82 K)	PS/KM	413, 23300 ± 650			68B003
	EPA (77 K)	PS/IV	417, 22000		λ_{\max} assumed from previous work; ϵ estimated by extrapolation to infinite excitation rate; $\tau_T = 19 \times 10^6 \mu s$	69E212
	EPA (77 K)	PS	417		Transition assigned as $^3B_{1g} \leftarrow ^3B_{2u}$	69B001
	EPA (77 K)	FP/SD	414.0, 40000 ± 6000		$\tau_T = 1.35 \times 10^7 \mu s$	69F389
	EtOH/Et ₂ O (77 K)	MOD/KM	417, 10000 ± 3000		Glass was 2:1 EtOH to Et ₂ O; temperature was not explicitly stated, but 77 K was inferred from the context	719059
	EtOH/MeOH (113 K)	FP/TD	575 ^a , 110 ^a 549 ^a , 70 ^a 529 ^a , 220 ^a 508 ^a , 130 ^a 493 ^a , 240 ^a 472 ^a , 140 ^a 457 ^a , 280 ^a 448 ^a , 170 ^a 439 ^a , 320 ^a		Glass was 3:1 EtOH to MeOH; shoulders at 595 ^a , 562 ^a , 515 ^a , 485 ^a , and 431 ^a nm	717460
	EtOH/MeOH (101 K)	PS	413 ^a 400 ^a 395 ^a 380 ^a 370 ^a		Glass was 3:1 EtOH to MeOH	75B002
	Hexamethylbenzene (77 K)	PS	415.3 393.2		Single crystal; relative intensities (100:48)	67B008
	PMMA (293 K)	PS	414.5 392.1 372		Relative intensities (100:44:4)	67B008
770.	Naphthalene/Chloranil Liquid paraffin	MOD	405		Mull	71E361
771.	Naphthalene/Pyromellitic dianhydride 1,2-Dichloroethane (293 K)	LP	420		Charge-transfer complex only was excited; naphthalene-PMDA charge-transfer complex	757413
	Dipropyl ether (100 K)	FP	426 ^a		Shoulder at 403 ^a nm; $\tau_T = 1.8 \times 10^6 \mu s$	696073
772.	Naphthalene/Tetrachlorophthalic anhydride Dipropyl ether (98 K)	FP	424 ^a		Shoulder at 407 ^a nm	696073
773.	Naphthalene- <i>d</i> ₈ /Tetrachlorophthalic anhydride EPA (77 K)	PS	414.8 390.5			67B009
774.	1-Naphthalenecarbothioic acid, <i>O</i> -ethyl ester Benzene	FP	490		†Triplet ET to triplet acceptors; the solvent was not specified explicitly	757534
775.	2-Naphthalenesulfonate ion Water	FP	413 390		†Rises with fluorescence decay; 50 μs delay; pH 6; rise time of 23 ns	767189
776.	1-Naphthalenesulfonic acid Liquid paraffin	FP	421 405 373.5		Solvent viscosity was 0.19 N·s/m ² ; relative intensities (100:88:54)	58E001
777.	<i>anti</i> -[2.2](1,4)-Naphthalenophane 2-MTHF (77 K)	LP	447.5			79B050
778.	<i>syn</i> -[2.2](1,4)-Naphthalenophane 2-MTHF (77 K)	LP	612.5 465			79B050

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
779.	2-(1-Naphthalenyl)-1,3-dioxane					
	1,2-Dichloroethane	FP	420	$\tau_T = 108 \mu s$		80E720
	Hexane	FP	415	$\tau_T = 63 \mu s$		80E720
780.	2-(2-Naphthalenyl)-1,3-dioxane					
	1,2-Dichloroethane	FP	417	$\tau_T = 142 \mu s$		80E720
	Hexane	FP	412	$\tau_T = 104 \mu s$		80E720
781.	1-Naphthoate ion					
	Water	FP	439	pH 13		61E008
782.	2-Naphthoate ion					
	Water	FP	414	pH 13		61E008
783.	Naphtho[1,2,3,4-def]chrysene					
	Benzene	MOD	610 522 442	Relative intensities (100:64:41); "1,2,4,5-dibenzpyrene"; $\tau_T = 140 \mu s$		71E361
784.	1-Naphthoic acid					
	Water	FP	446	$pK_a = 3.8 \pm 0.5$; pH 1		61E008
785.	2-Naphthoic acid					
	Liquid paraffin	FP	430 412			61E008
	Water	FP	428	$pK_a = 4.0$; pH 1		61E008
786.	1-Naphthol					
	EPA (77 K)	PS	483 463 455	‡Phosphorescence decay		51E001
	Hexane	FP	520 389.5 374.5 347.5 335	All bands except 520 nm were assigned to the second electronic transition; relative intensities (16:100:67:49:24)		58E001
	Isooctane	FP	435 400	‡Similarity to 1-naphthyl diisopropyl borate spectrum		717266
	Isopentane/MCH/Et ₂ O (77 K)	PS	481.0 455.0 431.5 412.0	‡Phosphorescence decay; glass was 6:1:0.7 isopentane to MCH to Et ₂ O; 455.0 and 431.5 nm were the most intense peaks		71B003
	PMMA	FP	485 460 430 400	‡Similarity of spectrum to that of 1-naphthyl diisopropyl borate; there was a weak band extending from 540 - 510 nm		717266
787.	2-Naphthol					
	Cyclohexane	LP	433 ^a 413 ^a 385 ^a			737287
	Cyclohexane	FP	467 ^a 429 ^a 408 ^a 384 ^a 355 ^a	‡Oxygen quenching and triplet ET to perylene; most intense peak at 429 nm; $\tau_T = 67 \mu s$		737113
	Cyclohexane	LP/ET	435, 6500	Triplet ET from perylene; ϵ relative to perylene in cyclohexane ($\epsilon_{430} = 13000 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 67 \mu s$		777391
	Gelatin	FP	443	‡Phosphorescence decay; $\tau_T = 4.7 \times 10^5 \mu s$		64E016
	Heptane	FP	430 ^a	Lifetimes suggest solution not fully deaerated; $\tau_T = 36 \pm 3 \mu s$		77E663
	Hexane	FP	429.0 404.0 381.0	Another band at 467 nm was identified as the 2-naphthoxyl radical; 429.0 nm was the most intense peak		71B003

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Isooctane	FP	433 405		†Similarity to 2-naphthyl diisopropyl borate spectrum	717266
	Isopentane/MCH/Et ₂ O (77 K)	PS	443.0 419.0 395.5		†Phosphorescence decay; glass was 6:1:0.7 isopentane to MCH to Et ₂ O; 443.0 nm was the most intense peak	71B003
	Liquid paraffin	FP	433 411 381.5			61E008
	PMMA	FP	525 438 405		†Similarity to 2-naphthyl diisopropyl borate spectrum; 438 nm was the most intense peak	717266
	Water	FP	432		Another maximum near 350 nm and a shoulder near 390 nm; pK _a = 8.1; pH 4.6	61E008
	Water	LP	431 ^a		Solution saturated with N ₂ O; [HClO ₄] = 0.2 mol L ⁻¹ ; shoulder at ~410 ^a nm; delay 30 ns; pH Acidic	737287
788.	1-Naphthol/Triethylamine Isopentane/MCH/Et ₂ O (77 K)	PS	488 ^a 461 ^a 435 ^a 417 ^a		H-bonded complex; glass was 6:1:0.7 isopentane to MCH to Et ₂ O; shoulder at 395 ^a nm; 461 nm was the most intense peak	71B003
789.	2-Naphthol/Triethylamine Isopentane/MCH/Et ₂ O (77 K)	PS	446 ^a 424 ^a 388 ^a		H-bonded complex; glass was 6:1:0.7 isopentane to MCH to Et ₂ O; 446 nm was the most intense peak; shoulder at 405 ^a nm	71B003
790.	9-Naphthoylanthracene EPA (77 K)	FP	427		†Phosphorescence decay; $\tau_T = 3.07 \times 10^4 \mu s$	82E338
791.	1-Naphthyl acetate MCH (77 K)	MOD/KM	436 ^b , 960 405 ^b , 1400 366 ^b , 560			82E648
792.	2-(1-Naphthyl)benzoxazole Pentane	LP/TD	530, 24200 ± 250 489 ^a 422 ^a		Shoulder at 454 ^a nm; $\tau_T = 0.52 \pm 0.05 \mu s$	82E632
793.	8-(1-Naphthyl)-6,7-dihydro-5H-benzocycloheptene Benzene	LP-ET	507 ^a 390 ^a		†Triplet ET from xanthone	84B007
794.	1-Naphthyl diisopropyl borate Triisopropyl borate	FP	900 535 502.5 476 450 420 397 377			717266
795.	2-Naphthyl diisopropyl borate Triisopropyl borate	FP	525 425 401 380		425 nm was the most intense peak	717266
796.	trans-1-(1-Naphthyl)-2-(2-naphthyl)ethylene Benzene	LP/ET	500 ± 3, 19000 ± 2900		†Quenching by oxygen, a nitroxide free radical, and azulene; ϵ relative to fluorenone in benzene ($\epsilon_{435} = 6000 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 0.56 \pm 0.08 \mu s$; $k_{et} = (6.9 \pm 1.0) \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	84E237

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Bromobenzene	LP	505 ± 3		$\tau_T = 0.43 \pm 0.06 \mu s$	84E237
797.	1-Naphthyloxy ion EtOH/MeOH (77 K)	PS	505 ^a 476 ^a 452 ^a		Glass was 5:1 EtOH to MeOH; there was a shoulder at 422 ^a nm; the solution was 0.11 mol L ⁻¹ KOH; 505 nm was the most intense peak	71B003
798.	2-Naphthyloxy ion Water	FP	460		Another maximum in the range 350 - 400 nm; pH 10.6	61E008
	Water	LP	455 ^a		Solution saturated with N ₂ O; delay 30 ns; pH 13	737287
799.	2-(2-Naphthyloxy)-3,5,5-trimethyl-2-cyclohexen-1-one MCH	LP	430 ^a		†Oxygen quenching (1.7×10^9 L mol ⁻¹ s ⁻¹), decay measured at 410 nm and spectrum obtained by extrapolation to zero time; $\tau_T = 0.53 \mu s$	81F111
300.	1-(1-Naphthyl)-1-phenylethylene Benzene	LP-ET	500 ^a 403 ^a		†Triplet ET from xanthone; $\tau_T = 0.040 \mu s$	84B007
801.	trans-1-(1-Naphthyl)-2-phenylethylene Benzene	LP/ET	490 ± 3, 17000 ± 2600		†Quenching by oxygen, a nitroxide free radical, and azulene; ϵ relative to fluorenone in benzene ($\epsilon_{435} = 6000$ L mol ⁻¹ cm ⁻¹); $\tau_T = 0.39 \pm 0.06 \mu s$; $k_{et} = (6.7 \pm 1.0) \times 10^9$ L mol ⁻¹ s ⁻¹	84E237
	Bromobenzene	LP	495 ± 3		$\tau_T = 0.25 \pm 0.04 \mu s$	84E237
802.	trans-1-(2-Naphthyl)-2-phenylethylene Benzene	LP-ET	530 ^a 410 ^a		†Triplet ET from xanthone and benzophenone, oxygen and azulene quenching; 50 ns delay; lifetimes are 0.08 μs (530 nm band) and 0.15 μs (410 nm band); possible triplet conformers; $E_T = 206$ kJ mol ⁻¹	81E490
	Benzene	LP/ET	395 ± 3, 8000 ± 1200		ϵ relative to fluorenone in benzene ($\epsilon_{435} = 6000$ L mol ⁻¹ cm ⁻¹); $\tau_T = 0.14 \pm 0.02 \mu s$; $k_{et} = (9.0 \pm 1.3) \times 10^9$ L mol ⁻¹ s ⁻¹	84E237
	Bromobenzene EPA (77 K)	LP LP	400 ± 3 510 497 492 470		$\tau_T = 0.09 \pm 0.01 \mu s$	84E237 747376
803.	2-(1-Naphthyl)-5-phenyl-1,3,4-oxadiazole Benzene	LP/SD	580, 67000		†Oxygen quenching (1.6×10^9 L mol ⁻¹ s ⁻¹); 100 ns delay; "α-NPD"; $\tau_T = 0.300 \mu s$	777265
	Cyclohexane	FP/TD	696, 6200 ^a 496, 5900 ^a 454, 4800 ^a		Only peaks reported by authors given - extensive vibronic structure observed; oscillator strength = 0.13, 0.02, 0.04	78B081
	Cyclohexane	FP/TD	550 ^a , 11000 ^a 495, 5000 ^a 455, 5000 ^a		Only principal maximum of the band(s) around 500 nm reported, extensive vibrational structure observed; ϵ method assumed from earlier work; oscillator strength = 0.13, 0.02, 0.04	79B119
804.	2-(2-Naphthyl)-5-phenyl-1,3,4-oxadiazole Cyclohexane	FP/TD	611, 6200 ^a 508, 4000 ^a 413, 14000 ^a		Only peaks reported by authors given; oscillator strength = 0.04, 0.02, 0.09	78B081
	Cyclohexane	FP/TD	611, 6200 ^a 508, 4000 ^a 488, 3800 ^a 441, 4200 ^a 423, 14100 ^a		ϵ method assumed from earlier work; oscillator strength = 0.03, <0.01, <0.01, 0.01, 0.09	79B119

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
805.	2-(1-Naphthyl)-5-phenyloxazole 3-MP (77 K) Benzene	CWL LP/SD	580 550, 100000		" α -NPO" †Oxygen quenching (2.3×10^9 L mol ⁻¹ s ⁻¹); 100 ns delay; $\tau_T = 0.215$ μ s	74B003 777265
806.	Neo-alloocimene Toluene	PR/ET	340 ^a , 930 ^a 315, 1840 ^a		†Triplet ET to perylene; ϵ relative to perylene in benzene ($\epsilon_{490} = 14300$ L mol ⁻¹ cm ⁻¹); $\tau_T = 0.33$ μ s; $k_{et} = 9.1 \times 10^9$ L mol ⁻¹ s ⁻¹	83E258
807.	Neomethylene Blue cation Water	FP/SD	840 ^b , 4000		†Triplet ET from naphthalene, oxygen and I ⁻ quenching; $pK_a = 6.5$; structure of the spectrum could not be discerned because it was not corrected for SD; pH 9	70E293
808.	<i>all-trans</i> -Neurosporene Cyclohexane	PR/ET	489, 274000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{360} = 42800$ L mol ⁻¹ cm ⁻¹); $\tau_T = 6.6$ μ s; $k_{et} = 1.22 \times 10^9$ L mol ⁻¹ s ⁻¹	83B121
809.	Neutral Red cation Water	FP/SD	580, 7300 380, 5800		†Oxygen quenching; pK_a 4.5; $\tau_T = 63 \pm 12$ μ s; pH 9.8	84B027
810.	Neutral Red cation, conjugate diacid Water	FP/SD	680, 12000 390, 8000		†Oxygen quenching; pK_a 4.8; $\tau_T = 63 \pm 12$ μ s; pH 2.8	84B027
811.	5-Nitroacenaphthene EPA (77 K)	PS/KM	850, 250 \pm 30 600, 500 \pm 60 370, 7100 \pm 850		†Phosphorescence decay; $\tau_T = (2.6 \pm 0.3) \times 10^5$ μ s	78E057
812.	4-Nitroaniline EPA (113 K) EPA (100 K)	FP FP	595 600 ^a		†Phosphorescence decay; $\tau_T = 250$ μ s Lifetime measured at 93 K; $\tau_T = 4.5 \times 10^4$ μ s	776104 80E318
813.	3-Nitroanisole CF ₃ CH ₂ OH	LP	400 ^a		$\tau_T = 3.5$ μ s	84E033
814.	4-Nitroanisole Glycerol triacetate (198 K) Glycerol triacetate	LP LP	560 418 550 414		418 nm peak was the more intense 550 nm peak was the more intense	78B088 78B088
815.	9-Nitroanthracene Benzene (293 K) EPA (77 K) EtOH EtOH (77 K)	LP FP LP FP	435 ^a 440 ^a 430 ^a 435 ^a		170 ps delay; growth followed at 452 nm; authors later [84F385] quote $\lambda_{\max} = 433$ nm, with a shoulder at 410 nm; rise time of $(76 \pm 6) \times 10^{-3}$ ns $\tau_T = 1.77 \times 10^4$ μ s Rise time measured at 452 nm; 100 ps delay; rise time of 0.086 ± 0.006 ns $\tau_T = 1.77 \times 10^4$ μ s	80B084 80B084 80E234 80E234
816.	4-Nitrobiphenyl Benzene EPA (77 K) Glycerol triacetate (198 K) Glycerol triacetate	LP FP LP LP	540 550 520 530 ~375 526 ~370		†Phosphorescence decay 530 nm peak was the more intense 526 nm peak was the more intense	747022 747022 78B088 78B088

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
817.	2-[2-(5-Nitro-2-furanyl)]ethenylquinoline 2-PrOH/Water	FP	590		†Triplet energy transfer to 3,3'-diethylthiacarbocyanine iodide (in BuOH); solvent was 1:1 2-PrOH to water; 50 μ s delay; τ_T = 670 \pm 90 μ s	79A403
818.	5-Nitro-2-furoic acid					
	Acetone (295 K)	LP/HAT	490, 19000		Hydrogen atom transfer from 2,4,6-tri- <i>tert</i> -butylphenol (1.71×10^9 L mol ⁻¹ s ⁻¹); ϵ relative to 2,4,6-tri- <i>tert</i> -butylphenoxyl radical (ϵ_{620} = 400 L mol ⁻¹ cm ⁻¹); τ_T = 0.2036 μ s	81A140
	Acetone (295 K)	LP/ELT	490, 20900		†Triplet ET to 1,4-benzoquinone, azulene and ferrocene, oxygen quenching (1.68×10^9 L mol ⁻¹ s ⁻¹); electron transfer from triphenylamine (3.3×10^9 L mol ⁻¹ s ⁻¹); ϵ relative to triphenylamine radical cation (ϵ_{660} = 28750 L mol ⁻¹ cm ⁻¹); τ_T = 0.2036 μ s; E_T = 240 \pm 4 kJ mol ⁻¹	81A140
	Acetone (295 K)	LP/ELT	490, 20500		Electron transfer from tris(4-bromophenyl)amine (8.94×10^9 L mol ⁻¹ s ⁻¹); ϵ relative to tris(4-bromophenyl)amine radical cation (ϵ_{725} = 20800 L mol ⁻¹ cm ⁻¹); τ_T = 0.290 μ s	81A140
	Acetone (295 K)	LP/ELT	490, 20700		Electron transfer from TMPD (5.98×10^9 L mol ⁻¹ s ⁻¹); ϵ relative to TMPD radical cation (ϵ_{565} = 12470 L mol ⁻¹ cm ⁻¹); τ_T = 0.290 μ s	81A140
	Acetonitrile (295 K)	LP/ELT	490, 22500		Electron transfer from iodide ion (6.65×10^9 L mol ⁻¹ s ⁻¹); ϵ relative to iodine molecular anion (ϵ_{400} = 12900 L mol ⁻¹ cm ⁻¹); τ_T = 0.290 μ s	81A140
	DMF (295 K)	LP	490		τ_T = 0.0766 μ s	81A140
	Ethyl acetate (295 K)	LP	490		τ_T = 0.070 μ s	81A140
	Methylene chloride (295 K)	LP	495		τ_T = 0.070 μ s	81A140
	Water (295 K)	LP	486		Lifetime increases to 34.2 ns in deuterated water; τ_T = 0.0272 μ s	81A140
819.	1-Nitronaphthalene					
	EPA (77 K)	PS	580			78E057
	EtOH	LP	580		†Phosphorescence decay at 77 K in PMMA and triplet ET to perylene and tetracene	747236
	EtOH	LP	580		†Oxygen quenching (3.3×10^9 L mol ⁻¹ s ⁻¹); τ_T = 4.9 μ s	81B064
	Hexane	LP	525		†Phosphorescence decay at 77 K in PMMA and triplet ET to perylene and tetracene	747236
	Hexane	LP	525		†Oxygen quenching (1.3×10^9 L mol ⁻¹ s ⁻¹); τ_T = 0.93 μ s	81B064
	N-Methylformamide	LP	615		†Phosphorescence decay at 77 K in PMMA and triplet ET to perylene and tetracene	747236
820.	2-Nitronaphthalene					
	Acetonitrile	LP	480		†Phosphorescence decay in EPA at 77K, triplet ET to tetracene; τ_T = 2.90 μ s; k_{et} = 7.1×10^9 L mol ⁻¹ s ⁻¹	767269
	EPA	LP	470		Radical or radical anion seen at λ_{\max} = 400 nm on exposure to a long (5 μ s) flash	767269
	EPA (77 K)	PS/KM	760, 510 \pm 25 475, 750 \pm 40 360, 3600 \pm 180		†Phosphorescence decay; τ_T = (2.47 \pm 0.02) $\times 10^5$ μ s	78E057
	EtOH	LP	470		†Phosphorescence decay in EPA at 77K, triplet ET to tetracene; τ_T = 1.70 μ s	767269
	EtOH	LP	470		†Oxygen quenching (1.6×10^9 L mol ⁻¹ s ⁻¹); τ_T = 1.7 μ s	81B064
	Formamide	LP	520 ^a		†Phosphorescence decay in EPA at 77K, triplet ET to tetracene	767269
	Hexane	LP	425		†Phosphorescence decay in EPA at 77K, triplet ET to tetracene; τ_T = 0.530 μ s	767269

TABLE 6 Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Hexane	LP	425		†Oxygen quenching (1.7×10^9 L mol ⁻¹ s ⁻¹); $\tau_T = 0.53$ μ s	81B064
	Water	LP	485		†Phosphorescence decay in EPA at 77K, triplet ET to tetracene; $\tau_T = 7.60$ μ s	767269
821.	4-Nitropyridine- <i>N</i> -oxide	FP	550 380		†Oxygen quenching; $\tau_T = 30$ μ s	737469
822.	4-Nitroquinoline- <i>N</i> -oxide	LP	530		†Triplet ET to tetracene; $k_{et} = 8 \times 10^9$ L mol ⁻¹ s ⁻¹	84B128
	Benzene	LP	590		Phosphate buffer; $\tau_T = 50$ μ s; pH 7	84B128
	Water	LP	410			
823.	<i>trans</i> -3-Nitrostilbene	LP	365			78B088
	Glycerol triacetate	LP	382			78B088
	Glycerol triacetate (198 K)					
824.	<i>trans</i> -4-Nitrostilbene	LP	540		†Oxygen quenching in benzene; $\tau_T = 0.063$ μ s	747022
	Cyclohexane	FP	585 ^a		Shoulder at 555 ^a	747022
	EPA (77 K)	LP	590		Solvent was 4:1 EtOH to MeOH; 450 nm peak was the more intense	78B088
	EtOH/MeOH (103 K)	LP	450			
	EtOH/MeOH	LP	585		Solvent was 4:1 EtOH to MeOH; 440 nm peak was the more intense	78B088
			440			
	Glycerol triacetate	LP	540		540 nm peak was the more intense	78B088
			425			
	Glycerol triacetate	LP	550		550 nm peak was the more intense	78B088
	(191 K)		426			
825.	4-Nitro- <i>p</i> -terphenyl	PS/KM	800, 14500 \pm 725 600, 25500 \pm 1275 340, 38500 \pm 1925		†Phosphorescence decay; $\tau_T = (3.14 \pm 0.16) \times 10^5$ μ s	78E057
826.	2-Nitrothiophene	LP/ELT	545 ^b , 8800 544		ϵ relative to (C ₆ H ₅) ₂ N \cdot in MTHF glass at 77 K ($\epsilon_{\max} = 3900$ L mol ⁻¹ cm ⁻¹); protonation was thought to take place after ELT; $\tau_T = 0.118$ μ s	82A153
	1-ProH (273 K)					
	Acetone (273 K)	LP	549		$\tau_T = 0.227$ μ s	82A153
	Acetonitrile (273 K)	LP/ELT	548 545 ^b , 10900		†Oxygen quenching; ϵ relative to 3,3',3''-tribromotriphenylamine radical cation in methylene chloride ($\epsilon_{\max} = 20800$ L mol ⁻¹ cm ⁻¹), assuming no solvent effect; ϵ was to be an upper bound since assumptions included complete ELT, no back-reaction, and no reaction with solvent; $\tau_T = 0.30$ μ s	82A153
	Acetonitrile (273 K)	LP/ELT	548 545 ^b , 8800		†Oxygen quenching; ϵ relative to TMPD ⁺ in water ($\epsilon_{\max} = 12470$ L mol ⁻¹ cm ⁻¹), assuming no solvent effect; ϵ was to be an upper bound since assumptions included complete ELT, no back-reaction, and no reaction with solvent; $\tau_T = 0.30$ μ s	82A153
	Acetonitrile (273 K)	LP/ELT	548 545 ^b , 11200		†Oxygen quenching; ϵ relative to triphenylamine radical cation in EtOH ($\epsilon_{\max} = 28750$ L mol ⁻¹ cm ⁻¹), assuming no solvent effect; ϵ was to be an upper bound since assumptions included complete ELT, no back-reaction, and no reaction with solvent; $\tau_T = 0.30$ μ s	82A153
	CCl ₄ (273 K)	LP	549		$\tau_T = 0.022$ μ s	82A153
	EtOH (273 K)	LP	545		$\tau_T = 0.129$ μ s	82A153
	MeOH (273 K)	LP	545		$\tau_T = 0.282$ μ s	82A153
	Methylene chloride (273 K)	LP	545		$\tau_T = 0.173$ μ s	82A153
	Water (273 K)	LP	545			82A153
	Water	LP	545		Phosphate buffer used; pH 7.0 \pm 0.1	82A154

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
827.	Octaethylporphinatopalladium(II) Methyl methacrylate	LP	497 ^a 417 ^a		417 nm peak was the more intense	747346
828.	Octaethylporphinatotin(IV) dichloride 1,2-Dichloroethane	LP/SD	584 ^b , 2400 566 ^b , 3500 551 ^b , 5300 433 ^a 418 ^b , 18000			747346
829.	8,9,10,11,12,13,14,15-Octahydro-5,7:16,18-dietheno-2,21-octanocyclopentadeca[1,2- α :1,15- α']diindene 2-MTHF (77 K)	PS/KM	397 ^a , 35900 ^a		Compound "X" in paper	83E383
830.	(all-E)-3,7,11,15,20,24,28,32-Octamethyl-1,34-bis(2,6,6-trimethyl-1-cyclohexen-1-yl)-1,3,5,7,9,11,13,15,17,19,21,23,25,27,29,31,33-tritriacontahexadecaene Benzene (296 K)	LP-ET	620		Triplet ET from chlorophyll <i>a</i> ; oxygen quenching (5.6×10^9 L mol ⁻¹ s ⁻¹); $\tau_T = 2.5$ μ s; $k_{et} = 1.8 \times 10^9$ L mol ⁻¹ s ⁻¹	73E347
831.	Orotate ion Water	FP/SD	315, 22000		\uparrow Oxygen quenching; pK_a 9.4; $\tau_T = 137 \pm 10$ μ s; pH 7.0	71E367
832.	Orotic acid Water	FP/SD	310, 11000 280, 14000		\uparrow Oxygen quenching; pK_a 4.6; $\tau_T = 137 \pm 10$ μ s; pH 3.0	71E367
833.	7-Oxa-2,3-dibenzoylbicyclo[2.2.1]hepta-2,5-diene Benzene	LP/RA	410 \pm 5, 5640 \pm 1400		\uparrow Oxygen and azulene quenching; ϵ relative to benzophenone in benzene ($\epsilon_{332} = 7600$ L mol ⁻¹ cm ⁻¹), assuming $\Phi_T = 1$ for benzophenone and taking $\Phi_T = 0.70$ for the compound in benzene; $\tau_T = 0.45 \pm 0.07$ μ s	84A355
	MeOH	LP	405 \pm 5		\uparrow Oxygen quenching; $\tau_T = 0.41 \pm 0.06$ μ s	84A355
834.	7-Oxa-2,3-dibenzoylbicyclo[2.2.1]hept-2-ene Benzene	LP/RA	450 \pm 5 ^b , 2700 \pm 700 342 ^a		\uparrow Oxygen and azulene quenching; ϵ relative to benzophenone in benzene ($\epsilon_{332} = 7600$ L mol ⁻¹ cm ⁻¹), assuming $\Phi_T = 1$ for benzophenone and taking $\Phi_T = 0.74$ for the compound in benzene; $\tau_T = 1.4 \pm 0.2$ μ s	84A355
835.	Oxonine cation MeOH	FP-ET/SD	750 ^a , 15000 ^a 675 ^a , 5000 ^a		\uparrow Triplet ET from 9,10-dibromoanthracene, oxygen quenching; ϵ assumes triplet does not absorb where singlet depletion is followed; $\tau_T = 140$ μ s; pH Basic	767246
836.	Oxonine cation, conjugate monoacid MeOH	FP-ET/SD	650 ^a , 16000 ^a 380 ^a , 13000 ^a		\uparrow Triplet ET from 9,10-dibromoanthracene; ϵ assumes triplet does not absorb where singlet depletion is followed; $\tau_T = 55$ μ s; pH Acidic	767246
837.	Oxotitanium(IV) tetraphenylporphyrin EtOH	LP	465		\uparrow Oxygen quenching; delay 200 ns; triplet species likely to be in equilibrium with EtOH-coordinated triplet; $\tau_T = 35 \pm 1$ μ s	84B008
838.	Palladium(II) etioporphyrin I DMF	FP/SD	410, 70000 365, 51000		\uparrow Phosphorescence decay; $\tau_T = 300$ μ s	73E345
839.	Palladium(II) tetrabenzoporphyrin Pyridine	FP	480 385		\uparrow Phosphorescence decay; $\tau_T = 190$ μ s	73E345

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
840.	Palladium(II) tetrakis(carboxyphenyl)porphyrin Water	LP	850 720			82E622
841.	Palladium(II) tetrakis(4-N-methylpyridyl)porphyrin Water	LP/TD	940, 6400 ± 600 470 ^b , 33400 ± 3300 450 ^b , 52000 ± 5200		Shoulder at 820 nm	82E622
842.	Palladium(II) tetrakis(p-sulfonatophenyl)porphyrin Water	LP/TD	830, 7600 ± 800 460 ^b , 28000 ± 2800		Shoulder at 700 nm	82E622
843.	Palladium(II) tetraphenylporphyrin MCH	FP/TD	450, 47000 385, 30000		†Oxygen quenching; $\tau_T = 380 \mu s$	81E271
	Pyridine	FP/SD	450, 47800 385, 35000			73E345
844.	[2.2]Paracyclophane 2-MTHF (77 K)	LP	550 380 340		Most intense peak at 380 nm; polarization also measured	79B050
	2-MTHF (77 K)	PS	380		Solvent uncertain; polarization also measured; $\tau_T = > 10^6 \mu s$	81F201
845.	Pentacene 1-Chloronaphthalene	FP/TD	505, 9900			727073
	Benzene (323 K)	FP/TD	505, 120000 472, 48000 392, 3000 374, 6500 340, 6900 305, 560000		Peak at 305 nm was the values measured in cyclohexane; shoulders at 440, 418, 355, and 292 nm, with the 292 nm figure for cyclohexane; 5 electronic transitions were assigned with 0-0 bands at 505 nm (³ B _{1g}), 392 nm (³ A _{1g}), 374 nm (³ B _{1g}), 340 nm (³ B _{1g}), and 305 nm (³ B _{1g}); T ₁ was ³ B _{1g} ; $\tau_T = 69.9 \pm 3.4 \mu s$; oscillator strength = 0.7, <0.005, 0.03, <0.005, 2.1	727348
	Hexane	FP/SD	525, 30000 490, 205000 457, 107000 383, 60000 305, 630000		Bands were assigned to 4 separate electronic transitions; bands at 490 and 457 nm were assigned to the 2nd electronic transition; oscillator strength = 0.2, 2.4, 1.4, 2.0	58E001
	Hexane	FP	492 460 385 300		Relative intensities (3:2:1:10); $\tau_T = 110 \pm 12 \mu s$	61E005
846.	Pentacyclo[18.2.2.2 ^{9,12} .0 ^{6,15} .0 ^{6,17}]hexacos-4,6(17),9,11,15,20,22,23,25-nonane-10,25-dicarbonitrile Acetonitrile	LP	775 ^a 505 ^a		†Triplet ET to piperylene, rises with fluorescence decay, oxygen quenching; 200 ns delay; "layered cyclophane"	776187
	Toluene	LP	760 ^a 525 ^a		†Triplet ET to piperylene, rises with fluorescence decay, oxygen quenching; 200 ns delay	776187
847.	Pentahelicene MCH/Isohexane	FP	630 420 ^a		†Rise time identical to fluorescence decay, oxygen quenching ($5.7 \times 10^8 \text{ L mol}^{-1} \text{ s}^{-1}$); relative intensities (3:1); solvent was 2:1 MCH to isohexane; $\tau_T = 0.31 \pm 0.02 \mu s$	79A237
	MCH/Isohexane (93 K)	FP	630 420 ^a		†Phosphorescence decay, ODMR signal; relative intensities (3:1); solvent was 2:1 MCH to isohexane; $\tau_T = 2 \times 10^6 \mu s$	79A237
848.	Pentaphene Cyclohexane	LP/SD	493, 45900 ^a 440, 37300 ^a		Delay 100 ns	84E390

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
849.	<i>cis</i> -Perinaphthothioindigo					
	MCH/Toluene (93 K)	LP	661 ^a		Glass was 1:1 MCH to toluene; shoulder at 715 ^a ; 661 nm peak was the more intense	82F295
	MCH/Toluene (93 K)	LP	590 ^a			
	MCH/Toluene (93 K)	LP	641 ^a		Shoulder at 716 ^a nm; glass was 1:1 MCH to toluene; $\tau_T = \sim 10 \mu s$	82A190
	MCH/Toluene (93 K)	LP	591 ^a			
850.	<i>trans</i> -Perinaphthothioindigo					
	MCH/Toluene (93 K)	LP	733 ^a		Glass was 1:1 MCH to toluene	82F295
	MCH/Toluene (93 K)	LP	733 ^a		†Oxygen quenching at 295 K; glass was 1:1 MCH to toluene; $\tau_T = \sim 10 \mu s$	82A190
851.	1,4-Perinaphthylidyl					
	EPA (77 K)	PS	512 ^a		†ESR, T ⁻¹ ESR signal intensity; triplet 0-0 bands only; apparent triplet ground state; spectrum probably contains singlet transitions	79E175
			345 ^a			
			250 ^a			
852.	Perylene					
	Benzene	PR/ET	490, 14300		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700 \text{ L mol}^{-1} \text{ cm}^{-1}$); ϵ obtained from a simultaneous least squares fit of data from several compounds making use of cyclohexane to benzene ϵ_{\max} ratios of 1.83 for naphthalene and 1.45 for anthracene	71E360
	Benzene	LP-ET	500		Triplet ET from 1-nitronaphthalene; $k_{et} = 4.2 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	747236
	Cyclohexane	LP/ET	460			
			480, 13000		Triplet ET from anthracene; ϵ relative to anthracene in cyclohexane ($\epsilon_{425} = 64700 \text{ L mol}^{-1} \text{ cm}^{-1}$)	777391
	Hexane	FP	485.9			54E001
	Hexane	FP	488		The two bands were assigned to different electronic transitions	58E001
			280			
	Polystyrene	PR	495		$G\epsilon = 6500 \text{ L mol}^{-1} \text{ cm}^{-1} / (100 \text{ eV absorbed})$; half-life = 6000 μs	672086
853.	Perylene/Chloranil					
	Liquid paraffin	MOD	475		Mull	71E361
	PMMA	PS	595		Charge transfer complex; most intense peak at 480 nm	766652
			580			
			563			
			544			
			505			
			480			
			463			
854.	Phenanthrene					
	1-BuOH/Isopentane (77 K)	PS	493.1		A distinction between a 1:7 and a 3:7 glass was not made; all 3 vibronic bands were assigned to the $^3B_2 \leftarrow ^3A_1$ transition	63B001
			460.2			
			131.0			
	3-MP (77 K)	PS/ESR	490, 41500		†ESR; only most intense visible peak reported	696115
	Benzene	PR/ET	492.5, 15700		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700 \text{ L mol}^{-1} \text{ cm}^{-1}$); ϵ obtained from a simultaneous least squares fit of data from several compounds making use of cyclohexane to benzene ϵ_{\max} ratios of 1.83 for naphthalene and 1.45 for anthracene	71E360
	Benzophenone (77 K)	PS/COM	500 ^a , 2100 ^c \pm 1000		Mixed crystals; shoulders at 495 ^a , 488 ^a , 478 ^a , 469 ^a , and 455 ^a nm; ϵ measured by a technique involving reabsorption of phosphorescence	68E106
			474 ^a , 1200 ^{a,c}			
			467 ^a , 1300 ^{a,c}			
			441 ^a , 500 ^a			
			435 ^a , 500 ^a			
	Butane/Isopentane (77 K)	PS/IV	492.5, 27000 \pm 5000		Glass was 3:7 butane to isopentane; ϵ_T was computed from $OD = \epsilon_T \tau_{pl_0}(1 - \Phi_f)$ after the optical density, OD, was extrapolated to zero intensity of the monitoring beam (namely $l_0 \rightarrow 0$), here τ was lifetime of triplet, and Φ_f was fluorescence quantum yield; oscillator strength = 0.11	65E031
			459 ^a , \sim 11000			

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Cyclohexane	FP	480		$\tau_T = 145 \mu\text{s}$	62E009
	Cyclohexane	PR/ET	482.5, 21000 \pm 5250		ϵ relative to benzophenone ketyl radical in water ($\epsilon_{537.5} = 3220 \text{ L mol}^{-1} \text{ cm}^{-1}$), assuming this value for $\epsilon_{542.5}$ in cyclohexane	680727
	Cyclohexane	LP	485 460 425		†Rise time of transient was the same as the decay time of the singlet	68B006
	Cyclohexane	PR/ET	482.5, 25200		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700 \text{ L mol}^{-1} \text{ cm}^{-1}$); reference ϵ obtained by starting from $\epsilon_{\max} = 3220 \text{ L mol}^{-1} \text{ cm}^{-1}$ for this ketyl radical in water and assuming the f of the ketyl radical is independent of solvent; final ϵ obtained from a simultaneous least squares fit to data from several compounds	71E360
	Dipropyl ether (100 K)	FP	493 ^a 463 ^a 437 ^a		Shoulder at 474 ^a nm; $\tau_T = 3.8 \times 10^6 \mu\text{s}$	696073
	Dipropyl ether (110 K)	PS	498 ^a 465 ^a 439 ^a		Relative intensities (4:2:1); polarization also measured	70B002
	EPA (77 K)	PS/IV	493.1, 1560 ^c \pm 190 459.6, 880 ^c \pm 80		†Phosphorescence decay; ϵ 's are too low; 1st transition assigned as $^3L_a \rightarrow ^3C_0$	51E001
	EPA (77 K)	PS	493.1 460.2 431.0		†Phosphorescence decay; relative intensities (100:45:14); $\tau_T = (3.3 \pm 0.2) \times 10^6 \mu\text{s}$	54B001
	EPA (77 K)	PS	828 735 656 599 493		†Phosphorescence decay; $\tau_T = (3.9 \pm 0.8) \times 10^6 \mu\text{s}$	67E106
	EPA (77 K)	PS/KM	499 ^a , 18500 ^a 466 ^a , 9200 ^a 431 ^a , 2800 ^a		Solvent, temperature and extinction method assumed from earlier work; polarization also measured	69E213
	EPA (77 K)	FP/SD	492.5, 38000 \pm 6000 461 ^a , 17000 ^a 431 ^a , 5700 ^a		$\tau_T = 3.4 \times 10^6 \mu\text{s}$; oscillator strength = 0.15	69F389
	EPA (77 K)	PS/KM	493, 20400		λ_{\max} assumed from previous work	69E212
	EtOH (293 K)	FP	482			68E098
	EtOH (77 K)	MOD/KM	489, 38000 \pm 9500 457.5, 16000 \pm 4000 428, 5400 \pm 1300 293, 13000 \pm 3300			737055
	EtOH (77 K)	MOD	490		Halfwidth 550 cm ⁻¹	777538
	EtOH	LP	480 ^a 450 ^a		Relative intensities (3:2); solution contains Ag ions; 1.8 μs delay	78E554
	EtOH	LP	482 ^a 452 ^a 427 ^a		Solvent absolute EtOH	82B052
	EtOH/MeOH (104 K)	PS	489 459		Solvent was 3:1 EtOH to MeOH; polarization also measured	68E102
	EtOH/MeOH (101 K)	PS	490 ^a 459 ^a		Glass was 3:1 EtOH to MeOH	75B002
	Ethylene glycol	FP	485 453 428 400		Relative intensities (100:60:25:10); $\tau_T = 910 \pm 83 \mu\text{s}$	61E005

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Heptane (77 K)	PS	829 723 650 496 492 485 474 465 455 443 435 428 405			69E211
	Hexamethylbenzene (130 K)	PS	496 465		Single crystal; relative intensities (100:61)	67B008
	Hexane	FP	481.0 452.9 425.0			54E001
	Hexane	FP	480 450 428 399		Relative intensities (100:60:25:10); $\tau_T = 93 \pm 7 \mu s$	61E005
	Hexane (77 K)	PS	497.5 485.9 471.9 460.0 435.9 430.1 410.0		Relative intensities (10:4:2:)	69E211
	Liquid paraffin	FP	480		Viscosity of solvent was 0.167 N·s/m ² ; $\tau_T = 2400 \mu s$	62E009
	Liquid paraffin	FP/TD	480 ^b , 24000 \pm 2000			67E031
	PMMA (293 K)	PS	488.8 456 426		Relative intensities (100:55:10)	67B008
	PMMA	LP	520 510 481 454			70E288
	PMMA (15 K)	PS	816 720 ^a 489			81B094
	Polystyrene	PR	495 ^a 458 ^a		$G_{\epsilon_{495}} = 11000^a$ L mol ⁻¹ cm ⁻¹ / (100 eV absorbed); $G_{\epsilon_{458}} = 9200^a$ L mol ⁻¹ cm ⁻¹ / (100 eV absorbed)	70I073
	Polyvinylbutyral (77 K)	LP	493 460 430 ^a			80E867
	Polyvinylbutyral (77 K)	FP	493 460 425 ^a		Relative intensities (5:2:1); polymer film; $\tau_T = 4.4 \times 10^6 \mu s$	82E129
	SDS	LP	490 ^a 460 ^a 420 ^a 400 ^a		Most intense peak at 490 nm; aqueous micelle	81N016
	Toluene (77 K)	MOD	500		Halfwidth 670 cm ⁻¹	77F538
855.	Phenanthrene- <i>d</i> ₁₀					
	3-MP (77 K)	PS	825 727 650 594			67E106
	3-MP (77 K)	PS/ESR	489, 42900 457 ^a , 19100 ^a 429 ^a , 7900 ^a		†ESR; band was assigned as ³ A _g ← ³ B ₂ ; oscillator strength = 0.23	68D211

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
3-MP (77 K)	PS		490 ^a 456 ^a 429 ^a			69E210
3-MP (77 K)	PS		827 ^a 728 ^a 648 ^a 586 ^a 495		Band at 495 nm assigned as the 0-0 of the $^3A_1^- \leftarrow ^3B_2^+$ and was the most intense band, ~ 4 times as strong as the bands to the red; 495 nm was the strongest vibronic band; shoulder at 426 ^a nm	74B002
? (77 K)	PS		454 ^a 488 459 432 ^a		†Phosphorescence decay and ESR lifetime; glass used was either 2-MTHF or 3-MP; $\tau_T = 1.36 \times 10^7$ μ s	67E108
Biphenyl (4 K)	PS		498 ^a		Maximum is taken from the c'-axis absorption spectrum	67B008
Cyclohexane	LP		520 510 481 454			70E288
EPA (77 K)	PS/RA		830, 675 734 658 598 493 459		†Phosphorescence decay; ratio of intensity at 493 nm to that at 830 nm was 40; ϵ relative to phenanthrene- <i>h</i> ₁₀ in 3:7 butane/isopentane glass at 77 K ($\epsilon_{492.5} = 27000$ L mol ⁻¹ cm ⁻¹); the 830 nm transition was assigned as $^3L_0 \leftarrow ^3L_{01}$; $\tau_T = (1.63 \pm 0.15) \times 10^7$ μ s	67E106
EPA (82 K)	PS/KM		487, 20400 \pm 1100			68B003
EPA (77 K)	PS		493		Transition assigned as $^3A_1^- \leftarrow ^3B_2^+$	69B001
EPA (77 K)	FP/SD		492.5, 31000 \pm 5000		$\tau_T = 1.22 \times 10^7$ μ s	69F389
EPA (77 K)	PS/KM		823 ^a , 1000 ^a 723 ^a , 1400 ^a 651 ^a , 1100 ^a 597 ^a , 960 ^a 495, 20400 460 ^a , 9500 ^a 426 ^a , 2800 ^a		Two electronic transitions were assigned, the 2nd starting at 495 nm; shoulder at 926 ^a nm; ϵ -method assumed on basis of earlier work by authors, oscillator strength = 0.02, 0.15	747347
Heptane (77 K)	PS		830 770 725 650 497 494 487 475 467.5 457 443 435 426 409 400		Relative intensities (14:13:19:12:100:88:59:35:43:30:19:24:20:11:10)	69F211
Hexamethylbenzene (130 K)	PS		495 465		Single crystal; relative intensities (100:65)	67B008
Hexane	PS		830 770 725 650		Weak bands in the red only studied	67B005
PMMA (293 K)	PS		486.6 454 424		Relative intensities (100:65:10)	67B008
856. Phenanthrene/Pyromellitic dianhydride Dipropyl ether (100 K)	FP		493 ^a 437 ^a			696073

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
857.	Phenanthrene/Tetrachlorophthalic anhydride Dipropyl ether (100 K)	FP	498 ^a 469 ^a		Shoulder at 435 ^a nm; $\tau_T = 1.1 \times 10^6 \mu s$	696073
858.	Phenanthrene- <i>d</i> ₁₀ /Tetrachlorophthalic anhydride EPA (77 K)	PS	492 458 431			67B009
859.	Phenanthridine EtOH/Et ₂ O (77 K)	PS	735 660 568 529 493 463 439		Solvent was 3:2 EtOH to Et ₂ O; relative intensities (17:25:70:82:84:100:65); $E_T = 265 \text{ kJ mol}^{-1}$	80B130
	PMMA (293 K)	FP	440		$\tau_T = 2.5 \times 10^4 \mu s$	70E291
	PMMA (77 K)	PS	560 520 465			70E291
860.	6(5 <i>H</i>)-Phenanthridinone EtOH (300 K)	FP	425			81E649
861.	1,10-Phenanthroline 2-PrOH	FP	445 422		\uparrow Oxygen quenching, pentadiene quenching; $\tau_T = 33 \pm 2 \mu s$	777201
	Benzene	PR	440 420		$\tau_T = 3.3 \pm 0.6 \mu s$	82A259
	Cyclohexane	FP	442 422		\uparrow Oxygen quenching, pentadiene quenching; $\tau_T = 26 \pm 3 \mu s$	777201
	EtOH	FP	445 425		\uparrow Oxygen quenching, pentadiene quenching; $\tau_T = 35 \pm 4 \mu s$	777201
	EtOH/Et ₂ O (77 K)	PS	450 427		Solvent was 3:2 EtOH to Et ₂ O; relative intensities (100:93); $E_T = 264 \text{ kJ mol}^{-1}$	80B130
	EtOH/MeOH	LP	434 ^a		Solvent was 4:1 EtOH to MeOH; shoulders at 456 ^a and 410 ^a nm	81E786
	Heptane	FP	440 420		\uparrow Oxygen quenching, pentadiene quenching; $\tau_T = 19 \pm 2 \mu s$	777201
	Isopentane	FP	440 420		\uparrow Oxygen quenching, pentadiene quenching; $\tau_T = 17 \pm 2 \mu s$	777201
	PMMA (77 K)	PS	450 421		$\tau_T = 8.7 \times 10^5 \mu s$	70E291
	PMMA (293 K)	FP	~440		Decay was nonexponential, and lifetime was calculated from a terminal 1st-order rate constant; $\tau_T = 4.3 \times 10^5 \mu s$	70E291
	Water	FP	450 425		\uparrow Oxygen quenching, pentadiene quenching; $\tau_T = 42 \pm 4 \mu s$	777201
862.	1,7-Phenanthroline EtOH/Et ₂ O (77 K)	PS	476 441		Solvent was 3:2 EtOH to Et ₂ O; relative intensities (92:100); $E_T = 267 \text{ kJ mol}^{-1}$	80B130
863.	1,8-Phenanthroline EtOH/Et ₂ O (77 K)	PS	481 457 426		Solvent was 3:2 EtOH to Et ₂ O; relative intensities (87:100:52); $E_T = 262 \text{ kJ mol}^{-1}$	80B130
864.	1,9-Phenanthroline EtOH/Et ₂ O (77 K)	PS	482 454 426		Solvent was 3:2 EtOH to Et ₂ O; relative intensities (100:65:43); $E_T = 262 \text{ kJ mol}^{-1}$	80B130

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
865.	4,7-Phenanthroline EtOH/Et ₂ O (77 K)	PS	488 458 435		Solvent was 3:2 EtOH to Et ₂ O; relative intensities (100:66:32); $E_T = 267$ kJ mol ⁻¹	80B130
866.	Phenazine Benzene	FP	440		$\tau_T = \sim 130$ μ s	717154
	CF ₃ CH ₂ OH	FP/SD	456 ^a , 14100 ^a 355 ^a , 36900 ^a		†Oxygen quenching; no absorption could be seen in the 360 - 400 nm region, possibly due to SD masking it; lifetime was concentration dependent, and the value quoted was the longest measured and was measured at the lowest concentration; $\tau_T = 170$ μ s	716169
	Cyclohexane	FP	511 ^a 445 ^a		Shoulder at 434 ^a nm; 445 nm was the more intense peak; delay 30 μ s	707337
	Isooctane	LP	500 ^a 440		Relative intensities (1:2); 50 ps delay; shoulders at 465, 455 and 425 nm; rise time of 140 ns	766469
	MCH	FP/SD	440 ^a , 15100 ^a 378 ^a , 36700 ^a 356 ^a , 38400 ^a		†Oxygen quenching; ϵ 's at wavelengths shorter than 420 nm were more uncertain than the 440 nm band because of the SD region; $\tau_T = 42 \pm 3$ μ s	716169
	MeOH	FP	508 ^a 447 ^a 391 ^a		Shoulder at 441 ^a nm; 447 nm was the most intense peak; delay 65 μ s	707337
	MeOH	LP	500 ^a 440		Relative intensities (1:2); rise time of 200 ns	766469
	PMMA (293 K)	FP	460 440 426		$\tau_T = 9000$ μ s	70E291
	Toluene (293 K)	FP	515 ^a 430 ^a		†Phosphorescence decay in EPA at 77 K, oxygen quenching	80E778
	Water	FP	506 ^a 445 ^a		Shoulder at 416 ^a nm; 445 nm was the more intense peak; pH ~ 6	707337
867.	Phenazine, conjugate monoacid MeOH	FP	460		Solution was 0.1 mol L ⁻¹ acetic acid and 0.1 mol L ⁻¹ sodium acetate; delay 60 μ s; pK_a 3.8; $\tau_T = 340 \pm 40$ μ s	687213
868.	Phenol Water (298 K)	LP	~ 250		Lifetime was measured at pH 7.5; spectrum is the difference between spectra at 20 ns and 15 μ s; $\tau_T = 3.3$ μ s; pH 7.7	757161
869.	Phenosafranin Water	FP	664 ^a		Any maxima in the range of 400 - 600 nm were masked by SD; pH 3-5	697061
870.	Phenothiazine 3-MP (77 K)	PS	465			67E106
	EPA (77 K)	PS	467			67E106
	MeOH	LP/ELT	460, 27000		ϵ relative to phenothiazine radical cations in MeOH ($\epsilon_{520} = 9300$ L mol ⁻¹ cm ⁻¹); ELT measurement used Eu ⁺³ ions as electron acceptors; 2nd-order rate constant in ELT = 4.7×10^9 L mol ⁻¹ s ⁻¹	757353
871.	Phenoxazine 3-MP (77 K)	PS	460		$\tau_T = (2.7 \pm 0.3) \times 10^6$ μ s	67E106
	Benzene	FP	465		†Phosphorescence decay in PMMA at 90 K; $\tau_T = 26$ μ s	707186
	EPA (77 K)	PS	458		†Phosphorescence decay; $\tau_T = (2.7 \pm 0.2) \times 10^6$ μ s	67E106
	EtOH	FP	465		†Phosphorescence decay in PMMA at 90 K; $\tau_T = 44$ μ s	707186
	Hexane	FP	465		†Phosphorescence decay in PMMA at 90 K; $\tau_T = 26$ μ s	707186
	MCH	FP	465		†Phosphorescence decay in PMMA at 90 K; $\tau_T = 32$ μ s	707186

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{max} , /nm	ϵ_{max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
872.	3-Phenylacetylacetone EtOH/MeOH (118 K)	FP	490		Solvent was 3:1 EtOH to MeOH	68B005
873.	Phenylacetylene Isopentane/2-PrOH (77 K)	PS	255 ^a		†ESR; glass was 7:3 isopentane to 2-propanol by volume; by assuming that radical formation occurs only from T* and that it is wavelength-independent, the author devise an ESR method that determines the shape of TTA even in the SD region	68B007
874.	9-Phenylacridan EtOH (97 K)	FP/?	520, 10000 ^a		Shoulder at 625 nm	697225
875.	N-Phenylacridan EtOH (93 K)	PS	525 ^a			69E214
876.	Phenylalanine Water (298 K)	LP	316 ^a 244 ^a		Spectrum is the difference between spectra at 10 μ s and 20 ns; $\tau_T = 2.4 \pm 0.2 \mu$ s; pH 0.3	757162
	Water (298 K)	LP	~ 310 ~ 240		†Oxygen quenching; spectrum is the difference between spectra at 10 μ s and 20 ns; $\tau_T = 3.1 \pm 0.3 \mu$ s; pH 7.5	757162
877.	4-(Phenylamino)-7H-benz[de]anthracen-7-one Benzene	LP	638 ^a 590 ^a		†Oxygen quenching (1.7×10^9 L mol ⁻¹ s ⁻¹); half-life = $\sim 0.28 \mu$ s	757427
878.	9-Phenylanthracene 2-PrOH	FP	428 ^a			65F031
	EtOH	FP	430		†Quenching by heavy atom containing molecules	67E109
	EtOH	FP/ET	440 ^b , 9100 428, 13500 424 ^b , 12500		Triplet ET from eosin and proflavine; ϵ relative to eosin in EtOH ($\epsilon_{580} = 9400$ L mol ⁻¹ cm ⁻¹) and proflavine in EtOH ($\epsilon_{550} = 11000$ L mol ⁻¹ cm ⁻¹)	716235
	EtOH	FP-ET/RA	694 ^b , 28		ϵ relative to 9-phenylanthracene in EtOH ($\epsilon_{430} = 14000$ L mol ⁻¹ cm ⁻¹)	78E019
	Ethylene glycol	FP	432		†Quenching by heavy atom containing molecules	67E109
	Liquid paraffin	FP	434			65F031
879.	4-Phenylbenzophenone Alcohol/Ether (77 K)	MOD	540		Glass was 2:1 alcohol to ether	76E682
880.	9-Phenyl-9H-9-bisfluorene MCH/Isopentane (77 K)	FP	387 ^a		†Phosphorescence lifetime; glass was 3:1 MCH to isopentane; shoulder at 369 ^a nm; $\tau_T = 1000 \mu$ s	81E648
881.	1-Phenylcyclohexene Cyclohexane	LP-ET/RA	325 ^b , 4400		ϵ relative to naphthalene in cyclohexane ($\epsilon_{413} = 24000$ L mol ⁻¹ cm ⁻¹), but in ref. cited ($\epsilon_{415} = 24500$ L mol ⁻¹ cm ⁻¹); triplet ET from thioxanthone; $\tau_T = \sim 0.065 \mu$ s	82E181
882.	1-Phenyldeoxybenzoin Benzene	LP	425 338		†Oxygen quenching ($> 10^9$ L mol ⁻¹ s ⁻¹), piperylene quenching; relative intensities (1:3); $\tau_T = 0.27 \mu$ s	79A028
883.	2,2'-(1,4-Phenylene)bisbenzoxazole Toluene	LP/TD	480, 18600 \pm 250 420		Shoulders at 562 ^a , 528 ^a , 451 ^a , and 405 ^a nm; $\tau_T = 0.48 \pm 0.05 \mu$ s	82E632
884.	2,2'-(1,4-Phenylene)bis[5-(4-butoxyphenyl)oxazole] 2-MTHF (77 K)	CWL	590		"DibutoxyPOPOP"	74B003
885.	2,2'-(1,4-Phenylene)bis[5-phenyloxazole] 2-MTHF (77 K)	CWL	~ 500		"POPOP"	74B003
	Dioxane	LP	550 ^a 360 ^a		Weak very broad bands	80E438

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
886.	<i>O</i> -(2-Phenylethyl) 4-(dimethylamino)benzenecarbothioate Cyclohexane	LP/SD	450, 20000 ± 10000		†Oxygen quenching and triplet ET to tetracene; $\tau_T = 0.5 \mu\text{s}$	737318
887.	<i>O</i> -(2-Phenylethyl) 4-methoxybenzenecarbothioate Cyclohexane	LP	430		†Oxygen quenching; $\tau_T = 0.3 \mu\text{s}$	737318
888.	<i>O</i> -(2-Phenylethyl) 1-naphthalenecarbothioate Cyclohexane	LP	480		†Oxygen quenching and triplet ET to tetracene; $\tau_T = 1 \mu\text{s}$	737318
889.	<i>O</i> -(2-Phenylethyl) 2-naphthalenecarbothioate Cyclohexane	LP	440		†Oxygen quenching	737318
890.	2-Phenylindene Benzene (298 K)	LP-ET	~390		†Triplet ET from benzophenone, oxygen quenching ($4 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); $\tau_T = > 3 \mu\text{s}$; $k_{et} = 6 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	81E214
	Benzene (298 K)	FP	392		$\tau_T = > 100 \mu\text{s}$	81E214
	Benzene (298 K)	LP-ET	392		†Triplet ET from xanthone, oxygen quenching ($3.8 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); $\tau_T = > 5 \mu\text{s}$; $k_{et} = 6 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	81E214
	EPA (77 K)	FP	390 368 352		Most intense peak at 390 nm; $\tau_T = 40 \mu\text{s}$	737069
	EPA (298 K)	LP	380 360			81E214
891.	β -Phenyl-4'-methoxypropiophenone Benzene	LP	381 ^a			84E018
892.	1-[[[(Phenylmethyl)sulfonyl]methyl]naphthalene Acetonitrile	LP	430 ^a			84B022
893.	1-Phenylnaphthalene 3-MP (77 K)	PS/ESR	495, 18400 462 ^a , 10300 ^a 441 ^a , 5700 ^a		†ESR; shoulder at 418 nm; oscillator strength = 0.16	69B002
	Cyclohexane	LP/ET	480, 17600 ± 1000		†Triplet ET from TMPD; ϵ relative to TMPD in cyclohexane ($\epsilon_{570} = 11900 \text{ L mol}^{-1} \text{ cm}^{-1}$)	757282
894.	2-Phenylnaphthalene 3-MP (77 K)	PS/ESR	476, 2400 ^a 433, 34600 410 ^a , 19300 ^a 382 ^a , 12400 ^a		†ESR; oscillator strength = 0.31	69B002
	Cyclohexane	LP/ET	425, 48300 ± 5000		†Triplet ET from phenanthrene; ϵ relative to phenanthrene in cyclohexane ($\epsilon_{482.5} = 25200 \text{ L mol}^{-1} \text{ cm}^{-1}$)	757282
895.	<i>N</i> -Phenyl-2-naphthylamine EtOH (77 K)	MOD/KM	520, 9000 ± 2300 331, 7000 ± 1700 292, 21500 ± 5400			737055
896.	2-Phenyl-2-norbornene Cyclohexane	LP-ET/RA	325 ^b , 17900		ϵ relative to naphthalene in cyclohexane ($\epsilon_{413} = 24000 \text{ L mol}^{-1} \text{ cm}^{-1}$), but in ref. cited ($\epsilon_{415} = 24500 \text{ L mol}^{-1} \text{ cm}^{-1}$); triplet ET from thioxanthone; $\tau_T = > 2.0 \mu\text{s}$	82E181
897.	9-Phenyl-9 <i>H</i> -9-phosphafluorene MCH/Isopentane (77 K)	FP	387 ^a		†Phosphorescence lifetime; glass was 3:1 MCH to isopentane; shoulder at 378 nm; $\tau_T = 1.2 \times 10^6 \mu\text{s}$	81E648

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
898.	9-Phenylproflavine, conjugate monoacid EtOH/Et ₂ O (90 K)	PS/SD	1190, 41000 ^a 690, 8000 ^a 575, 15000 ^a 395, 12000 ^a 280, 47500 ^a 240, 18000 ^a		Solvent was 2:1 EtOH to Et ₂ O	57B001
899.	9-Phenyl-9H-9-stibafluorene MCH/Isopentane (77 K)	FP	378 ^a		†Phosphorescence lifetime; glass was 3:1 MCH to isopentane; $\tau_T = 1.2 \times 10^4 \mu s$	81E648
900.	1-Phenylthio-3,4-dihydronaphthalene MCH	FP	550 ^a 430 ^a		†Oxygen quenching; $\tau_T = 0.46 \mu s$	78A237
	MCH (110 K)	FP	560 ^a		†Oxygen quenching; $\tau_T = 2.0 \times 10^3 \mu s$	78A237
901.	Pheophytin a Benzene	FP	480			55E003
	Benzene	FP	500		430 nm band was judged to be of "doubtful significance" by the experimenters; benzene was wet	58R002
	EtOH (298 K)	FP/SD	430 687 ^a , 4700 ^a 667 ^a , 4700 \pm 500 528 ^a , 18100 ^a 407 ^a , 62800 ^a 332 ^a , 47100 ^a		$\tau_T = 750 \pm 100 \mu s$	70E296
	Pyridine	FP	430		Maximum was judged to be of "doubtful significance" by the experimenters	58R002
902.	Pheophytin b Benzene	FP	510 460 430		430 nm maximum was judged to be of "doubtful significance" by the experimenters; 460 nm was the most intense peak; benzene was wet	58R002
	EtOH (298 K)	FP/SD	667 ^a , 4700 \pm 500 481 ^a , 31400 ^a 423 ^a , 71200 ^a		Shoulders at 508 nm and 662 nm; $\tau_T = 1050 \pm 100 \mu s$	70E296
	Pyridine	FP	520 460 410		Maxima at 410 and 520 nm were judged to be of "doubtful significance" by the experimenters	58R002
903.	Photoprotoporphyrin isomer "A", dimethyl ester Benzene	LP/TD	475 ^b , 35500 471 ^a 420 ^a 349		†Oxygen quenching (2.3×10^9 L mol ⁻¹ s ⁻¹); relative intensities (8:4:5); $\tau_T = 83 \mu s$	80E200
	Benzene	PR/ET	475 ^b , 39200 471 ^a 420 ^a 349		†Oxygen quenching (2.6×10^9 L mol ⁻¹ s ⁻¹); ϵ relative to biphenyl in benzene ($\epsilon_{360} = 27100$ L mol ⁻¹ cm ⁻¹); relative intensities (8:4:5); $\tau_T = 83 \mu s$	80E200
904.	Photoprotoporphyrin isomer "B", dimethyl ester Benzene	PR/ET	477 ^a 475 ^b , 37000 411 ^a 349 ^a		†Oxygen quenching (2.1×10^9 L mol ⁻¹ s ⁻¹); ϵ relative to biphenyl in benzene ($\epsilon_{360} = 27100$ L mol ⁻¹ cm ⁻¹); peaks roughly of equal intensity; $\tau_T = 89 \mu s$	80E200
905.	Phthalazine 1-BuOH/Isopentane (77 K)	FP/SD	420.5, 4450 \pm 200		Glass was 3:7 1-BuOH to isopentane; oscillator strength = 0.11	726177
	Benzene	FP	396 ^a			747093
	EPA (77 K)	FP	403 ^a		†Phosphorescence decay	747093
	Et ₂ O	FP	387 ^a		Shoulder at 423 ^a nm	747093
	EtOH	FP	385 ^a		Shoulder at 425 ^a nm	747093
	EtOIL/Acetic acid	FP	387 ^a			747093
	Hexane	FP	389 ^a		Shoulder at 427 ^a nm	747093

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
906.	Phthalocyaninatobis(pyridine)ruthenium(II) Methylene chloride	LP	~500			82C018
907.	Phthalocyanine 1-Chloronaphthalene	LP/TD	480, 28500 ± 2000		†Triplet ET to β -carotene; authors prefer this value to those obtained by either SD or E1 methods; $\tau_T = 130 \pm 10 \mu\text{s}$; $E_T = 120 \pm 10 \text{ kJ mol}^{-1}$; $k_{et} = 1.25 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	78A378
	1-Chloronaphthalene	PR/ET	480, 36000			78A378
	1-Chloronaphthalene	LP/SD	480, 38300		†Authors reject this ϵ value since residual triplet and excited singlet absorptions are neglected	78A378
	1-Chloronaphthalene	LP/TD	480, 37000 ± 4000		ϵ method inferred from text	81E457
	1-Chloronaphthalene	LP/SM	480, 18000 ± 2000		†Triplet ET to β -carotene; $\tau_T = 125 \mu\text{s}$	81E457
908.	15-cis-Phytoene Cyclohexane	LP-ET/ET	320, ~20000		†Triplet ET from fluorenone; ϵ relative to fluorenone ($\epsilon_{415} = 5500 \text{ L mol}^{-1} \text{ cm}^{-1}$), no solvent specified for this unpublished relative reference; $\tau_T = \sim 1 \mu\text{s}$; $E_T = 176\text{--}201 \text{ kJ mol}^{-1}$; $k_{et} = \sim 1 \times 10^{10} \text{ L mol}^{-1} \text{ s}^{-1}$	761035
909.	all-trans-Phytoene Cyclohexane	LP-ET/ET	320, ~20000		†Triplet ET from naphthalene; ϵ relative to naphthalene in cyclohexane ($\epsilon_{415} = 24500 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = \sim 1 \mu\text{s}$; $E_T = 176\text{--}201 \text{ kJ mol}^{-1}$; $k_{et} = \sim 1 \times 10^{10} \text{ L mol}^{-1} \text{ s}^{-1}$	761035
910.	Picene 2-MTHF (77 K)	PS/ESR	636, 75500 575 ^a , 20100 ^a 535 ^a , 11000 ^a 467 ^a , 5300 ^a		†ESR; the band was assigned as $^3A_1 \leftarrow ^3B_2$; oscillator strength = 0.40	68D211
	Benzene	MOD	560		$\tau_T = 160 \mu\text{s}$	71E361
	Benzene	LP	685 667 654 629 571 552 535 515 493 452		Delay 370 ns	737463
	EPA (77 K)	PS/SD	630, 62000 ± 6200			68E105
	EPA (77 K)	PS/KM	623 ^a , 21300 ^a 575 ^a , 7200 ^a 533 ^a , 3700 ^a		Solvent, temperature and extinction method assumed from earlier work; polarization also measured	69E213
	EPA (77 K)	PS/IV	620, 23000		λ_{\max} assumed from previous work; ϵ estimated by extrapolation to infinite excitation rate	69E212
	EtOH (77 K)	MOD	622		Halfwidth 570 cm ⁻¹	777538
	Toluene (77 K)	MOD	631		Halfwidth 570 cm ⁻¹	777538
911.	2-Piperidinoanthraquinone Benzene	PR/ET	581 ^a , 14400 ^a		†Triplet ET from triplet donors and triplet ET to triplet acceptors; ϵ relative to naphthalene in benzene ($\epsilon_{425} = 13200 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = \sim 20 \mu\text{s}$	720392
912.	Pivalophenone 2-PrOH	LP	500 ^a 450 ^a 400 ^a 360		†Triplet ET to naphthalene; relative intensities (1:2:3:5); maxima inferred from text; $\tau_T = 0.5 \mu\text{s}$	79P066
	Benzene	LP	500 ^a 450 ^a 400 ^a 335		†Triplet ET to naphthalene; relative intensities (1:2:3:5); $\tau_T = 0.37 \pm 0.03 \mu\text{s}$; $k_{et} = (2.0 \pm 0.4) \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	79P066

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
913.	Poly[oxy[2-(1-pyrenylmethyl)-1,3-propanediyl]oxy(1,4-dioxo-1,4-butanediyl)] DMF	LP	425 ^a		†Rises with fluorescence decay, oxygen quenching	80B076
914.	Poly[oxy[2-(1-pyrenylmethyl)-1,3-propanediyl]oxy(1,4-dioxo-1,4-butanediyl)]/1,4-Dicyanobenzene DMF	LP	425 ^a		†Rises with fluorescence decay, oxygen quenching	80B076
915.	Poly[oxy[2-(1-pyrenylmethyl)-1,3-propanediyl]oxy(1,4-dioxo-1,6-hexanediyl)] DMF	LP	430 ^a		†Rises with fluorescence decay, oxygen quenching	80B076
916.	Poly[oxy[2-(1-pyrenylmethyl)-1,3-propanediyl]oxy(1,4-dioxo-1,6-hexanediyl)]/1,4-Dicyanobenzene DMF	LP	435 ^a		†Rises with fluorescence decay, oxygen quenching	80B076
917.	Poly(phenylisopropenyl ketone) Benzene	LP	450		†Triplet ET to naphthalene, oxygen quenching (6×10^8 L mol ⁻¹ s ⁻¹); $\tau_T = 0.1$ μ s; $k_{et} = (1.5 \pm 1) \times 10^9$ L mol ⁻¹ s ⁻¹	79P066
918.	Poly(4-vinylbenzophenone) Benzene	LP	535		$\tau_T = 2.27$ μ s	79A171
919.	Poly(2-vinylnaphthalene) Benzene	PR/ET	426 ^a , 12000		†Triplet ET from biphenyl; ϵ relative to biphenyl in benzene ($\epsilon_{367} = 27100$ L mol ⁻¹ cm ⁻¹); degree of polymerization = 318 (monomer units); $\tau_T = 13.2$ μ s; $k_{et} = 1.8 \times 10^8$ L mol ⁻¹ s ⁻¹	79E666
	Benzene	PR/ET	426 ^a , 14000		†Triplet ET from biphenyl; ϵ relative to biphenyl in benzene ($\epsilon_{367} = 27100$ L mol ⁻¹ cm ⁻¹); degree of polymerization = 2690 (monomer units); $\tau_T = 10.6$ μ s; $k_{et} = 0.8 \times 10^8$ L mol ⁻¹ s ⁻¹	79E666
	Benzene	PR/ET	426 ^a , 11000		†Triplet ET from biphenyl; ϵ relative to biphenyl in benzene ($\epsilon_{367} = 27100$ L mol ⁻¹ cm ⁻¹); degree of polymerization = 688 (monomer units); $\tau_T = 12.2$ μ s; $k_{et} = 1.4 \times 10^8$ L mol ⁻¹ s ⁻¹	79E666
	Benzene	PR/ET	426 ^a , 11000		†Triplet ET from biphenyl; ϵ relative to biphenyl in benzene ($\epsilon_{367} = 27100$ L mol ⁻¹ cm ⁻¹); degree of polymerization = 109 (monomer units); $\tau_T = 9.7$ μ s; $k_{et} = 3.4 \times 10^8$ L mol ⁻¹ s ⁻¹	79E666
920.	21H,23H-Porphine Dimethyl phthalate	FP/SD	752 ^a , 4890 ^a 419 ^a , 98600 ^a 384 ^a , 35400 ^a		Shoulders at 795 nm and 440 nm	74B007
921.	Proflavine EtOH Water (296 K)	FP/SD FP/TD	550, 11000 549, 8000 \pm 800		pH Basic	716235 81E147
922.	Proflavine, conjugate diacid SDS	LP	1160 560		†Oxygen quenching; 1 μ s delay; aqueous micelle; $pK_a = 4.0$; cation radical (λ_{\max} 820 nm) also formed; $\tau_T = 7$ μ s; pH 3	80N112
923.	Proflavine, conjugate monoacid EtOH EtOH/Et ₂ O (90 K)	FP/TD PS/SD	455, 6000 1100, 63000 ^a 917 ^a , 12500 ^a 675, 10000 ^a 550, 12500 ^a 400, 4000 ^a 350, 6500 ^a 280, 47000 ^a 235, 11000 ^a 210, 10000 ^a		pH Acidic Solvent was 2:1 EtOH to Et ₂ O	727073 57B001

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Glycerol (293 K)	FP	650 ^a 600 ^a 540 ^a		Solvent has added glucose	776171
	SDS	LP	1080		1 μ s delay; aqueous micelle; $\tau_T = \sim 35 \mu$ s; pH 8.1	80N112
	Water	FP/SD	560 445 ^b , <10000 350		\dagger Oxygen quenching; decay was second order with $2k \sim 10^9$ L mol ⁻¹ s ⁻¹ ; pH 4	677029
	Water (296 K)	FP/TD	549, 5700		Light saturation; pH 4.1	80B057
	Water	LP	1160 930 670 550		\dagger Oxygen quenching; radical dication ($\lambda_{\max} = 810$ nm) also observed; $\tau_T = 20 \mu$ s; pH 3.0	80F373
924.	Promazine 2-PrOH	LP	465		\dagger Triplet ET to β -carotene, oxygen quenching; $\tau_T = 22.8 \mu$ s	767023
925.	9-Propionylantracene EPA (77 K)	FP	430		$\tau_T = 3.21 \times 10^4 \mu$ s	82E338
926.	Propiophenone 2-PrOH	LP	325			84E018
	Acetonitrile/Water	LP	325		Solvent was 9:1 acetonitrile to water	84E018
	Benzene	LP	418 ^a		Much more intense peak below 300 nm; shoulder at 457 ^a nm	84E018
927.	N-Propylphthalimide EtOH	FP	590 ^a 350 ^a		\dagger Oxygen quenching (3.5×10^9 L mol ⁻¹ s ⁻¹), diene quenching; 100 ns delay; relative intensities (1:5); $\tau_T = 40 \mu$ s	79A147
928.	1-Propynylbenzene Isopentane/2-PrOH (77 K)	PS	255 ^a		\dagger ESR; glass was 7:3 isopentane to 2-propanol by volume; by assuming that radical formation occurs only from T [*] and that it is wavelength-independent, the author devises an ESR method that determines the shape of TTA even in the SD region	68B007
929.	Protoporphyrin MeOH	FP	530 440		Maximum at 530 nm was judged to be of "doubtful significance" by the experimenters	58R002
	Pyridine	FP	420 350			58R002
930.	Protoporphyrin IX Dioxane	LP	742 ^a		\dagger Triplet quenching by <i>cis</i> -piperylene; oxygen quenching (1.6×10^9 L mol ⁻¹ s ⁻¹); TTA of globin complexes also studied	747630
	Water	LP	500 ^a 450 ^a		\dagger Oxygen quenching (1.6×10^9 L mol ⁻¹ s ⁻¹); difference spectrum only reported; $\tau_T = \sim 150 \mu$ s; pH 0	78E011
931.	Protoporphyrin IX, dimethyl ester Benzene	PR/ET	710 ^a , 9000 ^a		\dagger Triplet ET from biphenyl and to β -carotene, oxygen quenching (2.7×10^9 L mol ⁻¹ s ⁻¹); ϵ relative to biphenyl in benzene ($\epsilon_{550} = 27100$ L mol ⁻¹ cm ⁻¹); more intense peak below 450 nm; $E_T = 150$ kJ mol ⁻¹	771078
	Benzene	LP/TD	450 ^b , 35000 \pm 1500 425 320		\dagger First order decay, oxygen quenching (1.5×10^9 L mol ⁻¹ s ⁻¹); $\tau_T = 550 \mu$ s	80B017
932.	Protoporphyrin IX, dimethyl ester, conjugate diacid Benzene/Trifluoroacetic acid	LP/TD	560 470 ^b , 32000 \pm 1500 415		\dagger Triplet ET to β -carotene, oxygen quenching (4×10^9 L mol ⁻¹ s ⁻¹); $\tau_T = 71 \mu$ s; $k_{et} = 1.3 \times 10^{10}$ L mol ⁻¹ s ⁻¹	80B017

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
933.	Protoporphyrin IX, dimethyl ester, conjugate monoacid MeOH/Acetic acid	FP/TD	450 ^b , 27500 ± 1500		†First order decay, oxygen quenching; solvent was 1:1 MeOH to acetic acid by volume; $\tau_T = 110 \mu\text{s}$	80B017
934.	Psoralen Benzene	PR/ET	450 ^b , 8100		†Triplet ET to β -carotene; ϵ relative to biphenyl in benzene ($\epsilon_{367} = 27100 \text{ L mol}^{-1} \text{ cm}^{-1}$); $k_{et} = 5.0 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	78E157
	Benzene	LP	450 ^a			78E157
	EtOH	LP/ET	440, 30100		†Triplet ET to retinol; ϵ relative to retinol in hexane ($\epsilon_{405} = 80000 \text{ L mol}^{-1} \text{ cm}^{-1}$) assuming ϵ independent of solvent; $\tau_T = 5 \mu\text{s}$; $k_{et} = 1.1 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	79E678
	Water	LP	450 ^a		$\tau_T = 7.1 \mu\text{s}$	78E157
	Water	LP/TD	460, 10000		†Oxygen quenching; $\tau_T = 12 \mu\text{s}$; pH 8	83B068
935.	4',5'-Psoralen-thymine photo adduct MeOH	LP/ET	500, 8500		†Triplet ET to <i>all-trans</i> -retinol; ϵ relative to <i>all-trans</i> -retinol in hexane ($\epsilon_{405} = 80000 \text{ L mol}^{-1} \text{ cm}^{-1}$), assumed unchanged in MeOH; 100 ns delay; $\tau_T = 10.6 \mu\text{s}$; $k_{et} = 2.2 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	80A032
	Water	LP	500 ^a		100 ns delay; $\tau_T = 2.5 \mu\text{s}$	80A032
936.	Purine Acetonitrile	FP	410 ^a 380 ^a		†Oxygen quenching; 20 μs delay	80B077
	Water	FP	390 360 300		†Oxygen quenching; 20 μs delay; pH 5.9	80B077
	Water (77 K)	PS/KM	455 ^a 412 ^a 390, 4100 380 ^a		†Phosphorescence decay; $\tau_T = (3.6 \pm 0.3) \times 10^6 \mu\text{s}$; pH Basic	80B077
937.	Pyranthrene Benzonitrile	LP/RF	591 ^a 502 ^a , 8000 ± 500		ϵ relative to pyranthrene in toluene ($\epsilon_{300} = 14000 \text{ L mol}^{-1} \text{ cm}^{-1}$)	83F075
	Toluene	LP/ET	584 ^a 500 ^a , 14000 ± 1500		ϵ relative to anthracene in toluene ($\epsilon_{428.5} = 42000 \text{ L mol}^{-1} \text{ cm}^{-1}$)	83F075
938.	Pyrazine MeOH/EtOH (~ 100 K)	FP/?	820 ^a , 5000 ^a 260, 1000 ^a		†Phosphorescence decay; glass was 1:1 MeOH to EtOH; shoulder at 300 nm assignment $^3B_{3u} \rightarrow ^3B_{2g}$; $\tau_T = (11 \pm 1) \times 10^3 \mu\text{s}$	80B011
	Water	LP/HAT	810, 1000 ± 200 700, 1100 ± 200 ~260, 4900 ± 1500 230, 3500 ± 1100		Shoulders at ~640 and ~295 nm; ϵ relative to the neutral pyrazyl radical in water ($\epsilon_{310} = 6700 \text{ L mol}^{-1} \text{ cm}^{-1}$), assuming that the quenching of the triplet by 2-PrOH produced stoichiometric equivalents of the neutral pyrazyl radical and the dihydro radical cation of pyrazine; $\tau_T = 3.4 \mu\text{s}$; pH 5.4	747233
	Water (298 K)	LP/HAT	810, 1000 700, 1100 ~260, 4900 230, 3500		†Oxygen quenching; shoulders at ~640 and ~295 nm; ϵ relative to dihydropyrazyl radical cation in water ($\epsilon_{315} = 6900 \text{ L mol}^{-1} \text{ cm}^{-1}$), assuming H-abstraction from 2-PrOH was complete; N_2O was used to scavenge hydrated electrons; lifetime was measured at pH 7.1 and was for infinite dilution; $\tau_T = 4.5 \mu\text{s}$; pH 5.4	757309
939.	Pyrene 3-MP (77 K)	PS	518 487 ^a 455 ^a 411 ^a 387 ^a		The first band was assigned to one electronic transition, the 2nd two bands to a forbidden transition, and last 2 bands were assigned to a 3rd transition; this assignment was questioned in [70E294] where 518, 487, & 455 nm were all assigned to a single electronic transition; assumed 77 K	70B004
	Acetonitrile	FP	412		Pyrene radical cation ($\lambda_{\max} = 445 \text{ nm}$) also present	767071

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Benzene	PR/ET	420, 20900		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700$ L mol ⁻¹ cm ⁻¹); ϵ obtained from a simultaneous least squares fit of data from several compounds making use of cyclohexane to benzene ϵ_{\max} ratios of 1.83 for naphthalene and 1.45 for anthracene	71E360
	Cyclohexane	LP	510 490 415		†Rise time of transient was the same as the decay time of the singlet	68B006
	Cyclohexane	LP	520 483 416			70E288
	Cyclohexane	LP	525 480 415		Delay 1 μ s; 415 nm was the most intense peak	707346
	Cyclohexane	MOD	517 508 412 409		Relative intensities (26:18:100:75); $\tau_T = 180$ μ s	71E361
	Cyclohexane	PR/ET	412.5, 30400		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700$ L mol ⁻¹ cm ⁻¹); reference ϵ obtained by starting from $\epsilon_{\max} = 3220$ L mol ⁻¹ cm ⁻¹ for this ketyl radical in water and assuming the f of the ketyl radical is independent of solvent; final ϵ obtained from a simultaneous least squares fit to data from several compounds	71E360
	Cyclohexane	PR	515 490 440 410 385 365			720206
	EPA (77 K)	PS	525 489 415.5 372		Relative intensities (20:15:100:10); unresolved shoulder at 515 nm; $\tau_T = (7 \pm 2) \times 10^3$ μ s	54B001
	EPA (77 K)	PS/IV	415, 48200		λ_{\max} assumed from previous work; ϵ estimated by extrapolation to infinite excitation rate	69E212
	EPA (117 K)	FP	520 ^a 479 ^a 412 ^a 389 ^a		†Oxygen quenching; 412 nm was the most intense peak	716243
	EtOH (293 K)	FP	408			68E098
	EtOH	LP	500 ^a 450 ^a 400 ^a		Relative intensities (1:1:5); 1.8 μ s delay; solution contains Ag ions	78E554
	Hexane	FP	520.0 411.0 387.0 369.0			54E001
	Hexane (300 K)	MOD/SD	522 ^a , 6300 ^a 490 ^a , 2100 ^a 410 ^a , 34400 ^a 389 ^a , 14500 ^a 360 ^a , 2170 ^a 333 ^a , 3200 ^a 318 ^a , 3900 ^a 290, 12000 260, 14500			69E208
	Hexane	MOD	516 ^a 508 ^a 412 ^a 409 ^a		Shoulder at 484 ^a nm; 412 nm was the most intense peak; $\tau_T = 180$ μ s	70E295

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Liquid paraffin	FP	519.5 483 416 396.5 371.5		Solvent viscosity was 0.19 N·s/m ² ; the bands at 519.5 and 483 nm belong to the 1st electronic transition, and the other 3 bands belong to the 2nd electronic transition; relative intensities (24:19:100:48:26)	58E001
	MeOH	LP	500 ^a 414 ^a 371 ^a		Delay 1 μ s	707346
	PMMA	CWL/RF	971, ~100 901 870 781 741 613, 900 562 521, 12500 485 455 413, 48200 392 370		ϵ relative to pyrene in EPA ($\epsilon_{\max} = 48200$ L mol ⁻¹ cm ⁻¹); four electronic transitions were assigned, and the 0-0 bands of these transitions are identified by their ϵ ; T ₁ is ³ B _{2u} ⁺ , and the other states are ³ B _{3g} ⁻ , ³ A _g ⁺ , ³ B _{3g} ⁻ , and ³ A _g ⁻ in increasing energy; $\tau_T = 5 \times 10^5$ μ s; oscillator strength = $\sim 10^{-3}$, ~ 0.01 , 0.14, 0.31	70E294
	PMMA	I.P	520 483 416			70E288
	PMMA	PS	520 ^a 485 ^a 410 ^a 390 ^a		Relative intensities (7:5:21:10); polarization also measured; phase sensitive detection	78E152
	Pyrene (425 K)	LP	935 909 870 800 709 690 671 645 613 571 559 524 461		Liquid at 425 K; delay 350 ns; another maximum in 740 - 755 nm range	737101
	SDS	LP	414		Aqueous micelle	82N081
	Toluene	LP	529 ^a 421 ^a		Shoulder at 488 ^a nm; 421 nm was the more intense peak	717126
	Toluene	LP	520 410		50 ns delay	766528
	Water	LP	530 ^a 480 ^a 415 ^a		Solubilized by 1.5×10^{-5} mol L ⁻¹ bovine serum albumin; oxygen quenching (1.3×10^8 L mol ⁻¹ s ⁻¹); 1 μ s delay	773047
	Water	LP	414		Solubilized by caffeine; 2 μ s delay	81E149
	Water/ <i>tert</i> -BuOH	FP	410 ^a		Solvent mixture contains "1-5%" <i>tert</i> -BuOH for solubility; pH ~ 6	767189
940.	Pyrene- <i>d</i> ₁₀ 2-MTHF (77 K)	PS/ESR	416, 43600 394 ^a , 18100 ^a 415 ^a , 38600 ^a 391 ^a , 20000 ^a		\uparrow ESR; shoulder at 408 ^a nm; the band was assigned as ³ B _{3g} \leftarrow ³ B _{2u} ; oscillator strength = 0.21 $\tau_T = 3.7 \times 10^6$ μ s	68D211 696019
	PMMA (77 K)	PS/SD	415 ^a , 39800 ^a 391 ^a , 18700 ^a		$\tau_T = 2.4 \times 10^6$ μ s	696019
941.	Pyrene/Chloranil Liquid paraffin	MOD	405		Mull	71E361

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
942.	Pyrene/Diethylaniline Toluene	LP	500		50 ns delay; possible triplet exciplex; $\tau_T = 0.107 \mu\text{s}$	766528
943.	Pyrene/Tetracyanobenzene, 1:1 EPA (95 K)	FP	513 ^a 480 ^a 427 ^a 404 ^a 394 ^a 383 ^a		{Oxygen quenching and an accompanying phosphorescence; 404 nm was the most intense peak	716243
944.	Pyrene/Tetracyanobenzene, 2:1 EPA (117 K)	FP	435 ^a 415 ^a 400 ^a		{Oxygen quenching and an accompanying phosphorescence; 415 nm was the most intense peak	716243
945.	1-Pyrenecarboxaldehyde					
	2-PrOH (295 K)	LP/SD	440 ± 3, 18100 ± 1800	$\tau_T = 37 \mu\text{s}$		83E387
	Acetonitrile (295 K)	LP/SD	438 ± 3, 18700 ± 1900	$\tau_T = 30 \mu\text{s}$		83E387
	Acetone (295 K)	LP/SD	440 ± 3, 20100 ^c ± 2000	$\tau_T = 20 \mu\text{s}$		83E387
	Benzene (295 K)	LP/SD&PR/ ET	443 ± 3, 21000 ^c ± 2100	In ET method ϵ relative to biphenyl in benzene ($\epsilon_{361} = 42800 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 35 \mu\text{s}$		83E387
	CCl ₄ (295 K)	LP/SD	443 ± 3, 18800 ± 1900	$\tau_T = 49 \mu\text{s}$		83E387
	CF ₃ CH ₂ OH (295 K)	LP/SD	438 ± 3, 16900 ^c ± 1700	$\tau_T = 58 \mu\text{s}$		83E387
	Cyclohexane (295 K)	LP/SD&TD	440 ± 3, 18900 ± 1900	$\tau_T = 50 \mu\text{s}$		83E387
	Cyclohexene (295 K)	LP/SD	440 ± 3, 17700 ± 1800	$\tau_T = 70 \mu\text{s}$		83E387
	EtOH (295 K)	LP/SD	440 ± 3, 18200 ± 1800	$\tau_T = 38 \mu\text{s}$		83E387
	Ethylene glycol (295 K)	LP/SD	440 ± 3, 18100 ± 1800	$\tau_T = 300 \mu\text{s}$		83E387
	Hexane- <i>d</i> ₁₄ (295 K)	LP	428 ± 3	$\tau_T = 34 \mu\text{s}$		83E387
	MeOH (295 K)	LP/SD	437 ± 3, 18300 ± 1800	$\tau_T = 26 \mu\text{s}$		83E387
	MeOH/Water (295 K)	LP/SD	443 ± 3, 18500 ± 1900	Solvent was 1:1 MeOH to water; $\tau_T = 107 \mu\text{s}$		83E387
	THF (295 K)	LP/SD	440 ± 3, 19100 ± 1900	$\tau_T = 59 \mu\text{s}$		83E387
946.	Pyrene-1-sulfonate Water	LP	520 420		Delay 1 μs	84E533
947.	Pyrene-3-sulfonate Benzene/DDDAB	MOD	525 491 429		Inverse micelle; relative intensities (3:2:5); $\tau_T = 3000 \mu\text{s}$	79N036
	Water	MOD	422		Principal maximum only reported	79N036
948.	[2,2](2,7)-Pyrenophane 2-MTHF (77 K)	LP	625 480 380			79B050
949.	2-(1-Pyrenylmethyl)-1,3-propandiol diacetate DMF	LP	415 ^a		{Rises with fluorescence decay, oxygen quenching	80B076
950.	2-(1-Pyrenylmethyl)-1,3-propandiol diacetate/1,4-Dicyanobenzene DMF	LP	420		{Rises with fluorescence decay, oxygen quenching; pyrene radical cation also observed (460 nm)	80B076
951.	Pyrido[2,1,6- <i>de</i>]quinolizine Hexane	FP	380		{Ground state recovery, molecular orbital calculations; solvent uncertain; "cyl[3.3.3]azine"; $\tau_T = \sim 0.1 \mu\text{s}$; $E_T = \sim 90 \text{ kJ mol}^{-1}$	80E405
952.	Pyrimidine Water (298 K)	LP/HAT	600, 530 260 ^b , ~ 3000		{Oxygen quenching; ϵ relative to dihydropyrazyl radical cation in water ($\epsilon_{315} = 6900 \text{ L mol}^{-1} \text{ cm}^{-1}$), assuming H-abstraction from 2-PrOH was complete; N ₂ O and <i>tert</i> -BuOH were present as scavengers; lifetime was for infinite dilution; $\tau_T = 1.4 \mu\text{s}$; pH 7.1	757309

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
953.	Pyromellitic dianhydride CCl ₄	FP	560 ^a 335 ^a		†Oxygen and piperylene quenching; 335 nm peak more intense; shoulder at 530 ^a nm; delay 10 μ s; "PMDA"	737098
	Et ₂ O/Isopentane (77 K)	FP	496 ^a		†Phosphorescence decay; glass was 1:1 Et ₂ O to isopentane, shoulder or maximum at \sim 350 nm; $\tau_T = 6.7 \times 10^5 \mu$ s	747587
954.	Pyromellitic dianhydride/Et₂O Et ₂ O	FP	561 ^a 529 ^a		†Oxygen and piperylene quenching; there was a maximum or shoulder near 340 nm; delay 10 μ s; "PMDA"-Et ₂ O complex	737098
955.	Pyronine Water	FP	482 ^a		pH 7	747229
956.	4-Pyrrolidinopyridine EPA (77 K)	FP	510		†Phosphorescence decay; $\tau_T = 1.1 \times 10^6 \mu$ s	83B112
957.	Pyruvic acid Acetonitrile	FP	695 ^a		†Triplet ET to naphthalene, oxygen quenching; $\tau_T = 0.5 \mu$ s	81F070
958.	Quantacure SKS anion Water	LP	631 ^a 477 ^a		†Oxygen quenching; 447 nm peak was the more intense; anion of sodium (4-sulfomethylphenyl)-phenylethanedione salt; $\tau_T = 0.50 \pm 0.02 \mu$ s; pH 6	82P060
959.	[1,1':4',1'':4'',1''':4''']-Quaterphenyl]-4,4'''-diamine 2-MTHF (77 K)	CWL	650			74B003
960.	p-Quaterphenyl 2-MTHF (77 K)	CWL	530		Shoulder at 498 ^a nm	74B003
	BuOH (\sim 80 K)	FP	525 ^a 495 ^a		Relative intensities (5:2)	67B010
	Toluene/EtOH (77 K)	MOD	563 ^a 395 ^a		Glass was 19:1 toluene to EtOH; 563 nm peak was the more intense	719059
961.	Quinoline 1-BuOH/Isopentane (77 K)	FP/SD	458 ^a , 2700 ^a 425 ^a , 7000 ^a 400.0, 7100 \pm 700		Glass was 3:7 1-BuOH to isopentane; shoulder at 384 ^a nm; oscillator strength = 0.13	707240
	3-MP (77 K)	MOD/SD	485 ^a 455 ^a 422 ^a 398, 6000 238, 25000		Shoulder at 382 ^a nm	73B001
	EPA (77 K)	PS	487 458 427 402		Relative intensities (40:59:98:100); unresolved shoulders at 475 nm and 380 nm; $\tau_T = (5 \pm 2) \times 10^3 \mu$ s	54B001
	EtOH (77 K)	MOD/KM	425, 6500 \pm 1600			737055
	Isopentane/MCH (77 K)	PS	526 485 450 420 395		Relative intensities (4:22:42:97:100); unresolved shoulder at 373 nm	54B001
	Water	FP	418		pH Basic	61E008
962.	Quinoline, conjugate acid Water	FP	465		pK _a = 6.0; pH Acidic	61E008
963.	Quinoxaline 1-BuOH/Isopentane (77 K)	FP/SD	425.0, 8100 \pm 800		Glass was 3:7 1-BuOH to isopentane; oscillator strength = 0.15	707240

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	3-MP (77 K)	PS	414			67E106
	EPA (77 K)	PS	417			67E106
	EtOH/MeOH (113 K)	FP/TD	410, >2500		Solvent was 3:1 EtOH to MeOH; lower limit assumes total ground state depletion	67B004
	Hexane	LP	430		100 ps delay; rise time of 0.04 ns	79B007
	Isooctane	LP	425		130 ps delay, at earlier times overlapping $S_n \leftarrow S_1$ absorption	79B076
	PMMA (293 K)	FP	~415		Decay was nonexponential, and lifetime was calculated from a terminal 1st-order rate constant; $\tau_T = \sim 5 \times 10^4 \mu s$	70E291
	Water (298 K)	LP	432 413 ~270		\uparrow Oxygen quenching; shoulder at ~660 nm; a limit was set on ϵ_{413} as ≤ 7400 L mol ⁻¹ cm ⁻¹ , based on radical formation; $\tau_T = 29.4 \mu s$; pH 7.1	757309
	Water	FP	426 ^a 410 ^a		pK _b = 9.0 ± 0.7 ; pII 12.6	766206
964.	Quinoxaline-1,4-dioxide Water	FP	498 ^a 469 ^a		\uparrow Triplet ET from 1,5-naphthalene disulfonate and to Methylene Blue; shoulder at 429 ^a nm; $\tau_T = 77 \mu s$; pH Neutral	83F096
965.	Retinal Toluene (299 K)	FP	450		Whether isomer was all-trans or 11-cis was not specified; $\tau_T = 13 \mu s$	62E006
966.	11-cis-Retinal Benzene	LP-ET	455		Triplet ET from anthracene; $k_{et} = 2.0 \times 10^{10}$ L mol ⁻¹ s ⁻¹	777434
	Benzene	LP	455		$\tau_T = 8.70 \mu s$	777434
	DMSO	LP/RF	480, 27000		\uparrow Triplet ET to β -carotene; ϵ determined assuming oscillator strength independent of solvent, bands are symmetrical (in energy) and relative to measured value (unstated) in hexane	78E467
	EPA (77 K)	LP	445		$\tau_T = 41.3 \pm 1.7 \mu s$	79B077
	Hexane	PR/ET	450, 75000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹ , note literature value at 361.3 nm); $\tau_T = 8.3 \pm 0.7 \mu s$	741013
	Hexane	LP/ET	445, 57000 \pm 14000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹); original reference quoted (440 \pm 5) nm as λ_{\max} , stated λ_{\max} from [78Z194]; $\tau_T = 8.3 \pm 0.7 \mu s$	747633
	Hexane	PR/ET	445, 62000 \pm 6000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹); original reference quoted (440 \pm 5) nm as λ_{\max} , stated λ_{\max} from [78Z194]; $\tau_T = 8.3 \pm 0.7 \mu s$	747633
	Hexane	LP/SD	445, 50000 ^a		ϵ relative to anthracene in cyclohexane ($\epsilon_{423} = 64700$ L mol ⁻¹ cm ⁻¹)	78E208
	MeOH	LP/RF	450, 27000		\uparrow Triplet ET to β -carotene; ϵ determined assuming oscillator strength independent of solvent, bands are symmetrical (in energy) and relative to measured value (unstated) in hexane	78E467
967.	13-cis-Retinal EPA (77 K)	LP	465		$\tau_T = 48.3 \pm 2.8 \mu s$	79B077
	Hexane	PR/ET	445 \pm 5, 56000 \pm 6000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹); $\tau_T = 8.3 \pm 0.7 \mu s$	747633
	Hexane	LP/ET	445 \pm 5, 44000 \pm 11000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹); $\tau_T = 8.3 \pm 0.7 \mu s$	747633
	Hexane	PR/ET	445, 68000 \pm 10200		\uparrow Triplet ET from biphenyl; ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹), assuming authors used standard for cyclohexane; $\tau_T = 9.1 \pm 0.9 \mu s$; $k_{et} = (2.0 \pm 0.20) \times 10^{10}$ L mol ⁻¹ s ⁻¹	776412
	MeOH	LP/RF	450, 24300		\uparrow Triplet ET to β -carotene; ϵ determined assuming oscillator strength independent of solvent, bands are symmetrical (in energy) and relative to measured value (unstated) in hexane	78E467

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
968.	7,9-cis-Retinal					
	Benzene	LP-ET	455		Triplet ET from anthracene; $k_{et} = 2.8 \times 10^{10}$ L mol ⁻¹ s ⁻¹	777434
	Benzene	LP	455		$\tau_T = 8.47$ μ s	777434
969.	7-cis-Retinal					
	Benzene	LP-ET	455		Triplet ET from anthracene; $k_{et} = 2.0 \times 10^{10}$ L mol ⁻¹ s ⁻¹	777434
	Benzene	LP	455		$\tau_T = 8.70$ μ s	777434
	EPA (77 K)	LP	455 425		Decay wavelength dependent; lifetime measured at 455 nm; $\tau_T = 44.2 \pm 0.6$ μ s	79B077
970.	9-cis-Retinal					
	EPA (77 K)	LP	465		Concentration unspecified; $\tau_T = 58.8 \pm 2.4$ μ s	79B077
	Hexane	LP/ET	445 \pm 5, 63000 \pm 15000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹); $\tau_T = 8.3 \pm 0.7$ μ s	747633
	Hexane	PR/ET	445 \pm 5, 56000 \pm 6000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹); $\tau_T = 8.3 \pm 0.7$ μ s	747633
	MeOH	LP/RF	460, 24300		†Triplet ET to β -carotene; ϵ determined assuming oscillator strength independent of solvent, bands are symmetrical (in energy) and relative to measured value (unstated) in hexane; $\tau_T = \sim 0.17$ μ s	78E467
971.	all-trans-Retinal/(CF ₃) ₂ CHOH	LP	475		$\tau_T = 11.1$ μ s	82A288
972.	all-trans-Retinal					
	Acetonitrile	LP/RF	470, 67000		ϵ relative to retinal in hexane ($\epsilon_{445} = 78000$ L mol ⁻¹ cm ⁻¹) assuming oscillator strength independent of solvent; $\tau_T = 12.0$ μ s	79E546
	Acetonitrile	LP/SD	470, 59000		ϵ is upper limit; $\tau_T = 12.0$ μ s	79E546
	Benzene	LP	455		$\tau_T = 9.09$ μ s	777434
	Benzene	LP-ET	455		Triplet ET from anthracene; $k_{et} = 3.0 \times 10^{10}$ L mol ⁻¹ s ⁻¹	777434
	Benzene	PR/RF	460, 62000		ϵ relative to retinal in hexane ($\epsilon_{445} = 78000$ L mol ⁻¹ cm ⁻¹) assuming oscillator strength independent of solvent; $\tau_T = 9.1$ μ s	79E546
	Benzene	PR/ET	460, 67000		ϵ relative to biphenyl in benzene ($\epsilon_{367.5} = 27100$ L mol ⁻¹ cm ⁻¹); $\tau_T = 9.1$ μ s	79E546
	Benzene	PR	460		†Triplet ET from naphthalene; $\tau_T = 8.62$ μ s; $k_{et} = 5.5 \times 10^9$ L mol ⁻¹ s ⁻¹	81B008
	Cyclohexane	LP	445		$\tau_T = 9.3$ μ s	82A288
	DFMeOH	LP-ET	550		†Triplet ET from anthracene; $\tau_T = 24.2$ μ s	82A288
	EPA (77 K)	LP	465		$\tau_T = 41.0 \pm 3.2$ μ s	79B077
	Hexane	PR/ET	450, 74000		†Triplet ET to β -carotene; ϵ relative to biphenyl in cyclohexane ($\epsilon_{361.3} = 42800$ L mol ⁻¹ cm ⁻¹), assuming no solvent effect on biphenyl	733002
	Hexane	PR/ET	445 \pm 5, 78000 \pm 8000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹); $\tau_T = 8.3 \pm 0.7$ μ s	747633
	Hexane	PR-ET/ET	450, 75000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{365} = 42800$ L mol ⁻¹ cm ⁻¹ , note literature value at 361.3 nm); $\tau_T = 8.3 \pm 0.7$ μ s; $k_{et} = (3 \pm 1) \times 10^{10}$ L mol ⁻¹ s ⁻¹	741013
	Hexane	LP/ET	445 \pm 5, 69000 \pm 17000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹); $\tau_T = 8.3 \pm 0.7$ μ s	747633
	Hexane	PR/ET	445, 114000 ^c \pm 17100		†Triplet ET from biphenyl; ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹), assuming authors used standard for cyclohexane; $\tau_T = 9.1 \pm 0.9$ μ s; $k_{et} = (2.0 \pm 0.20) \times 10^{10}$ L mol ⁻¹ s ⁻¹	776412
	Hexane	LP/SD	445, 70000 \pm 7000		ϵ relative to anthracene in cyclohexane ($\epsilon_{423} = 64700$ L mol ⁻¹ cm ⁻¹)	78E208
	MCH	FP	440		†Oxygen quenching; half-life = 9.9 μ s	59E006
	MCH	FP/?	450, 75800		Lifetime measurement was made at 293 K with retinal concentration 2.18×10^{-5} mol L ⁻¹ ; $\tau_T = 11$ μ s	62E007
	MCH (296 K)	LP/SD	448, 76000		†Oxygen quenching and triplet ET to β -carotene; $\tau_T = 11$ μ s	747334

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	MeOH	FP/?	465, 67000		Lifetime measurement was made at 293 K with retinal concentration 2.30×10^{-5} mol L ⁻¹ ; $\tau_T = 17$ μ s	62E007
	MeOH (296 K)	LP/SD	450, 68900		{Oxygen quenching and triplet ET to β -carotene; $\tau_T = 16$ μ s	747334
	MeOH	LP/RF	460, 74000		{Triplet ET to β -carotene; ϵ determined assuming oscillator strength independent of solvent, bands are symmetrical (in energy) and relative to measured value (unstated) in hexane	78E467
	THF	FP/?	448, 78600		Lifetime measurement was made at 293 K with retinal concentration 2.28×10^{-5} mol L ⁻¹ ; $\tau_T = 14$ μ s	62E007
973.	<i>all-trans</i> -Retinoic acid					
	Hexane	PR/ET	440, 73000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹)	82A205
	MeOH	LP	440		{Oxygen quenching; $\tau_T = 1.6$ μ s	82A205
974.	<i>all-trans</i> -Retinol					
	Cyclohexane	LP	405		{Triplet ET from biphenyl and oxygen enhancement of triplet yields	737185
	Hexane (298 K)	FP-ET	400 ^a 383 ^a		{Triplet ET from benz[a]anthracene; delay 50 μ s; $\tau_T = 17$ μ s; $k_{et} = 5 \times 10^9$ L mol ⁻¹ s ⁻¹	69E217
	Hexane	PR/ET	405, 57500 \pm 500		{Triplet ET from biphenyl; ϵ relative to biphenyl (ϵ_{ref} unstated); $k_{et} = 3 \times 10^9$ L mol ⁻¹ s ⁻¹	746499
	Hexane	LP/ET	405, 57500 \pm 500		{Triplet ET from TMPD-biphenyl; ϵ relative to biphenyl (ϵ_{ref} unstated)	746499
	Hexane	PR/ET	405, 80000 \pm 12000		{Triplet ET from biphenyl; ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹), assuming authors used standard for cyclohexane; $\tau_T = 15 \pm 1.5$ μ s; $k_{et} = (2.3 \pm 0.23) \times 10^{10}$ L mol ⁻¹ s ⁻¹	776412
975.	Retinyl acetate					
	Hexane	LP/ET	405, 64000 \pm 1000		{Triplet ET from TMPD-biphenyl; ϵ relative to biphenyl (ϵ_{ref} unstated)	746499
	Hexane	PR/ET	405, 64000 \pm 1000		{Triplet ET from biphenyl; ϵ relative to biphenyl (ϵ_{ref} unstated)	746499
	Hexane	PR/ET	395, 79000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹)	82A205
	MeOH	LP	395		{Oxygen quenching (1.0×10^9 L mol ⁻¹ s ⁻¹); $\tau_T = 8.3$ μ s	82A205
976.	<i>N</i> -11- <i>cis</i> -Retinylidene- <i>n</i> -butylamine					
	Hexane	LP-ET/ET	\sim 435, 50000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361,3} = 42800$ L mol ⁻¹ cm ⁻¹); ET was from an exciplex of TMPD and biphenyl which decayed to triplet biphenyl	747189
977.	<i>N</i> -13- <i>cis</i> -Retinylidene- <i>n</i> -butylamine					
	Hexane	LP-ET	\sim 440		ET was from an exciplex of TMPD and biphenyl which decayed to triplet biphenyl; the maximum was determined by smoothing the literature data	747189
978.	<i>N</i> -9- <i>cis</i> -Retinylidene- <i>n</i> -butylamine					
	Hexane	LP-ET	\sim 435		ET was from an exciplex of TMPD and biphenyl which decayed to triplet biphenyl	747189
979.	<i>N</i> - <i>all-trans</i> -Retinylidene- <i>n</i> -butylamine					
	Hexane	LP-ET/ET	\sim 435, 50000		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361,3} = 42800$ L mol ⁻¹ cm ⁻¹); ET was from an exciplex of TMPD and biphenyl which decayed to triplet biphenyl	747189
	Hexane	PR/ET	430, 158000 \pm 23700		{Triplet ET from biphenyl; ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹), assuming authors used standard for cyclohexane; $\tau_T = 20 \pm 2$ μ s; $k_{et} = (1.5 \pm 0.15) \times 10^{10}$ L mol ⁻¹ s ⁻¹	776412
	MCH (296 K)	LP/SD	435, 45000		{Oxygen quenching and triplet ET from triplet donor, anthracene; $\tau_T = 17$ μ s	747334

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
980.	<i>N</i> -11- <i>cis</i> -Retinylidene- <i>n</i> -butylamine, conjugate acid EtOH	LP-ET	~550		†Triplet ET from phenanthrene; 550 nm was apparent maximum in difference spectrum; $\tau_T = 50 \mu\text{s}$; pH Acidic	757090
981.	<i>N</i> -all- <i>trans</i> -Retinylidene- <i>n</i> -butylamine, conjugate acid MCH/Trichloroacetic acid (296 K)	LP-ET/SD	570, 150000		†Triplet ET from anthracene; $\tau_T = 20 \mu\text{s}$; $k_{et} = 1.2 \times 10^{10} \text{ L mol}^{-1} \text{ s}^{-1}$	747334
982.	Rhodamine 110 EtOH	FP/SD	606 ^a , 11000 ^a 510, 15800 391 ^a , 8460 ^a		†Triplet ET from naphthalene; shoulder at 460 ^a nm; long-lived photoproduct also observed; ϵ units uncertain	777316
983.	Rhodamine B 1-PrOH	FP/SD	1000 640, 13000 450		†Triplet ET to anthracene sulfonic acid and from naphthalene, oxygen quenching; 250 μs delay	78A304
	EtOH	FP/SD	560, 23000 419 ^a , 12100 ^a		†Triplet ET from naphthalene; shoulder at 640 ^a nm; long-lived photoproduct also observed; ϵ units uncertain	777316
	Water	FP	464 ^a		pH 7	747229
	Water	FP/SD	1000 640 625 ^b , 10000 450		250 μs delay; blue band decays with same lifetime as others	777041
	Water	FP	1000 640 450		†Triplet ET to anthracene sulfonic acid and from naphthalene, oxygen quenching; 250 μs delay; pH 7.0	78A304
984.	Rhodamine 6G 1-PrOH	FP/SD	950 630, 13000 415		†Triplet ET to anthracene sulfonic acid and from naphthalene, oxygen quenching; 250 μs delay	78A304
	Bromoform	FP	651 ^a		$\tau_T = 190 \mu\text{s}$	72E277
	DMSO	FP-ET	624 ^a		†Triplet ET from anthracene; shoulder at ~705 ^a nm	72E277
	Deuterium oxide	FP/SD	1120, 42000 637, 20000 410, 28000 308, 28000		Shoulders at 578 and 308 nm	82B106
	EtOH	FP/ET	600, 18000 \pm 3000 411 ^a , 18000 ^a		ϵ relative to naphthalene in EtOH ($\epsilon_{415} = 40000 \text{ L mol}^{-1} \text{ cm}^{-1}$); strong SD masked spectrum between 490 and 550 nm; spectrum corrected for a photoproduct absorbing in 350 - 490 nm region; $\tau_T = 3400 \pm 120 \mu\text{s}$	747050
	Glycerol (293 K)	FP	1120 610 415			82B106
	PMMA (200 K)	FP	1100 633 415		†Phosphorescence decay	82B106
	Water	FP-ET	633 ^a		†Triplet ET from naphthalene; shoulder at ~740 ^a nm; $\tau_T = 770 \mu\text{s}$	72E277
	Water	FP/SD	950 630 596 ^b , 8500 415		250 μs delay; blue band decays with same lifetime as others	777041
	Water	FP-ET	950		Triplet ET from naphthalene; no bands observed in the visible; pH 8.8	78A304
	Water	FP	950 635 410		Maximum at 635 nm has apparent contribution from triplet of conjugate acid (authors)	79F576

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Water	FP/SD	637, 20000 410, 24000 308, 28000 256, 42000 227, 44000		Shoulders at 578 and 308 nm; pH 7	82B106
985.	Rhodamine 6G cation, conjugate monoacid					
	Water	FP	950 630 415		†Triplet ET to anthracene sulfonic acid and from naphthalene, oxygen quenching; 250 μ s delay; $pK_a = 7.3$; pH 7.0	78A304
	Water	FP	1000 420 355 313 260		pH 1.5	82B106
986.	Rhodamine 6G dimer					
	BuOH/CCl ₄	LP/ET	620 ^a , 16500 ^a		†Triplet ET to anthracene; dimer species; solvent was 2:3 BuOH to carbon tetrachloride; ϵ relative to anthracene (ϵ_{ref} unstated) assuming unit transfer efficiency	75E530
	CCl ₄	LP/ET	650 ^a , 10000 ^a		†Triplet ET to anthracene; aggregate species; solvent contains 0.2% BuOH; ϵ relative to anthracene (ϵ_{ref} unstated) assuming unit transfer efficiency	75E530
987.	Riboflavin, conjugate monoacid					
	Water	LP/RA	670 ^a , 5000 ^a 415 ^a , 7900 ^a 350 ^a , 7000 ^a		lifetime measured at 660 nm; ϵ relative to anthracene in cyclohexane ($\Phi_T = 0.71$, $\epsilon_{423} = 64700$ L mol ⁻¹ cm ⁻¹) and using $\Phi_T = 0.40$ for riboflavin in water; shoulder around 300 nm; $\tau_T = 19$ μ s; pH 2.2	77617
988.	Rose Bengal					
	Acetonitrile/Water	LP	820		Solvent was 3.5:1.5 acetonitrile to water; $\tau_T = 30$ μ s	84E216
	PrOH	FP	610 ^a 450 ^a		Relative intensities (1:1); $\tau_T = 500$ μ s	79E848
989.	Rubrene					
	Acetonitrile	FP-ET	498 ^a 473 ^a 450 ^a		†Triplet ET from biacetyl; delay 50 μ s; $\tau_T = \sim 80$ μ s	68E103
	Benzene	FP-ET	505 472		†Triplet ET from benzophenone; $\tau_T = 100 \pm 20$ μ s; $E_T = 100 - 124$ kJ mol ⁻¹	776262
	Benzene	FP/ET	450 ^b , 3000		†Triplet ET from benzophenone; ϵ_{ref} unstated; $\tau_T = 115 \pm 3$ μ s; $E_T = 110 \pm 2$ kJ mol ⁻¹	81E346
	Benzene	PR-ET	~ 475		†Triplet ET from biphenyl; $\tau_T = 109$ μ s	83E223
	Benzene	FP	496 ^a 467 ^a 438 ^a 413 ^a		496 nm was the most intense peak	84F248
	Hexane	FP	487.6			54E001
	Hexane	FP	536 519 418.5		Relative intensities (58:100:25)	58E001
	Toluene (295 K)	LP-ET/ET	495, 26000 \pm 2000 468 ^a , 19000 ^a 440 ^a , 15000 ^a 414 ^a , 10000 ^a		†Oxygen quenching, triplet ET from anthracene; ϵ relative to anthracene in benzene ($\epsilon_{431} = 42000$ L mol ⁻¹ cm ⁻¹ , assuming ϵ_{\max} in toluene at 429 nm was the same); $\tau_T = 140$ μ s; $k_{et} = (2.45 \pm 0.3) \times 10^9$ L mol ⁻¹ s ⁻¹	81E716
990.	Safranin cation					
	Water	FP/SD	420, 10000		† Oxygen quenching; unprotonated species; spectrum between 500-600 nm uncharacterized; $\tau_T = 59$ μ s; pH 10.4	81B032

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
991.	Safranine cation, conjugate diacid					
	Water	FP	698 ^a 657 ^a 384 ^a		‡Oxygen quenching; pH 3-5	677322
	Water	FP/SD	660, 21500 390, 17500		‡Oxygen quenching; pK _a = 7.5; spectrum between 500-600 nm uncharacterized; τ_T = 280 μ s; pH 3.0	81B032
992.	Safranine cation, conjugate monoacid					
	Water	FP	377 ^a		‡Oxygen quenching and KI quenching; another maximum > 800 nm with a shoulder at 743 ^a nm; τ_T = 67 \pm 13 μ s; pH 7-9	677322
	Water	FP/SD	> 800 800 ^b , 15500		‡Oxygen quenching; pK _a = 9.2; spectrum between 500-600 nm uncharacterized; τ_T = 44 μ s; pH 8.3	81B032
993.	Selenine cation					
	Water	FP/SD	770 ^a , 17000 ^a 690 ^a , 10500 ^a 425 ^a , 12000 ^a		ϵ assumes triplet does not absorb where singlet depletion is followed; lifetimes measured in MeOH; τ_T = 21 μ s; pH 8.5	767246
994.	Selenine cation, conjugate monoacid					
	Water	FP/SD	650 ^a , 15500 ^a 385 ^a , 13500 ^a		ϵ assumes triplet does not absorb where singlet depletion is followed; lifetimes measured in MeOH; pK _a = 6.4 \pm 0.2; τ_T = 0.39 μ s; pH 4.6	767246
995.	Selenopyronine					
	Water	FP	690 480 400		‡Triplet ET to benzophenone; τ_T = 167 μ s; pH 7.2	82E680
996.	<i>all-trans</i> -Spheroidene					
	Cyclohexane	PR/ET	510, 309000		ϵ relative to biphenyl in cyclohexane (ϵ_{360} = 42800 L mol ⁻¹ cm ⁻¹); τ_T = 5.5 μ s; k_{et} = 8.96 \times 10 ⁹ L mol ⁻¹ s ⁻¹	83B121
997.	<i>all-trans</i> -Spheroidenone					
	Cyclohexane	PR/ET	550, 60600		ϵ relative to biphenyl in cyclohexane (ϵ_{360} = 42800 L mol ⁻¹ cm ⁻¹); τ_T = 3.8 μ s; k_{et} = 3.22 \times 10 ¹⁰ L mol ⁻¹ s ⁻¹	83B121
998.	<i>all-trans</i> -Spirilloxanthin					
	Cyclohexane	PR/ET	550, 92000		ϵ relative to biphenyl in cyclohexane (ϵ_{360} = 42800 L mol ⁻¹ cm ⁻¹); τ_T = 3.7 μ s; k_{et} = 2.74 \times 10 ¹⁰ L mol ⁻¹ s ⁻¹	83B121
999.	Spiro[9,10-dihydro-9-oxoanthracene-10,2'-5',6'-benzindan]					
	Hexane	LP	423			84E077
1000.	Stilbene					
	Stilbene	PR	450 ^a		Triplet exciton; shoulders at 500 nm and 630 nm	80E230
1001.	<i>trans</i> -Stilbene-2,2'-d ₂					
	EPA (77 K)	FP	380 360 342		Most intense peak at 380 nm; τ_T = 2.2 \times 10 ⁴ μ s	707199
1002.	<i>trans</i> -Stilbene-2,3,4,5,6-d ₅					
	EPA (77 K)	FP	380 360 342		Most intense peak at 380 nm; τ_T = 2.4 \times 10 ⁴ μ s	707199
1003.	<i>trans</i> -Stilbene-4,4'-d ₂					
	EPA (77 K)	FP	380 360 344		Most intense peak at 380 nm; τ_T = 2.2 \times 10 ⁴ μ s	707199

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
1004.	<i>trans</i> -Stilbene-7,7'-d ₂ EPA (77 K)	FP	380 360 344		Most intense peak at 380 nm; $\tau_T = 5.3 \times 10^4 \mu s$	707199
1005.	<i>cis</i> -Stilbene Benzene Cyclohexane EtOH (108 K)	PR PR LP	360 360 380 360 342		{Oxygen quenching {Oxygen quenching; $\tau_T = 17 \pm 6 \mu s$ Triplet state absorption observed only after repetitive (~30) flashes	680379 680379 79E640
	MCH/Isohexane (93 K)	LP	380 360 341		Triplet state absorption observed only after repetitive (~30) flashes	79E640
1006.	<i>trans</i> -Stilbene Benzene Benzene Cyclohexane Cyclohexane EPA (77 K)	PR PR FP PR FP	360 360 325 360 379 360 343		{Oxygen quenching $G\epsilon_{100} = 2300 \pm 230 \text{ L mol}^{-1} \text{ cm}^{-1} / (100 \text{ eV absorbed})$; $\tau_T = 0.11 \mu s$ $\tau_T = 4 \times 10^4 \mu s$ {Oxygen quenching; $\tau_T = 14 \pm 6 \mu s$ 379 nm was the most intense peak; $\tau_T = 2.2 \times 10^4 \mu s$	680379 720447 600062 680379 677485
	EPA (103 K)	LP	381 362 344		Triplet absorption not observable above 107 K; lifetime constant below 88 K; $\tau_T = 1.1 \times 10^4 \mu s$	79E640
	EtOH (98 K)	LP	381 361 343		Relative intensities (3:2:1); triplet absorption not observable above 128 K; lifetime constant below 105 K; $\tau_T = 1.2 \times 10^4 \mu s$	79E640
	Glycerol (235 K)	LP	385 366 350		Triplet absorption not observable above 244 K; lifetime constant below 200 K; $\tau_T = 10^4 \mu s$	79E640
	Glycerol triacetate (213 K)	LP	380 360 340		Triplet absorption not observable above 235 K; lifetime constant below 201 K; $\tau_T = 10^4 \mu s$	79E640
	Isopentane/3-MP (77 K)	FP/IV	378, 34000 358 341		Glass was 6:1 isopentane : 3-MP; $\tau_T = 5300 \mu s$	687111
	MCH/Isohexane (98 K)	LP	380 360 342		Relative intensities (3:2:1); triplet absorption not observable above 109 K; reported lifetime constant below 95 K; $\tau_T = 1.4 \times 10^4 \mu s$	79E640
1007.	<i>trans</i> -Stilbene-d ₁₂ EPA (77 K)	FP	379 360 341		Main peak at 379 nm; $\tau_T = 9.2 \times 10^4 \mu s$	707199
	Isopentane/3-MP (77 K)	FP/IV	378, 28000 358 341		Glass was 6:1 isopentane to 3-MP; $\tau_T = 1.1 \times 10^4 \mu s$	687111
1008.	Styrene Cyclohexane	LP/RA-ET	325 ^b , 2200		ϵ relative to naphthalene in cyclohexane ($\epsilon_{413} = 24000 \text{ L mol}^{-1} \text{ cm}^{-1}$), but in ref. cited ($\epsilon_{415} = 24500 \text{ L mol}^{-1} \text{ cm}^{-1}$); triplet ET from thioxanthone; $\tau_T = \sim 0.025 \mu s$	82E181
1009.	Sulfacetamide Water	LP	450 330		{Oxygen quenching and triplet ET to β -carotene in EtOH; spectrum obtained from composite spectra; $\tau_T = 0.5 \pm 0.05 \mu s$; pH 7.0	82A138
1010.	4-Sulfomethylbenzophenone anion AOT/Decane/Water- /Cyclohexanol	LP	535		Microemulsion	84N125

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
1011.	[1,1':4',1''-Terphenyl]-4,4''-diamine 2-MTHF (77 K)	CWL	560			74B003
1012.	<i>m</i> -Terphenyl 2-PrOH/Isocetane/- Et ₂ O (81 K) BuOH (~80 K) BuOH (~80 K) EtOH/Et ₂ O (77 K) PVA (81 K) Toluene/EtOH (77 K)	FP FP/COM FP/IV MOD/KM FP MOD	404 ^a 401, 72 ^d 401, 63 ^d 436, 3900 ± 1200 412 ^a 341 ^a 441 ^a 400 ^a 364 ^a		{Phosphorescence decay; glass was 1:3:3 2-PrOH to isocetane to Et ₂ O ε estimated from numerical simulation of triplet state kinetics; $\tau_T = (4.65 \pm 0.2) \times 10^6 \mu s$ ε estimated by extrapolation to infinite excitation rate Glass was 2:1 EtOH to Et ₂ O; temp. temperature was not explicitly stated, but 77 K was inferred from the context {Phosphorescence decay Glass was 19:1 toluene to EtOH; most intense peak was at 441 nm	81B119 67B010 67B010 719059 81B119 719059
1013.	<i>o</i> -Terphenyl BuOH (~80 K) BuOH (~80 K) EtOH/Et ₂ O (77 K) Toluene/EtOH (77 K)	FP/COM FP/IV MOD/KM MOD	488, 80 ^d 488, 100 ^d 390, <1000 600 ^a 398 ^a 353 ^a		ε estimated from numerical simulation of triplet state kinetics; $\tau_T = (3.45 \pm 0.1) \times 10^6 \mu s$ ε estimated by extrapolation to infinite excitation rate Glass was 2:1 EtOH to Et ₂ O; temperature was not explicitly stated, but 77 K was inferred from the context Glass was 19:1 toluene to EtOH; most intense peak was at 398 nm	67B010 67B010 719059 719059
1014.	<i>p</i> -Terphenyl 2-MTHF (77 K) 2-PrOH/Isocetane/- Et ₂ O (81 K) Alcohol/Ether (77 K) Benzene BuOH (~80 K) BuOH (~80 K) EtOH/Et ₂ O (77 K) Hexane (300 K) MCH/Isopentane (77 K) Polystyrene Toluene/EtOH (77 K)	CWL FP MOD PK/EI FP/COM FP/IV MOD/KM MOD/SD PS PR MOD	460 459 459 460, 90000 449, 4880 ^d 449, 9920 ^d 460, 88000 ± 35000 444, 40000 455 ^a 470 467 ^a		Shoulder at 413 ^a nm {Phosphorescence decay; glass was 1:3:3 2-PrOH to isocetane to Et ₂ O; $\tau_T = 2.7 \times 10^6 \mu s$ Glass was 2:1 alcohol to ether ε relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700 \text{ L mol}^{-1} \text{ cm}^{-1}$); ε obtained from a simultaneous least squares fit of data from several compounds making use of cyclohexane to benzene ϵ_{\max} ratios of 1.83 for naphthalene and 1.45 for anthracene ε estimated from numerical simulation of triplet state kinetics; $\tau_T = (1.93 \pm 0.06) \times 10^6 \mu s$ ε estimated by extrapolation to infinite excitation rate Glass was 2:1 EtOH to Et ₂ O; temperature was not explicitly stated, but 77 K was inferred from the context $\tau_T = 450 \pm 40 \mu s$ Glass was 4:1 MCH to isopentane; shoulder at 425 nm Gε values increased with concentration; half-life = $2 \times 10^4 \mu s$ Glass was 19:1 toluene to EtOH; shoulder at 394 ^a nm	74B003 81B119 76E682 71E360 67B010 67B010 719059 69E208 74B006 672086 719059
1015.	<i>p</i> -Terphenyl- <i>d</i> ₁₄ 3-MP (77 K)	PS/ESR	461, 12900		{ESR; oscillator strength = 0.88	69B002
1016.	4-(4- <i>p</i> -Terphenylmethyl)benzophenone 2-MTHF (77 K)	PS/ESR	480 ^a , 110000 ± 20000			707036
1017.	2-([1,1':4',1''-Terphenyl]-4-ylmethyl)triphenylene 2-MTHF (77 K)	PS/ESR	471 ^a , 100000 ± 20000			707036
1018.	Testosterone EtOH	LP	305 ^a		{Triplet ET to piperylene, oxygen quenching ($2.2 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); solvent uncertain; $\tau_T = 0.044 \mu s$	80B055

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
1019.	Tetrabenzophenazine Toluene (293 K)	FP	495 ^a		†Phosphorescence decay in EPA at 77 K, oxygen quenching; isomer unspecified	80E778
1020.	Tetrabenzoporphine Pyridine	FP	440 400			73E345
1021.	Tetracene 2-MTHF (77 K)	CWL	485 468 440 412 385 360 320		The first 3 bands were assigned to $^3B_{1g} \leftarrow ^3B_{2g}$, and the last 4 were assigned to $^3A_{1g} \leftarrow ^3B_{2g}$	727000
	Acetonitrile	LP-ET	470 ^a		†Triplet ET from 2-nitronaphthalene; $k_{et} = 7.1 \times 10^9$ L mol ⁻¹ s ⁻¹	767269
	Acetonitrile	FP	460			78A163
	Benzene	PR/ET	465, 31200		ϵ relative to benzophenone ketyl radical in cyclohexane ($\epsilon_{\max} = 3700$ L mol ⁻¹ cm ⁻¹); ϵ obtained from a simultaneous least squares fit of data from several compounds making use of cyclohexane to benzene ϵ_{\max} ratios of 1.83 for naphthalene and 1.45 for anthracene	71E360
	Benzene	LP-ET	470 ^a		†Triplet ET from 2-nitronaphthalene; shoulder at 450 nm; $k_{et} = 7.4 \times 10^9$ L mol ⁻¹ s ⁻¹	767269
	EtOH/MeOH/2-MTHF (113 K)	FP/SD	962, 1000 \pm 300 935 ^a , 70 ^a 909 ^a , 80 ^a 893 ^a , 80 ^a 846.5, 790 ^a 806 ^a , 60 ^a 794 ^a , 70 ^a 769 ^a , 60 ^a 757, 300 ^a 694 ^a , 70 ^a 683, 90 ^a 671 ^a , 70 ^a		†Lifetime same as TTA bands in the blue; glass was 3:1:4 EtOH to MeOH to 2-MTHF; shoulders at 862 ^a , 752 ^a , and 741 ^a nm; $\tau_T = 1000 \pm 50$ μ s	69B006
	EtOH/MeOH (113 K)	FP/TD	962.0, 1000 931 ^a , 80 ^a 911 ^a , 90 ^a 888 ^a , 100 ^a 848 ^a , 750 ^a 807 ^a , 70 ^a 793 ^a , 80 ^a 774 ^a , 70 ^a 758 ^a , 290 ^a 699 ^a , 80 ^a 685 ^a , 100 ^a 497 ^a , 5600 ^a 481.5, 60000 463.3, 90000 438 ^a , 32900 ^a 413 ^a , 9150 ^a 368 ^a , 2300 ^a 339.1, 2000 318.0, 17000 305 ^a , 15600 ^a 285.8, 350000 244.0, 23000 235 ^a , 15800 ^a		†Phosphorescence decay; shoulders at 864 ^a , 752 ^a , 745 ^a , 669 ^a , 534 ^a , 488 ^a , 458 ^a , 451 ^a , 330 ^a , 294 ^a , and 291 ^a nm; 7 electronic transitions were assigned with 0-0's at 962.0, 481.5, 463.3, 339.1, 318.0, 285.8, and 244.0 nm; oscillator strength = 0.002, 0.2, 0.45, 0.004, 0.15, 0.85, \sim 0.2	72B001

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Liquid paraffin	FP/SD	476, 30000 460, 52600 430.5, 27300 400, 8000 312, 8000 285.5, 195000		Solvent had viscosity of 0.03 N·s/m ² ; 4 separate electronic transitions were assigned; bands between 460 and 400 nm were assigned to the second transition; oscillator strength = 0.1, 0.4, 0.05, 0.9	58E001
	MCH	LP	476 ^a 459 ^a 431 ^a		459 nm was the most intense peak	82F367
	THF	FP	480 ^a 460 ^a 435 ^a		460 nm was the most intense peak	757004
1022.	1,5,9,10-Tetrachloroanthracene Liquid paraffin	FP	440		Viscosity of solvent was 0.167 N·s/m ² ; $\tau_T = 310 \mu\text{s}$	62E009
1023.	1,2,4,5-Tetracyanobenzene Benzene	PR	1000		†Triplet ET to anthracene; charge transfer complex with solvent	776146
	EPA (77 K)	FP	516 477		†Phosphorescence decay, molecular orbital calculations; ϵ roughly 200 L mol ⁻¹ cm ⁻¹ (method unspecified); possible charge transfer complex with ether of solvent since radical anion detected on repeated flashing; $\tau_T = 3.1 \mu\text{s}$	776213
1024.	<i>N,N,N',N'</i> -Tetraethyloxonine cation EtOH	LP-ET/SD	820, 24000 500 ^a , 8600 ^a		†Triplet ET from 9,10-dibromoanthracene and oxygen quenching; solution 10 ⁻³ mol L ⁻¹ NH ₃ ; $\tau_T = 285 \mu\text{s}$	82E456
1025.	<i>N,N,N',N'</i> -Tetraethyloxonine cation, conjugate monoacid EtOH	LP-ET/SD	700, 28000 500 ^a , 9700 ^a		†Triplet ET from 9,10-dibromoanthracene and oxygen quenching; solution 0.2 mol L ⁻¹ H ₃ PO ₄ ; $\tau_T = 47 \mu\text{s}$; $k_{et} = 1.9 \times 10^9$ L mol ⁻¹ s ⁻¹	82E456
1026.	Tetraethylporphine Pyridine	FP	779 ^a 434 ^a		Shoulder at 451 nm	74B007
1027.	<i>N,N,N',N'</i> -Tetraethyl-[1,1':4',1'':4'',1'''-quaterphenyl]-4,4'''-diamine 2-MTHF (77 K)	CWL	710			74B003
1028.	<i>N,N,N',N'</i> -Tetraethyl-[1,1':4',1'':4'',1'''-terphenyl]-4,4'''-diamine 2-MTHF (77 K)	CWL	610			74B003
1029.	(<i>R</i>)-4,5,6,7-Tetrahydrodinaphtho[2,1- <i>b</i> :1',2'- <i>d</i>][1,6]dioxecin 2-MTHF (96 K)	PS	429 ^a 413 ^a		Compound "VIII" in paper	83E383
1030.	6,7,8,9-Tetrahydro-4-hydroxythiazolo[4,5- <i>h</i>]isoquinoline-7-carboxylate ion Water	LP/ET	370, 8000		†Oxygen quenching and triplet ET to crocetin; ϵ relative to crocetin in water ($\epsilon_{470} = 75000$ L mol ⁻¹ cm ⁻¹); ϵ considered an upper limit; $\tau_T = 54 \mu\text{s}$; pH 13; $k_{et} = 1.8 \times 10^{10}$ L mol ⁻¹ s ⁻¹	84A024
1031.	7,8,9,10-Tetrahydro-4-hydroxybenzo[<i>a</i>]pyrene Water	FP	415		Solution buffered; delay 50 μs	82R150
1032.	Tetrakis(2,6-dimethyl-4-sulfonatophenyl)porphine Water	LP/TD	780, 3200		$\tau_T = 1800 \mu\text{s}$	84E203
1033.	Tetrakis(2- <i>N</i> -methylpyridyl)porphine Water	LP/TD	790, 2600 \pm 300		$\tau_T = 1160 \pm 120 \mu\text{s}$; pH 6.0	84E346

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
1034.	Tetrakis(3- <i>N</i> -methylpyridyl)porphine Water	LP/TD	840, 3200 ± 300 710	$\tau_T = 390 \pm 39 \mu\text{s}$; pH 6.0		84E346
1035.	Tetrakis(4- <i>N</i> -methylpyridyl)porphine Water	LP/TD	450 ^b , 28600	†Oxygen quenching; tetra- <i>p</i> -toluenesulfonate salt; pH 7.4; half-life = 120 μs Shoulder at 820 nm		82A152
	Water	LP/TD	920, 7600 ± 800 470 ^b , 28000 ± 2800 450 ^b , 57600 ± 5800			82E622
	Water	LP/TD	920, 7600 ± 800 820	$\tau_T = 170 \pm 17 \mu\text{s}$; pH 6.0		84E346
1036.	Tetrakis(2-pyridyl)porphine Methylene chloride	LP/TD	790, 2600 ± 300 695	$\tau_T = 150 \pm 15 \mu\text{s}$		84E346
1037.	Tetrakis(3-pyridyl)porphine Methylene chloride	LP/TD	790, 3600 ± 400 700	$\tau_T = 170 \pm 17 \mu\text{s}$		84E346
1038.	Tetrakis(4-pyridyl)porphine Methylene chloride	LP/TD	790, 3800 ± 400 695	$\tau_T = 170 \pm 17 \mu\text{s}$		84E346
1039.	Tetrakis(2-pyridyl)porphine, conjugate diacid Water	LP	930 720	1 mol L ⁻¹ HCl; pH Acidic		84E346
1040.	Tetrakis(3-pyridyl)porphine, conjugate diacid Water	LP	890 720	1 mol L ⁻¹ HCl; pH Acidic		84E346
1041.	Tetrakis(4-pyridyl)porphine, conjugate diacid Water	LP	950 720	1 mol L ⁻¹ HCl; pH Acidic		84E346
1042.	Tetrakis(<i>p</i> -sulfonatophenyl)porphinatorhodate(III) ion Water	LP/TD	830			84E043
1043.	Tetrakis(<i>p</i> -sulfonatophenyl)porphine Water	LP/TD	460 ^b , 65000	†Oxygen quenching (1.9 × 10 ⁹ L mol ⁻¹ s ⁻¹); $\tau_T = 400 \mu\text{s}$; pH 7		81E084
	Water	LP/TD	790, 3100 ± 300 680			82E622
	Water	LP/TD	460 ^b , 130000 ± 13000 440 ^b , 44250	†Oxygen quenching; spectrum corrected for cation radical by monitoring e_{aq}^- absorption; pH 7.4		82A152
	Water	LP/TD	790, 3400	$\tau_T = 420 \mu\text{s}$		84E203
1044.	Tetrakis(trimethylaminophenyl)porphine Water	LP/TD	800, 3200 690	$\tau_T = 540 \mu\text{s}$		83E462
1045.	<i>N,N,N',N'</i> -Tetramethylbenzidine Acetonitrile	LP/SD	475, 36000 ± 4000	†Oxygen quenching (2.2 × 10 ¹⁰ L mol ⁻¹ s ⁻¹); $\tau_T = 10 \mu\text{s}$		84B066
	Acetonitrile	LP	475 ^a	†Quenching by 2,5-dimethyl-2,4-hexadiene; spectrum obtained by subtracting radical cation spectrum from a composite spectrum; $\tau_T = 5 \mu\text{s}$		84F074
	Cyclohexane	LP	475	†Triplet ET to naphthalene, oxygen quenching, intensity dependent lifetime; $E_T = 260 \pm 5 \text{ kJ mol}^{-1}$; $k_{\text{et}} = 2 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$		767177

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	EtOH/Water	LP	475		†Oxygen quenching; solvent was 1:1 EtOH to water; delay 250 ns	84E405
	MeOH	LP/ELT	475 ^b , 40000 ± 3000		†Triplet ET to naphthalene, oxygen quenching, intensity dependent lifetime; ϵ relative to cation radical in MeOH (ϵ_{475} = 41000 L mol ⁻¹ cm ⁻¹) obtained by electron transfer to Eu ³⁺ ions and noting no change in absorbance at 475 nm; τ_T = ~10 μ s; E_T = 260 ± 5 kJ mol ⁻¹	767177
	SDS	LP/ELT	475 ^b , 40000 ± 2000		†Triplet ET to naphthalene, oxygen quenching, intensity dependent lifetime; ϵ relative to cation radical in MeOH (ϵ_{475} = 41000 L mol ⁻¹ cm ⁻¹) obtained by electron transfer to Eu ³⁺ ions and noting no change in absorbance at 475 nm; aqueous micelle; E_T = 260 ± 5 kJ mol ⁻¹	767177
1046.	1,1',3,3'-Tetramethyldianthrone MCH/Isohexane (93 K)	FP	490		†ESR; shoulders at 450 nm and 530 nm; lifetime measured at 103 K; τ_T = 1200 μ s	70B006
	PMMA	FP/SD	485, <26000			677223
	PMMA (140 K)	FP/SD	485, 26000		†Phosphorescence decay; τ_T = 88000 μ s	677223
1047.	3,4,7,8-Tetramethyl-1,5-diazabicyclo[3,3,0]octa-3,7-diene-2,6-dione Acetonitrile	LP	420		†Variation of decay with viscosity; ϵ > 10000 L mol ⁻¹ cm ⁻¹ ; τ_T = 2.2 μ s	79E443
1048.	1,3,3,6-Tetramethyl-6'-nitrospiro-(2,2'-indolin[2H-1]benzopyran) Toluene (298 K)	FP	460		†Oxygen quenching, activationless decay; assigned to $\pi \rightarrow \pi^*$ transition in uncolored form	67B003
1049.	N,N,N',N'-Tetramethyl-p-phenylenediamine 3-MP (77 K)	FP/SD	613 ^a , 15300 ^a 610 ^b , 15000 570 ^c , 13600 ^a 330 ^d , 5500 ^c 299 ^e , 17500 ^a		†Phosphorescence decay and oxygen quenching; "TMPD"; shoulder at 532 ^a nm; spectrum was estimated in SD region, 340-360 nm; ϵ outside SD region was measured assuming integrated ϵ was equal to that of TMPD cation, and this was used in a modified SD method	69E215
	Acetonitrile	LP	560 ^a		Spectra extracted from composite spectra; τ_T = 0.5 μ s	84B061
	Benzene	PR/ET	605, 12200		ϵ relative to benzophenone ketyl radical in cyclohexane (ϵ_{\max} = 3700 L mol ⁻¹ cm ⁻¹); ϵ obtained from a simultaneous least squares fit of data from several compounds making use of cyclohexane to benzene ϵ_{\max} ratios of 1.83 for naphthalene and 1.45 for anthracene	71E360
	Cyclohexane	LP/ET	597 ^a 570 ^b , 11900 526 ^a		†Oxygen quenching and delayed fluorescence; ϵ relative to naphthalene in cyclohexane (ϵ_{415} = 22400 L mol ⁻¹ cm ⁻¹)	707061
	Dipropyl ether (77 K)	MOD/KM	615, 24000 ± 6000 570, 24000 ± 6000 297, 28000 ± 7000			737055
	EtOH	FP	620 ^a 568 ^a			707061
	Heptane	LP	596 ^a 562 ^a		Delay 150 ns; 562 nm peak was the more intense; τ_T = 1.4 μ s	82E474
	Hexane	LP	608 ^a 551 ^a		Delay 40 ns	84B061
	Liquid paraffin	FP	598 ^a 554 ^a		†Negative comparisons with TMPD ⁺ spectra and kinetics; shoulder at 524 ^a nm; another maximum < 300 nm, and this maximum was the more intense than the 2 listed maxima; τ_1 = 82 μ s	66B003
	MCH/Isopentane (77 K)	PS	615 ^a 600 ^a 578 ^a 565 ^a 536 ^a 523 ^a		†Phosphorescence decay; glass was 1:1 MCH to isopentane; the 6 peaks occurred in pairs, with the bands formed by each pair showing indications of additional structure	66B003

Table 1. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	PMMA	PR/ET	620, 10000 580, 11000		ϵ relative to naphthalene in cyclohexane ($\epsilon_{415} = 22000$ L mol ⁻¹ cm ⁻¹), assuming no solvent effects; reference ϵ was from a citation that actually reported $\epsilon_{412.5} = 22600$ L mol ⁻¹ cm ⁻¹	710278
	Polystyrene	PR/ET	630, 15000 580, 15000		ϵ relative to naphthalene in polystyrene ($\epsilon_{425} = 11000$ L mol ⁻¹ cm ⁻¹)	710278
	Water	LP	630 570		Water	707061
1050.	Tetramethylthiuram disulfide Hexane	FP	~400		†Oxygen quenching; $\tau_T = 90 \mu\text{s}$	78B108
1051.	Tetramethylthiuram monosulfide Hexane	FP	~400		†Oxygen quenching; $\tau_T = 90 \mu\text{s}$	78B108
1052.	1,1,4,4-Tetraphenyl-1,3-butadiene Toluene	PR	395 ^a		†Triplet ET to β -carotene and oxygen quenching; $\tau_T = 0.665 \mu\text{s}$	84E144
1053.	Tetraphenylchlorin Benzene	FP	520 470		Maximum at 520 nm was judged to be of "doubtful significance" by the experimenters; benzene was wet	58R002
	Dimethyl phthalate	FP/SD	441 ^a		Shoulder at 458 nm	74B007
1054.	Tetraphenyl- <i>p</i> -dioxin Benzene	LP	550 ^a 350 ^a		†Oxygen quenching (1.7×10^9 L mol ⁻¹ s ⁻¹), triplet ET to Di- <i>tert</i> -butylselenoketone; maxima assumed solvent independent (text); $\tau_T = 0.630 \mu\text{s}$; $k_{\text{et}} = 1.8 \times 10^8$ L mol ⁻¹ s ⁻¹	79A241
	MeOH	LP	550 ^a 350 ^a		†Oxygen quenching, triplet quenching; relative intensities (1:3); $\tau_T = 0.535 \mu\text{s}$	79A241
1055.	3,3,4,5-Tetraphenyl-2(3 <i>H</i>)-furanone Benzene	LP/ET	365 \pm 5, 13000 \pm 2600		†Oxygen quenching; ϵ relative to benzophenone in benzene ($\epsilon_{332} = 7600$ L mol ⁻¹ cm ⁻¹); $\tau_T = 12 \pm 2 \mu\text{s}$; $k_{\text{et}} = 2.5 \times 10^9$ L mol ⁻¹ s ⁻¹	84E520
1056.	Tetraphenylporphinatorrhodium(III) chloride MeOH	LP/TD	830			84E045
1057.	<i>meso</i> -Tetraphenylporphine Benzene	LP/TD	790, 6000		$\tau_T = 1500 \mu\text{s}$	84E203
	Toluene	FP/SD	780, 6000 690, 3500 430, 83000 390, 42000		$\tau_T = 1350 \mu\text{s}$	60E006
	Toluene	LP	780			80B101
1058.	Tetraphenylporphyrin dianion THF	FP	460		Electron photoejection ruled out by scavenging study; half-life = 1000 μs	79B114
1059.	1,3,6,8-Tetraphenylpyrene Cyclohexane	FP/TD	525.5, 14400 510, 19700 499, 18900 488.5, 18100 478.5, 16800 460, 13900 434, 16900 429.5, 16700 426, 16500		The 1st triplet level was assigned to $^3B_{2u}^+$; four electron transitions were assigned, with the 1st (band 1), 3rd (origin, band 6), and the 4th (origin, band 7) being $^3B_{2g}^-$, while the 2nd electronic transition (origin, band 2) being $^3A_g^-$	757439

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
1060.	Thiobenzophenone Benzene	LP/RA	515 ± 5, 4100 ± 800 400 ± 5, 4800 ± 1000		ϵ relative to benzophenone in benzene ($\epsilon_{532.5} = 7630$ L mol ⁻¹ cm ⁻¹), taking $\Phi_T = 0.6$ at 337 nm excitation and taking $\Phi_T = 1$ for benzophenone; $\tau_T = 1.7 \pm 0.2$ μ s	84A221
1061.	Thioflavine PMMA (77 K)	PS	1650 1300 1200 1100 950 750			65B004
1062.	Thioindigo 2-MTHF Benzene	LP LP	588 ^a 590 ^a 530 ^a 380 ^a		\uparrow Oxygen quenching \uparrow Oxygen quenching (3.2×10^9 L mol ⁻¹ s ⁻¹); presumably twisted triplet since spectrum independent of amount of <i>cis</i> or <i>trans</i> isomers used; $\tau_T = 0.279$ μ s	79E325 78F030
	Butyronitrile Chloroform Dichloromethane	LP LP LP	580 603 590 ^a 530 ^a 380 ^a		\uparrow Oxygen quenching \uparrow Oxygen quenching; presumably twisted triplet since spectrum independent of amount of <i>cis</i> or <i>trans</i> isomers used; $\tau_T = 0.158$ μ s	79E325 81E639 78F030
	Glycerol triacetate (203 K) Glycerol triacetate	LP LP	580 375 575 370		$\tau_T = 37$ μ s $\tau_T = 0.20$ μ s	79E543 79E543
	MCH/Toluene (77 K) MCH/Toluene Methylene chloride	LP LP LP	587 ^a 590 590 370		Solvent was 1:1 MCH to toluene \uparrow Oxygen quenching; solvent was 1:1 MCH to toluene $\tau_T = 0.15$ μ s	79E325 79E325 79E543
	PMMA	LP	580 375			79E543
1063.	Thionine cation Acetonitrile	FP	810 ^a 740 ^a 430 ^a			78A447
	MeOH	FP	780 420			747038
	MeOH	FP/SD	780 ^a , 13200 700 ^a , 7000 ^a 430 ^a , 12000 ^a		Upper bound on ϵ ; basic form of triplet; $\tau_T = 12$ μ s	777242
	Water	FP/SD	760 ^a , 9400 ^a 690 ^a , 6000 ^a 415 ^a , 12000 ^a		$\tau_T = 1.7$ μ s; pH 8.	64E017
	Water	FP/SD	775, 10000		$pK_a = 6.3 \pm 0.1$	67C001
	Water	FP	770		pK_a 8.95; pH 7.5-9.5	757148
	Water	FP	770 ^a		Shoulder at 730 nm	777315
1064.	Thionine cation, conjugate monoacid Sulfuric acid (295 K)	FP/SD	495 ^a , 5000 ^a		Absorption spectra extends to 700 nm and has at least one additional maximum which was more intense than at 495 nm, but is masked by bleaching of ground state; temperature ± 1 K; pH ~ 0.7	61A001
	Water	FP/SD	655 ^a , 15400 ^a 375 ^a , 14000 ^a		pH 1.	64E017
	Water	FP	650		$\tau_T = 16 \pm 2$ μ s; pH 2.5	67C001
	Water	FP	645 ^a			777315
	Water	LP/SD	650, 16000 380, 14000		SD assuming no triplet absorption at 630 nm; solution 0.01 mol L ⁻¹ acid; pH 2	82E232
	Water/Acetonitrile	LP/SD	650, 20000 380, 15500		SD assuming no triplet absorption at 630 nm; solution 0.01 mol L ⁻¹ acid; solvent was 1:1 water to acetonitrile	82E232
	Water/EtOH	LP/SD	650, 20500 380, 16500		SD assuming no triplet absorption at 630 nm; solution 0.01 mol L ⁻¹ acid; solvent was 1:1 water to EtOH	82E232

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
1065.	Thiopyronine Water	FP	690 470-480 440 400 370 340		Solution was phosphate buffered; $\tau_T = 420 \mu\text{s}$; pH 7.4	687142
1066.	Thiopyronine cation Acetonitrile MeOH	FP/? FP/?	660 ^b , 4560 720 ^a 660 ^b , 4200 470 ^a 420 ^a		ϵ method unspecified; $\tau_T = \sim 80 \mu\text{s}$ Radical ($\lambda_{\max} = 425 \text{ nm}$) and radical ion ($\lambda_{\max} = 470 \text{ nm}$) contributions in the blue unsubtracted; ϵ method unspecified; $\tau_T = \sim 80 \mu\text{s}$	80A369 80A369
	Water	FP	704 ^a 344 ^a		$\tau_T = 420 \mu\text{s}$; pH 7.4	697009
	Water	FP/?	660 ^b , 4350		ϵ method unspecified; $\tau_T = \sim 80 \mu\text{s}$	80A369
1067.	4-Thiouridine Acetonitrile	LP/ET	520, 50000		ϵ relative to retinol in hexane ($\epsilon_{405} = 80000 \text{ L mol}^{-1} \text{ cm}^{-1}$)	83E625
	Water	LP/ET	520, 50000		ϵ relative to retinol in hexane ($\epsilon_{405} = 80000 \text{ L mol}^{-1} \text{ cm}^{-1}$)	83E625
1068.	9H-Thioxanthene-9-thione Benzene	LP/RA	505 \pm 5, 2500 \pm 500 350 \pm 5, 10600 \pm 2100		ϵ relative to benzophenone in benzene ($\epsilon_{532.5} = 7630 \text{ L mol}^{-1} \text{ cm}^{-1}$), taking $\Phi_T = 0.8$ at 337 nm excitation and taking $\Phi_T = 1$ for benzophenone; SD method gave similar ϵ ; $\tau_T = 0.83 \pm 0.08 \mu\text{s}$	84A221
1069.	Thioxanthene-9-one Acetonitrile Acetonitrile Benzene	FP LP LP/RA	625 \pm 5 620 650, 30000 \pm 5000		†Triplet ET to triplet acceptors; $\tau_T = 73 \mu\text{s}$ ϵ relative to anthracene in liquid paraffin ($\epsilon_{424} = 71000 \text{ L mol}^{-1} \text{ cm}^{-1}$), assuming $\Phi_T = 0.7$ for anthracene in benzene and $\Phi_T = 1.0$ for thioxanthene-9-one and assuming ϵ independent of solvent; wavelength assumed to be a maximum	737190 84B116 79E099
	Benzene	LP/ET	650, 30000 \pm 5000		ϵ relative to anthracene in liquid paraffin ($\epsilon_{424} = 71000 \text{ L mol}^{-1} \text{ cm}^{-1}$) assuming ϵ independent of solvent; wavelength assumed to be a maximum	79E099
	Benzene	FP	650-670		$\tau_T = 95 \mu\text{s}$	81A294
	tert-BuOH	FP	620 \pm 5		†Triplet ET to triplet acceptors	737190
1070.	Thymidine Acetonitrile	LP/ET	370 ^b , 3600		†Triplet ET to retinol, oxygen quenching; ϵ relative to retinol in hexane ($\epsilon_{405} = 80000 \text{ L mol}^{-1} \text{ cm}^{-1}$), assuming no change in extinction coefficient of standard in acetonitrile, and making kinetic corrections; $k_{et} = 6 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	79B087
	Water	LP/RF	370 ^b , 3600		†Oxygen quenching; ϵ relative to thymidine in acetonitrile ($\epsilon_{370} = 3600 \text{ L mol}^{-1} \text{ cm}^{-1}$), assuming oscillator strength independent of solvent; $\tau_T = 25 \mu\text{s}$	79B087
1071.	Thymidine 5'-monophosphate EtOH	LP/ET	370 ^b , 4000		†Triplet ET to retinol, oxygen quenching; ϵ relative to retinol in hexane ($\epsilon_{405} = 80000 \text{ L mol}^{-1} \text{ cm}^{-1}$), assuming no change in extinction coefficient of standard in EtOH, and making kinetic corrections; $\tau_T = 25 \mu\text{s}$; $k_{et} = 2 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	79B087
	Water	LP/RF	370 ^b , 4000		†Oxygen quenching; ϵ relative to thymidine 5'-monophosphate in EtOH ($\epsilon_{370} = 4000 \text{ L mol}^{-1} \text{ cm}^{-1}$), assuming oscillator strength independent of solvent	79B087

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
1072.	Thymine Acetonitrile	LP/ET	340 ^b , 4000		†Triplet ET to β -carotene and retinol; ϵ relative to retinol in hexane ($\epsilon_{405} = 80000$ L mol ⁻¹ cm ⁻¹), assuming no solvent effect on reference f ; no maximum was present, but 300 nm appeared to be a shoulder; $\tau_T = 2.5$ μ s; $k_{et} = \sim 6 \times 10^9$ L mol ⁻¹ s ⁻¹	757510
	Water	FP	340			717071
	Water	LP/ET	340 ^b , 4000		†Triplet ET to β -carotene and retinol; ϵ relative to retinol in hexane ($\epsilon_{405} = 80000$ L mol ⁻¹ cm ⁻¹), assuming no solvent effects on reference f on going to acetonitrile and on thymine on going from acetonitrile to water; shoulder at ~ 340 nm; $\tau_T = 0.6$ μ s	757510
1073.	Thymine, negative ion Water	FP	421 ^a		†Oxygen quenching; shoulder at 490 ^a nm; another maximum <330 nm; $pK_b \sim 3$; pH 12	727036
1074.	Toluene/Tetracyanobenzene Toluene/Et ₂ O (77 K)	MOD	1250 559		Glass was 1:1 toluene to Et ₂ O	72E276
1075.	<i>s</i> -Triazine Acetonitrile	LP/HAT	303 245, ~ 6000		Shoulders at ~ 450 and ~ 345 nm; ϵ relative to H adduct of <i>s</i> -triazine in acetonitrile ($\epsilon_{315} = 1900$ L mol ⁻¹ cm ⁻¹); $\tau_T = 0.91$ μ s	757066
1076.	Tribenzo[<i>a,c,h</i>]phenazine PMMA (77 K)	PS	525		†Phosphorescence decay (?); $\tau_T = 7.4 \times 10^5$ μ s	70E291
	PMMA (293 K)	FP	525		Decay was nonexponential, and lifetime was calculated from a terminal 1st-order rate constant; $\tau_T = 3.2 \times 10^5$ μ s	70E291
1077.	2,9,10-Trichloroanthracene Liquid paraffin	FP	419		Viscosity of solvent was 0.167 N·s/m ² ; $\tau_T = 400$ μ s	62E009
1078.	1,3,5-Tricyanobenzene Et ₂ O/Isopentane (77 K)	PS	320		Glass was 1:1 Et ₂ O to isopentane	75B004
1079.	Tricyclo[8.2.2.2 ^{4,7}]hexadeca-4,6,10,12,13,15-hexaene-5,15-dicarbonitrile Acetonitrile	LP	520 ^a		100 ns delay	776187
	Toluene	LP	520 ^a		100 ns delay	776187
1080.	1,1,1-Trifluoroacetylacetone EtOH/MeOH (118 K)	FP	380		Solvent was 3:1 EtOH to MeOH	68B005
1081.	4-(Trifluoromethyl)acetophenone Benzene	LP/RA	455 ^a , 2800 ^a 426 ^a , 2700 ^a		ϵ relative to benzophenone in cyclohexane ($\epsilon_{533} = 7630$ L mol ⁻¹ cm ⁻¹ , assuming no solvent effect from benzene), and taking all the triplet yields to be unity	737198
1082.	4-(Trifluoromethyl)benzophenone Acetonitrile/Water	LP/ET	530, 5800 320, 12400		Solvent was 9:1 acetonitrile to water; ϵ relative to 1-methylnaphthalene in acetonitrile/water ($\epsilon_{415} = 11200$ L mol ⁻¹ cm ⁻¹)	84B033
1083.	4,4,4-Trifluoro-1-phenyl-1,3-butanedionatosodium EPA	LP/SD	620 ^b , 11000		$\tau_T = 0.04$ μ s	84E232
	EPA	LP/ET	620 ^b , 12000		ϵ relative to naphthalene in cyclohexane ($\epsilon_{414} = 24500$ L mol ⁻¹ cm ⁻¹); $\tau_T = 0.03$ μ s; $k_{et} = 4.1 \times 10^9$ L mol ⁻¹ s ⁻¹	84E232
1084.	2,4,6-Triisopropylbenzophenone Acetonitrile	LP	750 305		†Triplet ET to <i>cis</i> -1,3-pentadiene; relative intensities (1:2); decays into biradical; $\tau_T = 0.255 \pm 0.013$ μ s	80B088

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued.

No.	Solvent	Method	λ_{\max} /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
1085.	2,3,5-Trimethyl-1,4-benzoquinone Water	LP	480 440		$\tau_T = 5.9 \mu s$	80B112
1086.	(<i>all-E</i>)-3,7,12-Trimethyl-1,14-bis(2,6,6-trimethyl-1-cyclohexen-1-yl)-1,3,5,7,9,11,13-tetradecaheptaene Benzene (296 K)	LP-ET	495		Triplet ET from chlorophyll <i>a</i> ; oxygen quenching (3.2×10^9 L mol ⁻¹ s ⁻¹); $\tau_T = 10 \mu s$; $k_{et} = 0.8 \times 10^9$ L mol ⁻¹ s ⁻¹	73E347
1087.	1,1,2-Trimethyl-2-(2-naphthyl)ethylene Benzene	LP-ET	390 ^a		†Triplet ET from xanthone and oxygen quenching (4.7×10^9 L mol ⁻¹ s ⁻¹); $\tau_T = 0.07 \mu s$	84B090
1088.	1,3,3-Trimethyl-6'-nitrospiro-(2,2'-indolin[2H-1]benzopyran) MCH	LP	430		†Oxygen quenching; ground state dimers form readily; $\tau_T = \sim 20 \mu s$	80N064
	Toluene (298 K)	FP	443		†Oxygen quenching, activationless decay; ϵ estimated > 10000 L mol ⁻¹ cm ⁻¹ ; assigned to $\pi \rightarrow \pi^*$ transition in uncolored form	67B003
1089.	3,5,5-Trimethyl-2-phenoxy-2-cyclohexene-1-one MCH	LP	380 ^a		†Oxygen quenching; spectrum obtained by extrapolation to zero time	81F111
1090.	4-[Tri(4-methylphenyl)porphyrinyl]-3-phenoxypropyl β - <i>apo</i> -6'-carotenatene Benzene	LP	~ 540 ~ 440		The 80 ns lifetime associated with 440 nm triplet, assigned to porphyrin chromophore; 2.4 μs lifetime and 540 nm band assigned to carotenoid chromophore; 440 nm species is precursor of 540 nm species; $\tau_T = 2.4$; 0.080 μs	82E174
1091.	4,5',8-Trimethylpsoralen EtOH (180 K)	FP	481 ^a 461 ^a 376 ^a		†Triplet ET from naphthalene; shoulder at 498 ^a nm	83B122
	MeOH	LP/ET	470, 33000 \pm 3000 370, 27000 ^a		†Triplet ET to β -carotene; ϵ relative to β -carotene in hexane ($\epsilon_{515} = 240000$ L mol ⁻¹ cm ⁻¹) assuming ϵ independent of solvent; $\tau_T = 7.1 \mu s$	79B042
1092.	Triphenylamine 3-MP (77 K)	PS	525		$\tau_T = (9 \pm 3) \times 10^5 \mu s$	67E106
	EPA (77 K)	PS	531		†Phosphorescence decay; $\tau_T = (9 \pm 2) \times 10^5 \mu s$	67E106
	MCH	LP	520			72B002
	Toluene/EtOH (77 K)	MOD	583 ^a 410 ^a		Glass was 19:1 toluene to EtOH; 583 nm peak was the more intense	719059
1093.	Triphenylamine, conjugate acid Acetic acid/MeOH/- Et ₂ O (77 K)	MOD/KM	579 ^b , 5900 546 ^b , 9900 436 ^b , 2300		Glass was 14:3:3 acetic acid to MeOH to Et ₂ O	82E648
1094.	1,3,5-Triphenylbenzene 2-PrOH/Isocetane/- Et ₂ O (81 K)	FP	500 346		†Phosphorescence decay; glass was 1:3:3 2-PrOH to isocetane to Et ₂ O; $\tau_T = 3.9 \times 10^6 \mu s$	81B119
	Alcohol/Ether (77 K)	MOD	494		Glass was 2:1 alcohol to ether	76E682

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
1095.	Triphenylene					
	2-MTHF (77 K)	PS/ESR	431, 16500 408 ^a , 9900 ^a		†ESR; oscillator strength = 0.085	68D211
	3-MP (77 K)	PS	721 657			67E106
	? (77 K)	PS	434 ^a 410 ^a		†Phosphorescence decay and ESR lifetime; glass used was either 2-MTHF or 3-MP; $\tau_T = 1.37 \times 10^7 \mu s$	67E108
	Benzene	FP	428 407		Relative intensities (10:9)	61E005
	Benzene	LP	428			70E288
	Benzene	LP/ET	435, 6000 \pm 500		†Triplet ET from benzophenone; ϵ relative to benzophenone in benzene ($\epsilon_{532.5} = 7630 \text{ L mol}^{-1} \text{ cm}^{-1}$)	757282
	Butane/Isopentane (77 K)	PS/IV	430, 7000 \pm 2000 408 ^a , ~3800		Glass was 3:7 butane to isopentane; ϵ_T was computed from $OD = \epsilon_T \tau_{p10}(1 - \Phi_f)$ after the optical density, OD, was extrapolated to zero intensity of the monitoring beam (namely $i_0 \rightarrow 0$), here τ was lifetime of triplet, and Φ_f was fluorescence quantum yield; oscillator strength = 0.05	65E031
	Cyclohexane	LP	430		†Rise time of transient was the same as the decay time of the singlet	68B006
	EPA (77 K)	PS	431		†Phosphorescence decay	51E001
	EPA (77 K)	PS	727 661		†Phosphorescence decay; $\tau_T = (1.34 \pm 0.3) \times 10^7 \mu s$	67E106
	EPA (82 K)	PS/KM	430, 16800 \pm 300			68B003
	EPA (77 K)	FP/SD	430, 15600 \pm 2300 409 ^a , 8200 ^a		$\tau_T = 1.38 \times 10^7 \mu s$; oscillator strength = 0.11	69F389
	EPA (77 K)	PS/KM	429 ^a , 17100 ^a 407 ^a , 11200 ^a 392 ^a , 7700 ^a		Solvent, temperature and extinction method assumed from earlier work; polarization also measured	69E213
	EPA (77 K)	PS/KM	430, 16800		λ_{\max} assumed from previous work	69E212
	EtOH (77 K)	MOD/KM	340 ^a , 8500 310 ^a , 2500 295 ^a , 3200 274.3 ^b , 17500 270, 36500 262.5 ^b , 27500 250 ^a , 45000			757594
	EtOH (77 K)	MOD	727 650 ^a 431 408 ^a		Relative intensities (2:1:4:1); halfwidth of principal maximum 680 cm ⁻¹	777538
	EtOH/MeOH (113 K)	FP/TD	350, > 13000		Solvent was 3:1 EtOH to MeOH; lower limit assumes total ground state depletion	67B004
	Ethylene glycol	FP	428 407		Relative intensities (10:9); $\tau_T = 1000 \pm 100 \mu s$	61E005
	Hexane	FP	428.1			54E001
	Hexane	FP/SD	428, 4100 404, 3500 346, 7000		Band at 346 nm was assigned to the second electronic transition; oscillator strength = 0.025, 0.06	58E001
	Hexane	FP	428 405		Relative intensities (10:9); $\tau_T = 55 \pm 9 \mu s$	61E005
	Hexane (77 K)	PS	431 407 385 360 350		First 3 bands were of medium intensity and were assigned to one electronic transition; other 2 bands were of strong intensity and were assigned to another electronic transition	63B002
	Hexane (77 K)	PS	432.6 409.0 381.2 361.8		Sample was a Shpolskii matrix; first two bands were assigned to one electronic transition, and the last two were assigned to another one	65B003
	Hexane	PS	722 655		Weak bands in the red only studied	67B005

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
	Hexane (77 K)	FS	722.0 654.9 432.0 416.7 408.5 396.0 381.0 369.90 362.1 354.6		Relative intensities (15:10:100:42:62:33:40:35)	69E211
	MCH (77 K)	PS/KM	432, 14400 ± 1000		Photoreaction with solvent measured; $\tau_T = 14 \times 10^6$ μ s	81F315
	MCH	LP	424 ^a 405 ^a		424 nm peak was the more intense	82F367
	PMMA	LP	428			70E288
	Toluene (77 K)	MOD	737 660 ^a 443		Relative intensities (4:2:3); halfwidth of principal maximum 2000 cm ⁻¹	77F538
1096.	Triphenylene- <i>d</i> ₁₂ EPA (77 K)	FP/SD	431, 12000 ± 1800		$\tau_T = 1.73 \times 10^7$ μ s	69F389
1097.	Triphenylene/Chloranil Liquid paraffin	MOD	415		Mull	71E361
1098.	Triphenylethylene Toluene	PR	366 ^a		†Triplet ET to β -carotene; maximum uncertain because of low intensity of monitoring light (private communication, P.K. Das, 1984); $\tau_T = 0.130$ μ s	84E144
1099.	3,3,5-Triphenyl-2(3 <i>H</i>)-furanone Benzene	LP/ET	330 ± 5, 13000 ± 2600		†Oxygen quenching; ϵ relative to benzophenone in benzene ($\epsilon_{332} = 7600$ L mol ⁻¹ cm ⁻¹); $\tau_T = 1.2 \pm 0.1$ μ s; $k_{et} = 2.1 \times 10^9$ L mol ⁻¹ s ⁻¹	84E520
1100.	Triphenylmethane MCH (77 K)	PS	427 ^a			727440
1101.	Triphenylmethyl cation Sulfuric acid/Acetic acid (77 K)	PS	340 ^a		†Oxygen quenching; solvent was 1:1 sulfuric acid to acetic acid; pH Acidic	78A180
	Sulfuric acid/Acetic acid	FP	580 ^a 340		†Oxygen quenching; solvent was 1:1 sulfuric acid to acetic acid; pH Acidic	78A180
1102.	1,3,3-Triphenylprop-2-en-1-one MeOH	LP	440		†Oxygen quenching; $\tau_T = 0.325$ μ s	84E520
1103.	Tris(2,2'-bipyridine)iridium(III) ion MeOH/Water	LP	567 ^a 519 ^a 500 ^a 463 ^a 451 ^a 438 ^a 424 ^a 404 ^a 379 ^a			79B098
1104.	Tris(2,2'-bipyridine)osmium(II) ion Water	LP/SM	467 ^a , 5000 ^a 365 ^a , 17700 ^a		$\tau_T = 0.019$ μ s	80E040

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
1105.	Tris(2,2'-bipyridine)ruthenium(II) ion					
	Acetonitrile	LP/SD	360 ^b , 27300 ± 1500		†Triplet ET to retinol; $\tau_T = 1.1 \pm 0.1 \mu\text{s}$	767180
	Water	LP/SD	360 ^b , 27300 ± 1500		†Oxygen quenching ($3.3 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$); $\tau_T = 0.67 \pm 0.02 \mu\text{s}$	767180
	Water	LP/TD	453 ^b , 7700 ^b 445 ^a , 8000 ^{a,c} 370 ^a , 13500 ^{a,c}		In estimation of ϵ , TD assumed from saturation of OD at intensities above $0.3 \times 10^{18} \text{ photons/cm}^2$; shoulder at 350 nm	79B161
	Water	LP/SM	452 ^b , 2100 370 ^a , 28900 ^a 315 ^a , 36600 ^a 252 ^a , 41600 ^a			79B032
	Water	LP/SM	430 ^a , 6500 ^a 364 ^a , 25400 ^a			80E040
	Water	LP	368 310 255		pH 12.7	82A365
	Water	LP/TD	541, 3000 444, 5000 376, 29000 320, 37000 250 ^b , 40000			83B098
1106.	Tris(1,10-phenanthroline)rhodium(III) ion					
	Water (298 K)	LP/RA	490, 4300 ^a		ϵ relative to benzophenone in benzene ($\epsilon_{332} = 7630 \text{ L mol}^{-1} \text{ cm}^{-1}$, $\Phi_T = 1$), Φ_T for compound measured as 1; units for ϵ are assumed since they were not specified in the paper; $\tau_T = 0.25 \pm 0.02 \mu\text{s}$	84B055
	Water/Ethylene glycol (77 K)	LP	490		†Phosphorescence decay; glass was 1:1 water to ethylene glycol; $\tau_T = (4.8 \pm 2) \times 10^4 \mu\text{s}$	84B055
1107.	Tris(4,4,4-trifluoro-1-phenyl-1,3-butanedionato)lanthanum(III)					
	EPA	LP/SD	620, 20000		$\tau_T = 2.6 \mu\text{s}$	84E232
	EPA	LP/ET	620, 27000		ϵ relative to naphthalene in cyclohexane ($\epsilon_{414} = 24500 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 2.7 \mu\text{s}$; $k_{et} = 1.0 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	84E232
1108.	Tris(4,4,4-trifluoro-1-phenyl-1,3-butanedionato)lutetium(III)					
	EPA	LP/SD	620 ^b , 28000		$\tau_T = 1.4 \mu\text{s}$	84E232
	EPA	LP/ET	620 ^b , 28000		ϵ relative to naphthalene in cyclohexane ($\epsilon_{414} = 24500 \text{ L mol}^{-1} \text{ cm}^{-1}$); $\tau_T = 1.5 \mu\text{s}$; $k_{et} = 1.0 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$	84E232
1109.	Tryptophan					
	PVA (293 K)	FP	480 ^a		†Phosphorescence decay	757469
	Water (77 K)	PS	464 ^a		†Phosphorescence decay; 4 mol L ⁻¹ CaCl ₂ aqueous glass	700011
	Water (298 K)	LP	450		†Triplet ET to anthracene in EtOH; lifetime was measured at pH 7.5; spectrum is the difference between spectra at 100 ns in O ₂ -free solution and 1 μs in O ₂ -containing solution; $\tau_T = 14.3 \pm 1.4 \mu\text{s}$; pH 5.4	757163
	Water	LP/COM	460, 5000		ϵ corrected for radical and hydrated electron absorption at 460 nm ϵ from quadratic fit to absorption decay curve assuming triplet-triplet annihilation with rate constant $10^{10} \text{ L mol}^{-1} \text{ s}^{-1}$; pH 7	777432
1110.	DL-Tryptophan					
	PVA	FP/COM	485 ^a , 3700 ± 400		†Phosphorescence decay; ϵ estimated based on $\Phi_T = 0.55$ and the assumption of no internal conversion	753103
1111.	L-Tryptophan					
	Ethylene glycol/Water (77 K)	PS/KM	450, 8000		†Phosphorescence decay; glass was 1:1 ethylene glycol to water by volume; $\tau_T = 6.6 \times 10^6 \mu\text{s}$	716330
	Water	FP	460		pH 5.4	727041

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
1112.	Tyrosine Water	LP	~575 295 250		Zwitterion; lifetime was measured at pH 6.0 and extrapolated to zero tyrosine concentration; pK _a 's of triplet are 2.5 and 9.7; spectrum is the difference between spectra at 20 ns and 15 μ s; $\tau_T = 5.5 \pm 1.5 \mu$ s; pH 7.5	757161
1113.	L-Tyrosylglycine Water	LP	~385 ^a		Another maxima below 250 nm that was at least 3 times as high as the maximum at 385 ^a nm; spectrum is the difference between spectra at 20 ns and 15 μ s; $\tau_T = 3.4 \pm 0.2 \mu$ s; pH 6.0	757161
1114.	Ubiquinone 30 Benzene	PR/ET	430 ^b , 13000		†Triplet ET from anthracene; ϵ relative to biphenyl in benzene ($\epsilon_{367} = 27100$ L mol ⁻¹ cm ⁻¹); this ϵ may be too high, see comment under ubiquinone [743062]; half-life = 0.45 μ s; $k_{et} = 5.1 \times 10^9$ L mol ⁻¹ s ⁻¹	723043
	Cyclohexane	LP/ET	440 \pm 5, 8000 \pm 1600		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹); listed value updates authors' earlier value, $\epsilon = 19000$ L mol ⁻¹ cm ⁻¹ , from [723043] which was in error because of spectral overlap with the donor	743062
	Cyclohexane	PR/ET	440 \pm 5, 7200 \pm 1600		ϵ relative to biphenyl in cyclohexane ($\epsilon_{361} = 42800$ L mol ⁻¹ cm ⁻¹)	743062
1115.	Uracil Acetonitrile	LP/ET	340, 2750		†Triplet ET to β -carotene and retinol; ϵ relative to retinol in hexane ($\epsilon_{405} = 80000$ L mol ⁻¹ cm ⁻¹), assuming no solvent effect on reference f ; $\tau_T = 0.6 \mu$ s; $k_{et} = \sim 8 \times 10^9$ L mol ⁻¹ s ⁻¹	757510
	Water	FP	380		†Oxygen quenching; $\tau_T = 6.1 \pm 0.5 \mu$ s	697253
	Water	LP/ET	360, 2600 ^a		†Triplet ET to β -carotene and retinol; ϵ relative to retinol in hexane ($\epsilon_{405} = 80000$ L mol ⁻¹ cm ⁻¹), assuming no solvent effects on reference f on going to acetonitrile and on uracil on going from acetonitrile to water; $\tau_T = 0.35 \mu$ s	757510
1116.	Uridine Acetonitrile	LP/ET	370 ^b , 6400		†Triplet ET to retinol, oxygen quenching; ϵ relative to retinol in hexane ($\epsilon_{405} = 80000$ L mol ⁻¹ cm ⁻¹), assuming no change in extinction coefficient of standard in acetonitrile, and making kinetic corrections; $k_{et} = 6 \times 10^9$ L mol ⁻¹ s ⁻¹	79B087
	Water	LP/RF	370 ^b , 6400		†Oxygen quenching; ϵ relative to uridine in acetonitrile ($\epsilon_{370} = 6400$ L mol ⁻¹ cm ⁻¹), assuming oscillator strength independent of solvent; $\tau_T = 20 \mu$ s	79B087
1117.	Uridine monophosphate EtOH	LP/ET	390, 9000		†Triplet ET to retinol, oxygen quenching; ϵ relative to retinol in hexane ($\epsilon_{405} = 80000$ L mol ⁻¹ cm ⁻¹), assuming no change in extinction coefficient of standard in EtOH, and making kinetic corrections; $k_{et} = 5 \times 10^9$ L mol ⁻¹ s ⁻¹	79B087
	Water	LP/RF	390 ^b , 9000		†Oxygen quenching (3×10^9 L mol ⁻¹ s ⁻¹); ϵ relative to uridine monophosphate in EtOH ($\epsilon_{390} = 9000$ L mol ⁻¹ cm ⁻¹), assuming oscillator strength independent of solvent; $\tau_T = 33 \mu$ s	79B087
1118.	Uroporphyrin I, octamethyl ester Benzene	PR/ET	440 ^b , 32000 405 ^a		†Oxygen quenching (1.5×10^9 L mol ⁻¹ s ⁻¹); ϵ relative to biphenyl in benzene ($\epsilon_{360} = 27100$ L mol ⁻¹ cm ⁻¹); $\tau_T = 270 \mu$ s	80E200
	Benzene	LP/TD	440 ^b , 26000 405 ^a		†Oxygen quenching (1.5×10^9 L mol ⁻¹ s ⁻¹); $\tau_T = 270 \mu$ s	80E200

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
1119.	2-Vinylnaphthalene					
	Benzene	PR/ET	426 ^a , 13200		†Triplet ET from biphenyl; ϵ relative to biphenyl in benzene ($\epsilon_{367} = 27100$ L mol ⁻¹ cm ⁻¹); $\tau_T = 60$ μ s; $k_{et} = 7.1 \times 10^9$ L mol ⁻¹ s ⁻¹	79E666
	Benzene	LP-ET	417 ^a 396 ^a		†Triplet ET from xanthone and oxygen quenching (3.9×10^9 L mol ⁻¹ s ⁻¹); there was a weak, broad maximum in the region of 600 nm; $\tau_T = 0.25$ μ s	84B090
1120.	9H-Xanthene-9-thione					
	Benzene	LP/RA	460 \pm 5, 5500 \pm 1100 345 \pm 5, 15400 \pm 3100		ϵ relative to benzophenone in benzene ($\epsilon_{332.5} = 7630$ L mol ⁻¹ cm ⁻¹), taking $\Phi_T = 0.8$ at 337 nm excitation and taking $\Phi_T = 1$ for benzophenone; SD method gave similar ϵ ; $\tau_T = 1.8 \pm 0.2$ μ s	84A221
1121.	Xanthone					
	1,1,2-Trichloro- trifluoroethane	LP	610		$\tau_T = 0.42$ μ s	767171
	2-ProOH (295 K)	LP	610 ^a		$\tau_T = 0.370$ μ s	80A338
	Benzene	LP/RA	610, 5300 \pm 700		†Triplet ET to naphthalene, oxygen quenching (5.6×10^9 L mol ⁻¹ s ⁻¹); ϵ relative to benzophenone in benzene ($\epsilon_{325} = 7630$ L mol ⁻¹ cm ⁻¹) assuming $\Phi_T = 1$ for both molecules; $\tau_T = 0.092$ μ s; $k_{et} = (9.5 \pm 1.0) \times 10^9$ L mol ⁻¹ s ⁻¹	767171
	Benzene	LP/RA	600, 8600 \pm 2000		ϵ relative to anthracene in liquid paraffin ($\epsilon_{424} = 71000$ L mol ⁻¹ cm ⁻¹), assuming $\Phi_T = 0.7$ for anthracene in benzene and $\Phi_T = 1.0$ for xanthone and assuming ϵ independent of solvent; wavelength assumed to be a maximum	79E099
	Benzene	LP	650		100 ps delay; rise time of 0.013 ns	79B007
	CCl ₄ (295 K)	LP	655		$\tau_T = 0.70$ μ s	80A338
	Cyclohexane	LP	~ 610		$\tau_T = \sim 0.02$ μ s	767171
	EtOH/Water	LP	590		†Oxygen quenching (2.4×10^9 L mol ⁻¹ s ⁻¹); solvent was "10 wt% aqueous ethanol"; $\tau_T = 17.9$ μ s	767171
	EtOH/Water	LP	605		†Triplet ET to naphthalene, oxygen quenching (2.8×10^9 L mol ⁻¹ s ⁻¹); solvent was "95%" ethanol; $\tau_T = 1.41$ μ s; $k_{et} = (5.6 \pm 0.6) \times 10^9$ L mol ⁻¹ s ⁻¹	767171
	Poly(styrenesulfonate) /DTB/Water	LP	600 ^a		Xanthone associated with aggregates in aqueous medium; spectrum averaged during first 40 ns; spectrum blue-shifts at longer times indicating a shift in position in the aggregate	84N163
	SDS	LP	607 ^a		Aqueous micelles	83N127
	SDS	LP	610		Aqueous micelles	84N197
1122.	o-Xylene					
	Cyclohexane (289.5 K)	LP	$\sim 340^a$		Delay 200 ns	83E483
1123.	Zeaxanthin					
	?	FP-ET	505 ^a		†Triplet ET from anthracene; it was not reported whether the solvent was hexane or benzene	733001
	Benzene	PR-ET	520		†Triplet ET from naphthalene; $\tau_T = 9.4$ μ s	80A143
1124.	Zinc(II) chlorophyll a					
	Benzene	FP	450		Benzene was wet	58R002
1125.	Zinc(II) cytochrome c					
	Water (288.7 K)	LP	462		†Phosphorescence decay, oxygen quenching; cytochrome c is from horse heart, type III; $\tau_T = 7000$ μ s	81E091
	Water/Glycerol (288.7 K)	LP	462		†Phosphorescence decay, oxygen quenching; cytochrome c is from horse heart, type III; solvent contains 33% glycerol; $\tau_T = 14100$ μ s	81E091
1126.	Zinc(II) etioporphyrin I					
	Benzene	FP	510 ^a 430 ^a		Shoulder at 462 ^a nm; 430 nm was the more intense peak	747293
	DMF	FP/SD	440, 99000 411 ^b , 45300			73E345

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
1127.	Zinc(II) etioporphyrin I/Hexachloroethane Benzene	FP	434 ^a		Shoulder at 458 ^a nm	747293
1128.	Zinc(II) etioporphyrin I/ <i>trans</i> -4-Nitrostilbene Benzene	FP	450		Triplet exciplex; assigned as a 1:2 complex of the porphyrin to the stilbene	717463
1129.	Zinc(II) etioporphyrin I/4-Nitrotoluene Benzene	FP	440		Triplet exciplex; assigned as a 1:2 complex of the porphyrin to 4-nitrotoluene	717463
	Benzene	FP	532 ^a 439 ^a		Shoulder at 462 ^a nm; 439 nm was the more intense peak	747293
1130.	Zinc(II) phthalocyanine 1-Chloronaphthalene	LP/TD	480, 34000 ± 4000		λ_{\max} and ϵ method inferred from text	81E457
	1-Chloronaphthalene	LP/SM	480, 16000 ± 3000		†Triplet ET to β -carotene; λ_{\max} inferred from text	81E457
	Benzene	FP/?	545, 44000 450, 51000		Method to be published	81E346
	Dimethylacetamide- /Water	LP/SD	490, 32500		Solvent was 7:3 dimethylacetamide to water	82A290
	PrOH	FP/SD	470, 33000			73E345
1131.	Zinc(II) porphyrin CTAC	LP/SM	840, 7000 ^a 740 ^a , 5000 ^a 460, 10000		50 ns delay; aqueous micelle; ϵ method uncertain; τ_T = 200 μ s	80N087
	DODAC	LP/SM	840, 7200 ^a 740 ^a , 5000 ^a 660 ^a , 3200 ^a		Cationic vesicle; ϵ method uncertain; τ_T = ~500 μ s	80N035
	Dimethyl phthalate	FP/SD	811 ^a , 11100 ^a 427 ^a , 167000 ^a 389 ^a , 46300 ^a		Shoulder at 487 nm	74B007
1132.	Zinc(II) protoporphyrin MeOH	FP	455			58R002
1133.	Zinc(II) tetrabenzoporphyrin Pyridine	FP/SD	490, 74000 410, 37000		†Phosphorescence decay; τ_T = 525 μ s	73E345
1134.	Zinc(II) tetraethylporphyrin Pyridine	FP/SD	845 ^a 790 ^a			74B007
1135.	Zinc(II) tetrakis(carboxyphenyl)porphyrin Water	LP	820 ^a 710 ^a 680 ^a		Most intense peaks at 710 nm and 680 nm; decay followed at 840 nm; pH 7	80A074
	Water	LP	830 730			82E622
1136.	Zinc(II) tetrakis(2,6-dimethyl-4-sulfonatophenyl)porphyrin Water	LP/TD	830, 6000		τ_T = 1300 μ s	84E203
1137.	Zinc(II) tetrakis(2- <i>N</i> -methylpyridyl)porphyrin Water	LP	850		Shoulder at 750 nm; τ_T = 1400 ± 140 μ s; pH 6.0	84E346
1138.	Zinc(II) tetrakis(3- <i>N</i> -methylpyridyl)porphyrin Water	LP	830		Shoulder at 750 nm; τ_T = 2000 ± 200 μ s; pH 6.0	84E346
	Water	LP/SD	575 ^b , 3900 545 ^b , 8300 460 440 ^b , 57000 390 ^b , 18000		†Oxygen quenching; τ_1 = 1800 μ s	84A264

TABLE 6. Spectral parameters for triplet-triplet absorption of organic molecules in condensed phases — Continued

No.	Solvent	Method	λ_{\max} , /nm	ϵ_{\max} /L mol ⁻¹ cm ⁻¹	Comment	Ref.
1139.	Zinc(II) tetrakis(4-N-methylpyridyl)porphyrin					
	Water	LP	1040 ^a 1000 ^a 950 ^a 890 ^a 830 ^a 790 ^a 730 ^a		Most intense peaks at 1040 nm and 840 nm; decay followed at 840 nm; $\tau_T = 1.3 \times 10^3 \mu\text{s}$; pH 7	80A074
	Water	LP/TD	1020, 7200 \pm 700 470 ^b , 60800 \pm 6100 465 ^b , 38400 \pm 3800		Shoulders at 949 and 860 nm	82E622
	Water	LP/TD	1020, 7200 \pm 700		Shoulder at 950 nm; $\tau_T = 2000 \pm 200 \mu\text{s}$; pH 6.0	84E346
1140.	Zinc(II) tetrakis(p-sulfonatophenyl)porphyrin					
	MeOH	LP	840		$\tau_T = \sim 80 \mu\text{s}$	82N068
	Water	LP	840 780 ^a 750 ^a 660 ^a		Most intense peak at 840 nm; decay followed at 840 nm; $\tau_T = 1.5 \times 10^3 \mu\text{s}$; pH 7	80A074
	Water	LP/TD	840, 6000 \pm 600 730 460 ^b , 55200 \pm 5500			82E622
	Water (293 K)	LP/TD	470 ^b , 45000 ^a		pH 7.0 \pm 0.1	82A306
	Water	LP/TD	840, 6000		$\tau_T = 1400 \mu\text{s}$	84E203
1141.	Zinc(II) tetrakis(trimethylaminophenyl)porphyrin					
	Water	LP/TD	840, 5000 730		$\tau_T = 1200 \mu\text{s}$	83E462
1142.	Zinc(II) tetraphenylchlorin					
	Dimethyl phthalate	FP/SD	447 ^a			74B007
1143.	Zinc(II) tetraphenylporphyrin					
	Benzene	LP/TD	845, 8200			84E203
	CTAC	LP	840 720 ^a 460		$\tau_T = > 2500 \mu\text{s}$	80N044
	MCH	FP/TD	470, 71000 400, 38000		†Oxygen quenching; $\tau_T = 1200 \mu\text{s}$	81E271
	SIIS	LP	840 640		Aqueous microemulsion, droplet radius 17 nm; $\tau_T = 1700 \mu\text{s}$	80N143
	Toluene	FP/SD	845, 8200 745, 5300 470, 74000 400, 42000		$\tau_T = 1250 \mu\text{s}$	60E006
	Toluene	LP	750			80B101
	Triton X-100	LP	840 460		Relative intensities (1:10); aqueous micelle; $\tau_T = 2500 \mu\text{s}$	81N034

^a Data have been obtained by computer-assisted digitization from spectrum in cited reference^b (Wavelength, extinction coefficient) pair is not necessarily related to a spectral peak^c Measurement violates Chauvenet's criterion, see Section 4.2.2^d Measurement not included in averages because of systematic error^e Measurement not included in averages since performed in mixed crystals, neat liquids, or micelles

† Evidence supporting the assignment of the transient as a triplet state

9. References to Table 6

- 41E001 Reversible photochemical processes in rigid media. A study of the phosphorescent state. Lewis, G.N.; Lipkin, D.; Magel, T.T. *J. Am. Chem. Soc.* **63**: 3005-18 (1941).
- 42B002 Reversible photochemical processes in rigid media: The dissociation of organic molecules into radicals and ions. Lewis, G.N.; Lipkin, D. *J. Am. Chem. Soc.* **64**: 2801-8 (1942).
- 51E001 Excited triplet states of some polyatomic molecules. I. McClure, D.S. *J. Chem. Phys.* **19**: 670-5 (1951).
- 53E001 Triplet states in solution. Porter, G.; Windsor, M.W. *J. Chem. Phys.* **21**: 2088 (1953).
- 54B001 The triplet-triplet absorption spectra of some aromatic hydrocarbons and related substances. Craig, D.P.; Ross, I.G. *J. Chem. Soc.* : 1589-1606 (1954).
- 54E001 Studies of the triplet state in fluid solvents. Porter, G.; Windsor, M.W. *Discuss. Faraday Soc.* **17**: 178-86 (1954).
- 55E002 Excited triplet states of polyatomic molecules. II. Flash-lamp studies on aromatic ketones. McClure, D.S.; Hanst, P.L. *J. Chem. Phys.* **23**: 1772-7 (1955).
- 55E003 Preliminary study of a metastable form of chlorophyll in fluid solutions. Livingston, R. *J. Am. Chem. Soc.* **77**: 2179-82 (1955).
- 55R001 Reversible flash bleaching of chlorophyll. Abrahamson, E.W.; Linschitz, H. *J. Chem. Phys.* **23**: 2198-9 (1955).
- 57B001 Die spektroskopisch nachweisbaren Elektronenzustände der stabilen und metastabilen Molekülform einiger Acridinfarbstoffe und des Fluoresceinkations. Zanker, V.; Miethke, E. *Z. Naturforsch., Teil A* **12A**: 385-95 (1957).
- 587005 Primary photoprocesses in quinones and dyes. I. Spectroscopic detection of intermediates. Bridge, N.K.; Porter, G. *Proc. Roy. Soc. London, Ser. A* **244**: 259-75 (1958).
- 58E001 The triplet state in fluid media. Porter, G.; Windsor, M.W. *Proc. R. Soc. London, Ser. A* **245**: 238-58 (1958).
- 58E002 The triplet state of anthracene in liquid solutions. Livingston, R.; Tanner, D.W. *Trans. Faraday Soc.* **54**: 765-71 (1958).
- 58R001 The absorption spectra and decay kinetics of the metastable states of chlorophyll A and B. Linschitz, H.; Sarkanen, K. *J. Am. Chem. Soc.* **80**: 4826-32 (1958).
- 58R002 Some properties of the ground triplet state of chlorophyll and related compounds. Livingston, R.; Fujimori, E. *J. Am. Chem. Soc.* **80**: 5610-3 (1958).
- 59B001 Contribution à l'étude de l'état métastable des molécules de colorants: Bandes d'absorption dans le domaine visible. Laffitte, E.; Nouchi, G. *C.R. Seances Acad. Sci.* **248**: 2746-8 (1959).
- 59B002 Absorption spectrum and decay kinetics of the metastable state of chlorophyll b. Claesson, S.; Lindqvist, L.; Holmstroem, B. *Nature* **183**: 661-2 (1959).
- 59E006 Reversible spectral changes in retinene solutions following flash illumination. Abrahamson, E.W.; Adams, R.G.; Wulff, V.J. *J. Phys. Chem.* **63**: 441-3 (1959).
- 600062 Transient benzyl radical reactions produced by high-energy radiation. McCarthy, R.L.; MacLachlan, A. *Trans. Faraday Soc.* **56**: 1187-1200 (1960).
- 60A001 A flash photolysis study of fluorescein. Lindqvist, L. *Ark. Kemi* **16**: 79-138 (1960).
- 60E005 Experimental study of the triplet state of anthracene in fluid solutions. Jackson, G.; Livingston, R.; Pugh, A.C. *Trans. Faraday Soc.* **54**: 1635-9 (1960).
- 60E006 Studies on metastable states of porphyrins. II. Spectra and decay kinetics of tetraphenylporphine, zinc tetraphenylporphine and bacteriochlorophyll. Pekkarinen, L.; Linschitz, H. *J. Am. Chem. Soc.* **82**: 2407-11 (1960).
- 61A001 The photoreduction of thionine by ferrous sulphate. Hatchard, C.G.; Parker, C.A. *Trans. Faraday Soc.* **57**: 1093-106 (1961).
- 61E005 Energy transfer from the triplet state. Porter, G.; Wilkinson, F. *Proc. R. Soc. London, Ser. A* **264**: 1-18 (1961).
- 61E008 Acidity constants in the triplet state. Jackson, G.; Porter, G. *Proc. R. Soc. London, Ser. A* **260**: 13-30 (1961).
- 62E006 Some flash-photolytic and photochemical studies of retinene and related compounds. Grellmann, K.-H.; Memming, R.; Livingston, R. *J. Am. Chem. Soc.* **84**: 546-8 (1962).
- 62E007 Population and decay of the lowest triplet state in polyenes with conjugated heteroatoms: Retinene. Dawson, W.; Abrahamson, E.W. *J. Phys. Chem.* **66**: 2542-7 (1962).
- 62E008 Observed electronic transitions of hexahelicene. Rhodes, W.; El-Sayed, M.F.A. *J. Mol. Spectrosc.* **9**: 42-9 (1962).
- 62E009 Radiationless conversion from the triplet state. Hoffman, M.Z.; Porter, G. *Proc. R. Soc. London, Ser. A* **268**: 46-56 (1962).
- 62E010 Lifetime of the triplet state of anthracene in lucite. Melhuish, W.H.; Hardwick, R. *Trans. Faraday Soc.* **58**: 1908-11 (1962).
- 63A001 Transient measurements of photochemical processes in dyes. II. The mechanism of the photosensitized oxidation of aqueous phenol by eosin. Zwicker, E.F.; Grossweiner, L.I. *J. Phys. Chem.* **67**: 549-55 (1963).
- 63B001 Polarization of the triplet-triplet absorption spectrum of some polyacenes by the method of photoselection. El-Sayed, M.A.; Pavlopoulos, T. *J. Chem. Phys.* **39**: 834-8 (1963).
- 63B002 Absorption et luminescence du triphénylène en solution étendue dans l'hexane à 77°K: Mise en évidence d'une fluorescence par échelon (transition $T^* \rightarrow T$). Dupuy, F.; Nouchi, G.; Rousset, A. *C. R. Seances Acad. Sci.* **256**: 2976-9 (1963).
- 63B003 Intermolecular electron transfer in tetrapyrrole pigments under pulsed illumination. Shakhverdov, P.A.; Terenin, A.N. *Dokl. Phys. Chem.* **150**: 557-60 (1963) Translated from: *Dokl. Akad. Nauk SSSR* **150**: 1311-4 (1963).
- 63B004 Dispositif expérimental pour la mesure des absorptions $T^* \leftarrow T$. Lochet, R.; Nouchi, G. *J. Phys. (Paris)* **24**: 765-7 (1963).
- 63E007 Decay kinetics of the 1-naphthaldehyde and benzophenone triplet states in benzene. Bell, J.A.; Linschitz, H. *J. Am. Chem. Soc.* **85**: 528-32 (1963).
- 63F019 The triplet state of fluorescein in sulfuric acid. Lindqvist, L. *J. Phys. Chem.* **67**: 1701-4 (1963).
- 63F021 One-electron photo-ionization of acridine dyes in rigid media: Evidence for the formation of transient radical ion. Lim, E.C.; Swenson, G.W. *J. Chem. Phys.* **39**: 2768-9 (1963).
- 64B003 Flash photolysis of methylene blue. I. Reversible photolysis in the absence of a reducing agent and in the presence of certain kinds of reducing agents. Matsumoto, S. *Bull. Chem. Soc. Jpn.* **37**: 491-8 (1964).
- 64B005 Polarized emission and triplet-triplet absorption spectra of aromatic hydrocarbons in benzophenone crystals. Hochstrasser, R.M.; Lower, S.K. *J. Chem. Phys.* **40**: 1041-6 (1964).
- 64B006 Formation of triplet states of solutes in the radiolysis of organic liquids. Dainton, F.S.; Kemp, T.J.; Salmon, G.A.; Keene, J.P. *Nature* **203**: 1050-2 (1964).
- 64E013 Studies of the transient intermediates in the photoreduction of methylene blue. Kato, S.; Morita, M.; Koizumi, M. *Bull. Chem. Soc. Jpn.* **37**: 117-24 (1964).
- 64E014 Triplet-triplet polarisation measurements in mixed crystals. Craig, D.P.; Fischer, G. *Proc. Chem. Soc.* **176** (1964).
- 64E015 Eine experimentelle Methode zur Ermittlung der Singulett-Triplett-Konversionswahrscheinlichkeit und der Triplett-Spektren von gelösten organischen Molekülen Messungen an 1,2-Benzanthracen. Labhart, H. *Helv. Chim. Acta* **47**: 2279-88 (1964).
- 64E016 Flash photolysis in thin films of gelatin and other polymers. Buettner, A.V. *J. Phys. Chem.* **68**: 3253-9 (1964).
- 64E017 Blitzlichtuntersuchungen ueber die Ausbleichreaktion von Thionin mit Allylthioharnstoff. Fischer, H. *Z. Phys. Chem. (Frankfurt)* **43**: 177-90 (1964).
- 650006 Pulse radiolysis of dioxane solutions. Baxendale, J.H.; Fielden, E.M.; Keene, J.P. *Science* **148**: 637-8 (1965).
- 65A001 Deactivation of the fluorescent and phosphorescent states of tetrapyrrole pigments in liquid solutions. Shakhverdov, P.A.; Terenin, A.N. *Dokl. Phys. Chem.* **160**: 163-5 (1965) Translated from: *Dokl. Akad. Nauk SSSR* **160**: 1141-3 (1965).
- 65A002 Transient species in the photochemistry of eosin. Kasche, V.; Lindqvist, L. *Photochem. Photobiol.* **4**: 923-33 (1965).

- 65B003 Quasilinear structure of luminescence spectra of triphenylene in frozen crystalline solutions. Ruziewicz, Z. *Acta Phys. Pol.* **28**: 389-406 (1965).
- 65B004 Contribution a l'etude de l'etat triplet des colorants: Spectres d'absorption dans le domaine des grandes longueurs d'onde, rouge et proche infrarouge. Lochet, R.; Nouchi, G. *C.R. Seances Acad. Sci.* **260**: 1897-1900 (1965).
- 65E031 Extinction coefficients of triplet-triplet transitions in aromatic compounds. Keller, R.A.; Hadley, S.G. *J. Chem. Phys.* **42**: 2382-7 (1965).
- 65E032 Untersuchung reversibler Photoprozesse durch Blitzlichtphotolyse. II. Triplett-Triplett-Annihilation von Anthracen. Wild, U.; Guenthard, Hs.H. *Helv. Chim. Acta* **48**: 1843-57 (1965).
- 65F031 Mechanism of fluorescence quenching in solution. Part 1. Quenching by bromobenzene. Medinger, T.; Wilkinson, F. *Trans. Faraday Soc.* **61**: 620-30 (1965).
- 65R023 Triplet-triplet absorption spectra of some tetrapyrrole pigments in liquid solution. Shakhverdov, P.A. *Opt. Spectrosc. (USSR)* **19**: 519-20 (1965) Translated from: *Opt. Spektrosk.*
- 66A002 Intermediates in the photoreduction of eosine as revealed by a flash-photolysis study. Ohno, T.; Kato, S.; Koizumi, M. *Bull. Chem. Soc. Jpn.* **39**: 232-9 (1966).
- 66A003 Studies of the photoreduction of acridine in ethanol by the flash technique. Kira, A.; Kato, S.; Koizumi, M. *Bull. Chem. Soc. Jpn.* **39**: 1221-7 (1966).
- 66B001 Triplet-triplet absorption spectra of solid solutions of certain organic compounds. Zhmyreva, I.A.; Kolobkov, V.P.; Volkov, S.V. *Opt. Spectrosc. (USSR)* **20**: 162-4 (1966) Translated from: *Opt. Spektrosk.*
- 66B003 The photo-ionization of molecules in solutions. III. Photoionization and recombination processes of N,N,N',N'-tetramethyl-p-phenylenediamine in various organic solvent. Yamamoto, N.; Nakato, Y.; Tsubomura, H. *Bull. Chem. Soc. Jpn.* **39**: 2603-8 (1966).
- 66E085 Second triplet state of anthracene. Kellogg, R.E. *J. Chem. Phys.* **44**: 411-2 (1966).
- 66E086 Radiationless deactivation of the fluorescent state of substituted anthracenes. Bennett, R.G.; McCartin, P.J. *J. Chem. Phys.* **44**: 1969-72 (1966).
- 66E087 A fast reacting intermediate in flash excited flavin solution. Tegner, L.; Holmstrom, D. *Photochem. Photobiol.* **5**: 223-6 (1966).
- 66E089 Direct evidence for the sensitized formation of a metastable state of β -carotene. Chessin, M.; Livingston, R.; Truscott, T.G. *Trans. Faraday Soc.* **62**: 1519-24 (1966).
- 66E090 Stabilization of reactive species in polymethyl methacrylate. Melhuish, W.H. *Trans. Faraday Soc.* **62**: 3384-92 (1966).
- 672086 Pulse radiolysis of polystyrene. Ho, S.K.; Siegel, S.; Schwarz, H.A. *J. Phys. Chem.* **71**: 4527-33 (1967).
- 676011 Pulse-radiolysis studies of the formation of triplet excited states in cyclohexane solutions of naphthalene and anthracene. Hunt, J.W.; Thomas, J.K. *J. Chem. Phys.* **46**: 2954-8 (1967).
- 677029 Photoreduction of proflavine in the aqueous solution. I. A flash photolysis study. Kikuchi, K.; Koizumi, M. *Bull. Chem. Soc. Jpn.* **40**: 736-43 (1967).
- 677223 Untersuchung reversibler Photoprozesse durch Blitzlichtphotolyse. IV. Triplettzustande des 1,3,6,8'-Tetramethyl-dehydrodianthrone. Huber, J.R.; Wild, U.; Guenthard, Hs.H. *Helv. Chim. Acta* **50**: 841-8 (1967).
- 677259 Photochemical reaction between acridine and acridan in the presence and in the absence of oxygen. II. Studies by the flash technique. Kira, A.; Koizumi, M. *Bull. Chem. Soc. Jpn.* **40**: 2486-92 (1967).
- 677322 Transient measurements of photochemical processes in dyes. I. Primary processes in the flash-photolysis of safranin T. Chibisov, A.K.; Skvortsov, B.V.; Karyakin, A.V.; Shvindt, N.N. *Khim. Vys. Energ.* **1**: 529-35 (1967).
- 677485 Flashphotolysis of stilbenes at 77°K; in rigid EPA. Heinrich, G.; Blume, H.; Schulte-Frohlinde, D. *Tetrahedron Lett.* **47**: 4693-94 (1967).
- 677498 Flash-photolytic detection of triplet acridine formed by energy transfer from biacetyl. Kellmann, A.; Lindqvist, L. *Proc. of Internat. Symposium on Triplet State, Beirut, Lebanon, 1967*, p.439-45.
- 67A001 The one-electron oxidation of triplet diphenyl-p-phenylenediamine by the diimine. Linschitz, H.; Ottolenghi, M.; Bensasson, R. *J. Am. Chem. Soc.* **89**: 4592-9 (1967).
- 67B003 Spectrographie a resolution dans le temps appliquee a l'etude d'especes a tres courte duree de vie dans les spiropyranes photochromiques. Mosse, M.; Metras, J.-C. *J. Chim. Phys. Phys.-Chim. Biol.* **64**: 691-4 (1967).
- 67B004 Existence de bandes d'absorption transitoire entre 5000 et 8800 Å dans les molecules aromatiques et heterocycliques. Astier, R.; Meyer, Y. *J. Chim. Phys. Phys.-Chim. Biol.* **64**: 919-21 (1967).
- 67B005 Absorption triplet-triplet de quelques hydrocarbures aromatiques: Mise en evidence de bandes de faible intensite dans le rouge. Nouchi, G. *C.R. Seances Acad. Sci., Ser. B* **264**: 152-5 (1967).
- 67B006 Transient absorption spectra of benzophenone studied by the flash excitation. Tsubomura, H.; Yamamoto, N.; Tanaka, S. *Chem. Phys. Lett.* **1**: 309-10 (1967).
- 67B007 Concentration dependence of the intensity of triplet-triplet absorption in some aromatic hydrocarbons. Bolotnikova, T.N.; Naumova, T.M. *Opt. Spectrosc. (USSR)* **23**: 206-8 (1967) Translated from: *Opt. Spektrosk.*
- 67B008 Triplet-triplet spectra of aromatic molecules in mixed crystals. Craig, D.P.; Fischer, G. *Trans. Faraday Soc.* **63**: 530-9 (1967).
- 67B009 Polarized triplet-triplet absorption in charge-transfer complexes. Eiselthal, K.B. *J. Chem. Phys.* **46**: 3268-9 (1967).
- 67B010 Triplet state studies of some polyphenyls in rigid glasses. Ramsay, I.A.; Munro, I.H. *The Triplet State*, Cambridge Univ. Press, Cambridge, England, 1967, p.415-23.
- 67C001 Etude en spectroscopie par eclat des colorants thiaziniques en solution aqueuse. Faure, J.; Bonneau, R.; Jousset-Dubien, J. *Photochem. Photobiol.* **6**: 331-9 (1967).
- 67E027 Quantum yields of triplet formation in solutions of chlorophyll. Bowers, P.G.; Porter, G. *Proc. R. Soc. London, Ser. A* **296**: 435-44 (1967).
- 67E031 Triplet state quantum yields for some aromatic hydrocarbons and xanthene dyes in dilute solution. Bowers, P.G.; Porter, G. *Proc. R. Soc. London, Ser. A* **299**: 348-53 (1967).
- 67E105 Triplet-triplet absorption spectra of benzophenone and its derivatives. Godfrey, T.S.; Hilpern, J.W.; Porter, G. *Chem. Phys. Lett.* **1**: 490-2 (1967).
- 67E106 Triplet-triplet absorption studies on aromatic and heterocyclic molecules at 77° K. Henry, B.R.; Kasha, M. *J. Chem. Phys.* **47**: 3319-27 (1967).
- 67E107 The laser photolysis of methylene blue. Danziger, R.M.; Bar-Eli, K.H.; Weiss, K. *J. Phys. Chem.* **71**: 2633-40 (1967).
- 67E108 Application of electron spin resonance in the study of triplet states. II. Effect of triplet-triplet reabsorption on quantitative phosphorescence measurements. Brinen, J.S.; Hodgson, W.G. *J. Chem. Phys.* **47**: 2946-50 (1967).
- 67E109 Solvent dependence of the quantum yield of triplet state production of 9-phenylanthracene. Horrocks, A.R.; Medinger, T.; Wilkinson, F. *Photochem. Photobiol.* **6**: 21-8 (1967).
- 67F505 Radiationless transitions in cyanine dyes. Buettner, A.V. *J. Chem. Phys.* **46**: 1398-401 (1967).
- 680379 Transient species in the radiolysis of solutions of stilbene. Dainton, F.S.; Peng, C.T.; Salmon, G.A. *J. Phys. Chem.* **72**: 3801-7 (1968).
- 680386 The pulse radiolysis of benzene-biacetyl solutions. Cundall, R.B.; Evans, G.B.; Griffiths, P.A.; Keene, J.P. *J. Phys. Chem.* **72**: 3871-7 (1968).
- 680553 Yield of triplet states in the pulse radiolysis of an aliphatic hydrocarbon glass containing naphthalene. Land, E.J.; Swallow, A.J. *J. Chem. Phys.* **49**: 5552-3 (1968).
- 680727 Extinction coefficients of triplet-triplet transitions. Land, E.J. *Proc. Roy. Soc. (London) Ser. A* **305**: 457-71 (1968).

- 686045 Fluorescence quenching and the charge transfer in the excited state. III. Flash photolytic studies. Yamashita, H.; Kokubun, H.; Koizumi, M. *Bull. Chem. Soc. Jpn.* **41**: 2312-19 (1968).
- 686058 Laser photolysis and spectroscopy: a new technique for the study of rapid reactions in the nanosecond time range. Novak, J.R.; Windsor, M.W. *Proc. Roy. Soc. (London) Ser. A* **308**: 95-110 (1968).
- 687111 The lowest triplet state of stilbene. Herkstroeter, W.G.; McClure, D.S. *J. Am. Chem. Soc.* **90**: 4522-7 (1968).
- 687142 Flash photolysis of thiopyronine. Morita, M.; Kato, S. *Z. Naturforsch. Teil B* **23**: 931-3 (1968).
- 687213 The photoreduction of phenazine in acidic methanol. Bailey, D.N.; Roe, D.K.; Hercules, D.M. *J. Am. Chem. Soc.* **90**: 6291-7 (1968).
- 68B002 Triplet-triplet absorption spectrum of N,N-dimethylaniline (DMA) measured by the flash photolysis technique. Kimura, K.; Arimitsu, S.; Yamamoto, N.; Tsubomura, H. *Bull. Chem. Soc. Jpn.* **41**: 1274 (1968).
- 68B003 Une methode cinetique pour mesurer les coefficients d'extinction moleculaire des transitions triplet-triplet. Lavalette, D. *C.R. Seances Acad. Sci., Ser. B* **266**: 279-82 (1968).
- 68B004 The polarization of the allowed triplet-triplet absorption in anthracene. Hochstrasser, R.M.; Marchetti, A.P. *Chem. Phys. Lett.* **1**: 597-8 (1968).
- 68B005 Absorption des β -dicetones sous forte excitation optique. Astier, R.; Meyer, Y.H. *J. Chim. Phys. Phys.-Chim. Biol.* **65**: 1407-9 (1968).
- 68B006 Nanosecond flash photolysis and the absorption spectra of excited singlet states. Porter, G.; Topp, M.R. *Nature* **220**: 1228-9 (1968).
- 68B007 Determination of "I-I" absorption spectra by means of two-quantum photosensitized reactions. Zhuravlyeva, T.S. *Opt. Spectrosc. (USSR)* **25**: 389-91 (1968) Translated from: *Opt. Spektrosk.*
- 68B008 Acidity constants of anthracene derivatives in singlet and triplet excited states. Vander Donckt, E.; Porter, G. *Trans. Faraday Soc.* **64**: 3218-23 (1968).
- 68D211 Application of electron spin resonance in the study of triplet states. III. Extinction coefficients of triplet-triplet transitions. Brinen, J.S. *J. Chem. Phys.* **49**: 586-90 (1968).
- 68E098 Triplet state formation efficiencies of aromatic hydrocarbons in solution. Horrocks, A.R.; Wilkinson, F. *Proc. R. Soc. London, Ser. A* **306**: 257-73 (1968).
- 68E100 A flash-photolysis investigation of flavin photosensitization of purine nucleotides. Knowles, A.; Roe, E.M.F. *Photochem. Photobiol.* **7**: 421-36 (1968).
- 68E102 Dichroisme de l'anthracene triplet. Meyer, Y.H.; Astier, R. *J. Phys. (Paris)* **29**: 1075-80 (1968).
- 68E103 Triplet-triplet fluorescence of rubrene in solution. Yildiz, A.; Kissinger, P.T.; Reilley, C.N. *J. Chem. Phys.* **49**: 1403-6 (1968).
- 68E104 Measurement of extinction coefficients of triplet absorption and quantum yields of triplet formation. Dawson, W.R. *J. Opt. Soc. Am.* **58**: 222-7 (1968).
- 68E105 Quantum efficiencies of triplet formation in aromatic molecules. Windsor, M.W.; Dawson, W.R. *Mol. Cryst.* **4**: 253-8 (1968).
- 68E106 Triplet-triplet absorption of naphthalene and phenanthrene impurities in benzophenone crystals. Gaevskii, A.S.; Nelipovich, K.I.; Faizyash, A.N. *Opt. Spectrosc. (USSR)* **24**: 554-5 (1968) Translated from: *Opt. Spektrosk.*
- 68E107 Singlet-singlet absorption and intersystem crossing from the 1B_u state of naphthalene. Bonneau, R.; Faure, J.; Jousset-Dubien, J. *Chem. Phys. Lett.* **2**: 65-7 (1968).
- 690087 Yields of excited states in the pulse radiolysis of solutions of substituted anthracenes. Kemp, T.J.; Roberts, J.P. *Trans. Faraday Soc.* **65**: 725-31 (1969).
- 690520 Absorption spectrum of the triplet state of duroquinone. Land, E.J. *Trans. Faraday Soc.* **65**: 2815-22 (1969).
- 692001 Nanosecond pulse radiolysis of polystyrene and poly(methylmethacrylate) Ho, S.K.; Siegel, S. *J. Chem. Phys.* **50**: 1142-52 (1969).
- 696019 Radiative and radiationless processes in aromatic molecules. Pyrene. Kropp, J.L.; Dawson, W.R.; Windsor, M.W. *J. Phys. Chem.* **73**: 1747-52 (1969).
- 696020 Radiationless deactivation and anomalous fluorescence of singlet 1,12-benzperylene. Dawson, W.R.; Kropp, J.L. *J. Phys. Chem.* **73**: 1752-58 (1969).
- 696073 Triplet excimers favoured by charge transfer in electron-donor-acceptor-complexes. Briegleb, G.; Schuster, H.; Herre, W. *Chem. Phys. Lett.* **4**: 53-8 (1969).
- 696115 Quantitative ESR studies of organic triplet states. Extinction coefficients of $T \rightarrow T'$ transitions. Brinen, J.S. *Molecular Luminescence*, Lim, E.C. (ed.), W.A. Benjamin, Inc., New York, 1969, p.333-49.
- 697009 Studies of the transient intermediates of a thiopyronine aqueous solution under flash excitation. Morita, M.; Kato, S. *Bull. Chem. Soc. Jpn.* **42**: 25-35 (1969).
- 697061 Investigation of transient states in the photochemical reactions of dyes. III. Flash-photolysis of phenosafranine and neutral red in aqueous solutions. Chibisov, A.K.; Skvortsov, B.V.; Karyakin, A.V.; Rygalov, L.N. *High Energy Chem.* **3**: 190-4 (1969) Translated from: *Khim. Vys. Energ.* **3**: 210-6 (1969).
- 697092 Flash photolysis of camphorquinone and biacetyl. Singh, A.; Scott, A.R.; Sopchysyn, F. *J. Phys. Chem.* **73**: 2633-43 (1969).
- 697225 Blitzlichtphotolyse an Acridin und Acridan im Temperaturbereich von +25 °C bis -180 °C. Zanker, V.; Prell, G. *Ber. Bunsenges. Phys. Chem.* **73**: 791-5 (1969).
- 697226 Die Untersuchung angeregter Zustände aromatischer Kohlenwasserstoffe mit Hilfe der Laser-Blitzspektroskopie. Mueller, A.; Sommer, U. *Ber. Bunsenges. Phys. Chem.* **73**: 819-26 (1969).
- 697253 Optical detection of the triplet state of uracil. Whillans, D.W.; Herbert, M.A.; Hunt, J.W.; Johns, H.E. *Biochem. Biophys. Res. Commun.* **36**: 912-8 (1969).
- 697272 The triplet state of chloranil. Kemp, D.R.; Porter, G. *Chem. Commun.* : 1029-30 (1969).
- 69B001 Polarized excitation spectrum of the triplet-triplet absorption of aromatic hydrocarbons. Lavalette, D. *Chem. Phys. Lett.* **3**: 67-70 (1969).
- 69B002 Experimental and theoretical study of triplet-triplet absorption in the phenylanthracenes. Brinen, J.S.; Orloff, M.K. *J. Chem. Phys.* **51**: 527-31 (1969).
- 69B003 Spectres d'absorption triplet-triplet de colorants derives du triphenylmethane. Nouchi, G.; Silvie, C. *C.R. Seances Acad. Sci., Ser. B* **268**: 546-8 (1969).
- 69B004 Low-energy triplet-triplet transitions in naphthalene and some monosubstituted derivatives. Melhuish, W.H. *J. Chem. Phys.* **50**: 2779 (1969).
- 69B005 Triplet absorption spectra of cyanine dyes. Pierce, R.A.; Berg, R.A. *J. Chem. Phys.* **51**: 1267 (1969).
- 69B006 Low-energy triplet state of tetracene. Astier, R.; Bokobza, A.; Meyer, Y.H. *J. Chem. Phys.* **51**: 5174-5 (1969).
- 69B007 Etude et interpretation des spectres d'absorption triplet-triplet. Classification des états triplets superieurs. I. Les matieres colorantes. Nouchi, G. *J. Chim. Phys. Phys.-Chim. Biol.* **66**: 548-53 (1969).
- 69E207 First allowed triplet-triplet transition in benzene. Astier, R.; Meyer, Y.H. *Chem. Phys. Lett.* **3**: 399-401 (1969).
- 69E208 Triplet-triplet spectra and triplet quantum yields of some aromatic hydrocarbons in liquid solution. Heinzelmann, W.; Labhart, H. *Chem. Phys. Lett.* **4**: 20-4 (1969).
- 69E209 The triplet state of lycopene. Sykes, A.; Truscott, T.G. *Chem. Commun.* : 274-6 (1969).
- 69E210 Polarization of electronic transitions of aromatic hydrocarbons. Gallivan, J.B.; Brinen, J.S. *J. Chem. Phys.* **50**: 1590-5 (1969).

- 69E211 Etude et interpretation des spectres d'absorption triplet-triplet. Classification des etats triplets superieurs. II. Les hydrocarbures aromatiques. Nouchi, G. J. Chim. Phys. Phys.-Chim. Biol. **66**: 554-65 (1969).
- 69E212 L'absorption triplet-triplet des hydrocarbures aromatiques alternants. II. Coefficients d'extinction moleculaire et forces d'oscillateur. Lavalette, D. J. Chim. Phys. Phys.-Chim. Biol. **66**: 1853-9 (1969).
- 69E213 L'absorption triplet-triplet des hydrocarbures aromatiques alternants. III. Polarisation des transitions. Lavalette, D. J. Chim. Phys. Phys.-Chim. Biol. **66**: 1860-4 (1969).
- 69E214 Absorptions-, Emissions- und Polarisationsgradmessungen an 9-C- und 10-N-substituierten Acridanen. Zanker, V.; Schneider, B. Z. Phys. Chem. (Frankfurt) **68**: 19-29 (1969).
- 69E215 Polarized triplet-triplet absorption spectra in three benzene derivatives. Cadogan, K.D.; Albrecht, A.C. J. Phys. Chem. **73**: 1868-77 (1969).
- 69E216 Radiative and radiationless processes in aromatic molecules. Coronene and benzcoronene. Dawson, W.R.; Kropp, J.L. J. Phys. Chem. **73**: 693-9 (1969).
- 69E217 A metastable state of retinol (vitamin A). Sykes, A.; Truscott, T.G. Chem. Commun. 929-30 (1969).
- 69E218 The photoreduction of fluorenone. Davis, G.A.; Carapellucci, P.A.; Szoc, K.; Gresser, J.D. J. Am. Chem. Soc. **91**: 2264-72 (1969).
- 69F389 Direct determination of the extinction coefficients for triplet \leftarrow triplet transitions in naphthalene, phenanthrene, and triphenylene. Hadley, S.G.; Keller, R.A. J. Phys. Chem. **73**: 4351-5 (1969).
- 69N001 Etude par spectroscopie d'eclairs du transfert d'energie chlorophylle-carotenoide. Mathis, P. Photochem. Photobiol. **9**: 55-63 (1969).
- 700011 Reactions of electrons photoejected from aromatic amino acids in frozen aqueous solutions of divalent metal salts. Santus, R.; Helene, A.; Helene, C.; Ptak, M. J. Phys. Chem. **74**: 550-61 (1970).
- 700114 Excitation transfer in the pulse radiolysis of naphthalene and benzophenone solutions. Holroyd, R.A.; Theard, L.M.; Peterson, F.C. J. Phys. Chem. **74**: 1895-900 (1970).
- 700277 Radiolysis transients in viscous liquids. Biphenyl in liquid paraffin. Fuller, J.; Peteleski, N.; Ruppel, D.; Tomlinson, M. J. Phys. Chem. **74**: 3066-73 (1970).
- 701073 Investigation of aromatic triplet molecules in polystyrene by the method of pulsed radiolysis. Shekk, Yu.B.; Alfimov, M.V.; Nikoiforov, V.I. Bull. Acad. Sci. USSR, Div. Chem. Sci. (Engl. Transl.) : 2309-14 (1970) Translated from: Izv. Akad. Nauk SSSR, Ser. Khim. : 2454-60 (1970).
- 703001 The metastable state of β -carotene excited by pulse radiolysis. Land, E.J.; Sykes, A.; Truscott, T.G. Chem. Commun. : 332 (1970).
- 706135 The fluorescence and phosphorescence of 1,2,5,6-dibenzacridine and 1,2,7,8-dibenzacridine in glassy and liquid solution. Kropp, J.L.; Lou, J.J. J. Phys. Chem. **74**: 3953-9 (1970).
- 706157 Polarized spectroscopy of excited states of substituted anthracenes on a nanosecond time scale. Kliger, D.S.; Albrecht, A.C. J. Chem. Phys. **53**: 4059-65 (1970).
- 707036 Triplet-triplet absorption via intramolecular energy transfer. Brinen, J.S.; Gerhardt, G.E.; Kazan, J. Chem. Phys. Lett. **5**: 150-2 (1970).
- 707061 Effect of solvent on the flash photolysis and laser photolysis of N,N,N',N'-tetramethyl-paraphenylene diamine in solution. Richards, J.T.; Thomas, J.K. Trans. Faraday Soc. **66**: 621-32 (1970).
- 707152 Mechanism of photodissociation of hydroquinone derivatives. Yamada, H.; Nakashima, N.; Tsubomura, H. J. Phys. Chem. **74**: 2897-903 (1970).
- 707186 Photochemistry of phenoxazine. A flash-photolytic study. Gegiou, D.; Huber, J.R.; Weiss, K. J. Am. Chem. Soc. **92**: 5058-62 (1970).
- 707199 Triplett-Lebensdauern von Diphenylpolyenen und deuterierten Stilbenen in eingefrorener Loesung bei 77 °K. Heinrich, G.; Holzer, G.; Blume, H.; Schulte-Frohlinde, D. Z. Naturforsch. Teil B **25**: 496-9 (1970).
- 707230 Polarization measurements on low-lying T-T absorptions with CW argon-ion laser photoselection. Langelaar, J.; Wegdam-Van Beek, J.; Van Voorst, J.D.W.; Lavalette, D. Chem. Phys. Lett. **6**: 460-4 (1970).
- 707240 Direct determination of triplet \leftarrow triplet absorption extinction coefficients. II. Quinoline, isoquinoline, quinoxaline. Hadley, S.G. J. Phys. Chem. **74**: 3551-2 (1970).
- 707305 Photochemical reactions of duroquinone. Part 2. Flash photolytic investigations. Nafisi-Movaghgar, J.; Wilkinson, F. Trans. Faraday Soc. **66**: 2268-78 (1970).
- 707337 Photoreduction of phenazine in alcohols and by EDTA in the aqueous solution. Iwaoka, T.; Niizuma, S.; Koizumi, M. Bull. Chem. Soc. Jpn. **43**: 2786-95 (1970).
- 707346 Formation of ions and excited states in the laser photolysis of solutions of pyrene. Richards, J.T.; West, G.; Thomas, J.K. J. Phys. Chem. **74**: 4137-41 (1970).
- 707364 A study of the intermediate compositions in the photochemical reactions of dyes. IV. Oxidation-reduction transformations of methylene blue and methylene green in pulsed photoexcitation. Chibisov, A.K.; Roitman, G.P.; Karyakin, A.V. High Energy Chem. **4**: 235-6 (1970) Translated from: Khim. Vys. Energ. **4**: 273-4 (1970).
- 707561 Transient species in the laser photolysis at 265 m μ of the halobenzenes in solution. Loeff, I.; Lutz, H.; Lindqvist, L. Isr. J. Chem. **8**: 141-6 (1970).
- 70B002 Die Uebergangsmomentrichtung von Triplett-Triplett-Absorptionsbanden. Eine neue Methode zur Bestimmung der Anisotropie der Absorption bei der Photoselektion. Kuball, H.-G.; Euing, W.; Karstens, T. Ber. Bunsenges. Phys. Chem. **74**: 316-23 (1970).
- 70B003 Polarization of the triplet-triplet absorption spectrum of 1,2-benzopyrene. Pavlopoulos, T.G. Ber. Bunsenges. Phys. Chem. **74**: 989-92 (1970).
- 70B004 Polarization of the triplet-triplet absorption spectrum of pyrene. Pavlopoulos, T.G. J. Chem. Phys. **52**: 3307-8 (1970).
- 70B005 Search for the first $^3A_g \leftarrow ^3B_{2u}$ transition in naphthalene. Pavlopoulos, T.G. J. Chem. Phys. **53**: 4230-3 (1970).
- 70B006 Dianthrone photochromism 1950-1970. Bercovici, T.; Korenstein, R.; Muszkat, K.A.; Fischer, E. Pure Appl. Chem. **24**: 531-65 (1970).
- 70E288 Nanosecond flash photolysis. Porter, G.; Topp, M.R. Proc. R. Soc. London, Ser. A **315**: 163-84 (1970).
- 70E290 Triplet-triplet absorption of biphenyl. Meyer, Y.H.; Astier, R.; Leclercq, J.M. Chem. Phys. Lett. **4**: 587-90 (1970).
- 70E291 Absorption and emission spectra and triplet decay of some aromatic and N-heterocyclic compounds in polymethylmethacrylate. West, M.A.; McCallum, K.J.; Woods, R.J.; Formosinho, S.J. Trans. Faraday Soc. **66**: 2135-47 (1970).
- 70E293 Spectral characteristics of the triplet states of cyanin and thiazine dyes. Chibisov, A.K.; Kuz'min, V.A.; Roitman, G.P.; Levkoev, I.I.; Karyakin, A.V. Bull. Acad. Sci. USSR, Phys. Ser. **34**: 1144-7 (1970) Translated from: Izv. Akad. Nauk SSR, Ser. Fiz.
- 70E294 The location, polarization and assignment of low lying excited triplet states in pyrene. Langelaar, J.; Wegdam-van Beek, J.; ten Brink, H.T.; van Voorst, J.D.W. Chem. Phys. Lett. **7**: 368-70 (1970).
- 70E295 A simple apparatus for measuring the absorption spectrum and lifetimes of photo-produced short lived species using commercial subunits. Slifkin, M.A.; Walmsley, R.H. J. Phys. E **3**: 160-2 (1970).
- 70E296 Pheophytin-a and pheophytin-b: Quantitative absorption spectra of the singlet- and triplet-state. Zanker, V.; Rudolph, E.; Prell, G. Z. Naturforsch., Teil B **25B**: 1137-43 (1970).
- 710186 Pulse radiolysis studies of acetone solutions of biphenyl and anthracene: Formation of ions and excited singlet state. Arai, S.; Kira, A.; Imamura, M. J. Chem. Phys. **54**: 5073-81 (1971).

- 710278 Aromatic solute molecular ion and triplet state formation in polystyrene and polymethylmethacrylate. Siegel, S.; Stewart, T. J. *Chem. Phys.* **55**: 1775-86 (1971).
- 710617 Yields of excited singlet and triplet states of some aromatics in the pulse radiolysis of cyclohexane solutions. Baxendale, J.H.; Wardman, P. *Trans. Faraday Soc.* **67**: 2997-3007 (1971).
- 713035 The in vitro photochemistry of biological molecules. II. The triplet states of β -carotene and lycopene excited by pulse radiolysis. Land, E.J.; Sykes, A.; Truscott, T.G. *Photochem. Photobiol.* **13**: 311-20 (1971).
- 716169 Flash spectroscopy and photoreduction of phenazine. Japar, S.M.; Abrahamson, E.W. *J. Am. Chem. Soc.* **93**: 4140-4 (1971).
- 716235 Studies on triplet energy transfer by means of an emission-absorption flash technique. II. Mixed triplet-triplet annihilation in ethanol. Kikuchi, K.; Kokubun, H.; Koizumi, M. *Bull. Chem. Soc. Jpn.* **44**: 1527-34 (1971).
- 716243 Triplet states in the pyrene-tetracyanobenzene charge-transfer system. Teitelbaum, Z.; Potashnik, R.; Ottolenghi, M. *Mol. Photochem.* **3**: 107-22 (1971).
- 716244 Temperature dependence of the deactivation of electronically excited indazoles in solution. Bircher, P.; Pantke, E.R.; Labhart, H. *Chem. Phys. Lett.* **11**: 347-9 (1971).
- 716330 Ionization processes in polar solution of tryptophan at 77 ° K. Moan, J. *Isr. J. Chem.* **9**: 637-43 (1971).
- 717009 Photoreduction of benzophenone in benzene. II. Flash photolysis study of primary photochemical reactions. Buettner, A.V.; Dedinas, J. J. *Phys. Chem.* **75**: 187-91 (1971).
- 717071 Properties of the triplet states of thymine and uracil in aqueous solution. Whillans, D.W.; Johns, H.E. *J. Am. Chem. Soc.* **93**: 1358-2 (1971).
- 717126 Intersystem crossing in the charge-transfer quenching of molecular fluorescence. Goldschmidt, C.R.; Potashnik, R.; Ottolenghi, M. *J. Phys. Chem.* **75**: 1025-31 (1971).
- 717154 Photoreduction of nitrogen heterocycles. I. The photoreduction of phenazine: Evidence for singlet-state reactivity. Davis, G.A.; Gresser, J.D.; Carapellucci, P.A. *J. Am. Chem. Soc.* **93**: 2179-82 (1971).
- 717171 The primary process of the photochemical dimerization of carbostyryl. Yamamuro, T.; Tanaka, I.; Hata, N. *Bull. Chem. Soc. Jpn.* **44**: 667-71 (1971).
- 717179 Nanosecond laser photolysis of acetophenone in organic solvents. Lutz, H.; Lindqvist, L. *Chem. Commun.* : 493-4 (1971).
- 717266 Triplet and radical formation as observed on flashing aromatic hydroxyl derivatives in liquid and solid solution. de Groot, R.L. *Mol. Photochem.* **3**: 1-22 (1971).
- 717460 Weak triplet-triplet absorption bands of naphthalene. Astier, R.; Meyer, Y.H. *Chem. Phys. Lett.* **11**: 523-5 (1971).
- 717463 Direct observation of triplet exciplexes by flash photconductivity and flash spectroscopy. Formation of three-component exciplexes between porphyrins and nitro aromatics. Roy, J.K.; Whitten, D.G. *J. Am. Chem. Soc.* **93**: 7093-4 (1971).
- 717489 Detection and lifetime of the triplet state of acetone in solution. Porter, G.; Yip, R.W.; Dunston, J.M.; Cessna, A.J.; Sugamori, S.E. *Trans. Faraday Soc.* **67**: 3149-54 (1971).
- 717520 Photochemistry of methylated p-benzoquinones. Kemp, D.R.; Porter, G. *Proc. Roy. Soc. (London) Ser. A* **326**: 117-30 (1971).
- 719059 Triplet-triplet absorption and energy transfer from high triplet states. Alfimov, M.V.; Batekha, I.G.; Sheck, Yu.B.; Gerko, V.I. *Spectrochim. Acta* **27**: 329-41 (1971).
- 71B001 Triplet-triplet absorption studies on coumarin and related molecules. Henry, B.R.; Hunt, R.V. *J. Mol. Spectrosc.* **39**: 466-70 (1971).
- 71B002 A flash photolysis study of the triplet state of acridine orange in basic solvents. Kellmann, A. *Photochem. Photobiol.* **14**: 85-93 (1971).
- 71B003 Observation and interpretation of triplet-triplet absorption spectra of substituted naphthalenes. Takemura, T.; Hara, K.; Baba, H. *Bull. Chem. Soc. Jpn.* **44**: 977-84 (1971).
- 71B005 The second triplet level of 1,5-dichloroanthracene in fluid solutions. Roberts, J.P.; Dixon, R.S. *J. Phys. Chem.* **75**: 845-8 (1971).
- 71B006 Use of a silicon photodiode for detection of weak transient absorption in flash photolysis experiments. Melhuish, W.H. *J. Phys. E* **4**: 60-1 (1971).
- 71E360 Triplet-triplet extinction coefficients via energy transfer. Bensasson, R.; Land, E.J. *Trans. Faraday Soc.* **67**: 1904-15 (1971).
- 71E361 Triplet states of polycyclic aromatic hydrocarbons in fluid solution and in the solid state. Slifkin, M.A.; Walmsley, R.H. *Photochem. Photobiol.* **13**: 57-65 (1971).
- 71E367 Flash photolysis studies of orotic acid. Herbert, M.A.; Johns, H.E. *Photochem. Photobiol.* **14**: 693-704 (1971).
- 720206 The radiation-induced formation of excited states of aromatic hydrocarbons in benzene and cyclohexane. II. Yields of excited singlet and triplet state solute molecules. Dainton, F.S.; Morrow, T.; Salmon, G.A.; Thompson, G.F. *Proc. Roy. Soc. (London) Ser. A* **328**: 457-79 (1972).
- 720243 Yields and reactions of lowest excited singlet and triplet states in the radiolysis of naphthalene and 1-methylnaphthalene. Holroyd, R.A.; Capellos, C. *J. Phys. Chem.* **76**: 2485-91 (1972).
- 720392 Pulse radiolysis of 9,10-anthraquinones. Part 2. Triplet excited states. Hulme, B.E.; Land, E.J.; Phillips, G.O. *J. Chem. Soc., Faraday Trans. 1* **68**: 2003-12 (1972).
- 720440 Excited states in the nanosecond pulse radiolysis and laser flash photolysis of N,N-dimethylaniline. Land, E.J.; Richards, J.T.; Thomas, J.K. *J. Phys. Chem.* **76**: 3805-12 (1972).
- 720447 Pulse radiolysis of solutions of stilbene. I. Evidence for triplet and singlet excited state formation. Dainton, F.S.; Robinson, E.A.; Salmon, G.A. *J. Phys. Chem.* **76**: 3897-904 (1972).
- 720464 Sensitized triplet-triplet absorption of biphenylene. Tetreau, C.; Lavalette, D.; Land, E.J.; Peradejordi, F. *Chem. Phys. Lett.* **17**: 245-7 (1972).
- 723043 Nanosecond irradiation studies of biological molecules. I. Coenzyme Q 6 (ubiquinone-30). Bensasson, R.; Chachaty, C.; Land, E.J.; Salet, C. *Photochem. Photobiol.* **16**: 27-37 (1972).
- 726156 Photochemical characteristics of cyanine dyes. Part 1. 3,3'-Diethyloxadicarbocyanine iodide and 3,3'-diethylthiadibocarbocyanine iodide. Dempster, D.N.; Morrow, T.; Rankin, R.; Thompson, G.F. *J. Chem. Soc., Faraday Trans. 2* **68**: 1479-96 (1972).
- 726177 Triplet \leftarrow triplet absorption and intersystem crossing in phthalazine. Alvarez, V.L.; Hadley, S.G. *J. Phys. Chem.* **76**: 3937-40 (1972).
- 727000 Measurement of the triplet-triplet absorption spectrum of tetracene using cw argon laser excitation. Pavlopoulos, T.G. *J. Chem. Phys.* **56**: 227-32 (1972).
- 727036 Photoreactions in aqueous solutions of thymine, pH 12. Whillans, D.W.; Johns, H.E. *J. Phys. Chem.* **76**: 489-93 (1972).
- 727041 Primary products in the flash photolysis of tryptophan. Santos, R.; Grossweiner, L.I. *Photochem. Photobiol.* **15**: 101-5 (1972).
- 727069 Transient species obtained by the flash photolysis of chloranil in various solutions. Kawai, K.; Shiota, Y.; Tsubomura, H.; Mikawa, H. *Bull. Chem. Soc. Jpn.* **45**: 77-81 (1972).
- 727073 Study of triplet quantum yields using a tunable dye laser. Soep, B.; Kellmann, A.; Martin, M.; Lindqvist, L. *Chem. Phys. Lett.* **13**: 241-4 (1972).
- 727098 Primary photochemical processes in aromatic molecules. Part 15. The photochemistry of aromatic carbonyl compounds in aqueous solution. Ledger, M.B.; Porter, G. *J. Chem. Soc., Faraday Trans. 1* **68**: 539-53 (1972).
- 727105 Primary photoprocesses in isoquinoline N-oxides. Lohse, C. *J. Chem. Soc., Perkin Trans. 2* : 229-33 (1972).
- 727348 Absorption spectrum and decay kinetics of triplet pentacene in solution, studied by flash photolysis. Hellner, C.; Lindqvist, L.; Roberge, P.C. *J. Chem. Soc., Faraday Trans. 2* **68**: 1928-37 (1972).
- 727374 Quenching of triplet states by Schiff base nickel(II) complexes. Adamczyk, A.; Wilkinson, F. *J. Chem. Soc., Faraday Trans. 2* **68**: 2031-41 (1972).
- 727440 Predissociation mechanism for the decomposition of electronically excited hydrocarbon molecules. Smirnov, V.A.; Plotnikov, V.G.; Zavyalov, Yu.A.; Alfimov, M.V. *Opt. Spectrosc. (Engl. Transl.)* **33**: 124-7 (1972) Translated from: *Opt. Spektrosk.* **33**: 230-6 (1972).

- 72A022 Photochemical reactions of aromatic ketones with nucleic acids and their components. I. Purine and pyrimidine bases and nucleosides. Charlier, M.; Helene, C. *Photochem. Photobiol.* **15**: 71-87 (1972).
- 72B001 Triplet-triplet spectroscopy of polyacenes. Meyer, Y.H.; Astier, R.; Leclercq, J.M. *J. Chem. Phys.* **56**: 801-15 (1972).
- 72B002 The light-induced conversion of triphenylamine to the excited triplet state of 11,12-dihydrocarbazole. Foerster, E.W.; Grellmann, K.H. *Chem. Phys. Lett.* **14**: 536-8 (1972).
- 72B003 Triplet-triplet absorption spectra of aromatic hydrocarbons. I. Naphthalene. Wild, U.P.; Kaenzig, H.; Ranald, U.B. *Helv. Chim. Acta* **55**: 2724-30 (1972).
- 72B004 Triplet-triplet absorption in 4-N-dimethylaminopyridine. Testa, A.C. *J. Chem. Phys.* **57**: 3019-20 (1972).
- 72E276 The triplet-triplet absorption observed for some tetracyanobenzene complexes. Matsumoto, S.; Nagakura, S.; Iwata, S.; Nakamura, J. *Chem. Phys. Lett.* **13**: 463-5 (1972).
- 72E277 Triplet state of rhodamine 6G molecules. Ketsle, G.A.; Levshin, L.V.; Slavnova, T.D.; Chibisov, A.K. *Sov. Phys. Doklady* **16**: 986-8 (1972) Translated from: *Dokl. Akad. Nauk SSSR* **201**: 60-3 (1971).
- 733001 The in vitro photochemistry of biological molecules. III. Absorption spectra, lifetimes and rates of oxygen quenching of the triplet states of β -carotene, retinal and related polyenes. Truscott, T.G.; Land, E.J.; Sykes, A. *Photochem. Photobiol.* **17**: 43-51 (1973).
- 733002 The triplet state of retinal: Is it involved in vision? Bensasson, R.; Land, E.J.; Truscott, T.G. *Photochem. Photobiol.* **17**: 53-5 (1973).
- 733184 Laser flash photolysis studies of chlorin and porphyrin systems. I. Energetics of the triplet state of bacteriochlorophyll. Connolly, J.S.; Gorman, D.S.; Seely, G.R. *Ann. N.Y. Acad. Sci.* **206**: 649-69 (1973).
- 736051 Photochemical characteristics of the mode-locking dyes 1,1'-diethyl-4,4'-carbocyanine iodide (cryptocyanine, DCI) and 1,1'-diethyl-2,2'-dicarbocyanine iodide (DDI). Dempster, D.N.; Morrow, T.; Rankin, R.; Thompson, G.F. *Chem. Phys. Lett.* **18**: 488-92 (1973).
- 736218 Blitzspektroskopische Untersuchung der Triplettbildung von Naphthalin, 1-Methoxynaphthalin und 1-Cyanonaphthalin in Loesung in Gegenwart der Fluoreszenzloescher Tetrachlorokohlenstoff oder Diaethylamin. Hermann, H.; Koltzenburg, G.; Schulte-Frohlinde, D. *Ber. Bunsenges. Phys. Chem.* **77**: 677-81 (1973).
- 737046 The primary process of the photochemical dimerization of carbostyryl (2-quinolone) in an aqueous solution. Yamamuro, T.; Hata, N.; Tanaka, I. *Bull. Chem. Soc. Jpn.* **46**: 29-34 (1973).
- 737055 Quantitative studies of biphotonic reactions. Bagdasaryan, Kh.S.; Kiryukhin, Yu.I.; Sinitina, Z.A. *J. Photochem.* **1**: 225-40 (1973).
- 737069 Lebensdauer der Triplett-Zustände von partiell deuterierten trans-Stilbenen und strukturverwandten Verbindungen. Heinrich, G.; Guesten, H.; Mark, F.; Olbrich, G.; Schulte-Frohlinde, D. *Ber. Bunsenges. Phys. Chem.* **77**: 103-8 (1973).
- 737098 Mechanism of photoinduced anion formation of pyromellitic dianhydride in n-donor solvents. Achiba, Y.; Kimura, K. *Chem. Phys. Lett.* **19**: 45-8 (1973).
- 737101 Laser induced transient absorption in liquid pyrene. Hodgkinson, K.A.; Munro, I.H. *Chem. Phys. Lett.* **19**: 260-2 (1973).
- 737113 Interaction between excited β -naphthol and pyridine. Hydrogen-bond formation in S* and H-transfer reaction in T. Kikuchi, K.; Watarai, H.; Koizumi, M. *Bull. Chem. Soc. Jpn.* **46**: 749-54 (1973).
- 737125 Reaction patterns and kinetics of the photoconversion of N-methyldiphenylamine to N-methylcarbazole. Foerster, E.W.; Grellmann, K.H.; Linschitz, H. *J. Am. Chem. Soc.* **95**: 3108-15 (1973).
- 737127 Flash photolysis study of 4-N-dimethylaminopyridine. Testa, A.C. *J. Am. Chem. Soc.* **95**: 3128-31 (1973).
- 737140 Quantum yield of triplet state formation of 9-cyanoanthracene in three solvents. Vander Donckt, E.; Barthels, M.R.; Delestinne, A. *J. Photochem.* **1**: 429-32 (1973).
- 737158 Flash photolysis studies of solute triplets: an anomalous heavy atom effect of xenon. Head, D.A.; Singh, A.; Cook, M.G.; Quinn, M.J. *Can. J. Chem.* **51**: 1624-33 (1973).
- 737185 Primary photoprocesses in retinol. Rosenfeld, T.; Alchahal, A.; Ottolenghi, M. *Chem. Phys. Lett.* **20**: 291-7 (1973).
- 737190 Triplet state of ketones in solutions. Quenching rate studies of thioxanthone triplets by flash absorption. Yip, R.W.; Szabo, A.G.; Tolg, P.K. *J. Am. Chem. Soc.* **95**: 4471-2 (1973).
- 737198 Effects of solvent and substituents on the absorption spectra of triplet acetophenone and the acetophenone ketyl radical studied by nanosecond laser photolysis. Lutz, H.; Breheret, E.; Lindqvist, L. *J. Phys. Chem.* **77**: 1758-62 (1973).
- 737287 Primary processes in excited β -naphthol systems: A pulsed laser photolysis study. Klaening, U.K.; Goldschmidt, Ch.R.; Ottolenghi, M.; Stein, G. *J. Chem. Phys.* **59**: 1753-9 (1973).
- 737318 Photolysis of aromatic thioacid O-esters: the nature of the reactive excited state. Wirz, J. *J. Chem. Soc., Perkin Trans. 2* **1307-12** (1973).
- 737439 Photochemistry of flavins. I. Conventional and laser flash photolysis study of alloxazine. Dekker, R.H.; Srinivasan, B.N.; Huber, J.R.; Weiss, K. *Photochem. Photobiol.* **18**: 457-66 (1973).
- 737463 Excited singlet and triplet states of aromatic hydrocarbons determined by a laser photolysis technique. Hodgkinson, K.A.; Munro, I.H. *J. Mol. Spectry.* **48**: 57-71 (1973).
- 737466 Photoreactions of aromatic compounds. XXIX. Pathway and intermediates of the photoreactions of 3,5-dinitroanisole with nucleophiles. de Gunst, G.P.; Havinga, E. *Tetrahedron* **29**: 2167-71 (1973).
- 737469 The primary photochemical processes of 4-nitropyridine N-oxide. III. Its photochemical behavior in an aqueous solution. Hata, N.; Ono, I.; Osaka, K. *Bull. Chem. Soc. Jpn.* **46**: 3363-6 (1973).
- 737473 Photoenolization of ortho-methyl-substituted acetophenones: Solvent effects on the triplet state reactivity. Lutz, H.; Breheret, E.; Lindqvist, L. *J. Chem. Soc., Faraday Trans. 1* **69**: 2096-102 (1973).
- 73B001 Triplet-triplet absorption spectra of quinoline and isoquinoline. Kreibich, U.T.; Wild, U.P. *J. Mol. Spectrosc.* **47**: 189-93 (1973).
- 73E284 Triplet-triplet absorption spectra; of organic molecules. Labhart, H.; Heinzelmann, W. *Organic Molecular Photophysics*, J.B. Birks (ed.), John Wiley and Sons, New York, N.Y., 1973, Vol. 1, p.297-355.
- 73E342 On the apparent absence of triplet-triplet absorption in pure organic molecular crystals. Priestley, E.B.; Robinson, G.W. *Mol. Phys.* **26**: 159-67 (1973).
- 73E344 Prediction of laser action properties of organic dyes from their structure and the polarization characteristics of their electronic transitions. Pavlopoulos, T.G. *IEEE J. Quantum Electron.* **QE-9**: 510-6 (1973).
- 73E345 Triplet-triplet absorption and phosphorescence of metal-porphyrin complexes in liquid solutions. Tsvirko, M.P.; Sapunov, V.V.; Solovov, K.N. *Opt. Spectrosc.* **34**: 635-8 (1973) Translated from: *Opt. Spektrosk.* **34**: 1094-100 (1973).
- 73E346 Measurement of molar triplet extinction coefficients of organic molecules by means of cw laser excitation. Pavlopoulos, T.G. *J. Opt. Soc. Am.* **63**: 180-4 (1973).
- 73E347 The triplet state of β -carotene and of analog polyenes of different length. Mathis, P.; Kleo, J. *Photochem. Photobiol.* **18**: 343-6 (1973).
- 741006 Equilibria between triplet states of aromatic hydrocarbons. Kira, A.; Thomas, J.K. *J. Phys. Chem.* **78**: 196-9 (1974).
- 741013 On the role of the triplet state in the photoisomerization of retinal isomers. Rosenfeld, T.; Alchahal, A.; Ottolenghi, M. *J. Phys. Chem.* **78**: 336-41 (1974).
- 743062 Triplet states of ubiquinone analogs studied by ultraviolet and electron nanosecond irradiation. Amouyal, E.; Bensasson, R.; Land, E.J. *Photochem. Photobiol.* **20**: 415-22 (1974).

- 746168 Interactions of triplet eosin with aromatics and intermolecular complexes of the ground state of eosin with aniline and pyridine in aqueous solutions and in ethanol. Chrysochoos, J. *Mol. Photochem.* **6**: 23-42 (1974).
- 746270 9,10-Diphenylanthracene as a fluorescence quantum yield standard. Heinrich, G.; Schoof, S.; Gusten, H. *J. Photochem.* **3**: 315-20 (1974).
- 746499 Intersystem crossing, ionic dissociation and cis-trans isomerization mechanisms in the photolysis of retinol and related molecules. Rosenfeld, T.; Alchalel, A.; Ottolenghi, M. *Excited States of Biological Molecules* (Proc. of the Int. Conf. at the Calouste Gulbenkian Foundation Centre, Lisbon, Portugal, April 18-24, 1974) J.B. Birks (ed.), Wiley, London, 1976, p.540-54.
- 747022 Laser flash photolysis of substituted stilbenes in solution. Bent, D.V.; Schulte-Frohlinde, D. *J. Phys. Chem.* **78**: 446-50 (1974).
- 747038 Increased quantum yield of photoreduction of dyes (a catalytic effect). Steiner, U.; Hafner, M.; Schreiner, S.; Kramer, H.E.A. *Photochem. Photobiol.* **19**: 119-28 (1974).
- 747049 Extinction coefficients for triplet-triplet absorption in ethanol solutions of anthracene, naphthalene, 2,5-diphenyloxazole, 7-diethylamino-4-methyl coumarin and 4-methyl-7-aminocarboxystyrene. Dempster, D.N.; Morrow, T.; Quinn, M.F. *J. Photochem.* **2**: 329-41 (1974).
- 747050 The photochemical characteristics of rhodamine 6G-ethanol solutions. Dempster, D.N.; Morrow, T.; Quinn, M.F. *J. Photochem.* **2**: 343-59 (1974).
- 747056 Time resolved molecular spectroscopy using high power solid state lasers in pulse transmission mode. A re-examination of the $S_0 \leftarrow S_1$ spectra of naphthalene and anthracene. Bebelar, D. *Chem. Phys.* **3**: 205-16 (1974).
- 747093 Flash-spectroscopic study of phthalazine. Kanamaru, N.; Nagakura, S.; Kimura, K. *Bull. Chem. Soc. Jpn.* **47**: 745-6 (1974).
- 747187 Primary photochemical processes of cationic acridine orange in aqueous solution studied by flash photolysis. Kellmann, A. *Photochem. Photobiol.* **20**: 103-8 (1974).
- 747189 Triplet states and cis-trans photoisomerization processes in the Schiff bases of retinal isomers. Rosenfeld, T.; Alchalel, A.; Ottolenghi, M. *Photochem. Photobiol.* **20**: 121-5 (1974).
- 747229 Formation and reactions of long lived xanthene dye radicals. II. Photochemical reduction of rhodamine-B and fluorescein derivatives. Krueger, U.; Memming, R. *Ber. Bunsenges. Phys. Chem.* **78**: 679-85 (1974).
- 747233 Quenching of triplet states of diazines by H-atom donors. Formation of azyl radicals. Bent, D.V.; Hayon, E.; Moorthy, P.N. *Chem. Phys. Lett.* **27**: 544-7 (1974).
- 747236 Triplet state of α -nitronaphthalene. Capellos, C.; Porter, G. *J. Chem. Soc., Faraday Trans. 2* **70**: 1159-64 (1974).
- 747293 Spectroscopic studies of formation and decay of triplet exciplexes. Evidence for a limited role of charge-transfer interactions in a nonpolar solvent. Roy, J.K.; Carroll, F.A.; Whitten, D.G. *J. Am. Chem. Soc.* **96**: 6349-55 (1974).
- 747334 Laser photolysis of retinal and its protonated and unprotonated n-butylamine Schiff base. Fisher, M.M.; Weiss, K. *Photochem. Photobiol.* **20**: 423-32 (1974).
- 747346 Picosecond flash photolysis: transient absorption in Sn(IV), Pd(II), and Cu(II) porphyrins. Magde, D.; Windsor, M.W.; Holtz, D.; Gouterman, M. *Chem. Phys. Lett.* **29**: 183-8 (1974).
- 747347 Triplet-triplet absorption of biphenyl and related compounds. Lavalette, D.; Tetreau, C. *Chem. Phys. Lett.* **29**: 204-9 (1974).
- 747376 Laser photolysis study of trans \rightarrow cis photoisomerization of trans-1-phenyl-2-(2-naphthyl)ethylene. Sumitani, M.; Nagakura, S.; Yoshihara, K. *Chem. Phys. Lett.* **29**: 410-3 (1974).
- 747389 Kinetics and mechanism of photochemical solvolysis of arylmethylacetates. Ivanov, V.B.; Ivanov, V.L.; Kuzmin, M.G. *Mol. Photochem.* **6**: 125-32 (1974).
- 747390 The triplet state of phenylpyridyl ketones in perfluorocarbon solvent. Bianchi, J.-P.; Watkins, A.R. *Mol. Photochem.* **6**: 133-42 (1974).
- 747417 The primary photochemical process of diphenylacetylene. Ota, K.; Murofushi, K.; Hoshi, T.; Inoue, H. *Tetrahedron Lett.* **14**: 1431-4 (1974).
- 747587 Ionic photodissociation in excited singlet and triplet states of the 1-methylnaphthalene-pyromellitic dianhydride charge-transfer complex. Achiba, Y.; Kimura, K. *Oyo Denki Kenkyusho Hokoku* **26**: 114-23 (1974, Publ. 1975).
- 747630 Laser study of triplet porphyrin quenching by oxygen in porphyrin-globins. Alpert, B.; Lindqvist, L. *Excited States of Biological Molecules* (Proc. of the Int. Conf. at the Calouste Gulbenkian Foundation Centre, Lisbon, Portugal, April 18-24, 1974) J.B. Birks (ed.), Wiley, London, 1976, p.425-33.
- 747633 Singlet-triplet quantum efficiencies of retinal isomers. Azerad, R.; Bensasson, R.; Cooper, M.B.; Dawe, E.A.; Land, E.J. *Excited States of Biological Molecules* (Proc. of the Int. Conf. at the Calouste Gulbenkian Foundation Centre, Lisbon, Portugal, April 18-24, 1974) J.B. Birks (ed.), Wiley, London, 1976, p.531-9.
- 74B002 Location of the first $^3B_1 \leftarrow ^3B_2^+$ transition in phenanthrene. Pavlopoulos, T.G. *Chem. Phys. Lett.* **27**: 245-8 (1974).
- 74B003 Spectroscopic studies of some laser dyes. Pavlopoulos, T.G.; Hammond, P.R. *J. Am. Chem. Soc.* **96**: 6568-79 (1974).
- 74B004 Polarization measurement and configuration analysis of the triplet-triplet absorption spectra of substituted naphthalenes. Hara, K.; Takemura, T.; Baba, H. *J. Mol. Spectrosc.* **50**: 90-105 (1974).
- 74B006 Chemische Gesellschaft Fribourg. Wild, U.P. *Chimia* **28**: 125-7 (1974).
- 74B007 Triplet-triplet absorption of porphyrins and their derivatives. Sapunov, V.V.; Solov'ev, K.N.; Tsvirko, M.P. *Zh. Prikl. Spektrosk.* **21**: 667-71 (1974).
- 751124 Nanosekunden-Pulsradiolyse und Laserphotolyse von Benzophenon in Benzol und Cyclohexan. Brede, O.; Helmstret, W.; Mehnert, R. *Z. Phys. Chem. (Leipzig)* **256**: 505-12 (1975).
- 753103 Kinetic studies of the tryptophan triplet state in solid film and in wool keratin. Ghiggino, K.P.; Nicholls, C.H.; Pailthorpe, M.T. *Photochem. Photobiol.* **22**: 169-73 (1975).
- 756077 Flash photolysis and phosphorescence of benzoic acid at 77-K. Acuna, A.U.; Ceballos, A.; Garcia Dominguez, J.A.; Molera, M.J. *An. Quim.* **71**: 22-7 (1975).
- 756162 Electron transfer reaction in the triplet state. Role of ferrocene as an electron donor. Kikuchi, K.; Kokubun, H.; Kikuchi, M. *Bull. Chem. Soc. Jpn.* **48**: 1378-81 (1975).
- 756270 Triplet-triplet absorption studies of the intersystem crossing mechanism of 2-aminopurines. Wierzchowski, K.L.; Berens, K.; Szabo, A.G. *J. Lumin.* **10**: 331-43 (1975).
- 756293 Excited singlet yield in T-T annihilation. A comparative study of naphthalene and anthracene in solution. Tfibel, F.; Lindqvist, L. *Chem. Phys.* **10**: 471-8 (1975).
- 757004 Flash photolytic kinetic studies of the quenching of tetracene triplets by tetracene radical anions. Levin, G.; Szwarc, M. *Chem. Phys. Lett.* **30**: 116-9 (1975).
- 757066 Formation and reactions of the triplet state of sym-triazine in solution studied by laser spectroscopy. Bent, D.V.; Hayon, E. *Chem. Phys. Lett.* **31**: 325-7 (1975).
- 757078 Determination of the pK values of the lumiflavin triplet state by flash photolysis. Schreiner, S.; Steiner, U.; Kramer, H.E.A. *Photochem. Photobiol.* **21**: 81-4 (1975).
- 757090 Triplet-sensitized cis-trans isomerization of the protonated Schiff base of retinal isomers. Alchalel, A.; Honig, B.; Ottolenghi, M.; Rosenfeld, T. *J. Am. Chem. Soc.* **97**: 2161-6 (1975).
- 757112 Activation-controlled hydrogen abstraction by benzophenone triplet. Topp, M.R. *Chem. Phys. Lett.* **32**: 144-9 (1975).
- 757141 The determination of molar absorption coefficients of metastable states. Ranald, U.B.; Kaenzig, H.; Wild, U.P. *J. Photochem.* **4**: 97-107 (1975).
- 757148 Mechanism of photoreduction of thiazine dyes by EDTA studied by flash photolysis. III. Consequences of a newly found pK_T of thionine on the mechanism in basic solutions. Bonneau, R.; Pereyre, J. *Photochem. Photobiol.* **21**: 173-7 (1975).

- 757161 Excited state chemistry of aromatic amino acids and related peptides. I. Tyrosine. Bent, D.V.; Hayon, E. *J. Am. Chem. Soc.* **97**: 2599-606 (1975).
- 757162 Excited state chemistry of aromatic amino acids and related peptides. II. Phenylalanine. Bent, D.V.; Hayon, E. *J. Am. Chem. Soc.* **97**: 2606-12 (1975).
- 757163 Excited state chemistry of aromatic amino acids and related peptides. III. Tryptophan. Bent, D.V.; Hayon, E. *J. Am. Chem. Soc.* **97**: 2612-9 (1975).
- 757167 Photochemical rearrangements of indazoles. Investigation of the triplet excited states of 1- and 2-methylindazole. Ferris, J.P.; Prabhu, K.V.; Strong, R.L. *J. Am. Chem. Soc.* **97**: 2835-9 (1975).
- 757282 Determination of triplet quantum yields by laser flash absorption spectroscopy. Amand, B.; Bensasson, R. *Chem. Phys. Lett.* **34**: 44-8 (1975).
- 757309 Chemistry of the triplet state of diazines in solution studied by laser spectroscopy. Bent, D.V.; Hayon, E.; Moorthy, P.N. *J. Am. Chem. Soc.* **97**: 5065-71 (1975).
- 757353 Laser photoionization of phenothiazine in alcoholic and aqueous micellar solution. Electron transfer from triplet states to metal ion acceptors. Alkaitis, S.A.; Beck, G.; Graetzel, M. *J. Am. Chem. Soc.* **97**: 5723-9 (1975).
- 757361 Photosensitizing properties of N-formylkynurenine. Walrant, P.; Santus, R.; Grossweiner, L.I. *Photochem. Photobiol.* **22**: 63-5 (1975).
- 757413 Rate constants of triplet state ionic photodissociation of weak charge-transfer complexes formed between pyromellitic dianhydride (PMDA) and naphthalenes. Achiba, Y.; Kimura, K. *J. Phys. Chem.* **79**: 2626-8 (1975).
- 757427 The photochemistry of benz(de)anthracen-7-ones. Part II. Flash photolysis. Bentley, P.; McKellar, J.F.; Phillips, G.O. *J. Chem. Soc., Perkin Trans. 2* : 1259-62 (1975).
- 757439 Phenyl substitution effects on triplet-triplet absorption spectra. I. Study of 1,3,6,8-tetraphenylpyrene. Rulliere, C.; Colson, E.C.; Roberge, P.C. *Can. J. Chem.* **53**: 3269-75 (1975).
- 757441 Study of the intermediate states in photochemical reactions of dyes. VII. Electron transfer in redox reactions of thiocarbocyanines. Lifanov, Yu.I.; Korobov, V.E.; Karvakin, A.V.; Chibisov, A.K. *High Energy Chem.* **9**: 232-6 (1975). Translated from: *Khim. Vys. Energ.* **9**: 265-70 (1975).
- 757469 An electro-optic shutter to assist flash photolysis studies of solid films. Ghigino, K.P.; Nicholls, C.H.; Pailthorpe, M.T. *J. Phys. E* **8**: 900-2 (1975).
- 757485 Photodynamic sensitization by 8-methoxypsoralen via the singlet oxygen mechanism. Poppe, W.; Grossweiner, L.I. *Photochem. Photobiol.* **22**: 217-9 (1975).
- 757510 Studies on thymine and uracil triplet excited state in acetonitrile and water. Salet, C.; Bensasson, R. *Photochem. Photobiol.* **22**: 231-5 (1975).
- 757522 Triplet absorption of phthalimide solutions. Borisevich, N.A.; Bolot'ko, L.M.; Staneva, T.G. *Zh. Prikl. Spektrosk.* **23**: 633-7 (1975).
- 757534 An efficient photosensitizer: O-ethyl-1-thionaphthoate. Gisin, M.; Wirz, J. *Helv. Chim. Acta* **58**: 1768-71 (1975).
- 757573 Microsecond and nanosecond impulse photolysis of carbazoles. Two-quantum reaction in a liquid medium. Borovkova, V.A.; Bagdasar'yan, Kh.S.; Pikel'ni, V.F.; Kolosov, V.A.; Kiryukhin, Yu.I. *Dokl. Phys. Chem. (Engl. Transl.)* **224**: 973-6 (1975). Translated from: *Dokl. Akad. Nauk SSSR* **224**: 616-9 (1975).
- 757594 Absolute efficiency of the two-quanta sensitized decomposition of benzylamine. Sinitsyna, Z.A.; Kiryukhin, Yu.I.; Bagdasar'yan, Kh.S. *Dokl. Phys. Chem. (Engl. Transl.)* **225**: 1212-4 (1975). Translated from: *Dokl. Akad. Nauk SSSR* **225**: 361-3 (1975).
- 75B002 Anisotropy of the T-T absorption spectra of naphthalene and phenanthrene. Kuball, H.-G.; Euing, W. *Chem. Phys. Lett.* **30**: 457-62 (1975).
- 75B003 Matrix effect on the linewidth of the $^3B_{2u} \rightarrow ^3B_{1g}$ absorption spectrum of naphthalene. Haertel, H.; Khelladi, F.Z.; Ostertag, R. *Chem. Phys. Lett.* **30**: 472-6 (1975).
- 75B004 The triplet-triplet absorption spectra and electronic structures of some cyano-substituted benzenes. Morita, H.; Matsumoto, S.; Nagakura, S. *Bull. Chem. Soc. Jpn.* **48**: 420-3 (1975).
- 75E530 Degradation of the energy of electronic excitation of associated molecules. The triplet states of associates of rhodamine 6G. Chibisov, A.K.; Ketsle, G.A.; Levshin, L.V.; Slavnova, T.D. *Opt. Spectrosc.* **38**: 45-8 (1975). Translated from: *Opt. Spektrosk.* **38**: 83-8 (1975).
- 761024 Energy transfer from an alkene triplet state during pulse radiolysis. Barwise, A.J.G.; Gorman, A.A.; Rodgers, M.A.J. *Chem. Phys. Lett.* **38**: 313-6 (1976).
- 761030 Studies of higher triplet exciton bands through T-T absorption in naphthalene single crystal. Higuchi, M.; Nakayama, T.; Itoh, N. *J. Phys. Soc. Jpn.* **40**: 250-7 (1976).
- 761035 Triplet states of carotenoids from photosynthetic bacteria studied by nanosecond ultraviolet and electron pulse irradiation. Bensasson, R.; Land, E.J.; Maudinas, B. *Photochem. Photobiol.* **23**: 189-93 (1976).
- 761069 Intermediates in the nanosecond pulse radiolysis of dimethylaniline solutions in cyclohexane. Zador, E.; Warman, J.M.; Hummel, A. *J. Chem. Soc., Faraday Trans. 1* **72**: 1368-76 (1976).
- 761080 Pulse radiolysis of ethyl acetate and its solutions. Ramanam, G. *J. Phys. Chem.* **80**: 1553-8 (1976).
- 761088 The triplet state of 1,6-diphenyl-1,3,5-hexatriene and 1,8-diphenyl-1,3,5,7-octatetraene. Bensasson, R.; Land, E.J.; Lafferty, J.; Sinclair, R.S.; Truscott, T.G. *Chem. Phys. Lett.* **41**: 333-5 (1976).
- 761122 Triplet excited states of 1,4-disubstituted anthraquinones: Possible evidence for association of quinones in solution. Land, E.J.; McAlpine, E.; Sinclair, R.S.; Truscott, T.G. *J. Chem. Soc., Faraday Trans. 1* **72**: 2091-100 (1976).
- 761168 The triplet excited state of bilirubin. Land, E.J. *Photochem. Photobiol.* **24**: 475-7 (1976).
- 766206 Excited states of six-membered aza-aromatic rings. X. Acid-base equilibria of electronically excited quinoxaline as an example of anomalous diazines. Grabowska, A.; Herbach, J.; Kirkorkaminska, E.; Pakula, B. *J. Lumin.* **11**: 403-12 (1976).
- 766377 The hydrogen bonding effect of pyridine on the deactivation of the excited states of acridone. Fushimi, K.; Kikuchi, K.; Kokubun, H. *J. Photochem.* **5**: 457-68 (1976).
- 766403 Properties of benzopyrene-DNA complexes investigated by fluorescence and triplet flash photolysis techniques. Geacintov, N.E.; Prusik, T.; Khosrofi, J.M. *J. Am. Chem. Soc.* **98**: 6444-52 (1976).
- 766421 Photochemical behaviour and spectroscopical properties of anilino naphthalenes. Fratev, F.; Polansky, O.E.; Zander, M. *Z. Naturforsch. Teil A* **31A**: 987-9 (1976).
- 766464 Picosecond time-resolved spectroscopy and the intersystem crossing rates of anthrone and fluorenone. Kobayashi, T.; Nagakura, S. *Chem. Phys. Lett.* **43**: 429-34 (1976).
- 766469 Intersystem crossing to the lowest triplet state of phenazine following singlet excitation with a picosecond pulse. Hirata, Y.; Tanaka, I. *Chem. Phys. Lett.* **43**: 568-70 (1976).
- 766528 Evidence for a triplet exciplex state from pulsed laser photolysis. Bell, I.P.; Rodgers, M.A.J. *Chem. Phys. Lett.* **44**: 249-52 (1976).
- 766652 The use of charge transfer complexing to elucidate some triplet-triplet spectra. Slifkin, M.A.; Al-Chalabi, A.O. *Spectrochim. Acta, Part A* **32A**: 661-2 (1976).
- 767023 Photochemistry of chlorpromazine (2-chloro-N-(3-dimethylaminopropyl)phenothiazine) in propan-2-ol solution. Davies, A.K.; Navaratnam, S.; Phillips, G.O. *J. Chem. Soc., Perkin Trans. 2* : 25-9 (1976).
- 767071 Direct and sensitized photooxidation of cyanine dyes. Byers, G.W.; Gross, S.; Henrichs, P.M. *Photochem. Photobiol.* **23**: 37-43 (1976).

- 767094 Mechanism of quenching of the triplet states of organic compounds by tris-(β -diketonato) complexes of iron(III), ruthenium(III) and aluminium(III). Wilkinson, F.; Farmilo, A. J. Chem. Soc., Faraday Trans. 2 72: 604-18 (1976).
- 767144 Duroquinone triplet reduction, in cyclohexane, ethanol and water, and by durohydroquinone. Amouyal, E.; Bensasson, R. J. Chem. Soc., Faraday Trans. 1 72: 1274-87 (1976).
- 767147 Triplet-triplet extinction coefficients of anthracene and 9-bromoanthracene determined by a ground state depletion method. Ledger, M.B.; Salmon, G.A. J. Chem. Soc., Faraday Trans. 2 72: 883-6 (1976).
- 767159 Behaviors of the triplet molecule in neat liquid of naphthalene derivatives at room temperature. Ohno, T.; Kato, S. Chem. Lett. (Tokyo): 263-6 (1976).
- 767171 Laser photolysis studies of the triplet state of xanthone and its ketyl radical in fluid solution. Garner, A.; Wilkinson, F. J. Chem. Soc., Faraday Trans. 2 72: 1010-20 (1976).
- 767177 Laser photoionization and light-initiated redox reactions of tetramethylbenzidine in organic solvents and aqueous micellar solution. Alkaitis, S.A.; Graetzel, M. J. Am. Chem. Soc. 98: 3549-54 (1976).
- 767180 Laser flash spectroscopy of tris(2,2'-bipyridine)ruthenium(II) in solution. Bensasson, R.; Salet, C.; Balzani, V. J. Am. Chem. Soc. 98: 3722-4 (1976).
- 767189 Quenching of triplet states by inorganic ions. Energy transfer and charge transfer mechanisms. Treinin, A.; Hayon, E. J. Am. Chem. Soc. 98: 3884-91 (1976).
- 767201 A laser-photolytic study of 9-fluorenone. Preliminary observations of transient absorption spectra. Nakajima, A. Mol. Photochem. 7: 251-62 (1976).
- 767246 Photochemical investigations of oxazine, thiazine and selenazine dyes. The reactivity of protolytic triplet forms in electron transfer reactions. Vogelmann, E.; Kramer, H.E.A. Photochem. Photobiol. 23: 383-90 (1976).
- 767261 Build-up of T-T absorption of acridine following singlet excitation with a picosecond pulse. Hirata, Y.; Tanaka, I. Chem. Phys. Lett. 41: 336-8 (1976).
- 767269 Triplet excited states of nitronaphthalenes. II. b-Nitronaphthalene. Capellos, C.; Suryanarayanan, K. Int. J. Chem. Kinet. 8: 529-39 (1976).
- 767270 Triplet excited states of nitronaphthalenes. III. 1,4-Dinitronaphthalene. Capellos, C.; Suryanarayanan, K. Int. J. Chem. Kinet. 8: 541-8 (1976).
- 767488 Photochemical reactions of bifluorenylidene. Part 2. Flash photolytic investigations. Van Sinooy, A.; Vander Donckt, E. J. Chem. Soc., Faraday Trans. 1 72: 2312-23 (1976).
- 767526 Low temperature flash photolysis of 2-N-dimethylaminopyridine. Wolleben, J.; Testa, A.C. Mol. Photochem. 7: 277-85 (1976).
- 767556 Electron transfer in the photochemistry of substituted benzoate and naphthoate esters. Coyle, J.D.; Kingston, D.H. J. Chem. Soc., Perkin Trans. 2: 1475-9 (1976).
- 767574 The photoreduction of methylene blue by amines. I. A flash photolysis study of the reaction between triplet methylene blue and amines. Kayser, R.H.; Young, R.H. Photochem. Photobiol. 24: 395-401 (1976).
- 767582 Excited state chemistry of indigoid dyes. IV. Triplet state spectra and lifetimes from flash photolysis at low temperatures. Schulte-Frohlinde, D.; Herrmann, H.; Wyman, G.M. Z. Phys. Chem. (Frankfurt Am Main) 101: 115-21 (1976).
- 767626 Deactivation of triplet 2-naphthol and 1-anthrol by aromatic N-heterocycles. Yamamoto, S.; Kikuchi, K.; Kokubun, H. Bull. Chem. Soc. Jpn. 49: 2950-1 (1976).
- 767647 Participation of triplet states in the photocoloration of photochromic spiropyranes of the phenanthridine series. Kuz'min, V.A.; Malkin, Ya.N.; Martynova, V.P.; Zakhs, E.R.; Efros, L.S. Dokl. Phys. Chem. (Engl. Transl.) 228: 422-5 (1976) Translated from: Dokl. Akad. Nauk SSSR 228: 127-30 (1976).
- 767764 Configuration of photoisomers of merocyanine dyes. Kuz'min, V.A.; Vinogradov, A.M.; Romanov, N.N.; Al'perovich, M.A.; Levkoev, I.I.; Babichev, F.S. Bull. Acad. Sci. USSR, Div. Chem. Sci. 25(8): 1755-7 (1976) Translation from: Izv. Akad. Nauk SSSR, Ser. Khim. (8): 1864-6 (1976).
- 767803 Photoreduction of methylene blue by aromatic amines. Nanosecond pulse photolysis. Romashov, L.V.; Kiryukhin, Yu.I.; Bagdasar'yan, Kh.S. Dokl. Phys. Chem. 230: 961-4 (1976) Translated from: Dokl. Akad. Nauk SSSR 230: 1145-8 (1976).
- 76B012 Phosphorescence and triplet-triplet absorption spectra of anthracene-d₁₀ in polymethylmethacrylate under high pressures. Shaw, R.W.; Nicol, M. Chem. Phys. Lett. 39: 108-12 (1976).
- 76E682 Features of the triplet-triplet absorption spectra of composite molecules. Gerko, V.I.; Alifimov, M.V.; Popov, L.S.; Kovalenko, N.P. Dokl. Phys. Chem. 230: 811-3 (1976) Translated from: Dokl. Akad. Nauk SSSR 230: 125-7 (1976).
- 771021 Triplet state quenching involving charge transfer interactions between N-methyl indole and aromatic carbonyl compounds in benzene solution. Wilkinson, F.; Garner, A. J. Chem. Soc., Faraday Trans. 2 73: 222-33 (1977).
- 771048 Radiation induced racemization of 1,1'-binaphthyl in tetrahydrofuran and toluene. Irie, M.; Yorozu, T.; Yoshida, K.; Hayashi, K. J. Phys. Chem. 81: 973-6 (1977).
- 771059 Ion recombination processes in rigid solutions. Miller, J.R. Can. J. Chem. 55: 1867-75 (1977).
- 771078 Excited states of protoporphyrin IX dimethyl ester: Reaction of the triplet with carotenoids. Chantrell, S.J.; McAuliffe, C.A.; Munn, R.W.; Pratt, A.C.; Land, E.J. J. Chem. Soc., Faraday Trans. 1: 858-65 (1977).
- 773047 A pulsed laser study of excited states of aromatic molecules absorbed in globular proteins. Cooper, M.; Thomas, J.K. Radiat. Res. 70: 312-24 (1977).
- 776081 Intersystem crossing in althracene-N,N-diethylaniline exciplex. Nishimura, T.; Nakashima, N.; Mataga, N. Chem. Phys. Lett. 46: 334-8 (1977).
- 776104 Charge transfer triplet state of p-nitroaniline. Wolleben, J.; Testa, A.C. J. Phys. Chem. 81: 429-31 (1977).
- 776146 Photophysical processes in the molecular complexes of 1,2,4,5-tetracyanobenzene with aromatic donors. Craig, B.B.; Rodgers, M.A.J.; Wood, B. J. Chem. Soc., Faraday Trans. 2 73: 349-55 (1977).
- 776171 Probabilities of intramolecular radiative and radiationless transitions of some amino derivatives of acridine. Morozov, Yu.V.; Savenko, A.K. Mol. Photochem. 8: 1-43 (1977).
- 776187 The fluorescent state of cyano-substituted layered cyclophanes. Masuhara, H.; Mataga, N.; Yoshida, M.; Tatsumitsu, H.; Sakata, Y.; Misumi, S. J. Phys. Chem. 81: 879-83 (1977).
- 776190 The effect of chemical structure on the photosensitizing efficiencies of porphyrins. Cauzzo, G.; Gennari, G.; Jori, G.; Spikes, J.D. Photochem. Photobiol. 25: 389-95 (1977).
- 776194 Triplet excited states of nitronaphthalenes. IV. 1,2- and 1,8-dinitronaphthalenes. Capellos, C.; Suryanarayanan, K. Int. J. Chem. Kinet. 9: 399-407 (1977).
- 776195 The triplet state of 4-nitronaphthylamine. Capellos, C.; Lang, F. Int. J. Chem. Kinet. 9: 409-15 (1977).
- 776213 Weak triplet-triplet absorption spectra of cyanobenzenes. Achiba, Y.; Kimura, K. Chem. Phys. Lett. 48: 107-10 (1977).
- 776222 Luminescence and triplet absorption of o-, m-, and p-methylbenzoic acids. Acuna, A.U.; Ceballos, A.; Molera, M.J. J. Phys. Chem. 81: 1090-3 (1977).
- 776227 Intersystem crossing processes of acridine. Kikuchi, K.; Uji-ie, K.; Miyashita, Y.; Kokubun, H. Bull. Chem. Soc. Jpn. 50(4): 879-82 (1977).
- 776258 Intersystem crossing and internal conversion quantum yields of acridine in polar and nonpolar solvents. Kellmann, A. J. Phys. Chem. 81: 1195-8 (1977).
- 776262 P-type delayed fluorescence from rubrene. Liu, D.K.K.; Faulkner, L.R. J. Am. Chem. Soc. 99: 4594-9 (1977).

- 776279 Polarized triplet-triplet absorption spectra using stretched polymer films: Biphenyl and fluorene. Ota, K.; Murofushi, K.; Hoshi, T.; Shibuya, E.; Yoshino, J. *Z. Phys. Chem. (Frankfurt am Main)* **104**: 181-8 (1977).
- 776335 Studies of the interaction of excited chloranil with acrylonitrile, methyl methacrylate, and styrene by means of laser photolysis. Kobashi, H.; Gyoda, H.; Morita, T. *Bull. Chem. Soc. Jpn.* **50**: 1731-8 (1977).
- 776412 Singlet \rightarrow triplet intersystem crossing quantum yields of photosynthetic and related polyenes. Bensasson, R.; Dawe, E.A.; Long, D.A.; Land, E.J. *J. Chem. Soc., Faraday Trans. 1* **73**: 1319-25 (1977).
- 776431 Proton-induced radiationless processes in excited 9,9'-bianthryl. Shizuka, H.; Ishii, Y.; Morita, T. *Chem. Phys. Lett.* **51**: 40-4 (1977).
- 776464 Two emitting states of excited p-(9-anthryl)-N,N-dimethylaniline derivatives in polar solvents. Siemiarz, A.; Grabowski, Z.R.; Krowczynski, A.; Asher, M.; Ottolenghi, M. *Chem. Phys. Lett.* **51**: 315-20 (1977).
- 777036 Triplet states of cyanine dyes and reactions of electron transfer with their participation. Chibisov, A.K. *J. Photochem.* **6**: 199-214 (1976/77).
- 777037 Laser photolysis of indole in cyclohexane. Pernot, C.; Lindqvist, L. *J. Photochem.* **6**: 215-20 (1976/77).
- 777041 Triplet state of rhodamine dyes and its role in production of intermediates. Korobov, V.E.; Shubin, V.V.; Chibisov, A.K. *Chem. Phys. Lett.* **45**: 498-501 (1977).
- 777045 Laser photolysis studies of duroquinone triplet state electron transfer reactions. Scheerer, R.; Graetzel, M. *J. Am. Chem. Soc.* **99**: 865-71 (1977).
- 777062 Anion and solvent effects on the rate of reduction of triplet excited thiazine dyes by ferrous ions. Wildes, P.D.; Lichtin, N.N.; Hoffman, M.Z.; Andrews, L.; Linschitz, H. *Photochem. Photobiol.* **25**: 21-5 (1977).
- 777088 Photoionization and recombination luminescence of N,N'-disubstituted dihydropyrazines in 3-methylpentane at 77 K. Bruehlmann, U.; Huber, J.R. *J. Phys. Chem.* **81**: 386-91 (1977).
- 777201 Photoreduction of 1,10-phenanthroline. Bandyopadhyay, B.N.; Harriman, A. *J. Chem. Soc., Faraday Trans. 1* **73**: 663-74 (1977).
- 777238 Reversible photochemistry and photocyclization of 10H,10'H-bianthrylidene. Korenstein, R.; Muszkat, K.A.; Fischer, E. *J. Chem. Soc., Perkin Trans. 2* **5**: 564-9 (1977).
- 777241 Laser photolysis study of the photoracemization of 1,1'-binaphthyl. Irie, M.; Yoshida, K.; Hayashi, K. *J. Phys. Chem.* **81**: 969-72 (1977).
- 777242 Investigation of physical triplet quenching by electron donors. Steiner, U.; Winter, G.; Kramer, H.E.A. *J. Phys. Chem.* **81**: 1104-10 (1977).
- 777265 Laser investigation of some oxazole and oxadiazole derivatives. Fouassier, J.-P.; Lougnot, D.-J.; Wieder, F.; Faure, J. *J. Photochem.* **7**(1): 17-28 (1977).
- 777315 Photoredox reactions of thionine. Ferreira, M.I.C.; Harriman, A. *J. Chem. Soc., Faraday Trans. 1* **73**: 1085-92 (1977).
- 777316 Triplet-triplet absorption spectra and the spectra of the photo-reduced states of rhodamine B and rhodamine 110. Dunne, A.; Quinn, M.F. *J. Chem. Soc., Faraday Trans. 1* **73**: 1104-10 (1977).
- 777345 Lifetime enhancement of the first electronic triplet state of 3,5-dinitroanisole by hydrogen bonding to clusters in mixed aqueous solvent systems. Varma, C.A.G.O.; Plantenga, F.L.; van den Ende, C.A.M.; van Zeyl, P.H.M.; Tamminga, J.J.; le Cornélisse, J. *Chem. Phys.* **22**: 475-83 (1977).
- 777388 Photoionization of aromatic compounds in boric acid glass. Andreev, O.M.; Smirnov, V.A.; Alfimov, M.V. *J. Photochem.* **7**: 149-56 (1977).
- 777391 Mechanisms of deactivation of triplet 1-anthrol and 2-naphthol by aromatic N-heterocycles. Yamamoto, S.-A.; Kikuchi, K.; Kokubun, H. *J. Photochem.* **7**: 177-84 (1977).
- 777432 Flash photolysis of tryptophan and N-acetyl-L-tryptophanamide. The effect of bromide on transient yields. Volkert, W.A.; Kuntz, R.R.; Ghiron, C.A.; Evans, R.F.; Santus, R.; Bazin, M. *Photochem. Photobiol.* **26**: 3-9 (1977).
- 777434 Photochemistry of polyenes. XI. Flash photolytic studies of hindered isomers of retinal. Harriman, A.; Liu, R.S.H. *Photochem. Photobiol.* **26**: 29-32 (1977).
- 777455 Laser flash photolysis and quantum yields of the direct photochemical cis-trans isomerization of 4-nitro-4'-methoxystilbene in cyclohexane solutions in presence of quenchers. Model for the photoisomerization of nitrostilbenes. Goerner, H.; Schulte-Frohlinde, D. *Ber. Bunsenges. Phys. Chem.* **81**: 713-20 (1977).
- 777538 Abnormal broadening in triplet-to-triplet absorption spectra caused by aromatic solvents. Alfimov, M.V.; Gerko, V.I.; Popov, L.S.; Razumov, V.F. *Chem. Phys. Lett.* **50**: 398-401 (1977).
- 777602 Interaction of duroquinone lowest triplet with amines. Amouyal, E.; Bensasson, R. *J. Chem. Soc., Faraday Trans. 1* **73**: 1561-8 (1977).
- 777603 Photochemistry of Michler's ketone in cyclohexane and alcohol solvents. Brown, R.G.; Porter, G. *J. Chem. Soc., Faraday Trans. 1* **73**: 1569-73 (1977).
- 777617 Photochemistry of flavins. II. Photophysical properties of alloxazines and isoalloxazines. Grodowski, M.S.; Veyret, B.; Weiss, K. *Photochem. Photobiol.* **26**: 341-52 (1977).
- 777635 Picosecond-laser study of triplet state population of carbonyl derivatives of anthracene. Hirayama, S.; Kobayashi, T. *Chem. Phys. Lett.* **52**: 55-8 (1977).
- 77A178 Direct evidence of photodissociation of carbazole via the higher excited triplet state by a double excitation method. Yamamoto, S.; Kikuchi, K.; Kokubun, H. *Chem. Lett.* **6**: 1173-6 (1977).
- 77A196 Intermediates from the photolysis of 2,3-dihydroxynaphthalene. Khudyakov, I.V.; Tatikolov, A.S.; Kuz'min, V.A. *Bull. Acad. Sci. USSR, Div. Chem. Sci.* **26**: 1197-9 (1977) Translated from: *Izv. Akad. Nauk SSSR, Ser. Khim.* **1300**-3 (1977).
- 77A203 The electron transfer reaction between triplet methylene blue and aromatic compounds. Kikuchi, K.; Tamura, S.-I.; Iwanaga, C.; Kokubun, H.; Usui, Y. *Z. Phys. Chem. (Frankfurt am Main)* **106**: 17-24 (1977).
- 77A216 The triplet state of 4-nitro-N,N-dimethylnaphthylamine. Capellos, C.; Lang, F. *Int. J. Chem. Kinet.* **9**: 943-52 (1977).
- 77B012 Two-quantum photoionization in liquid solutions. Quantitative investigation by means of nanosecond photolysis. Borovkova, V.A.; Bagdasar'yan, Kh.S. *Dokl. Phys. Chem.* **235**: 649-52 (1977) Translated from: *Dokl. Akad. Nauk SSSR* **235**: 132-5 (1977).
- 77B022 Triplet-triplet absorption of some carbonyl-containing compounds. Alfimov, M.V.; Buben, N.Ya.; Glagolev, V.L.; Kuyumdzhii, E.S.; Pomazan, Yu.V.; Shamshev, V.N. *Opt. Spectrosc.* **42**: 267-70 (1977) Translated from: *Opt. Spektrosk.* **42**: 476-82 (1977).
- 77C001 Dimerisation de la chlorophylle dans les solvants apolaires. Etude de l'équilibre et de l'état triplet. Journeaux, R.; Chene, G.; Viovy, R. *J. Chim. Phys. Phys.-Chim. Biol.* **74**: 1203-10 (1977).
- 77E543 Excited singlet state decay pathways in the fluorescence probe molecule 1,8-anilidonaphthalene sulphonate (ANS). Fleming, G.R.; Porter, G.; Robbins, R.J.; Synowiec, J.A. *Chem. Phys. Lett.* **52**: 228-32 (1977).
- 77E663 Mechanism of the degradation of electronic excitation energy in hydrogen-bonded ion pairs. Ivanov, V.L.; Martynov, I.Yu.; Uzhinov, B.M.; Kuz'min, M.G. *High Energy Chem.* **11**: 361-4 (1977) Translated from: *Khim. Vys. Energ.* **11**: 327-31 (1977).
- 77E797 Etude par spectroscopie laser de dérivés de benzoïne. Fouassier, J.-P.; Lougnot, D.-J.; Faure, J. *C.R. Seances Acad. Sci., Ser. C* **284**: 643-6 (1977).
- 78A163 Photooxidation of aromatic hydrocarbons by europium(III) salts. Levin, G. *J. Phys. Chem.* **82**: 1584-8 (1978).

- 78A180 Deactivation of the electronic excitation of triarylcarbonium ions in acid media. Ivanov, V.L.; Al'-Ainen, S.A.; Kuz'min, M.G. *High Energy Chem.* **12**: 39-42 (1978) Translated from: *Khim. Vys. Energ.* **12**: 48-52 (1978).
- 78A195 Hydrogen atom abstraction by p-chloranil triplet from tetrachlorohydroquinone and the presence of a competitive deactivation process. Kobashi, H.; Nagumo, T.; Morita, T. *Chem. Phys. Lett.* **57**: 369-72 (1978).
- 78A237 Flash photolytic studies on the photochemical formation of five-membered sulfur heterocycles. Wolff, T. *J. Am. Chem. Soc.* **100**: 6157-9 (1978).
- 78A268 Transient absorption in the sensitized photoisomerization of stilbene. Sumitani, M.; Yoshihara, K.; Nagakura, S. *Bull. Chem. Soc. Jpn.* **51**: 2503-7 (1978).
- 78A304 Primary processes in the photochemistry of rhodamine dyes. Korobov, V.E.; Chibisov, A.K. *J. Photochem.* **9**: 411-24 (1978).
- 78A324 Electron transfer from aromatic molecules to dimethylmercury via a triplet exciplex. Vander Donckt, E.; Van Vooren, C. *J. Chem. Soc., Faraday Trans. 1* **74**: 827-36 (1978).
- 78A345 Blitzlichtspektroskopische Untersuchungen des Tripletzustandes und der photolytisch gebildeten Radikalkationen des 3,4-Benzpyrens in waessrigen Natriumdodecylsulfat- und Coffeinloesungen. Bauer, H.; Reske, G. *J. Photochem.* **9**: 43-54 (1978).
- 78A368 Quantitative study of the photodissociation of carbazoles in the higher excited triplet state by a double excitation method. Yamamoto, S.-A.; Kikuchi, K.; Kokubun, H. *Z. Phys. Chem. (Frankfurt am Main)* **109**: 47-58 (1978).
- 78A378 Triplet states of copper and metal-free phthalocyanines. McVie, J.; Sinclair, R.S.; Truscott, T.G. *J. Chem. Soc., Faraday Trans. 2* **74**: 1870-9 (1978).
- 78A386 Laser photolysis of dicarbocyanine dyes. Darmanyan, A.P.; Kuz'min, V.A. *Bull. Acad. Sci. USSR, Div. Chem. Sci.* **27**: 506-10 (1978) Translated from: *Izv. Akad. Nauk SSSR, Ser. Khim.* : 587-92 (1978).
- 78A411 Pulse radiolysis of benzene solutions of indolinispiropyrans. Vannikov, A.V.; Kryukov, A.Yu. *High Energy Chem.* **12**: 280-4 (1978) Translated from: *Khim. Vys. Energ.* **12**: 334-8 (1978).
- 78A447 The electron transfer reactions between triplet dyes and aromatic compounds in acetonitrile. Tamura, S.-I.; Kikuchi, K.; Kokubun, H.; Usui, Y. *Z. Phys. Chem. (Wiesbaden)* **111**: 7-18 (1978).
- 78B071 Radiationless deactivation of benzophenone triplets in aromatic solvents. Characterization of transient intermediates in the presence of diphenyl ether. Small, R.D.; Scaiano, J.C. *J. Phys. Chem.* **82**: 2064-6 (1978).
- 78B072 Mechanism of the photochemical reaction of diphenylamines with carbon tetrachloride. Wyrzykowska, K.; Grodowski, M.; Weiss, K.; Latowski, T. *Photochem. Photobiol.* **28**: 311-8 (1978).
- 78B081 Triplet-triplet absorption of some aryl-substituted 1,3,4-oxadiazoles. Rulliere, C.; Rayez, J.-C.; Roberge, P.C. *Can. J. Chem.* **56**: 2781-5 (1978).
- 78B088 Tripletspectra of 4-nitrostilbenes in polar and non-polar solvents at different temperatures and the configuration of the triplet state. Goerner, H.; Schulte-Frohlinde, D. *Ber. Bunsenges. Phys. Chem.* **82**: 1102-7 (1978).
- 78B089 Formation and decay of triplet excimers as revealed by their absorption spectra: Intramolecular excimers of α,α -dinaphthylalkanes. Subudhi, P.C.; Lim, E.C. *Chem. Phys. Lett.* **56**: 59-61 (1978).
- 78B108 Mechanisms of antioxidant action. A photochemical study of transition metal dibutyl dithiocarbamates (and their photostabilizing role in a polyolefin matrix). Scott, G.; Sheena, H.H.; Harriman, A.M. *Eur. Polym. J.* **14**: 1071-7 (1978).
- 78B112 Dynamic behaviour of excited charge transfer systems in polar solvents. Masuhara, H.; Saito, T.; Maeda, Y.; Mataga, N. *J. Mol. Struct.* **47**: 243-59 (1978).
- 78B129 Polarized triplet-triplet absorption spectra using stretched polymer film technique: 4-Hydroxy- and 4,4'-dihydroxy-biphenyl. Hoshi, T.; Oka, M.; Komuro, M.; Yoshino, J.; Ota, K.; Murofushi, K. *Z. Phys. Chem. (Wiesbaden)* **112**: 129-37 (1978).
- 78E011 Triplet state of protoporphyrin IX. Lafferty, J.; Truscott, T.G. *J. Chem. Soc., Chem. Commun.* : 51-2 (1978).
- 78E019 Nonradiative relaxation processes of the higher excited triplet states of anthracenes studied by a double excitation method. Kobayashi, S.; Kikuchi, K.; Kokubun, H. *Chem. Phys.* **27**: 399-407 (1978).
- 78E057 Triplet-triplet absorption spectroscopy of some highly dipolar unsaturated nitro compounds. Cowley, D.J. *Helv. Chim. Acta* **61**: 184-97 (1978).
- 78E062 Flash photolysis study of aminopyridines. Wolleben, J.; Testa, A.C. *J. Photochem.* **8**: 125-30 (1978).
- 78E152 Polarized photochromism in solid solutions. Pasternak, C.; Slifkin, M.A.; Shinitzky, M. *J. Chem. Phys.* **68**: 2669-73 (1978).
- 78E157 Triplet excited state of furocoumarins: Reactions with nucleic acid bases and amino acids. Bensasson, R.V.; Land, E.J.; Salet, C. *Photochem. Photobiol.* **27**: 273-80 (1978).
- 78E208 Laser photolysis study of the photoisomerization of retinals. Veyret, B.; Davis, S.G.; Yoshida, M.; Weiss, K. *J. Am. Chem. Soc.* **100**: 3283-90 (1978).
- 78E308 265 nm Laser flash photolysis of some ortho-substituted anilides and related N-formylkynurenine derivatives. Pileni, M.P.; Santus, R.; Land, E.J. *Photochem. Photobiol.* **27**: 671-81 (1978).
- 78E394 Determination of the extinction coefficient for triplet-triplet absorption from the kinetics of triplet-triplet energy transfer. Romashov, L.V.; Borovkova, V.A.; Kiryukhin, Yu.I.; Bagdasar'yan, Kh.S. *High Energy Chem.* **12**: 132-4 (1978) Translated from: *Khim. Vys. Energ.* **12**: 156-9 (1978).
- 78E414 Effect of the nature of the donor on the intersystem crossing in 9,10-dicyanoanthracene exciplexes in heptane. Soboleva, I.V.; Sadovalskii, N.A.; Kuz'min, M.G. *Dokl. Phys. Chem.* **238**: 70-3 (1978) Translated from: *Dokl. Akad. Nauk SSSR* **238**: 400-3 (1978).
- 78E467 Intersystem crossing efficiencies of retinal isomers in different solvents measured by laser flash absorption spectrophotometry. Bensasson, R.; Land, E.J. *Nouv. J. Chim.* **2**: 503-7 (1978).
- 78E495 Photophysical processes in fluorenone. Andrews, L.J.; Derouede, A.; Linschitz, H. *J. Phys. Chem.* **82**: 2304-9 (1978).
- 78E534 A luminescence and flash photolysis study of the photosensitizer N-methyl-2-benzoyl- β -naphthiazoline. Raemme, G. *J. Photochem.* **9**: 439-47 (1978).
- 78E538 Interband transitions in molecular crystals. Triplet-triplet absorptions in ketone crystals. Morris, J.M.; Yoshihara, K. *Mol. Phys.* **36**: 993-1003 (1978).
- 78E554 Kinetic spectroscopy of triplet states of benzenoid hydrocarbons using modulated Hg resonance radiation. Sime, M.E.; Phillips, D. *Chem. Phys. Lett.* **56**: 138-42 (1978).
- 78E721 Triplet excited states of carbonyl-containing polyenes. Becker, R.S.; Bensasson, R.V.; Lafferty, J.; Truscott, T.G.; Land, E.J. *J. Chem. Soc., Faraday Trans. 2* : 2246-55 (1978).
- 78F030 Mechanism of the photochemical cis \rightleftharpoons trans isomerization of thioindigo and 6,6'-diethoxy-thioindigo in solution. Grellmann, K.H.; Hentzschel, P. *Chem. Phys. Lett.* **53**: 545-51 (1978).
- 78Z194 Physical properties of excited states: A general method for measuring triplet-triplet extinction coefficients, singlet triplet intersystem crossing efficiencies, and related parameters. Bensasson, R.; Land, E.J. *Photochemical and Photobiological Reviews*, K.C. Smith (ed.), Plenum Press, New York, N.Y., 1978, Vol. 3, p.163-91.
- 79A028 Blitzphotolyse von Desoxybenzoinen in Loesung. Amirzadeh, G.; Kuhlmann, R.; Schnabel, W. *J. Photochem.* **10**: 133-44 (1979).
- 79A114 The triplet state of 8-methoxypsoralen. Sloper, R.W.; Truscott, T.G.; Land, E.J. *Photochem. Photobiol.* **29**: 1025-9 (1979).

- 79A147 Reversible photorearrangement of N-substituted phthalimides: a flash photolysis study. Coyle, J.D.; Harriman, A.; Newport, G.L. *J. Chem. Soc., Perkin Trans. 2* : 799-802 (1979).
- 79A171 Polymer effect on the photochemical reaction of benzophenone with tetrahydrofuran and with diphenylamine in benzene. Kamachi, M.; Kikuta, Y.; Nozakura, S. *Polym. J.* **11**: 273-7 (1979).
- 79A177 Steady-state and nanosecond spectroscopic studies of triplet sensitized reaction of K-region arene oxides. Itoh, M.; Murata, K.; Tokumura, K.; Shudo, K.; Miyata, N.; Okamoto, T. *Tetrahedron* **35**: 1059-63 (1979).
- 79A178 Hydrogen atom abstraction of p-chloranil triplet in 1,4-dioxane in the presence and absence of tetrachlorohydroquinone. Kobashi, H.; Tomioka, Y.; Morita, T. *Bull. Chem. Soc. Jpn.* **52**: 1568-72 (1979).
- 79A230 Laser flash-photolysis of methylene blue solutions. Rehak, V.; Poskocil, J. *Collect. Czech. Chem. Commun.* **44**: 2015-23 (1979).
- 79A237 The photophysics and photochemistry of pentahelicene. Grellmann, K.-H.; Hentzschel, P.; Wismontski-Knittel, T.; Fischer, E. *J. Photochem.* **11**: 197-213 (1979).
- 79A241 Photochemistry and photooxidation of tetraphenyl-p-dioxin. George, M.V.; Kumar, Ch.V.; Scaiano, J.C. *J. Phys. Chem.* **83**: 2452-5 (1979).
- 79A289 Photochemical α -cleavage and hydrogen abstraction in deoxybenzoin: a laser spectroscopy investigation. Fouassier, J.-P.; Merlin, A. *Can. J. Chem.* **57**: 2812-7 (1979).
- 79A346 Flash photolysis of tricarboyanine dyes. Shvedova, L.A.; Tatikolov, A.S.; Makin, S.M.; Romanov, N.N.; Kuz'min, V.A. *Bull. Acad. Sci. USSR, Div. Chem. Sci.* **28**: 696-701 (1979) Translated from: *Izv. Akad. Nauk SSSR, Ser. Khim.* : 749-56 (1979).
- 79A403 Mechanism for photoreduction of 2-[2'-5'-nitrofuryl-2'']-vinyl]quinoline. Luzhkov, V.B.; Khudyakov, I.V. *Bull. Acad. Sci. USSR, Div. Chem. Sci.* **28**: 608-11 (1979) Translated from: *Izv. Akad. Nauk SSSR, Ser. Khim.* : 653-6 (1979).
- 79B007 Picosecond transient spectroscopy of molecules in solution. Greene, B.I.; Hochstrasser, R.M.; Weisman, R.B. *J. Chem. Phys.* **70**: 1247-59 (1979).
- 79B032 Optical absorption spectrum of excited ruthenium tris-bipyridyl ($\text{Ru}(\text{bpy})_3^+$). Lachish, U.; Infelta, P.P.; Graetzel, M. *Chem. Phys. Lett.* **62**: 317-9 (1979).
- 79B037 Determination of the triplet and excited singlet absorption spectra of chlorophyll a with a tunable dye laser. Baugher, J.; Hindman, J.C.; Katz, J.J. *Chem. Phys. Lett.* **63**: 159-62 (1979).
- 79B042 A laser flash photolysis study of the reactivities of the triplet states of 8-methoxypsoralen and 4,5',8-trimethylpsoralen with nucleic acid bases in solution. Beaumont, P.C.; Parsons, B.J.; Phillips, G.O.; Allen, J.C. *Biochim. Biophys. Acta* **562**: 214-21 (1979).
- 79B043 The infrared and visible absorption spectra of the lowest triplet state of acridine isolated in an argon matrix. Mitchell, M.B.; Smith, G.R.; Jansen, K.; Guillory, W.A. *Chem. Phys. Lett.* **63**: 475-8 (1979).
- 79B044 Laser photolysis study of the spectral-kinetic characteristics of short-lived triplet exciplexes and the mechanism of atomic hydrogen transfer in the course of 2,6-diphenyl-1,4-benzoquinone triplet quenching with aromatic amines. Kuzmin, V.A.; Darmanyan, A.P.; Levin, P.P. *Chem. Phys. Lett.* **63**: 509-14 (1979).
- 79B050 Transannular interaction in the excited triplet states of [2.2]paracyclophane and related compounds. Ishikawa, S.; Nakamura, J.; Iwata, S.; Sumitani, M.; Nagakura, S.; Sakata, Y.; Misumi, S. *Bull. Chem. Soc. Jpn.* **52**: 1346-50 (1979).
- 79B061 Laserflash-photolysis of the p-chloranil/naphthalene system: Characterization of the naphthalene radical cation in a fluid medium. Gschwind, R.; Haselbach, E. *Helv. Chim. Acta* **62**: 941-55 (1979).
- 79B076 Picosecond relaxation processes in excited state quinoxaline. Anderson, R.W., Jr.; Damschen, D.E.; Scott, G.W.; Talley, L.D. *J. Chem. Phys.* **71**: 1134-40 (1979).
- 79B077 A study of isomeric retinal triplets by low temperature spectroscopic and kinetic flash photolysis. Grodowski, M.; Liu, R.S.H.; Herkstroeter, W.G. *Chem. Phys. Lett.* **65**: 42-5 (1979).
- 79B086 Kynurenic acid. I. Spectroscopic properties. Pileni, M.P.; Giraud, M.; Santus, R. *Photochem. Photobiol.* **30**: 251-6 (1979).
- 79B087 Triplet excited states of pyrimidine nucleosides and nucleotides. Salet, C.; Bensasson, R.; Becker, R.S. *Photochem. Photobiol.* **30**: 325-9 (1979).
- 79B098 A study on electronic excited states of iridium(III) complexes containing bipyridine and phenanthroline ligands. Solvent effect on triplet-triplet absorption spectra. Ohashi, Y.; Kobayashi, T. *Bull. Chem. Soc. Jpn.* **52**: 2214-7 (1979).
- 79B104 A previously unreported intense absorption band and the pK_a of protonated triplet methylene blue. Ohno, T.; Osif, T.L.; Lichtin, N.N. *Photochem. Photobiol.* **30**: 541-6 (1979).
- 79B114 Chemistry and photochemistry of the disodium salts of free-base porphyrin. Lee, C.Y.; Levin, G. *J. Phys. Chem.* **83**: 3165-8 (1979).
- 79B119 Photophysics of aryl substituted 1,3,4-oxadiazoles. II. SCF CI calculations in different molecular configurations and assignment of the $T_1 \rightarrow T_n$ spectra of α NPD and β NPD. Rulliere, C.; Rayez, J.-C.; Roberge, P.C. *Chem. Phys.* **44**: 337-47 (1979).
- 79B141 Determination of the photophysical constants of benzophenone by a nanosecond excitation method. Kiryukhin, Yu.I.; Sinitsyna, Z.A.; Bagdasar'yan, Kh.S. *High Energy Chem.* **13**: 432-7 (1979) Translated from: *Khim. Vys. Energ.* **13**: 515-21 (1979).
- 79B161 Absorption spectrum and quantum yield of formation of the lowest triplet excited state of tris(2,2'-bipyridine)ruthenium(II) in aqueous solution. Bensasson, R.V.; Salet, C.; Balzani, V. C. R. *Hebd. Seances Acad. Sci., Ser. B* **289**: 41-3 (1979).
- 79E099 Triplet deactivation in benzotriazole-type ultraviolet stabilizers. Werner, T. *J. Phys. Chem.* **83**: 320-5 (1979).
- 79E136 The triplet excited states of biliverdin and biliverdin dimethyl ester. Land, E.J. *Photochem. Photobiol.* **29**(3): 483-7 (1979).
- 79E175 Biradicaloid intermediates in photochemistry: Spectroscopic and kinetic study of 1,4-perinaphthadiyl and related 1,8-naphthoquinodimethanes. Gisin, M.; Rommel, E.; Wirz, J.; Burnett, M.N.; Pagni, R.M. *J. Am. Chem. Soc.* **101**: 2216-8 (1979).
- 79E219 Reactivity of acridine dye triplet states in electron transfer reactions. Vogelmann, E.; Rauscher, W.; Kramer, H.E.A. *Photochem. Photobiol.* **29**: 771-6 (1979).
- 79E243 Flash photolysis of cyanine dyes. Pinacyanol chloride (1,1'-diethyl-2,2'-carbocyanine chloride). Arvis, M.; Mialocq, J.-C. *J. Chem. Soc., Faraday Trans. 2* **75**: 415-21 (1979).
- 79E282 Triplet excited state of coumarin and 4',5'-dihydropsoralen: Reaction with nucleic acid bases and amino acids. Land, E.J.; Truscott, T.G. *Photochem. Photobiol.* **29**: 861-6 (1979).
- 79E297 Photophysics of aryl substituted 1,3,4-oxadiazoles. I. SCF CI calculations in different molecular configurations and assignment of the $T_1 \rightarrow T_n$ spectrum of 2,5-di-(1-naphthyl)-1,3,4-oxadiazole. Rulliere, C.; Rayez, J.-C.; Roberge, P.C. *Chem. Phys.* **39**: 1-12 (1979).
- 79E304 Studies on the photo-oxidation mechanism of polymers-IX. The photo-sensitized oxidation of cis-1,4-polybutadiene by N-methyl-2-benzoyl- β -naphthiazoline. Rabek, J.F.; Raemme, G.; Canbaeck, G.; Ranby, B.; Kagiya, V.T. *Eur. Polym. J.* **15**: 339-48 (1979).
- 79E325 The triplet state in trans \rightarrow cis-photoisomerization of thioindigo. Karstens, T.; Kobs, K.; Memming, R. *Ber. Bunsenges. Phys. Chem.* **83**: 504-10 (1979).
- 79E378 Upper excited triplet state mechanism in the trans \rightarrow cis photoisomerization of 4-bromostilbene. Goerner, H.; Schulte-Frohlinde, D. *J. Am. Chem. Soc.* **101**: 4388-90 (1979).
- 79E443 Picosecond spectroscopic measurement of very fast intersystem crossing for 9,10-dioxo-anti-bimanes. Huppert, D.; Dodiuk, H.; Kanety, H.; Kosower, E.M. *Chem. Phys. Lett.* **65**: 164-8 (1979).
- 79E543 Laser flash studies of thioindigo and indigo dyes. Evidence for a trans configuration of the triplet state. Goerner, H.; Schulte-Frohlinde, D. *Chem. Phys. Lett.* **66**: 363-9 (1979).

- 79E546 Triplet state photophysical properties and intersystem crossing quantum efficiencies of homologues of retinals in various solvents. Das, P.K.; Becker, R.S. *J. Am. Chem. Soc.* **101**: 6348-53 (1979).
- 79E640 Trans \rightarrow cis photoisomerization of stilbene and 4-halogenated stilbenes. Evidence for an upper excited triplet pathway. Goerner, H.; Schulte-Frohlinde, D. *J. Phys. Chem.* **83**: 3107-18 (1979).
- 79E666 Lowest triplet properties of poly(2-vinylnaphthalene) in solution. Bensasson, R.V.; Ronfard-Haret, J.C.; Land, E.J.; Webber, S.E. *Chem. Phys. Lett.* **68**: 438-42 (1979).
- 79E678 Some furocoumarins and analogs: Comparison of triplet properties in solution with photobiological activities in yeast. Sa E Melo, M.T.; Averbek, D.; Bensasson, R.V.; Land, E.J.; Salet, C. *Photochem. Photobiol.* **30**: 645-51 (1979).
- 79E690 Laser photolysis study of the exciplex between triplet benzil and triethylamine. Encinas, M.V.; Scaiano, J.C. *J. Am. Chem. Soc.* **101**: 7740-1 (1979).
- 79E799 Excited states of six-membered azaaromatic rings. Part XII. Acid-base equilibria of "double" azines. Bulska, H.; Kotlicka, J. *Pol. J. Chem.* **53**: 2103-15 (1979).
- 79E828 Etude par spectroscopie picoseconde et par photolyse éclair conventionnelle de polyméthine-cyanines. Mialocq, J.C.; Goujon, P.; Arvis, M. *J. Chim. Phys. Phys.-Chim. Biol.* **76**: 1067-75 (1979).
- 79E848 Mixed annihilation delayed fluorescence of Bengal Pink and anthracene. Ketsle, G.A.; Levshin, L.V.; Sokolova, L.K. *Opt. Spectrosc. (USSR)* **47**: 494-7 (1979) Translated from: *Opt. Spektrosk.* **47**: 893-9 (1979).
- 79E964 Systematic classification of molecules in terms of their luminescence spectroscopic properties. IV. Acridinone and its derivatives. Shcherbo, S.N.; Val'kova, G.A.; Shigorin, D.N.; Sorokina, R.S.; Rybakova, L.F. *Russ. J. Phys. Chem.* **53**: 318-320 (1979) Translated from: *Zh. Fiz. Khim.* **53**: 562-5 (1979).
- 79E965 Systematic classification of molecules in terms of luminescence spectroscopic properties. V. Linear quinacridinone and its derivatives. Shcherbo, S.N.; Val'kova, G.A.; Shigorin, D.N.; Moiseeva, Z.Z.; Bir, E.Sh. *Russ. J. Phys. Chem.* **53**: 320-2 (1979) Translated from: *Zh. Fiz. Khim.* **53**: 566-9 (1979).
- 79F576 Protonation of the ground and triplet states of rhodamine 6G in aqueous solutions. Kuznetsov, V.A.; Shamraev, V.N.; Nurmukhametov, R.N. *Opt. Spectrosc. (USSR)* **47**: 157-60 (1979) Translated from: *Opt. Spektrosk.* **47**: 279-83 (1979).
- 79N005 Light-initiated electron transfer in functional surfactant assemblies. 1. Micelles with transition metal counterions. Moroi, Y.; Braun, A.M.; Graetzel, M. *J. Am. Chem. Soc.* **101**: 567-72 (1979).
- 79N036 Absorption and emission studies of solubilization in micelles. Part 5. Pyrene-3-sodium sulphonate solubilised in dodecyltrimethylammonium bromide inverted micelles in benzene. Hunter, T.F.; Younis, A.I. *J. Chem. Soc., Faraday Trans. 1* **75**: 550-60 (1979).
- 79P066 Laser flash photolysis studies on homo- and copolymers of phenylisopropenyl ketone and of pivalophenone in solution. Naito, I.; Kuhlmann, R.; Schnabel, W. *Polymer* **20**: 165-70 (1979).
- 79R026 The effect of DNA binding on initial 8-methoxypsoralen photochemistry. Goyal, G.C.; Grossweiner, L.I. *Photochem. Photobiol.* **29**: 847-50 (1979).
- 80A030 Photodimerization of benzyl 9-anthroate in the presence of triethylamine. Role of exciplex and triplet formation. Costa, S.M. de B.; Melo, E.C.C. *J. Chem. Soc., Faraday Trans. 2* **76**: 1-13.
- 80A032 Triplet excited state of the 4'5' photoadduct of psoralen and thymine. Bensasson, R.V.; Salet, C.; Land, E.J.; Rushton, F.A.P. *Photochem. Photobiol.* **31**: 129-33 (1980).
- 80A051 Laser flash photolysis of aromatic azo-hydrazone systems. McVie, J.; Mitchell, A.D.; Sinclair, R.S.; Truscott, T.G. *J. Chem. Soc., Perkin Trans. 2*: 286-90 (1980).
- 80A074 Light induced redox reactions of water soluble porphyrins, sensitization of hydrogen generation from water by zinc-porphyrin derivatives. Kalyanasundaram, K.; Graetzel, M. *Helv. Chim. Acta* **63**: 478-85 (1980).
- 80A143 Kinetic and spectroscopic features of some carotenoid triplet states: Sensitization by singlet oxygen. Rodgers, M.A.J.; Bates, A.L. *Photochem. Photobiol.* **31**: 533-7 (1980).
- 80A196 Photochemistry of 5-methylphenazinium salts in aqueous solution. 2. Optical flash photolysis and fluorescence results and a proposed mechanism. Chew, V.S.F.; Bolton, J.R.; Brown, R.G.; Porter, G. *J. Phys. Chem.* **84**: 1909-16 (1980).
- 80A206 Dynamics of electron transfer in amine photooxidation. Peters, K.S.; Freilich, S.C.; Schaeffer, C.G. *J. Am. Chem. Soc.* **102**: 5701-2 (1980).
- 80A222 Chemistry of fluorenylidene. Direct observation of, and kinetic measurements on, a singlet and a triplet carbene at room temperature. Zupancic, J.J.; Schuster, G.B. *J. Am. Chem. Soc.* **102**: 5958-60 (1980).
- 80A235 The temperature dependence of the triplet lifetimes of some aryl iodides and bromides. Grieser, F.; Thomas, J.K. *J. Chem. Phys.* **73**: 2115-9 (1980).
- 80A338 Solvent effects in the photochemistry of xanthone. Scaiano, J.C. *J. Am. Chem. Soc.* **102**: 7747-53 (1980).
- 80A369 Heavy atom substituents as molecular probes for solvent effects on the dynamics of short-lived triplet exciplexes. Winter, G.; Steiner, U. *Ber. Bunsenges. Phys. Chem.* **84**: 1203-14 (1980).
- 80B001 Transient absorption spectra of 2-hydroxybenzophenone photostabilizers. Merritt, C.; Scott, G.W.; Gupta, A.; Yavrouian, A. *Chem. Phys. Lett.* **69**: 169-73 (1980).
- 80B010 Triplet sensitized cycloreversion of dinaphthyl dimethoxycarbonyl cyclobutane as studied by laser flash photolysis. Takamuku, S.; Schnabel, W. *Chem. Phys. Lett.* **69**: 399-402 (1980).
- 80B011 Triplet-triplet absorption of pyrazine related to the $n\pi^*(B_{3u})-n\pi^*(B_{2g})$ splitting. Inoue, A.; Webster, D.; Lim, E.C. *J. Chem. Phys.* **72**: 1419-20 (1980).
- 80B017 Triplet states of protoporphyrin IX and protoporphyrin IX dimethyl ester. Sinclair, R.S.; Tait, D.; Truscott, T.G. *J. Chem. Soc., Faraday Trans. 1* **76**: 417-25 (1980).
- 80B021 The triplet-triplet absorption of cyclohexadiene-1,3 and ergosterol. Sensitized total ground state depletion of dienes by triplet energy transfer from the ruby laser excited 2-fluorenylphenylketone. Toth, M. *Chem. Phys.* **46**: 437-43 (1980).
- 80B023 Nanosecond laser photolysis of the benzene monomer and eximer. Nakashima, N.; Sumitani, M.; Olmuci, I.; Yoshihara, K. *J. Chem. Phys.* **72**: 2226-30 (1980).
- 80B030 Laser flash photolysis investigation of primary processes in the sensitized polymerization of vinyl monomers. IV. Experiments with hydroxy alkylphenones. Eichler, J.; Herz, C.P.; Naito, I.; Schnabel, W. *J. Photochem.* **12**: 225-34 (1980).
- 80B055 Transient species in photochemistry of enones. The orthogonal triplet state revealed by laser photolysis. Bonneau, R. *J. Am. Chem. Soc.* **102**: 3816-22 (1980).
- 80B057 Protolytic reactions of acridine in the triplet state. Nishida, Y.; Kikuchi, K.; Kokubun, H. *J. Photochem.* **13**: 75-81 (1980).
- 80B066 Mechanistic studies of photochemical reactions of N-ethylphthalimide with olefins. Hayashi, H.; Nagakura, S.; Kubo, Y.; Maruyama, K. *Chem. Phys. Lett.* **72**: 291-4 (1980).
- 80B076 Photophysics and ionic photodissociation of polyesters with pendant 1-pyrenyl groups in solution. Masuhara, H.; Ohwada, S.; Seki, Y.; Mataga, N.; Sato, K.; Tazuke, S. *Photochem. Photobiol.* **32**: 9-15 (1980).
- 80B077 Intermediates in the room temperature flash photolysis and low temperature photolysis of purine solutions. Arce, R.; Jimenez, L.A.; Rivera, V.; Torres, C. *Photochem. Photobiol.* **32**: 91-5 (1980).
- 80B084 Nonradiative relaxation process of the higher excited states of meso-substituted anthracenes. Hamanoue, K.; Hirayama, S.; Nakayama, T.; Teranishi, H. *J. Phys. Chem.* **84**: 2074-8 (1980).

- 80B087 Direct observation of the 1,4-biradical of benzophenone with cis-3-methyl-2-pentene excited by the fourth harmonic of a Nd:YAG laser. Hayashi, H. *Bull. Chem. Soc. Jpn.* **53**: 2201-4 (1980).
- 80B088 Laser-photolysis study of biradical formation from the triplet state of 2,4,6-triisopropylbenzophenone. Hayashi, H.; Nagakura, S.; Ito, Y.; Umehara, Y.; Matsuura, T. *Chem. Lett.* : 939-42 (1980).
- 80B090 Extinction coefficients and quantum yields for triplet-triplet absorption using laser flash photolysis. Compton, R.H.; Grattan, K.T.V.; Morrow, T. *J. Photochem.* **14**: 61-6 (1980).
- 80B101 Laser flash photolysis study of substituted azobenzenes. Evidence for a triplet state in viscous media. Goerner, H.; Gruen, H.; Schulte-Frohlinde, D. *J. Phys. Chem.* **84**: 3031-9 (1980).
- 80B112 Assignment of transient species observed on laser flash photolysis of p-benzoquinone and methylated p-benzoquinones in aqueous solution. Ronfard-Haret, J.-C.; Bensasson, R.V.; Amouyal, E. *J. Chem. Soc., Faraday Trans. 1* **76**: 2432-6 (1980).
- 80B130 Triplet-triplet absorption of aza- and diazaphenanthrenes. Petrov, S.N.; Bocherkov, A.I.; Shatrov, V.D.; Kuznetsov, V.S.; Batekha, I.G.; El'tsov, A.V. *Zh. Prikl. Spektrosk.* **33**: 675-9 (1980).
- 80C005 Effect of pH on the behavior of duroquinone triplets. Scaiano, J.C.; Neta, P. *J. Am. Chem. Soc.* **102**: 1608-11 (1980).
- 80E023 Two-photon stepwise dissociation of carbazole in solution. Martin, M.; Breheret, E.; Tfibel, F.; Lacourbas, B. *J. Phys. Chem.* **84**: 70-2 (1980).
- 80E040 Lifetimes, spectra, and quenching of the excited states of polypyridine complexes of iron(II), ruthenium(II), and osmium(II). Creutz, C.; Chou, M.; Netzel, T.L.; Okumura, M.; Sutin, N. *J. Am. Chem. Soc.* **102**: 1309-19 (1980).
- 80E137 Visual pigments. 11. Spectroscopy and photophysics of retinoic acids and all-trans-methyl retinoate. Takemura, T.; Chihara, K.; Becker, R.S.; Das, P.K.; Hug, G.L. *J. Am. Chem. Soc.* **102**: 2604-9 (1980).
- 80E156 Photophysical properties of aminomethylpsoralen in presence and absence of DNA. Salet, C.; De Sa E Melo, T.M.; Bensasson, R.; Land, E.J. *Biochim. Biophys. Acta* **607**: 379-83 (1980).
- 80E200 Triplet states of porphyrin esters. Bonnett, R.; Charalambides, A.A.; Land, E.J.; Sinclair, R.S.; Tait, D.; Truscott, T.G. *J. Chem. Soc., Faraday Trans. 1* **76**: 852-9 (1980).
- 80E230 Spectral and electrical investigations of triplet excitons on pulsed electron excitation of molecular crystals. Kostin, A.K.; Vannikov, A.V. *Radiat. Phys. Chem.* **15**: 613-6 (1980).
- 80E234 Intersystem crossing of nitroanthracene derivatives studied by picosecond spectroscopy. Hamanoue, K.; Hirayama, S.; Nakayama, T.; Teranishi, H. *Chem. Lett.* : 407-10 (1980).
- 80E318 Excited state interaction between 2-N,N-dimethylamino-pyridine and p-nitroaniline. Wolleben, J.; Testa, A.C. *J. Photochem.* **13**: 215-21 (1980).
- 80E361 Circular dichroism of chiral molecules in the lowest excited triplet state. Tetreau, C.; Lavalette, D. *Nouv. J. Chim.* **4**: 423-5 (1980).
- 80E375 Excited state properties of 5H-dibenzo(a,d)cycloheptene. Watkins, A.R.; Bayrakceken, F. *J. Lumin.* **21**: 239-46 (1980).
- 80E405 Low-lying electronically excited states of cycl[3.3.3]azine, a bridged 12 π -perimeter. Leupin, W.; Wirz, J. *J. Am. Chem. Soc.* **102**: 6068-75 (1980).
- 80E416 Benzophenone triplet properties in acetonitrile and water. Reduction by lactams. Bensasson, R.V.; Gramain, J.-C. *J. Chem. Soc., Faraday Trans. 1* **76**: 1801-10 (1980).
- 80E438 Studies of excited state absorption of the dye p-phenylene-bis-(5-phenyl-2-oxazole) in liquid solution and in the vapour phase. Marowsky, G.; Schomburg, H. *J. Photochem.* **14**: 1-10 (1980).
- 80E439 Quenching of excited 2,5-diphenyloxazole by CCl₄. Takahashi, T.; Kikuchi, K.; Kokubun, H. *J. Photochem.* **14**: 67-76 (1980).
- 80E441 Laser photochemistry of poly(N-vinylcarbazole) in solution. Masuhara, H.; Ohwada, S.; Mataga, N.; Itaya, A.; Okamoto, K.; Kusabayashi, S. *J. Phys. Chem.* **84**: 2363-8 (1980).
- 80E593 The photophysics of bonellin: A chlorin found in marine animals. Matthews, J.I.; Braslavsky, S.E.; Camilleri, P. *Photochem. Photobiol.* **32**: 733-8 (1980).
- 80E642 On the photolysis of hydroxy alkylphenones and O-substituted derivatives. Laser flash photolysis and photocuring studies. Eichler, J.; Herz, C.P.; Schnabel, W. *Angew. Makromol. Chem.* **91**: 39-54 (1980).
- 80E720 Polymeric charge-transfer complexes based on poly(vinyl alcohol acetal)s. 2. Geissler, U.; Schulz, R.C. *Makromol. Chem.* **181**: 1483-94 (1980).
- 80E778 Kinetic investigation of triplet exciplexes of phenazine and its mono-, di-, and tetrabenzod derivatives with an electron donor. Osipov, V.V.; Usacheva, M.N.; Dilung, I.I. *Dokl. Phys. Chem.* **254**: 765-8 (1980) Translated from: *Dokl. Akad. Nauk SSSR* **254**: 407-10 (1980).
- 80E867 Migration of electron excitation energy with involvement of excited acceptor molecules. Gurinovich, G.P.; Zen'kevich, E.I.; Sagun, E.I. *Bull. Acad. Sci. USSR, Phys. Ser.* **44**: 7-13 (1980) Translated from: *Izv. Akad. Nauk SSSR, Ser. Fiz.* **44**: 693-701 (1980).
- 80E884 Nanosecond laser photolysis and the time resolved absorption spectra of organic molecules. Nakayama, T.; Miyake, T.; Okamoto, M.; Hamanoue, K.; Teranishi, H. *Mem. Fac. Ind. Arts, Kyoto Tech. Univ. Sci. Technol.* **29**: 35-45 (1980).
- 80F001 Laser-induced two-photon ionization of acridine in aqueous solution. Kellmann, A.; Tfibel, F. *Chem. Phys. Lett.* **69**: 61-5 (1980).
- 80F299 Singlet and triplet states in the cis-trans photoisomerization of 4-cyanostilbenes in solution. Goerner, H. *J. Photochem.* **13**: 269-94 (1980).
- 80F372 Photochemical transformations of cis-1,2-dibenzoylalkenes. Lahiri, S.; Dabral, V.; Chauhan, S.M.S.; Chakachery, E.; Kumar, C.V.; Scaiano, J.C.; George, M.V. *J. Org. Chem.* **45**: 3782-90 (1980).
- 80F373 Role of proflavin as a photosensitizer for the light-induced hydrogen evolution from water. Kalyanasundaram, K.; Dung, D. *J. Phys. Chem.* **84**: 2251-6 (1980).
- 80N033 Interfacial electron transfer involving radical ions of carotene and diphenylhexatriene in micelles and vesicles. Almgren, M.; Thomas, J.K. *Photochem. Photobiol.* **31**: 329-35 (1980).
- 80N035 Zinc-porphyrin sensitized reduction of simple and functional quinones in vesicle systems. Pileni, M.-P. *Chem. Phys. Lett.* **71**: 317-21 (1980).
- 80N044 Light-driven redox processes in functional micellar units. III. Zn-tetraphenylporphyrin sensitized reactions in methyl viologene surfactant assemblies. Pileni, M.-P.; Braun, A.M.; Graetzel, M. *Photochem. Photobiol.* **31**: 423-7 (1980).
- 80N057 Chlorophyll a sensitized redox processes in microemulsion systems. Kiwi, J.; Graetzel, M. *J. Phys. Chem.* **84**: 1503-7 (1980).
- 80N064 Laser photolysis studies of the light-induced formation of spiropyran-merocyanine complexes in solution. Krongauz, V.; Kiwi, J.; Graetzel, M. *J. Photochem.* **13**: 89-97 (1980).
- 80N087 Zinc porphyrin sensitized reduction of simple and functional quinones in micellar systems. Pileni, M.-P.; Graetzel, M. *J. Phys. Chem.* **84**: 1822-5 (1980).
- 80N112 Light-induced redox reactions of proflavin in aqueous and micellar solution. Pileni, M.-P.; Graetzel, M. *J. Phys. Chem.* **84**: 2402-6 (1980).
- 80N143 Zinc porphyrin sensitized redox processes in microemulsions. Pileni, M.-P. *Chem. Phys. Lett.* **75**: 540-4 (1980).
- 81A024 The flavin sensitized photooxidation of ascorbic acid: A continuous and flash photolysis study. Heelis, P.F.; Parsons, B.J.; Phillips, G.O.; McKellar, J.F. *Photochem. Photobiol.* **33**: 7-13 (1981).
- 81A140 Triplet state of 5-nitro-2-furoic acid by laser flash photolysis. Spectrum, lifetime and reactivity. Kemp, T.J.; Martins, L.J.A. *J. Chem. Soc., Faraday Trans. 1* **77**: 1425-35 (1981).

- 81A174 Laser flash photolysis study of the reactions of carbonyl triplets with phenols and photochemistry of p-hydroxypropiophenone. Das, P.K.; Encinas, M.V.; Scaiano, J.C. *J. Am. Chem. Soc.* **103**: 4154-62 (1981).
- 81A232 The photophysical and photochemical processes of tryptophan in interaction with polynucleotides: Laser flash photolysis study. Doan, T.L.; Toulme, J.J.; Santus, R.; Helene, C. *Photochem. Photobiol.* **34**: 309-13 (1981).
- 81A294 On the photoinitiation of free radical polymerization. Laser flash photolysis investigations on thioxanthone derivatives. Amirzadeh, G.; Schnabel, W. *Makromol. Chem.* **182**: 2821-35 (1981).
- 81A314 Quenching, radical formation, and disproportionation in the photoreduction of 4-carboxybenzophenone by 4-carboxybenzhydrol, hydrazine, and hydrazinium ion. Inbar, S.; Linschitz, H.; Cohen, S.G. *J. Am. Chem. Soc.* **103**: 7323-8 (1981).
- 81A345 Mechanism of photoreduction of 1- and 4-azoniaphenanthrene by dimethylformamide and phenol. Ivanov, V.L.; Al'-Ainen, S.A.; Kuz'min, M.G. *J. Org. Chem. USSR* **17**: 720-5 (1981) Translated from: *Zh. Org. Khim.* **17**: 822-7 (1981).
- 81A347 Photoinduced electron ejection from methylene blue in water and acetonitrile. Kamat, P.V.; Lichtin, N.N. *J. Phys. Chem.* **85**: 3864-8 (1981).
- 81A390 Investigation of photochemical processes involving acridone and its derivatives in ethanol solutions by flash photolysis. Shcherbo, S.N.; Val'kova, G.A.; Shigorin, D.N. *Russ. J. Phys. Chem.* **55**: 286-7 (1981) Translated from: *Zh. Fiz. Khim.* **55**: 509-10 (1981).
- 81A404 Time-resolved spectroscopy of carbonyl compounds. Lougnot, D.J.; Merlin, A.; Salvin, R.; Fouassier, J.P.; Faure, J. *Nuovo Cimento Soc. Ital. Fis. B* **63B**: 284-94 (1981).
- 81A434 Structure and properties of 5-deazaflavin radicals as compared to natural flavosemiquinones. Goldberg, M.; Pecht, I.; Kramer, H.E.A.; Traber, R.; Hemmerich, P. *Biochim. Biophys. Acta* **673**: 570-93 (1981).
- 81B008 Time-resolved resonance Raman spectroscopy: The triplet state of all-trans-retinal. Wilbrandt, R.; Jensen, N.-H. *J. Am. Chem. Soc.* **103**: 1036-41 (1981).
- 81B032 Transient photochemistry of Safranin-O. Baumgartner, C.E.; Richtol, H.H.; Aikens, D.A. *Photochem. Photobiol.* **34**: 17-22 (1981).
- 81B064 Time-resolved spectroscopy and chemical reactivity of energetic transient species of nitroaromatics. Capellos, C. *J. Photochem.* **17**: 213-25 (1981).
- 81B094 Triplet-triplet electronic absorption spectroscopy using a Fourier transform infrared spectrometer. Baiardo, J.; Mukherjee, R.; Vala, M. *Appl. Spectrosc.* **35**: 510-1 (1981).
- 81B097 Vibrational spectroscopy of the electronically excited state. 4. Nanosecond and picosecond time-resolved resonance Raman spectroscopy of carotenoid excited states. Dallinger, R.F.; Farquharson, S.; Woodruff, W.H.; Rodgers, M.A.J. *J. Am. Chem. Soc.* **103**: 7433-40 (1981).
- 81B111 Near infra-red absorption spectra of the chlorophyll a cations and triplet state in vitro and in vivo. Mathis, P.; Setif, P. *Isr. J. Chem.* **21**: 316-20 (1981).
- 81B115 Laser flash photolysis of the hematoporphyrin- β -carotene system. Poletti, A.; Murgia, S.M.; Cannistraro, S. *Photobiophys.* **2**: 167-72 (1981).
- 81B119 Nature of the triplet-triplet transitions in various many fragment molecules. Glagolev, V.L.; Pomazan, Yu.V.; Shamshev, V.N. *Opt. Spectrosc. (USSR)* **50**: 86-9 (1981) Translated from: *Opt. Spektrosk.* **50**: 165-70 (1981).
- 81E015 "Fast" triplet excimer formation in fluid solutions of dinaphthylalkanes. Webster, D.; Baugher, J.F.; Lim, B.T.; Lim, E.C. *Chem. Phys. Lett.* **77**: 294-8 (1981).
- 81E016 Enhanced phosphorescence in intramolecular exciplex systems. Hatano, Y.; Yamamoto, M.; Nishijima, Y. *Chem. Phys. Lett.* **77**: 299-303 (1981).
- 81E042 Photophysical properties of nucleic acid components. II. Effect of N,N' dimethyl substitution on intersystem crossing in pyrimidine bases. Becker, R.S.; Bensasson, R.V.; Salet, C. *Photochem. Photobiol.* **33**: 115-6 (1981).
- 81E082 Photophysics of indole. Comparative study of quenching, solvent, and temperature effects by laser flash photolysis and fluorescence. Klein, R.; Tatischeff, I.; Bazin, M.; Santus, R. *J. Phys. Chem.* **85**: 670-7 (1981).
- 81E084 Properties of excited tetrakis(sulfonatophenyl)porphyrin in aqueous solutions. Photoredox reactions with quenchers. Nahor, G.S.; Rabani, J.; Grieser, F. *J. Phys. Chem.* **85**: 697-702 (1981).
- 81E091 Triplet absorption and phosphorescence emission in zinc cytochrome c. Dixit, S.N.; Waring, A.J.; Vanderkooi, J.M. *FEBS Lett.* **125**: 86-8 (1981).
- 81E098 The electronic triplet state of a peralkylated cyclobutadiene. Wirz, J.; Krebs, A.; Schmalstieg, H.; Angliker, H. *Angew. Chem., Int. Ed. Engl.* **20**: 192-3 (1981).
- 81E147 Relaxation mechanism of excited acridine in nonreactive solvents. Kasama, K.; Kikuchi, K.; Yamamoto, S.; Uji-ie, K.; Nishida, Y.; Kokubun, H. *J. Phys. Chem.* **85**: 1291-6 (1981).
- 81E149 Reactions of the excited singlet state of pyrene with metal ions. Intermolecular electron transfer in caffeine-solubilized aqueous solution. Nosaka, Y.; Kira, A.; Imamura, M. *J. Phys. Chem.* **85**: 1353-8 (1981).
- 81E151 Photophysical and photosensitizing properties of 2-amino-4-pteridinone: A natural pigment. Chahidi, C.; Aubailly, M.; Monzikoff, A.; Bazin, M.; Santus, R. *Photochem. Photobiol.* **33**: 641-9 (1981).
- 81E214 Observation of the triplet state of stilbene in fluid solution. Determination of the equilibrium constant (${}^1t \rightleftharpoons {}^3p$) and of the rate constant for intersystem crossing (${}^3p \rightarrow {}^1p$). Goerner, H.; Schulte-Frohlinde, D. *J. Phys. Chem.* **85**: 1835-41 (1981).
- 81E222 Intersystem crossing and lowest triplet states of 4-chromanone, chromone, and flavone. Hamanoue, K.; Nakayama, T.; Miyake, T.; Teranishi, H. *Chem. Lett.* **10**: 39-42 (1981).
- 81E271 Luminescence of porphyrins and metalloporphyrins. Part 3. Heavy-atom effects. Harriman, A. *J. Chem. Soc., Faraday Trans. 2* **77**: 1281-91 (1981).
- 81E309 Picosecond triplet state dynamics and photodissociation in 2- and 9-iodoanthracene. Pineault, R.L.; Morgante, C.G.; Struve, W.S. *J. Photochem.* **17**: 435-49 (1981).
- 81E346 The triplet state energies of rubrene and diphenylisobenzofuran. Herkstroeter, W.G.; Merkel, P.B. *J. Photochem.* **16**: 331-41 (1981).
- 81E442 Photochemistry in solution. XX. Triplet reactivity of aliphatic aldehydes. Kossanyi, J.; Sabbah, S.; Chaquin, P.; Ronfart-Haret, J.C. *Tetrahedron* **37**: 3307-15 (1981).
- 81E457 Laser flash photolysis of phthalocyanines in solution and microemulsion. Jacques, P.; Braun, A.M. *Helv. Chim. Acta* **64**: 1800-6 (1981).
- 81E490 Analysis of the decay of 1-phenyl-2-(2-naphthyl)ethene triplets. A nanosecond laser pulse study. Goerner, H.; Eaker, D.W.; Saltiel, J. *J. Am. Chem. Soc.* **103**: 7164-9 (1981).
- 81E508 Quenching of the triplet state of poly(2-vinylnaphthalene) in solution: Effect of molecular weight. Pratte, J.F.; Noyes, W.A., Jr.; Webber, S.E. *Polym. Photochem.* **1**: 3-13 (1981).
- 81E552 Deactivation mechanism of excited acridine and 9-substituted acridines in water. Kasama, K.; Kikuchi, K.; Nishida, Y.; Kokubun, H. *J. Phys. Chem.* **85**: 4148-53 (1981).
- 81E639 Investigation of the mechanism of photoisomerization by the trans-thioindigo method of picosecond flash photolysis. Krysanov, S.A.; Alfimov, M.V. *Dokl. Phys. Chem.* **258**: 460-2 (1981) Translated from: *Dokl. Akad. Nauk SSSR* **258**: 665-8 (1981).
- 81E648 Spectroscopic study of the triplet states of organoclemental derivatives of fluorene. Davydov, S.N.; Rodionov, A.N.; Shigorin, D.N.; Syutkina, O.P.; Krasnova, T.L. *Russ. J. Phys. Chem.* **55**: 444-5 (1981) Translated from: *Zh. Fiz. Khim.* **55**: 784-7 (1981).

- 81E649 Determination of the energies of the highly excited triplet states of derivatives of acridinone, linear quinacridinone, and phenanthridinone. Shcherbo, S.N.; Val'kova, G.A.; Shigorin, D.N. Russ. J. Phys. Chem. **55**: 452-3 (1981) Translated from: Zh. Fiz. Khim. **55**: 795-7 (1981).
- 81E650 The influence of the relative positions of the chromophoric groups on the luminescence-spectroscopic and photochemical properties of compounds with heteroatoms. Val'kova, G.A.; Sakhno, T.V.; Shigorin, D.N.; Andrievskii, A.M.; Poplavskii, A.N. Russ. J. Phys. Chem. **55**: 459-60 (1981) Translated from: Zh. Fiz. Khim. **55**: 803-5 (1981).
- 81E715 Effect of structure of donor and acceptor on kinetic spectral characteristics of triplet exciplexes. Levin, P.P. Bull. Acad. Sci. USSR, Div. Chem. Sci. **30**: 1970-2 (1981) Translated from: Izv. Akad. Nauk SSSR, Ser. Khim. : 2390-2 (1981).
- 81E716 Laser photolysis of rubrene. Darmanyan, A.P.; Kuz'min, V.A. Dokl. Phys. Chem. **260**: 938-41 (1981) Translated from: Dokl. Akad. Nauk SSSR **260**: 1167-70 (1981).
- 81E786 A study of the electronic states of iridium(III) complexes containing 1,10-phenanthroline ligands. The contribution of a dd excited state to the lowest triplet state. Ohashi, Y. Bull. Chem. Soc. Jpn. **54**: 3673-6 (1981).
- 81F070 The mechanism of the photo-induced decarboxylation of pyruvic acid in solution. Davidson, R.S.; Goodwin, D.; Fornier de Violet, Ph. Chem. Phys. Lett. **78**: 471-4 (1981).
- 81F111 Flash photolytic studies on the photoconversion of aryl vinyl ethers to dihydrofurans. Wolff, T. J. Org. Chem. **46**: 978-83 (1981).
- 81F121 Probing of chlorophyll *a* with a pulsed tunable laser: Monomer and dimer excited state lifetimes and their time resolved fluorescence spectra. Asano, M.; Koningstein, J.A. Chem. Phys. **57**: 1-10 (1981).
- 81F130 Photochemistry of anthraquinone-2,6-disodium sulphate in aqueous solution. Harriman, A.; Mills, A. Photochem. Photobiol. **33**: 619-25 (1981).
- 81F201 Photodissociation reaction from higher excited triplet states of [2.2]paracyclophane. Ishikawa, S.; Nakamura, J.; Nagakura, S. Bull. Chem. Soc. Jpn. **54**: 685-6 (1981).
- 81F275 Observation of a triplet state of azulene generated by energy transfer in solution. Goerner, H.; Schulte-Frohlinde, D. J. Photochem. **16**: 169-77 (1981).
- 81F315 Mechanism and efficiency of photochemical attachment of aliphatic hydrocarbons to triphenylene in rigid solutions at 77 K. Lamotte, M. J. Phys. Chem. **85**: 2632-6 (1981).
- 81F390 Photochemical and thermal rearrangements of a benzoyl-naphthobarrelene-like system. Demuth, M.; Amrein, W.; Bender, C.O.; Braslavsky, S.E.; Burger, U.; George, M.V.; Lemmer, D.; Schaffner, K. Tetrahedron **37**: 3245-61 (1981).
- 81N016 Static and dynamic phenomena in the quenching of phenanthrene triplets by conjugated dienes in anionic micelles. Selwyn, J.C.; Scaiano, J.C. Can. J. Chem. **59**: 663-8 (1981).
- 81N034 Reaction photoredox de la tetraphenyl porphyrine de zinc en solution dans le triton. Pradevan, G.-O.; Pileni, M.-P. J. Chim. Phys. Phys.-Chim. Biol. **78**: 203-5 (1981).
- 81N070 Dynamics and statistics of triplet-triplet annihilation in micellar assemblies. Rothenberger, G.; Infelta, P.P.; Graetzel, M. J. Phys. Chem. **85**: 1850-6 (1981).
- 81N156 Photochemical processes of benzophenone in micro-heterogeneous systems. Braun, A.M.; Krieg, M.; Turro, N.J.; Aikawa, M.; Gould, I.R.; Graf, G.A.; Lee, P.C.-C. J. Am. Chem. Soc. **103**: 7312-6 (1981).
- 81P245 Laser photochemistry of polymers having 1,2-trans-dicarbazolylecyclobutane groups in solution. Masuhara, H.; Shioyama, H.; Mataga, N.; Inoue, T.; Kitamura, N.; Tanabe, T.; Tazuke, S. Macromolecules **14**: 1738-42 (1981).
- 82A006 The triplet state of anthracene photodimers and the wavelength dependence of the photodissociation process. Yamamoto, S.; Grellmann, K.-H. Chem. Phys. Lett. **85**: 73-80 (1982).
- 82A082 Anion-induced triplet quenching of aromatic ketones by nanosecond laser photolysis. Shizuka, H.; Obuchi, H. J. Phys. Chem. **86**: 1297-302 (1982).
- 82A138 Primary processes in the photochemistry of aqueous sulphacetamide: A laser flash photolysis and pulse radiolysis study. Land, E.J.; Navaratnam, S.; Parsons, B.J.; Phillips, G.O. Photochem. Photobiol. **35**: 637-42 (1982).
- 82A152 Pulsed irradiation of water-soluble porphyrins. Bonnet, R.; Ridge, R.J.; Land, E.J.; Sinclair, R.S.; Tait, D.; Truscott, T.G. J. Chem. Soc., Faraday Trans. 1 **78**: 127-36 (1982).
- 82A153 Triplet state of 2-nitrothiophen. A laser flash-photolysis study. Martins, L.J.A.; Kemp, T.J. J. Chem. Soc., Faraday Trans. 1 **78**: 519-31 (1982).
- 82A154 Electron-transfer reactions of the 2-nitrothiophen triplet state studied by laser flash photolysis. Martins, L.J.A. J. Chem. Soc., Faraday Trans. 1 **78**: 533-43 (1982).
- 82A190 Cis \rightleftharpoons trans photoisomerization of thioindigoid dyes. Klages, C.P.; Kobs, K.; Memming, R. Chem. Phys. Lett. **90**: 46-50 (1982).
- 82A205 Primary intermediates in the pulsed irradiation of retinoids. Lo, K.K.N.; Land, E.J.; Truscott, T.G. Photochem. Photobiol. **36**: 139-45 (1982).
- 82A259 Triplet state of 1,10-phenanthroline and benzophenone in benzene studied by pulse radiolysis. Ieply, J.; Mehnert, R.; Brede, O.; Fojtik, A. Radiochem. Radioanal. Lett. **53**: 141-51 (1982).
- 82A288 Photokinetic aspects of specific hydrogen bonding in all-trans retinal at room temperature. Das, P.K.; Hug, G.L. Photochem. Photobiol. **36**: 455-61 (1982).
- 82A290 Electron transfer in the quenching of triplet states of zinc phthalocyanine and methylene blue by the use of Fe(III), Co(III), and organic oxidants. Ohno, T.; Kato, S.; Lichtin, N.N. Bull. Chem. Soc. Jpn. **55**: 2753-9 (1982).
- 82A292 Effect of conformation on ring opening of cis and trans dimers of acenaphthylene in the triplet state. Direct detection of hydrocarbon biradical, 1,1'-biacenaphthene-2,2'-diyl. Kobashi, H.; Ikawa, H.; Kondo, R.; Morita, T. Bull. Chem. Soc. Jpn. **55**: 3013-8 (1982).
- 82A297 Sensitized photoreduction of nitrate in homogeneous and micellar solutions. Frank, A.J.; Graetzel, M. Inorg. Chem. **21**: 3834-7 (1982).
- 82A306 Kinetics and mechanism of porphyrin-photosensitized reduction of methylviologen. Rougee, M.; Ebbesen, T.; Ghetti, F.; Bensasson, R.V. J. Phys. Chem. **86**: 4404-12 (1982).
- 82A365 Tris(2,2'-bipyridine)ruthenium(II)-sensitized photooxidation of phenols. Environmental effects on electron transfer yields and kinetics. Miedlar, K.; Das, P.K. J. Am. Chem. Soc. **104**: 7462-9 (1982).
- 82B014 Substituent effect on a photo-Smiles rearrangement by laser photolysis. Yokoyama, K.; Nakamura, J.; Mutai, K.; Nagakura, S. Bull. Chem. Soc. Jpn. **55**: 317-8 (1982).
- 82B043 Deazaflavin photocatalyzed methyl viologen reduction in water. A laser flash-photolysis study. Visser, A.J.W.G.; Fendler, J.H. J. Phys. Chem. **86**: 2406-9 (1982).
- 82B045 Photochemical reactions of triplet acetone with indole, purine, and pyrimidine derivatives. Kasama, K.; Takematsu, A.; Arai, S. J. Phys. Chem. **86**: 2420-7 (1982).
- 82B052 Comment on the publication of transient spectra obtained using flash techniques. Scaiano, J.C. J. Photochem. **18**: 395-6 (1982).
- 82B057 Detection of the triplet-triplet absorption of 1,3-dienes by pulse radiolysis. Gorman, A.A.; Gould, I.R.; Hamblett, I. J. Photochem. **19**: 89-94 (1982).
- 82B094 Mimicry of carotenoid function in photosynthesis: Synthesis and photophysical properties of a carotenopyropheophorbide. Liddell, P.A.; Nemeth, G.A.; Lehman, W.R.; Joy, A.M.; Moore, A.L.; Bensasson, R.V.; Moore, T.A.; Gust, D. Photochem. Photobiol. **36**: 641-5 (1982).
- 82B102 Laser flash studies of the photochemistry of anthrone. Scaiano, J.C.; Lee, C.W.B.; Chow, Y.L.; Buono-Core, G.E. J. Photochem. **20**: 327-34 (1982).
- 82B106 Triplet-triplet transitions of the rhodamine 6G molecule. Kuznetsov, V.A.; Shamraev, V.N.; Nurmukhametov, R.N. Opt. Spectrosc. (USSR) **52**: 501-3 (1982) Translated from: Opt. Spektrosk. **52**: 838-41 (1982).

- 82B118 A flash photolysis study of hematoporphyrin IX in neutral, aqueous and ethanolic solutions. Murgia, S.M.; Pasqua, A.; Poletti, A. *Photobiophys.* **4**: 329-36 (1982).
- 82B121 Laser system and data analysis techniques for adsorbing and non-adsorbing excited transients. O'Dowd, R.F.; O'Hare, A.; Cooke, J.; Taaffe, J.K. *J. Phys. E* **15**: 736-40 (1982).
- 82C018 Excited-state redox properties of ruthenium(II) phthalocyanine from electron-transfer quenching. Prasad, D.R.; Ferraudi, G. *J. Phys. Chem.* **86**: 4037-40 (1982).
- 82E041 Laser photolysis studies of spiropyran-merocyanine aggregate formation in solution. Kalisky, Y.; Williams, D.J. *Chem. Phys. Lett.* **86**: 100-4 (1982).
- 82E129 Radiationless intermolecular energy transfer with participation of acceptor excited triplet states. Gurinovich, G.P.; Zenkevich, E.I.; Sagun, E.I. *J. Lumin.* **26**: 297-317 (1982).
- 82E133 Some properties of the triplet excited state of the photosensitizing furocoumarin: 3-carbethoxypsoralen. Ronfard-Haret, J.C.; Averbeck, D.; Bensasson, R.V.; Bisagni, E.; Land, E.J. *Photochem. Photobiol.* **35**: 479-89 (1982).
- 82E174 Photoprotection by carotenoids during photosynthesis: Motional dependence of intramolecular energy transfer. Moore, A.L.; Joy, A.; Tom, R.; Gust, D.; Moore, T.A.; Bensasson, R.V.; Land, E.J. *Science* **216**: 982-4 (1982).
- 82E181 Intersystem crossing in styrene and styrene derivatives. Bonneau, R. *J. Am. Chem. Soc.* **104**: 2921-3 (1982).
- 82E204 Triplet states of phenylethylenes in solution. Energies, lifetimes, and absorption spectra of 1,1-diphenyl-, triphenyl-, and tetraphenylethylene triplets. Goerner, H. *J. Phys. Chem.* **86**: 2028-35 (1982).
- 82E232 Electron transfer in the quenching of protonated triplet thionine and methylene blue by ground state thionine. Kamat, P.V.; Lichtin, N.N. *J. Photochem.* **18**: 197-209 (1982).
- 82E280 Electronic spectra of hexacene in solution (ground state, triplet state, dication and dianion). Angliker, H.; Rommel, E.; Wirz, J. *Chem. Phys. Lett.* **87**: 208-12 (1982).
- 82E287 Intramolecular singlet and triplet excimers with diphenanthrylpropanes. Zachariasse, K.A.; Busse, R.; Schrader, U.; Kuehnle, W. *Chem. Phys. Lett.* **89**: 303-8 (1982).
- 82E303 Intersystem crossing and predissociation of haloaromatics. Huppert, D.; Rand, S.D.; Reynolds, A.H.; Rentzepis, P.M. *J. Chem. Phys.* **77**: 1214-24 (1982).
- 82E338 Effect of substituent on the behaviour of the excited singlet and triplet states in carbonyl derivatives of anthracene of the type 9-X-CO-A. Hirayama, S. *J. Chem. Soc., Faraday Trans. 1* **78**: 2411-21 (1982).
- 82E359 Nanosecond laser flash photolysis of 1-anilinonaphthalene. Nakamura, H.; Tanaka, J.; Nakashima, N.; Yoshihara, K. *Bull. Chem. Soc. Jpn.* **55**: 1795-7 (1982).
- 82E365 Photoprocesses in diphenylpolyenes. Oxygen and heavy-atom enhancement of triplet yields. Chattopadhyay, S.K.; Das, P.K.; Hug, G.L. *J. Am. Chem. Soc.* **104**: 4507-14 (1982).
- 82E373 Absorption triplet-triplet et cinétique de la désactivation du premier état triplet de dérivés de l'amino-2-Δ²-thiazoline. Corval, A.; Jardon, P.; Gautron, R. *J. Chim. Phys. Phys.-Chim. Biol.* **79**: 335-41 (1982).
- 82E429 Singlet mechanism for trans → cis photoisomerization of α,ω-diphenylpolyenes in solution: Laser flash study of the triplet states of diphenylbutadiene, diphenylhexatriene and diphenyloctatetraene. Goerner, H. *J. Photochem.* **19**: 343-56 (1982).
- 82E451 Generation of ¹O₂ and the mechanism of internal conversion in 9,10-diphenylanthracene. Darmanyan, A.P. *Chem. Phys. Lett.* **91**: 396-400 (1982).
- 82E456 Properties of the triplet state of N,N,N',N'-tetraethyloxonine. Kamat, P.V.; Lichtin, N.N. *Isr. J. Chem.* **22**: 113-6 (1982).
- 82E474 Time-resolved resonance Raman spectra of the lowest triplet state of N,N,N',N'-tetramethyl-p-phenylenediamine in solution. Yokoyama, K. *Chem. Phys. Lett.* **92**: 93-6 (1982).
- 82E475 Solution photochemistry of o-hydroxybenzophenone at low temperatures. Scaiano, J.C. *Chem. Phys. Lett.* **92**: 97-9 (1982).
- 82E497 Picosecond laser photolysis studies of deactivation processes of excited hydrogen-bonding complexes. 2. Dibenzocarbazole-pyridine systems. Martin, M.M.; Ikeda, N.; Okada, T.; Mataga, N. *J. Phys. Chem.* **86**(21): 4148-56 (1982).
- 82E509 Studies on triplet-singlet energy transfer with 1,3-dibromo-9,10-bis-[phenylethynyl]-anthracene. Fang, T.-S.; Lin, J.; Schneider, R.; Yamada, T.; Singer, L.A. *Chem. Phys. Lett.* **92**: 283-7 (1982).
- 82E527 Reactivity and decay pathways of photoexcited anilinonaphthalenes. Grellmann, K.H.; Schmitt, U. *J. Am. Chem. Soc.* **104**: 6267-72 (1982).
- 82E561 Relaxation mechanism of excited acridine in methyl methacrylate and poly(methyl methacrylate). Kasama, K.; Kikuchi, K.; Ujiie, K.; Yamamoto, S.-A.; Kokubun, H. *J. Phys. Chem.* **86**: 4733-7 (1982).
- 82E614 Quenching of duroquinone by aromatic donors of electron and hydrogen atom. Levin, P.P.; Tatikolov, A.S.; Kuz'min, V.A. *Bull. Acad. Sci. USSR, Div. Chem. Sci.* **31**: 890-4 (1982) Translated from: *Izv. Akad. Nauk SSSR, Ser. Khim.* : 1005-10 (1982).
- 82E622 Photophysical and redox properties of water-soluble porphyrins in aqueous media. Kalyanasundaram, K.; Neumann-Spallart, M. *J. Phys. Chem.* **86**: 5163-9 (1982).
- 82E632 Triplet-triplet absorption of organic compounds lasing efficiently in the ultraviolet. Bolotko, L.M.; Gruzinskii, V.V.; Danilova, V.I.; Kopylova, T.N. *Opt. Spectrosc. (USSR)* **52**: 379-81 (1982) Translated from: *Opt. Spektrosk.* **52**: 635-8 (1982).
- 82E648 Quenching of phosphorescence of aromatic molecules through highly excited triplet states in solutions at 77°K by electron acceptors. Skvortsov, V.I.; Alfimov, M.V. *Dokl. Phys. Chem.* **263**(1-3) 221-4 (1982) Translated from: *Dokl. Akad. Nauk SSSR* **263**(3) 652-5 (1982).
- 82E680 Photochemical primary processes of xanthene dyes. I. Investigations of the primary processes of selenopyronine by flash excitation. Ortmann, W.; Kassem, A.; Hinzmann, S.; Fanghaenel, E. *J. Prakt. Chem.* **324**: 1017-25 (1982).
- 82F150 Mechanistic and kinetic aspects of the photoinduced OCH₃ substitution in 3,5-dinitroanisole. A probe for solvent effects in thermal reactions. Varma, C.A.G.O.; Tamminga, J.J.; Cornelisse, J. *J. Chem. Soc., Faraday Trans. 2* **78**: 265-84 (1982).
- 82F295 Photocatalytic cis→trans-isomerization of thioindigoid dyes. Klages, C.P.; Kobs, K.; Memming, R. *Ber. Bunsenges. Phys. Chem.* **86**: 716-20 (1982).
- 82F367 The adiabatic photodissociation of the anthracene-tetracene photodimer. Yamamoto, S.; Grellmann, K.-H. *Chem. Phys. Lett.* **92**: 533-40 (1982).
- 82F477 Fluorescence and triplet yield quenching of β-apo-14'-carotenal (C₂₂ aldehyde) by aromatic molecules. Selvarajan, N.; Das, P.K.; Hug, G.L. *J. Photochem.* **20**: 355-66 (1982).
- 82F494 Effect of solvent polarity on the photoelimination of halogen in 9,10-dichloroanthracene by the action of amines. Soloveichik, O.M.; Klimakova, A.; Ivanov, V.L. *J. Org. Chem. USSR* **18**: 894-8 (1982) Translated from: *Zh. Org. Khim.* **18**: 1033-8 (1982).
- 82F498 Study of the trans→cis photoisomerization of 4-nitro-4'-dimethylaminostilbene in toluene solutions. Goerner, H.; Schulte-Frohlinde, D. *J. Mol. Struct.* **84**: 227-36 (1982).
- 82N068 Light-driven electron transfer from tetrathiafulvalene to porphyrins and Ru(bpy)₃³⁺. Charge separation by organized assemblies. Graetzel, C.K.; Graetzel, M. *J. Phys. Chem.* **86**: 2710-4 (1982).
- 82N081 Enhancement of the intersystem crossing of pyrene by metal ions in sodium dodecyl sulfate micelle solutions. Nakamura, T.; Kira, A.; Imamura, M. *J. Phys. Chem.* **86**: 3359-63 (1982).
- 82N210 Laser flash photolysis studies on hematoporphyrin IX in aqueous and micellar systems. Murgia, S.M.; Poletti, A.; Pasqua, A. *Med., Biol., Environ.* **10**: 279-84 (1982).

- 82P060 Novel and efficient water-soluble photoinitiator for polymerization. Bonamy, A.; Fouassier, J.P.; Loughnot, D.J.; Green, P.N. *J. Polym. Sci., Polym. Lett. Ed.* **20**: 315-20 (1982).
- 82R150 Probing the microenvironment of benzo[a]pyrene diol epoxide-DNA adducts by triplet excited state quenching methods. Poulos, A.T.; Kuzmin, V.; Geacintov, N.E. *J. Biochem. Biophys. Methods* **6**: 269-81 (1982).
- 83A102 Mechanism of porphyrin ion production from the triplet state of magnesium octaethylporphyrin. Smalley, J.F.; Feldberg, S.W.; Brunschwig, B.S. *J. Phys. Chem.* **87**: 1757-65 (1983).
- 83A213 A flash photolysis study of 1-methylindole. Pepmiller, C.; Bedwell, E.; Kuntz, R.R.; Ghiron, C.A. *Photochem. Photobiol.* **38**: 273-80 (1983).
- 83A218 Laser-photolysis study of the photochemical primary processes of hydrogermanes in solution. Hayashi, H.; Mochida, K. *Chem. Phys. Lett.* **101**: 307-11 (1983).
- 83A225 On the photochemical synthesis of N,N'-dimethylindolo-[2,3-c]carbazole and the mechanism of its formation from N,N'-dimethyl-N,N'-diphenyl-1,4-phenylenediamine. Weller, H.; Grellmann, K.-H. *J. Am. Chem. Soc.* **105**: 6268-73 (1983).
- 83A249 Hydrogen-atom transfer reaction from triplet 2-naphthylammonium ion to ground-state aromatic ketones. Shizuka, H.; Fukushima, M. *Chem. Phys. Lett.* **101**: 598-602 (1983).
- 83A355 Mechanism of photoisomerization of thioindigoid dyes. Krysanov, S.A.; Alfimov, M.V. *Dokl. Phys. Chem.* **272**: 696-9 (1983) Translated from: *Dokl. Akad. Nauk SSSR* **272**: 637-40 (1983).
- 83B030 Transient absorption study of the intramolecular excited-state and ground-state proton transfer in 3-hydroxyflavone and 3-hydroxychromone. Itoh, M.; Tanimoto, Y.; Tokumura, K. *J. Am. Chem. Soc.* **105**: 3339-40 (1983).
- 83B041 Photo-oxidation of 1,4-dihydropyridines by various electron acceptors: A laser flash photolysis study. Martens, F.M.; Verhoeven, J.W.; Varma, C.A.G.O.; Bergwerf, P. *J. Photochem.* **22**: 99-113 (1983).
- 83B054 Photochemistry of 9,10-anthraquinone-2-sulfonate in solution. I. Intermediates and mechanism. Loeff, I.; Treinin, A.; Linschitz, H. *J. Phys. Chem.* **87**: 2536-44 (1983).
- 83B067 Actinometry in monochromatic flash photolysis: The extinction coefficient of triplet benzophenone and quantum yield of triplet zinc tetraphenyl porphyrin. Hurley, J.K.; Sinai, N.; Linschitz, H. *Photochem. Photobiol.* **38**: 9-14 (1983).
- 83B068 The characterisation of the triplet state of crocetin, a water soluble carotenoid, by nanosecond laser flash photolyses. Craw, M.; Lambert, C. *Photochem. Photobiol.* **38**: 241-3 (1983).
- 83B098 The electronic absorption spectrum and structure of the emitting state of the tris(2,2'-bipyridyl)ruthenium(II) complex ion. Brateman, P.S.; Harriman, A.; Heath, G.A.; Yellowlees, L.J. *J. Chem. Soc., Dalton Trans.* : 1801-3 (1983).
- 83B112 Photoionization and triplet-triplet absorption of 4-pyrrolidinopyridine. Santos, O.; Testa, A.C.; O'Sullivan, M. *J. Photochem.* **23**: 369-77 (1983).
- 83B113 Two-step laser excitation fluorescence study of the ground- and excited-state proton transfer in 3-hydroxyflavone and 3-hydroxychromone. Itoh, M.; Fujiwara, Y. *J. Phys. Chem.* **87**: 4558-60 (1983).
- 83B117 Dynamics of the spiropyran-merocyanine conversion in solution. Kalisky, Y.; Orlowski, T.E.; Williams, D.J. *J. Phys. Chem.* **87**: 5333-8 (1983).
- 83B121 The triplet extinction coefficients of some bacterial carotenoids. Cogdell, R.J.; Land, E.J.; Truscott, T.G. *Photochem. Photobiol.* **38**: 723-5 (1983).
- 83B122 The role of a photochemical reaction on the deactivation of excited 4,5',8-trimethylpsoralen in ethanol. Ishikawa, M.; Kikuchi, K.; Kokubun, H. *Photochem. Photobiol.* **38**: 727-9 (1983).
- 83B137 Absorption spectra and dynamics of the triplet state of bis[1-(1-pyrenyl)ethyl]ethers. Masuhara, H.; Tanaka, J.A.; Mataga, N.; De Schryver, F.C.; Collart, P. *Polym. J.* **15**: 915-7 (1983).
- 83E016 Phosphorescence spectra and triplet lifetimes of halogenoanthraquinones. Hamaoune, K.; Kajiwara, Y.; Miyake, T.; Nakayama, T.; Hirase, S.; Teranishi, H. *Chem. Phys. Lett.* **94**: 276-80 (1983).
- 83E018 Excited-states decay dynamics of phenyllosazone D-glucose in the liquid phase. Maciejewski, A. *Chem. Phys. Lett.* **94**: 344-9 (1983).
- 83E025 Chemistry of singlet oxygen. 40. Enhanced 1O_2 formation in cyanoaromatic-sensitized photooxidations by substrate-enhanced intersystem crossing. Manning, L.E.; Gu, C.; Foote, C.S. *J. Phys. Chem.* **87**: 40-4 (1983).
- 83E026 Specific hydrogen-bonding effects in the photophysics of β -apo-14'-carotenal. Static and dynamic aspects of fluorescence and triplet yield quenching. Das, P.K.; Hug, G.L. *J. Phys. Chem.* **87**: 49-54 (1983).
- 83E088 Electron transfer reactions of the photoexcited triplet state of chloroaluminum phthalocyanine with aromatic amines, benzoquinones, and coordination compounds of iron(II) and iron(III). Ohno, T.; Kato, S.; Yamada, A.; Tanno, T. *J. Phys. Chem.* **87**: 775-81 (1983).
- 83E216 Photophysics of aryl substituted 1,3,4-oxadiazoles: The triplet state of 2,5-diphenyl-1,3,4-oxadiazole. Rulliere, C.; Roberge, P.C. *Chem. Phys. Lett.* **97**: 247-52 (1983).
- 83E223 Sensitization and quenching processes of alkylcobalt(III) compounds. Al-Saigh, H.Y.; Kemp, T.J. *J. Chem. Soc., Perkin Trans. 2* : 615-9 (1983).
- 83E258 Lifetime, triplet-triplet absorption spectrum and relaxation energy of an acyclic conjugated triene triplet: A pulse radiolysis study. Gorman, A.A.; Hamblett, I. *Chem. Phys. Lett.* **97**: 422-6 (1983).
- 83E262 Electron-transfer quenching vs. exciplex-mediated quenching of the low-lying excited states in phthalocyanines. Prasad, D.R.; Ferraudi, G. *Inorg. Chem.* **22**: 1672-4 (1983).
- 83E278 Conformations of the triplet state of aromatic olefins. Observation of the transoid triplet states of 3,3-dimethyl-1-(2-naphthyl)-1-butene and 3,4-dihydrophenanthrene as a rigid planar triplet model. Arai, T.; Sakuragi, H.; Tokumaru, K.; Sakaguchi, Y.; Nakamura, J.; Hayashi, H. *Chem. Phys. Lett.* **98**: 40-4 (1983).
- 83E281 Triplet-related photophysics of 9,10-diphenylanthracene. A kinetic study of reversible energy transfer from anthracene triplet by nanosecond laser flash photolysis. Chattopadhyay, S.K.; Kumar, Ch.V.; Das, P.K. *Chem. Phys. Lett.* **98**: 250-4 (1983).
- 83E311 Photosensitization of pyrimidines by 2-methylnaphthoquinone in water: A laser flash photolysis study. Fisher, G.J.; Land, E.J. *Photochem. Photobiol.* **37**: 27-32 (1983).
- 83E324 Some photophysical properties of 3-carboxypsoralen, 8-methoxypsoralen and 5-methoxypsoralen triplet states. Craw, M.; Bensasson, R.V.; Ronfard-Haret, J.C.; Sa E Melo, M.T.; Truscott, T.G. *Photochem. Photobiol.* **37**: 611-5 (1983).
- 83E347 Effect of a polar substituent on olefin triplet lifetime. Caldwell, R.A.; Singh, M. *J. Am. Chem. Soc.* **105**: 5139-40 (1983).
- 83E383 Circular dichroism in excited triplet states. 2. Chiral dimerlike molecules in the binaphthyl and spirobifluorene series. Tetreau, C.; Lavalette, D.; Cabaret, D.; Geraghty, N.; Welvert, Z. *J. Phys. Chem.* **87**: 3234-9 (1983).
- 83E387 A laser flash photolysis study of pyrene-1-aldehyde. Intersystem crossing efficiency, photoreactivity and triplet state properties in various solvents. Kumar, C.V.; Chattopadhyay, S.K.; Das, P.K. *Photochem. Photobiol.* **38**: 141-52 (1983).
- 83E392 The $T_3(\pi, \pi^*)$ state of acridine. Periasamy, N. *Chem. Phys. Lett.* **99**: 322-5 (1983).
- 83E427 The fluorescence of some dipolar N,N-dialkyl-4-(dichloro-1,3,5-triazinyl)anilines. Part 3. Intersystem crossing yields from intramolecular charge-transfer excited states and triplet state properties. Cowley, D.J.; Pasha, I. *J. Chem. Soc., Perkin Trans. 2* : 1139-45 (1983).

- 83E462 Photochemistry and sensitized evolution of hydrogen from water using water-soluble cationic porphyrins. Tetrakis-(trimethylaminophenyl)porphyrinatozinc and its free base. Kalyanasundaram, K. J. Chem. Soc., Faraday Trans. 2 **79**: 1365-74 (1983).
- 83E483 Transient absorption spectra of the excited states of triptycene and 3-acetyltriptycene. Sugawara, T.; Iwamura, H.; Nakashima, N.; Yoshihara, K.; Hayashi, H. Chem. Phys. Lett. **101**: 303-6 (1983).
- 83E509 Trans→cis photoisomerization of the quaternary iodides of 4-cyano- and 4-nitro-4'-azastilbene in ethanol solution: Singlet versus triplet mechanism. Goerner, H.; Schulte-Frohlinde, D. Chem. Phys. Lett. **101**: 79-85 (1983).
- 83E554 Excited and ionic states of polymers with pendant phenanthryl groups in solution. Model systems for photophysics in phenanthrene aggregates. Tamai, N.; Masuhara, H.; Mataga, N. J. Phys. Chem. **87**: 4461-7 (1983).
- 83E625 Studies on the triplet excited state of 4-thiouridine. Salet, C.; Bensasson, R.V.; Favre, A. Photochem. Photobiol. **38**: 521-5 (1983).
- 83E662 Absorption spectra and dynamics of some excited and ionic dicarbazolyl compounds with specific geometrical structures. Masuhara, H.; Tamai, N.; Mataga, N.; De Schryver, F.C.; Vandendriessche, J. J. Am. Chem. Soc. **105**: 7256-62 (1983).
- 83E667 Flash photolysis studies of hemato- and copro-porphyrins in homogeneous and microheterogeneous aqueous dispersions. Reddi, E.; Jori, G.; Rodgers, M.A.J.; Spikes, J.D. Photochem. Photobiol. **38**: 639-45 (1983).
- 83E737 Study of triplet states of some ketocyanine dyes, the polyenic bis- ω,ω' -dimethylamino ketones. Shvedova, L.A.; Borisevich, Yu.E.; Tatikolov, A.S.; Kuz'min, V.A.; Krasnaya, Zh.A. Bull. Acad. Sci. USSR, Div. Chem. Sci. **32**: 1290-3 (1983) Translated from: Izv. Akad. Nauk SSSR, Ser. Khim.: 1421-5 (1983).
- 83F075 Solvent effect on $^1\text{O}_2$ yield and the mechanism of polycyclic hydrocarbon triplet-state quenching by oxygen. Darmanyan, A.P. Chem. Phys. Lett. **96**: 383-9 (1983).
- 83F081 Primary processes in the photoisomerization of a nitrochromene studied by nanosecond laser photolysis. Kellmann, A.; Lindqvist, L.; Monti, S.; Tfibel, F.; Guglielmetti, R. J. Photochem. **21**: 223-35 (1983).
- 83F096 Studies of the photoreactions of heterocyclic N-dioxides: Identification of the oxaziridine intermediate of quinoxaline-1,4-dioxide. Kawata, H.; Kikuchi, K.; Kokubun, H. J. Photochem. **21**: 343-50 (1983).
- 83F334 Kinetics of photochemical H-abstraction by $^3\pi\pi^*$ -acridine in fluorene single crystals as studied by triplet-triplet absorption. Prass, B.; Fujara, F.; Stehlik, D. Chem. Phys. **81**: 175-84 (1983).
- 83F405 Sequential biphotonic processes in rhodium(III) phthalocyanines. Muralidharan, S.; Ferraudi, G. J. Phys. Chem. **87**: 4877-81 (1983).
- 83N056 Magnetic field effect on the photoinduced electron transfer reaction between duroquinone and aromatic amines in SDS micellar solution. Tanimoto, Y.; Shimizu, K.; Udagawa, H.; Itoh, M. Chem. Lett. **12**: 353-6 (1983).
- 83N062 Laser photolysis of methylene blue in aqueous micellar systems. Murgia, S.M.; Poletti, A. Photobiophys. **5**: 53-60 (1983).
- 83N073 Comparison of the photochemical processes of magnesium tetraphenylporphyrin in micellar and in microemulsion solution. Pileni, M.P.; Chevalier, S. J. Colloid Interface Sci. **92**: 326-31 (1983).
- 83N082 Thermo- and photochemical aspects of the valence isomerization of 9-tert-butylanthracene. Kraljic, I.; Mintas, M.; Klasinc, L.; Ranogajec, F.; Guesten, H. Nouv. J. Chim. **7**: 239-43 (1983).
- 83N127 Magnetic field effect on the hydrogen abstraction of xanthone from xanthene in SDS micelles. Tanimoto, Y.; Takashima, M.; Itoh, M. Chem. Phys. Lett. **100**: 442-4 (1983).
- 83N133 Magnetic field effects on the photolysis of p-benzoquinone derivatives in sodium dodecyl sulfate micelles. Tanimoto, Y.; Udagawa, H.; Katsuda, Y.; Itoh, M. J. Phys. Chem. **87**: 3976-82 (1983).
- 83P212 Triplet state of 4-biphenylcarbonyl and 2-naphthylcarbonyl chromophores in homopolymers, copolymers and model compounds. Hrdlovic, P.; Guyot, G.; Lemaire, J.; Lukac, I. Polym. Photochem. **3**: 119-29 (1983).
- 83Z042 Applications of holography in the investigation of photochemical reactions. Burland, D.M. Acc. Chem. Res. **16**: 218-24 (1983).
- 84A024 Photochemistry of benzothiazole models of pheomelanin. Lambert, C.; Sinclair, R.S.; Truscott, T.G.; Land, E.J.; Chedekel, M.R.; Liu, C.-T. Photochem. Photobiol. **39**: 5-10 (1984).
- 84A030 Photochemistry of alkyl esters of benzoylformic acid. Encinas, M.V.; Lissi, E.A.; Zanoeco, A.; Stewart, L.C.; Scaiano, J.C. Can. J. Chem. **62**: 386-91 (1984).
- 84A122 Electron-transfer reactions of excited phthalocyanines: Spin restriction on reaction rate of electron transfer and energy transfer to cobalt compounds. Ohno, T.; Kato, S. J. Phys. Chem. **88**: 1670-4 (1984).
- 84A218 Transient absorption and two-step laser excitation fluorescence studies of the excited-state proton transfer and relaxation in the methanol solution of 7-hydroxyflavone. Itoh, M.; Adachi, T. J. Am. Chem. Soc. **106**: 4320-4 (1984).
- 84A221 Aromatic thioketone triplets and their quenching behaviour towards oxygen and di-*t*-butylnitroxyl radical. A laser-flash-photolysis study. Kumar, C.V.; Qin, L.; Das, P.K. J. Chem. Soc., Faraday Trans. 2 **80**: 783-93 (1984).
- 84A222 Photochemistry of naphthylmethyl halides. Direct and sensitized paths to homolytic and heterolytic carbon-halogen bond cleavage. Slocum, G.H.; Schuster, G.B. J. Org. Chem. **49**: 2177-85 (1984).
- 84A227 A laser flash photolysis and pulse radiolysis study of amiloride in aqueous and alcoholic solution. Hamoudi, H.I.; Heelis, P.F.; Jones, R.A.; Navaratnam, S.; Parsons, B.J.; Phillips, G.O.; Vandenburg, M.J.; Currie, W.J.C. Photochem. Photobiol. **40**: 35-9 (1984).
- 84A253 Primary processes in the photoreduction of 1-chloro- and 1,8-dichloro-anthraquinones in ethanol studied by laser spectroscopy. Hamaoué, K.; Yokoyama, K.; Kajiwara, Y.; Nakajima, K.; Nakayama, T.; Teranishi, H. Chem. Phys. Lett. **110**: 25-8 (1984).
- 84A263 The triplet state of N-(n-butyl)-5-nitro-2-furamide by laser flash photolysis. Spectrum, lifetime, energy and electron-transfer reactions. Martins, L.J.A.; Kemp, T.J. J. Chem. Soc., Faraday Trans. 1 **80**: 2509-24 (1984).
- 84A264 Electron-transfer reactions of a photosensitized water-soluble zinc porphyrin. Le Roux, D.; Mialocq, J.-C.; Anitoff, O.; Folcher, G. J. Chem. Soc., Faraday Trans. 2 **80**: 909-20 (1984).
- 84A344 Laser flash photolysis study of photochemical ring opening of 2,3-di-2-naphthylloxiranes and resultant ylide behaviors. Das, P.K.; Griffin, G.W. J. Org. Chem. **49**: 3452-7 (1984).
- 84A355 Photochemical transformations and laser flash photolysis studies of 1,4- and 1,2-epoxy compounds containing 1,2-dibenzoylalkene moieties. Murty, B.A.R.C.; Kumar, C.V.; Dabral, V.; Das, P.K.; George, M.V. J. Org. Chem. **49**: 4165-71 (1984).
- 84B003 Photoenolization of 2-methylbenzophenone studied by picosecond and nanosecond laser spectroscopy. Nakayama, T.; Hamaoué, K.; Hidaka, T.; Okamoto, M.; Teranishi, H. J. Photochem. **24**: 71-8 (1984).
- 84B007 Triplet state of α -naphthylethylene derivatives. Structural effects on the equilibrium between planar and perpendicular triplet conformations. Lazare, S.; Bonneau, R.; Lapouyade, R. J. Phys. Chem. **88**: 18-23 (1984).
- 84B008 Photoexcited triplet state of oxotitanium(IV) tetraphenylporphyrin studied by ESR and laser photolysis. Proton-induced quenching of the triplet state. Hoshino, M.; Imamura, M.; Watanabe, S.; Hama, Y. J. Phys. Chem. **88**: 45-9 (1984).

- 84B017 Photoinduced electron transfer C-C bond cleavage reactions: oxidations and isomerizations. Reichel, L.W.; Griffin, G.W.; Muller, A.J.; Das, P.K.; Ege, S.N. *Can. J. Chem.* **62**: 424-36 (1984).
- 84B019 Laser photolysis studies of the anthracene dimer produced by photolytic dissociation of dianthracene in 2-methyltetrahydrofuran solution at 77 K. Hoshino, M.; Seki, H.; Imamura, M.; Yamamoto, S. *Chem. Phys. Lett.* **104**: 369-72 (1984).
- 84B022 Mechanistic studies of the photodecomposition of arylmethyl sulfones in homogeneous and micellar solutions. Gould, I.R.; Tung, C.; Turro, N.J.; Givens, R.S.; Matuszewski, B. *J. Am. Chem. Soc.* **106**: 1789-93 (1984).
- 84B025 Triplet, radical anion and radical cation spectra of furcoumarins. Bensasson, R.V.; Chalvet, O.; Land, E.J.; Ronfard-Haret, J.C. *Photochem. Photobiol.* **39**: 287-91 (1984).
- 84B026 Twenty-two carbon homologue of 11-cis retinal. Photophysical and photochemical properties. Das, P.K.; Becker, R.S. *Photochem. Photobiol.* **39**: 313-8 (1984).
- 84B027 Transient photochemistry of neutral red. Marks, G.T.; Lee, E.D.; Aikens, D.A.; Richtol, H.H. *Photochem. Photobiol.* **39**: 323-8 (1984).
- 84B028 Excited state absorption spectra and intersystem crossing kinetics in the naphthaldehydes. Boldridge, D.W.; Justus, B.L.; Scott, G.W. *J. Chem. Phys.* **80**: 3179-84 (1984).
- 84B033 A laser flash photolysis study of paraquat reduction by photo-generated aromatic ketyl radicals and carbonyl triplets. Baral-Tosh, S.; Chattopadhyay, S.K.; Das, P.K. *J. Phys. Chem.* **88**: 1404-8 (1984).
- 84B051 The laser-flash photolysis of propanone, butanone, and 2-pentanone. Study of triplet decay processes in aliphatic carbonyl compounds. Naito, I.; Schnabel, W. *Bull. Chem. Soc. Jpn.* **57**: 771-5 (1984).
- 84B055 Excited-state absorption of tris(phenanthroline)rhodium(III). A handle on the excited-state behavior of a powerful photochemical oxidant. Indelli, M.T.; Carioli, A.; Scandola, F. *J. Phys. Chem.* **88**: 2685-6 (1984).
- 84B061 The photoionization mechanism of N,N,N',N'-tetramethyl-p-phenylenediamine in acetonitrile. Nakamura, S.; Kanamaru, N.; Nohara, S.; Nakamura, H.; Saito, Y.; Tanaka, J.; Sumitani, M.; Nakashima, N.; Yoshihara, K. *Bull. Chem. Soc. Jpn.* **57**: 145-50 (1984).
- 84B066 Photoinduced electron-transfer processes involving substituted stilbene oxides. Das, P.K.; Muller, A.J.; Griffin, G.W. *J. Org. Chem.* **49**: 1977-85 (1984).
- 84B090 β -Naphthylethylene: A conformational study of the triplet state by laser flash photolysis. Lazare, S.; Lapouyade, R.; Robert, M.-P. *Nouv. J. Chim.* **8**: 407-9 (1984).
- 84B110 Photochemical debromination of meso-substituted bromoanthracenes studied by steady-state photolysis and laser photolysis. Hamanoue, K.; Tai, S.; Hidaka, T.; Nakayama, T.; Kimoto, M.; Teranishi, H. *J. Phys. Chem.* **88**: 4380-4 (1984).
- 84B116 Photoreduction of triplet thioxanthone by amines: Charge transfer generates radicals that initiate polymerization of olefins. Yates, S.F.; Schuster, G.B. *J. Org. Chem.* **49**: 3349-56 (1984).
- 84B120 Diffuse reflectance pulse radiolysis of opaque samples. Wilkinson, F.; Willsher, C.J.; Warwick, P.; Land, E.J.; Rushton, F.A.P. *Nature* **311**: 40-1 (1984).
- 84B128 Photochemical reactions of 4-nitroquinoline 1-oxide with DNA and related compounds. Kasama, K.; Takematsu, A.; Yamamoto, S.; Arai, S. *J. Phys. Chem.* **88**: 4918-21 (1984).
- 84B139 Time-resolved absorption and resonance Raman spectra of the lowest excited triplet state of all-trans-1,3,5-heptatriene. Langkilde, F.W.; Wilbrandt, R.; Jensen, N.-H. *Chem. Phys. Lett.* **111**: 372-8 (1984).
- 84B154 Time-dependent picosecond transient absorption spectra of 9-acetylanthracene, benzophenone and acridine in solution. Hamanoue, K.; Nakajima, K.; Hidaka, T.; Nakayama, T.; Teranishi, H. *Laser Chem.* **4**: 287-95 (1984).
- 84E018 Intramolecular quenching of carbonyl triplets by β -phenyl rings. Wismontski-Knittel, T.; Kilp, T. *J. Phys. Chem.* **88**: 110-5 (1984).
- 84E033 Picosecond and nanosecond kinetic spectroscopic investigations of the relaxation and the solute-solvent reaction of electronically excited 3,5-dinitroanisole. Varma, C.A.G.O.; Plantenga, F.L.; Huizer, A.H.; Zwart, J.P.; Bergwerf, Ph.; van der Ploeg, P.M. *J. Photochem.* **24**: 133-99 (1984).
- 84E036 Triplet states of isomers of the C₁₅-aldehyde and the C₁₈-ketone: Lower homologues of retinal. Bensasson, R.V.; Land, E.J.; Liu, R.S.H.; Lo, K.K.N.; Truscott, T.G. *Photochem. Photobiol.* **39**: 263-5 (1984).
- 84E043 Nanosecond time-resolved T_n \rightarrow T₁ fluorescence, T_n \leftarrow T₁ absorption, and resonance Raman scattering spectra in diphenylamine. Kobayashi, T.; Koshihara, S. *Chem. Phys. Lett.* **104**: 174-8 (1984).
- 84E044 Detection of triplet-triplet absorption in microcrystalline benzophenone by diffuse-reflectance laser flash photolysis. Wilkinson, F.; Willsher, C.J. *Chem. Phys. Lett.* **104**: 272-6 (1984).
- 84E045 Luminescence and triplet-triplet absorption spectra of rhodium(III) porphyrins. Kalyanasundaram, K. *Chem. Phys. Lett.* **104**: 357-62 (1984).
- 84E056 Laser induced excited state properties of rubreneperoxide in solution. Bayrakceken, F. *J. Lumin.* **29**: 111-8 (1984).
- 84E077 Time-resolved spectroscopy of intramolecular energy transfer in a rigid spiran. Maki, A.H.; Weers, J.G.; Hilinski, E.F.; Milton, S.V.; Rentzepis, P.M. *J. Chem. Phys.* **80**: 2288-97 (1984).
- 84E092 Polymeric photosensitizers: Effects of intramolecular energy migration on sensitization efficiencies. Urruti, E.H.; Kilp, T. *Macromolecules* **17**: 50-4 (1984).
- 84E111 Laser flash photolysis study of (2-anthryl)ethylenes. Triplet-related photophysical behaviors. Wismontski-Knittel, T.; Das, P.K. *J. Phys. Chem.* **88**: 1168-73 (1984).
- 84E144 Short-lived olefin triplets and energy transfer from them to β -carotene. Kumar, C.V.; Chattopadhyay, S.K.; Das, P.K. *Chem. Phys. Lett.* **106**: 431-6 (1984).
- 84E162 Photochemistry and photophysics of osazones. Part 3. Properties of D-arabino-hexulose phenylosazone in excited singlet and triplet states. Maciejewski, A.; Wojtczak, J.; Szymanski, M. *J. Chem. Soc., Faraday Trans. 2* **80**: 411-23 (1984).
- 84E203 Anomalous ortho effects in sterically hindered porphyrins: Tetrakis(2,6-dimethylphenyl)porphyrin and its sulfonato derivative. Lee, W.A.; Graetzel, M.; Kalyanasundaram, K. *Chem. Phys. Lett.* **107**: 308-13 (1984).
- 84E216 Photosensitization of phenylhydrazones derivatives by methylene blue and xanthene dyes. Grajcar, L.; Ivanoff, N.; Delouis, J.F.; Faure, J. *J. Chim. Phys. Phys.-Chim. Biol.* **81**: 33-8 (1984).
- 84E232 Electronic relaxation processes of rare-earth chelates of benzoyltrifluoroacetone. Tobita, S.; Arakawa, M.; Tanaka, I. *J. Phys. Chem.* **88**: 2697-702 (1984).
- 84E236 Photophysical behavior of exciplexes of 1,4-dicyanonaphthalene with methyl- and methoxy-substituted benzenes. Davis, H.F.; Chattopadhyay, S.K.; Das, P.K. *J. Phys. Chem.* **88**: 2798-803 (1984).
- 84E237 A laser flash photolysis study of triplets of trans-naphthylethylenes. Relative importance of planar and perpendicular forms. Wismontski-Knittel, T.; Das, P.K. *J. Phys. Chem.* **88**: 2803-8 (1984).
- 84E319 Role of geometric distortion in the quenching behaviors of all-trans 1,4-diphenyl-1,3-butadiene triplet. Chattopadhyay, S.K.; Kumar, C.V.; Das, P.K. *J. Photochem.* **26**: 39-47 (1984).
- 84E342 Photophysical behaviors of N-methylthioacridone triplet and efficiency of singlet oxygen generation from its quenching by oxygen. Kumar, C.V.; Davis, H.F.; Das, P.K. *Chem. Phys. Lett.* **109**: 184-9 (1984).
- 84E346 Photochemistry of water-soluble porphyrins: Comparative study of isomeric tetrapyrrolyl- and tetrakis(N-methylpyridinium)porphyrins. Kalyanasundaram, K. *Inorg. Chem.* **23**: 2453-9 (1984).

- 84E390 Excited singlet- and triplet-absorptions of pentaphene. Menzel, R.; Rapp, W. *Chem. Phys.* **89**: 445-55 (1984).
- 84E393 Experimental study of singlet-triplet energy transfer in liquid solutions. Darmanyan, A.P. *Chem. Phys. Lett.* **110**: 89-94 (1984).
- 84E405 Effect of environment on decay pathways of the singlet excited state of N,N,N',N'-tetramethylbenzidine. Hashimoto, S.; Thomas, J.K. *J. Phys. Chem.* **88**: 4044-9 (1984).
- 84E456 Acid-base properties in the triplet state of aromatic ketones studied by nanosecond laser flash photolysis. Shizuka, H.; Kimura, E. *Can. J. Chem.* **62**: 2041-6 (1984).
- 84E491 A pulse radiolysis study of the azulene triplet state. Gorman, A.A.; Hamblett, I.; Harrison, R.J. *J. Am. Chem. Soc.* **106**: 6952-5 (1984).
- 84E520 Photochemical and thermal transformations of 2(3H)-furanones and bis(benzofuranones). A laser flash photolysis study. Lohray, B.B.; Kumar, C.V.; Das, P.K.; George, M.V. *J. Am. Chem. Soc.* **106**: 7352-9 (1984).
- 84E529 Photophysics of chlorobenzene in cyclohexane. Previtali, C.M.; Ebbesen, T.W. *J. Photochem.* **27**: 9-15 (1984).
- 84E533 Fluorescence quenching mechanism of aromatic hydrocarbons by closed-shell heavy metal ions in aqueous and organic solutions. Masuhara, H.; Shioyama, H.; Saito, T.; Hamada, K.; Yasoshima, S.; Mataga, N. *J. Phys. Chem.* **88**: 5868-73 (1984).
- 84F039 Hydrogen atom transfer reaction from excited carbazole to pyridine. Kikuchi, K.; Yamamoto, S.; Kokubun, H. *J. Photochem.* **24**: 271-83 (1984).
- 84F068 Photoisomerization pathways of 8,16-methano[2.2]metacyclophane-1,9-diene. A model case for adiabatic electrocyclic ring closure in the excited singlet state. Wirz, J.; Persy, G.; Rommel, E.; Murata, I.; Nakasuji, K. *Helv. Chim. Acta* **67**: 305-17 (1984).
- 84F074 Photosensitization via charge transfer or reversible electron transfer. Oxirane isomerization and sulfur dioxide extrusion. Das, P.K.; Muller, A.J.; Griffin, G.W.; Gould, I.R.; Tung, C.-H.; Turro, N.J. *Photochem. Photobiol.* **39**: 281-5 (1984).
- 84F248 Primary photochemical and photophysical processes in 2,2,4-trimethyl-1,2-dihydroquinolines. Malkin, Ya.N.; Pirogov, N.O.; Kuzmin, V.A. *J. Photochem.* **26**: 193-202 (1984).
- 84F379 Quenching of triplet states of quinones by phenols. Kokrashvili, T.A.; Levin, P.P.; Kuz'min, V.A. *Bull. Acad. Sci. USSR, Div. Chem. Sci.* **33**: 702-8 (1984) Translated from: *Izv. Akad. Nauk SSSR, Ser. Khim.* : 765-71 (1984).
- 84F385 Photochemical reactions of nitroanthracene derivatives in fluid solutions. Hamanoue, K.; Amano, M.; Kimoto, M.; Kajiwara, Y.; Nakayama, T.; Teranishi, H. *J. Am. Chem. Soc.* **106**: 5993-7 (1984).
- 84N017 Triplet state properties of croconate dyes in homogeneous and polymer-containing solutions. Kamat, P.V.; Fox, M.A. *J. Photochem.* **24**: 285-92 (1984).
- 84N125 A water-soluble benzophenone in reverse micelles: Kinetics and spectroscopy. Lougnot, D.-J.; Scaiano, J.C. *J. Photochem.* **26**: 119-30 (1984).
- 84N163 Exploratory study of the effect of polyelectrolyte-surfactant aggregates on photochemical behavior. Abuin, E.B.; Scaiano, J.C. *J. Am. Chem. Soc.* **106**: 6274-83 (1984).
- 84N197 Magnetic field effect on the hydrogen abstraction reaction of xanthone in sodium dodecyl sulfate micellar solution. Tanimoto, Y.; Takashima, M.; Itoh, M. *J. Phys. Chem.* **88**: 6053-6 (1984).
- 84P005 Photophysical study on a new class of benzophenone containing photoinitiators of polymerization. Properties of poly(4-acryloxybenzophenone) and a model compound. Flamigni, L.; Barigelletti, F.; Bortolus, P.; Carlini, C. *Eur. Polym. J.* **20**: 171-5 (1984).
- 84P257 Excited-state annihilation processes in poly(vinylaromatics) in solution: Poly(2-vinylnaphthalene) and poly(4-vinylbiphenyl). Pratte, J.F.; Webber, S.E. *Macromolecules* **17**: 2116-23 (1984).
- 84R057 The triplet exciton of chlorophyll a and carotenoid in solution and in photosynthetic antenna proteins. Shepanski, J.F.; Williams, D.J.; Kalisky, Y. *Biochim. Biophys. Acta.* **766**: 116-25 (1984).

10. Indexes to Table 6

10.1. Compound Name Index

- 1(2*H*)-Aceanthrylenone, 2-methyl- 690.
 Acenaphthene 1.
 Acenaphthene, 5-nitro- 811.
 Acenaphthene, (1-phenylvinyl)- 4.
 Acenaphthylene, *cis*-photodimer 3.
 Acenaphthylene, *trans*-photodimer 3.
 1-Acenaphthyl-1-phenylethylene 4.
 Acetamide, *N*-[2-[(2-bromo-6-cyano-4-nitrophenyl)azo]-5-[(2-cyanoethyl)(2-hydroxyethyl)amino]-4-methoxyphenyl]- 218.
 Acetamide, *N*-[2-[(2-bromo-4,6-dinitrophenyl)azo]-5-[(2-cyanoethyl)(2-hydroxyethyl)amino]-4-methoxyphenyl]- 219.
 Acetamide, *N*-[2-[(2-chloro-4,6-dinitrophenyl)azo]-5-[(2-cyanoethyl)(2-hydroxyethyl)amino]-4-methoxyphenyl]- 270.
 Acetamide, *N*-[5-[(2-cyanoethyl)(2-hydroxyethyl)amino]-2-[(2,4-dinitrophenyl)azo]-4-methoxyphenyl]- 312.
 Acetic acid, 1-naphthyl ester 791.
 1'-Acetonaphthone 5.
 2'-Acetonaphthone 6.
 Acetone 7.
 Acetone, 1-benzoyl- 146.
 Acetone, 1-benzoyl-3,3,3-trifluoro- 158.
 Acetophenone 8.
 Acetophenone, 2-(*N*-acetylamino)- 16.
 Acetophenone, 2-(*N*-acetyl-*N*-methylamino)- 18.
 Acetophenone, conjugate acid 9.
 Acetophenone, 2-(*N*-formylamino)- 17.
 Acetophenone, 2-(*N*-formyl-*N*-methylamino)- 19.
 Acetophenone, α -hydroxy- α -phenyl- 124.
 Acetophenone, 3-methoxy- 661.
 Acetophenone, 4-methoxy- 662.
 Acetophenone, 3-methyl- 691.
 Acetophenone, 4-methyl- 692.
 Acetophenone, α -phenyl- 332.
 Acetophenone, 4-(trifluoromethyl)- 1081.
 2-Acetoxy 2-methyl-1-phenyl-1-propanone 10.
 Acetylacetone 11.
 Acetylacetone, 1,1,1,5,5,5-hexafluoro- 607.
 (*E*)-1-Acetyl-2(1-acetyl-1,3-dihydro-3-oxo-2*H*-indol-2-ylidene)-1,2-dihydro-3*H*-indol-3-one 334.
 2-(*N*-Acetylamino)acetophenone 16.
 1-Acetylanthracene 12.
 9-Acetylanthracene 13.
 4-Acetylbiphenyl 14.
 Acetylene, diphenyl- 527.
 Acetylene, 1-ethyl-2-phenyl- 237.
 Acetylene, 1-methyl-2-phenyl- 928.
 Acetylene, phenyl- 873.
 3-Acetyl-9,10-epoxy-9,10-dihydrophenanthrene 15.
 2-(*N*-Acetyl-*N*-methylamino)acetophenone 18.
N-(2-Acetylphenyl)acetamide 16.
N-(2-Acetylphenyl)formamide 17.
N-(2-Acetylphenyl)-*N*-methylacetamide 18.
N-(2-Acetylphenyl)-*N*-methylformamide 19.
N'-Acetylsulfanilamide 1009.
 3-Acetyltriptycene 20.
N-Acetyl-L-tryptophanamide 21.
 Acid Red 94 988.
 Acridan 22.
 Acridine 23.
 Acridine, 9-amino- 33.
 Acridine, 3,6-bis(dimethylamino)-, conjugate monoacid 25.
 Acridine, 3,6-bis(dimethylamino)- 26.
 Acridine, 3,6-bis(methylamino)-, conjugate monoacid 94.
 Acridine, conjugate acid 28.
 Acridine, 3,6-diamino-, conjugate diacid 922.
 Acridine, 3,6-diamino-, conjugate monoacid 923.
 Acridine, 3,6-diamino-, hydrochloride 921.
 Acridine, 3,6-diamino-2,7-dimethyl-, conjugate monoacid 27.
 Acridine, 3,6-diamino-2,7-dimethyl-9-phenyl-, monohydrochloride 119.
 Acridine, 3,6-diamino-9-phenyl-, conjugate monoacid 898.
 Acridine, 3,6-diamino-2,7,9-trimethyl-, conjugate monoacid 580.
 Acridine, 9,10-dihydro-9-phenyl- 874.
 Acridine, 9,10-dihydro-*N*-phenyl- 875.
 Acridine, 9-methyl- 693.
 Acridine-*d*₈ 24.
 Acridine Orange, conjugate monoacid 25.
 Acridine Orange, free base 26.
 Acridine Yellow, conjugate monoacid 27.
 2-Acridinesulfonate ion, 9,10-dihydro-9-oxo- 424.
 2-Acridinesulfonic acid, 9,10-dihydro-9-oxo- 425.
 9(10*H*)-Acridinethione, 10-methyl- 694.
 Acridinium, 3,6-diamino-10-methyl- 30.
 Acridinium ion 28.
 9(10*H*)-Acridinone, 10-ethyl- 569.
 9(10*H*)-Acridinone 29.
 9(10*H*)-Acridone 29.
 Acriflavine cation 30.
 Alloxazine 31.
 Amiloride 32.
 Aminacrine 33.
 9-Aminoacridine 33.
 1-Amino-4-anilinoanthraquinone 34.
 2-Aminoanthracene, conjugate acid 35.
 1-Aminoanthraquinone 36.
 2-Aminoanthraquinone 37.
 6-Amino-7*H*-benz[*de*]anthracen-7-one 38.
 6-Aminobenzanthrone 38.
 2-Aminobenzoic acid 39.
 4-Aminobenzoic acid 40.
 3-Aminobenzophenone 41.
 4-Aminobenzophenone 42.
 4-Aminobiphenyl 43.
 2-Aminofluorene 44.

- 2-Aminofluorene 44.
 α -Amino-2-(formylamino)- γ -oxobenzenebutanoic acid 599.
1-Amino-4-hydroxyanthraquinone 45.
4-Amino-4'-hydroxybiphenyl 46.
2-[6-Amino-3-imino-3*H*-xanthen-9-yl]benzoic acid, monohydrochloride 982.
2-Amino-3-(4-methoxy-6-benzothiazolyl)propionate ion 47.
1-Amino-4-(*N*-methylamino)anthraquinone 48.
7-Amino-4-methylcarbostyryl 49.
4-Amino-2-methyl-1*H*-isoindole-1,3(2*H*)-dione 50.
3-Amino-*N*-methylphthalimide 50.
4'-Aminomethyl-4,5',8-trimethylpsoralen 51.
2-Aminonaphthalene 766.
1-Amino-7-nitrofluorene 52.
1-Amino-4-nitronaphthalene 53.
trans-4-Amino-4'-nitrostilbene 54.
N-[2-(3-Amino-1-oxopropyl)phenyl]formamide 55.
9-Aminophenanthrene 56.
2-Amino-4-[3*H*]pteridinone 57.
2-Aminopurine 58.
2-Aminopyridine 59.
4-Aminopyridine 60.
Angelicin 640.
Aniline 61.
Aniline, 4-(4,6-dichloro-1,3,5-triazin-2-yl)-*N,N*-diethyl- 379.
Aniline, *N,N*-dimethyl- 470.
Aniline/Eosin 557.
4-Anilinobenzanthrone 877.
1-Anilinonaphthalene 62.
Anisole 63.
Anisole, 4-*tert*-butyl-3,5-dinitro- 233.
Anisole, 3,5-dinitro- 515.
Anisole, 3-nitro- 813.
Anisole, 4-nitro- 814.
Anthanthrene 344.
Anthanthrone 346.
Anthracene 64.
Anthracene, 1-acetyl- 12.
Anthracene, 9-acetyl- 13.
Anthracene, 2-amino-, conjugate acid 35.
Anthracene, 9-benzoyl- 148.
Anthracene, 9-benzoyl-10-bromo- 149.
Anthracene, 9-benzoyl-10-chloro- 150.
Anthracene, 10-benzoyl-9-cyano- 151.
Anthracene, 9-benzoyl-10-nitro- 154.
Anthracene, 9,10-bis(phenylethynyl)- 211.
Anthracene, 9-bromo- 215.
Anthracene, 9-butyryl- 239.
Anthracene, 9-cinnamoyl- 296.
Anthracene, 9-cyano- 308.
Anthracene, 9-cyano-10-nitro- 315.
Anthracene, 9,10-dibromo- 358.
Anthracene, 1,5-dichloro- 369.
Anthracene, 9,10-dichloro- 370.
Anthracene, 9,10-dicyano- 381.
Anthracene, 1,3-dimethyl- 472.
Anthracene, 9,10-dimethyl- 473.
Anthracene, 9-[4-(dimethylamino)phenyl]- 463.
Anthracene, 9,10-diphenyl- 529.
Anthracene, 1-hydroxy- 80.
Anthracene, 2-hydroxy- 81.
Anthracene, 9-hydroxy- 82.
Anthracene, 2-iodo- 632.
Anthracene, 9-iodo- 633.
Anthracene, 2-methyl- 699.
Anthracene, 9-methyl- 700.
Anthracene-*d*₁₂, 9-methyl- 701.
Anthracene, 9-naphthoyl- 790.
Anthracene, 9-nitro- 815.
Anthracene, 9-phenyl- 878.
Anthracene, 9-propionyl- 925.
Anthracene, 1,5,9,10-tetrachloro- 1022.
Anthracene, 2,9,10-trichloro- 1077.
Anthracene *d*₁₀ 65.
Anthracene/Chloranil 66.
9-Anthraceneacetic acid, methyl ester 67.
9-Anthracenecarbonitrile 308.
Anthracene-9-carboxaldehyde 68.
Anthracene-9-carboxamide 69.
Anthracene-2-carboxylate ion 79.
Anthracene-9-carboxylic acid, phenylmethyl ester 159.
Anthracene-9-carboxylic acid 70.
9,10-Anthracenedione, 1-amino-4-(methylamino)- 48.
9,10-Anthracenedione, 1,4-bis(methylamino)- 207.
Anthracene photodimer 71.
Anthracene-tetracene photodimer 72.
9-Anthracenol, 9,10-dihydro-10-(10-hydroxy-9(10*H*)-anthracenylidene)- 436.
9-Anthracenone, 10-(1,3-dimethyl-10-oxo-9-anthracenylidene)-2,4-dimethyl-, (*E*)- 1046.
[2.2](1,4)(9,10)-Anthracenophane 73.
anti-[2.2](1,4)-Anthracenophane 74.
syn-[2.2](1,4)-Anthracenophane 75.
9-Anthraldchyd 68.
Anthranilic acid 39.
Anthranol 82.
Anthraquinone 76.
Anthraquinone, 1-amino- 36.
Anthraquinone, 2-amino- 37.
Anthraquinone, 1-amino-4-anilino- 34.
Anthraquinone, 1-amino-4-hydroxy- 45.
Anthraquinone, 1-chloro- 263.
Anthraquinone, 2-chloro- 264.
Anthraquinone, 1,4-diamino- 335.
Anthraquinone, 1,8-dichloro- 371.
Anthraquinone, 2-piperidino- 911.
Anthraquinone-2,6-disulfonate ion 77.
9,10-Anthraquinone-2-sulfonate ion 78.
Anthraquinone-2-sulfonate ion 78.
2-Anthroate ion 79.
1-Anthrol 80.
2-Anthrol 81.
9-Anthrol 82.
Anthrone 83.

- 9-Anthronitrile, 10-nitro- 315.
 1-Anthryl diisopropyl borate 84.
 2-Anthryl diisopropyl borate 85.
 9-Anthryl diisopropyl borate 86.
 9-Anthryl ethenyl ketone 87.
 9-Anthryl ethyl ketone 925.
 1-Anthryl methyl ketone 12.
 1-(2-Anthryl)-2-(2-naphthyl)ethylene 88.
 9-Anthryl 1-naphthyl ketone 790.
 1-(2-Anthryl)-2-phenylethylene 89.
 9-Anthryl phenyl ketone 148.
 9-Anthryl propyl ketone 239.
 9-Anthryl styryl ketone 296.
 1-(2-Anthryl)-2-(2-thienyl)ethylene 90.
 Astacene 91.
 Astacin 91.
 Astaxanthin 92.
 Auramine O 93.
 Aurophosphine, conjugate monoacid 94.
 1-Azaphenanthrene 139.
 4-Azaphenanthrene 140.
 9-Azaphenanthrene 859.
 1-Azoniaphenanthrene 141.
 4-Azoniaphenanthrene 142.
 Azulene 95.
 Azulene, 1,4-dimethyl-7-isopropyl- 488.
 Azure A 96.
 Azure A cation 97.
 Azure B 98.
 Azure C 99.
 BBO 357.
 BBOT 189.
 Bacteriochlorophyll 100.
 Basic Red 1 984.
 Basic Violet 10 983.
 Basic Violet 3 307.
 Basic Yellow 2 93.
 1,2-Benzacridine 101.
 3,4-Benzacridine 102.
 Benz[a]acridine 101.
 Benz[c]acridine 102.
 Benz[b]acridin-12(5*H*)-one 103.
 Benzaldehyde 104.
 Benzaldehyde, 4-phenyl- 175.
 Benzamide 105.
 1,2-Benzanthracene 106.
 2,3-Benzanthracene 1021.
 Benz[a]anthracene 106.
 Benz[b]anthracene 1021.
 7*H*-Benz[de]anthracen-7-one, 6-amino- 38.
 7*H*-Benz[de]anthracen-7-one 107.
 Benzanthrone 107.
 Benzanthrone, 6-amino- 38.
 Benzanthrone, 4-anilino- 877.
 Benzanthrone, 6-hydroxy- 612.
 Benzanthrone, 3-methoxy- 663.
 Benz[a]azulene 108.
 Benzenamine, 4-[(2-bromo-4,6-dinitrophenyl)azo]-*N,N*-dimethyl- 220.
 Benzenamine,
 4-[(2-bromo-4-nitrophenyl)azo]-*N,N*-dimethyl- 226.
 Benzenamine, *N,N*-dimethyl-4-(2-nitroethenyl)- 465.
 Benzenamine, *N,N*-dimethyl-4-nitro- 497.
 Benzenamine, *N,N*-dimethyl-4-[(4-nitrophenyl)azo]- 498.
 Benzenamine, 4-[(2,4-dinitrophenyl)azo]-*N,N*-dimethyl- 520.
 Benzenamine, *N,N*-diphenyl-, conjugate acid 1093.
 Benzenamine, 4-nitro- 812.
 Benzene 109.
 Benzene, 1-butenyl- 578.
 Benzene, 1-butyryl- 237.
 Benzene, chloro- 265.
 Benzene, cyano- 128.
 Benzene, 1,2-dicyano- 382.
 Benzene, 1,3-dicyano- 383.
 Benzene, 1,4-dicyano- 384.
 Benzene, 1,2-dimethoxy-4-nitro- 452.
 Benzene, ethynyl- 873.
 Benzene, fluoro- 594.
 Benzene, methoxy- 63.
 Benzene, 1-methoxy-4-nitro- 814.
 Benzene, (1-methyl-1-butenyl)- 724.
 Benzene, 1-propynyl- 928.
 Benzene, 1,2,4,5-tetracyano- 1023.
 Benzene, 1,3,5-tricyano- 1078.
 Benzene, 1,3,5-triphenyl- 1094.
 Benzene-*d*₆ 110.
 Benzenoacetic acid, α -oxo-, butyl ester 232.
 Benzenobutanoic acid, α -amino-2-(formylamino)- γ -oxo- 599.
 Benzenecarbothioic acid, 4-(dimethylamino)-, *O*-(2-phenylethyl) ester 886.
 Benzenecarbothioic acid, 4-methoxy-, *O*-(2-phenylethyl) ester 887.
 1,4-Benzenedicarboxylic acid, 2-(9*H*-carbazoyl)ethyl methyl ester 665.
 1,4-Benzenedicarboxylic acid, 3-(9*H*-carbazoyl)propyl methyl ester 666.
 Benzenemethanesulfonate ion, 4-(oxophenylacetyl)- 958.
 1,2,4,5-Benzenetetracarbonitrile 1023.
 1,2,4,5-Benzenetetracarboxylic anhydride 953.
 Benzdine 111.
 Benzdine, *N,N,N',N'*-tetramethyl- 1045.
 Benzil 112.
 Benzil/Triethylamine 113.
 Benzo[a]carbazole 114.
 Benzo[b]carbazole 115.
 Benzo[c]carbazole 116.
 Benzo[b]chrysene 117.
 Benzo[a]coronene 118.
 5*H*-Benzocycloheptene, 8-(1-naphthyl)-6,7-dihydro- 793.
 Benzo[1,2-*c*:4,5-*c'*]dipyrrole-1,3,5,7(2*H*,6*H*)-tetrone, 2,6-dipentyl- 525.
 Benzoflavine 119.
 1,2-Benzofluorene 120.
 2,3-Benzofluorene 121.
 3,4-Benzofluorene 122.

- Benzo[*a*]fluorene 120.
 Benzo[*b*]fluorene 121.
 Benzo[*c*]fluorene 122.
 Benzoic acid 123.
 Benzoic acid, 4-(Dimethylamino)-, methyl ester 717.
 Benzoic acid, 2-amino- 39.
 Benzoic acid, 4-amino- 40.
 Benzoic acid, 2-(dimethylamino)ethyl ester 457.
 Benzoic acid, 2-[6-(ethylamino)-3-(ethylimino)-3*H*-xanthen-9-yl]-, monohydrochloride 400.
 Benzoic acid, 2-methyl- 703.
 Benzoic acid, 3-methyl- 704.
 Benzoic acid, 4-methyl- 705.
 Benzoic acid, 4-(methylamino)- 695.
 Benzoic acid, methyl ester 702.
 Benzoin 124.
 Benzo[*c*]1,5]naphthyridine 125.
 Benzo[*c*]1,6]naphthyridine 126.
 Benzo[*c*]1,7]naphthyridine 127.
 Benzonitrile 128.
 Benzonitrile, 4-(1-oxo-3-phenyl-2-propenyl)- 310.
 Benzo[*rst*]pentaphene 129.
 Benzo[*ghi*]perylene, 12*a*,12*b*-dihydro- 408.
 1,12-Benzoperylene 130.
 Benzo[*ghi*]perylene 130.
 Benzo[*c*]phenanthrene 131.
 Benzophenazine 132.
 Benzophenone 133.
 Benzophenone, 3-amino- 41.
 Benzophenone, 4-amino- 42.
 Benzophenone, 4,4'-bis(dimethylamino)- 190.
 Benzophenone, 4-carboxy- 249.
 Benzophenone, 4-chloro- 266.
 Benzophenone, conjugate acid 134.
 Benzophenone, 4,4'-dichloro- 372.
 Benzophenone, 4,4'-dimethoxy- 445.
 Benzophenone, 4[2-[4-(dimethylamino)phenyl]ethyl]- 464.
 Benzophenone, 4,4'-diphenyl- 530.
 Benzophenone, 4-fluoro- 595.
 Benzophenone, 2-hydroxy- 613.
 Benzophenone, 4-hydroxy- 614.
 Benzophenone, 4-methoxy- 664.
 Benzophenone, 2-methyl- 706.
 Benzophenone, 4-methyl- 707.
 Benzophenone, 4-phenyl- 879.
 Benzophenone, 4-sulfomethyl-, anion 1010.
 Benzophenone, 4-(trifluoromethyl)- 1082.
 Benzophenone, 2,4,6-triisopropyl- 1084.
 Benzophenone-(CH₃)₂-(*N,N*-dimethylaniline) 464.
 Benzo[*g*]pteridine-2,4(1*H*,3*H*)-dione, 7,8-dimethyl- 644.
 Benzo[*g*]pteridine-2,4-(3*H*,10*H*)-dione, 7,8,10-trimethyl- 645.
 Benzo[*g*]pteridine-2,4(1*H*,3*H*)-dione 31.
 4*H*-1-Benzopyran-4-one, 3-hydroxy- 616.
 2*H*-1-Benzopyran-2-one, 4-hydroxy-6-methyl- 623.
 1-Benzopyran-4(4*H*)-one 294.
 1,2-Benzopyrene 137.
 3,4-Benzopyrene 137.
 4,5-Benzopyrene 138.
 Benzo[*a*]pyrene/DNA 135.
 Benzo[*e*]pyrene/DNA 136.
 Benzo[*a*]pyrene 137.
 Benzo[*e*]pyrene 138.
 Benzo[*a*]pyrene-7,8,9,10-tetrol, 7,8,9,10-tetrahydro- 1031.
 1,2-Benzopyrone 303.
 5,6-Benzoquinoline 139.
 7,8-Benzoquinoline 140.
 Benzo[*f*]quinoline 139.
 Benzo[*h*]quinoline 140.
 Benzo[*f*]quinolinizinium 141.
 Benzo[*h*]quinolinizinium 142.
 1,4-Benzquinone, 2,3-dimethoxy- 446.
 1,4-Benzquinone, 2,5-dimethoxy- 447.
 1,4-Benzquinone, 2,3-dimethoxy-5-methyl- 450.
 1,4-Benzquinone, 2,5-dimethyl- 474.
 1,4-Benzquinone, 2,6-dimethyl- 475.
 1,4-Benzquinone, 2,6-diphenyl- 531.
 1,4-Benzquinone, methyl- 708.
 1,4-Benzquinone, tetramethyl- 552.
 1,4-Benzquinone, 2,3,5-trimethyl- 1085.
 1,4-Benzquinone 143.
p-Benzquinone 143.
 Benzothiazolium, 2-[4-(dimethylamino)phenyl]-3,6-dimethyl-, chloride 1061.
 Benzo[*b*]thiophene-3(2*H*)-one, 5-(1,1-dimethylpropyl)-2-[5-(1,1-dimethylpropyl)]-3-oxobenzo[*b*]thien-2(3*H*)-ylidene- 337.
 Benzo[*b*]thiophene-3(2*H*)-one, 5-(2,2-dimethylpropyl)-2-[5-(2,2-dimethylpropyl)]-3-oxobenzo[*b*]thien-2(3*H*)-ylidene- 514.
 Benzo[*b*]thiophene-3(2*H*)-one, 6-ethoxy-3-oxobenzo[*b*]thien-2(3*H*)-ylidene- 388.
 Benzo[*b*]thiophen-3(2*H*)-one, 2-(3-oxobenzo[*b*]thien-2(3*H*)-ylidene)- 1062.
 Benzo[*b*]triphenylene/Chloranil 144.
 Benzo[*b*]triphenylene 145.
 Benzoxazole, 2-biphenylphenyl- 177.
 Benzoxazole, 2-(4-biphenyl)- 178.
 Benzoxazole, 2-(1-naphthyl)- 792.
 Benzoxazole, 2,2'-(1,4-phenylene)bis- 883.
 1-Benzoylacetone 146.
 Benzoylamino-2-Δ²-thiazoline 147.
 9-Benzoylanthracene 148.
 4-Benzoylbenzoic acid 249.
 4-Benzoylbiphenyl 879.
 9-Benzoyl-10-bromoanthracene 149.
 9-Benzoyl-10-chloroanthracene 150.
 9-Benzoyl-10-cyanoanthracene 151.
 2-Benzoyl-*N*-methyl-β-naphthiazoline 152.
 8-Benzoylnaphtho[*de*-2.3.4]bicyclo[3.2.2]nona-2,6,8-triene 153.
 9-Benzoyl-10-nitroanthracene 154.
 2-Benzoylpyridine 155.
 3-Benzoylpyridine 156.
 4-Benzoylpyridine 157.

- 1-Benzoyl-3,3,3-trifluoroacetone 158.
 3,4-Benzphenanthrene 131.
 Benzyl 9-anthroate 159.
 Benzylideneacetophenone 258.
 1-Benzylisoquinoline-*N*-oxide 160.
 Benzyl phenyl ketone, conjugate acid 161.
 3-Benzyl-3-phenylphenanthro[9,10-*b*]furan-2(3*H*)-one 162.
 3-Benzyl-3,4,5-triphenyl-2(3*H*)-furanone 163.
 Bergapten 683.
 Biacetyl 164.
 9,9'-Bianthracene 165.
 9,9'-Bianthryl 165.
 9,9'-Bicarbazole 166.
 Bicyclo[2.2.1]hepta-2,5-diene, 7-oxa-2,3-dibenzoyl- 833.
 Bicyclo[2.2.1]hepta-2,5-diene-2,3-diylbis[phenylmethanone] 353.
 Bicyclo[2.2.1]hept-2-ene, 7-oxa-2,3-dibenzoyl- 834.
 Bicyclo[2.2.1]hept-2-ene, 2-phenyl- 896.
 Bicyclo[2.2.2]octa-2,5-diene, 2,3-dibenzoyl- 354.
 Bicyclo[2.2.2]octa-2,5-diene-2,3-diylbis[phenylmethanone] 354.
 Bifluorenylidene 167.
 21*H*-Biline-8,12-dipropanoic acid, 2,17-diethenyl-1,10,19,22,23,24-hexahydro-3,7,13,17-tetramethyl-1,19-dioxo- 168.
 21*H*-Biline-8,12-dipropanoic acid, 3,18-diethenyl-1,19,22,24-tetrahydro-2,7,13,17-tetramethyl-1,19-dioxo- 169.
 (15*E*)-21*H*-Biline-8,12-propanoic acid, 3,18-diethenyl-1,19,22,24-tetrahydro-2,7,13,17-tetramethyl-1,19-dioxo-, dimethyl ester 170.
 Bilirubin 168.
 Bilirubin IX 168.
 Biliverdin 169.
 Biliverdin, dimethyl ester 170.
 Biliverdin IX 169.
 1,1'-Binaphthyl, 3,4-dihydro- 409.
 1,1'-Binaphthyl, 3,4-dihydro-2'-methyl- 421.
 1,1'-Binaphthyl 171.
 2,2'-Binaphthyl 172.
p,p'-Biphenol 437.
 Biphenyl 173.
 Biphenyl, 4-acetyl- 14.
 Biphenyl, 4-amino- 43.
 Biphenyl, 4-amino-4'-hydroxy- 46.
 Biphenyl, 4-benzoyl- 879.
 Biphenyl, 4-bromo- 216.
 Biphenyl, 4-cyano- 309.
 Biphenyl, 4,4'-diamino- 111.
 Biphenyl, 4,4'-dibromo- 359.
 Biphenyl, 4,4'-dihydroxy- 437.
 Biphenyl, 4,4'-dimethoxy- 448.
 Biphenyl, 3,3'-dimethyl- 476.
 Biphenyl, 4-hydroxy- 615.
 Biphenyl, 4-iodo- 634.
 Biphenyl, 4-nitro- 816.
 Biphenyl, 2-(1-phenylethenyl)- 180.
 Biphenyl-*d*₁₀ 174.
 4-Biphenylamine 43.
 4-Biphenylcarbonitrile 309.
 4-Biphenylcarboxaldehyde 175.
 Biphenylene 176.
 2-Biphenylphenylbenzoxazole 177.
 2-(4-Biphenyl)benzoxazole 178.
 1-(4-Biphenyl)-3-chloro-1-propanone 179.
 2-[1,1'-Biphenyl]-4-ylphenylbenzoxazole 177.
 1-(2-Biphenyl)-1-phenylethylene 180.
 2-(4-Biphenyl)-5-phenyl-1,3,4-oxadiazole 181.
 2-(4-Biphenyl)-5-phenyloxazole 182.
 5'-[1,1'-Biphenyl]-4-yl-1,1':4',1'':3'',1''':4''',1''':4''''-quinquephenyl 183.
 6,6'-Biquinoline, conjugate diacid 185.
 2,2'-Biquinoline, conjugate monoacid 186.
 6,6'-Biquinoline, conjugate monoacid 187.
 2,2'-Biquinoline 184.
 2,5-Bis([1,1'-biphenyl]-4-yl)oxazole 357.
 Bis(2,2'-bipyridine)dichloroiridium(III) ion 188.
 2,5-Bis(5-*tert*-butyl-2-benzoxazolyl)thiophene 189.
 3,6-Bis(dimethylamino)acridine, conjugate monoacid 25.
 3,6-Bis(dimethylamino)acridine 26.
 4,4'-Bis(dimethylamino)benzophenone 190.
 6,6'-Bis(dimethylamino)-3,3'-diethyl-2,2'-thiacarbocyanine iodide 191.
 1,7-Bis(dimethylamino)-1,4,6-heptatrien-3-one 192.
 2,5-Bis[7-(dimethylamino)-2,4,6-heptatrienyldene]-cyclopentanone 193.
 3,7-Bis(dimethylamino)-4-nitrophenothiazin-5-ium 722.
 1,9-Bis(dimethylamino)-1,3,6,8-nonatetraen-5-one 194.
 1,9-Bis(dimethylamino)-1,4,6,8-nonatetraen-3-one 195.
all-trans-1,15-Bis(dimethylamino)-1,3,6,8,10,12,14-pentadecaheptaen-5-one 196.
 2,6-Bis[5-(dimethylamino)-2,4-pentadienyldene]-cyclohexanone 197.
 2,5-Bis[5-(dimethylamino)-2,4-pentadienyldene]-cyclopentanone 198.
 3,7-Bis(dimethylamino)phenothiazin-5-ium 719.
 2,6-Bis[3-(dimethylamino)-2-propenyldene]cyclohexanone 199.
 4,4'-Bis(dimethylamino)thiobenzophenone 200.
all-trans-1,13-Bis(dimethylamino)-1,3,5,8,10,12-tridecahexaen-7-one 201.
all-trans-1,13-Bis(dimethylamino)-1,3,6,8,10,12-tridecahexaen-5-one 202.
all-trans-1,11-Bis(dimethylamino)-1,3,6,8,10-undecapentaen-5-one 203.
 Bis(dimethylformamide)phthalocyaninoruthenium(II) 204.
 Bis(dimethylsulfoxide)phthalocyaninoruthenium(II) 205.
 Bis(dimethylsulfoxide)phthalocyaninato-ruthenium(II)/1,4-Dinitrobenzene 206.
 Bis(dimethylthiocarbamyl) disulfide 1050.
 9*H*-9-Bisfluorene, 9-phenyl- 880.
 3,6-Bis(methylamino)acridine, conjugate monoacid 94.
 1,4-Bis(methylamino)anthraquinone 207.
 1,1-Bis(1-naphthyl)ethylene 208.
trans-1,2-Bis(1-naphthyl)ethylene 209.

- trans*-1,2-Bis(2-naphthyl)ethylene 210.
 9,10-Bis(phenylethynyl)anthracene 211.
 1,4-Bis[2-(5-phenyloxazolyl)]benzene 885.
rac-Bis[1-(1-pyrenyl)ethyl]ether 212.
m,m'-Bitoluene 476.
 Bonellin 213.
 2,3-Bornanediol 243.
 Brilliant Sulfaflavine anion 214.
 9-Bromoanthracene 215.
 10-Bromo-9-anthryl phenyl ketone 149.
 4-Bromobiphenyl 216.
 4-Bromo-1-cyanoisoquinoline-*N*-oxide 217.
N-[2-[(2-Bromo-6-cyano-4-nitrophenyl)azo]-5-[(2-cyanoethyl)(2-hydroxyethyl)amino]-4-methoxyphenyl]acetamide 218.
N-[2-[(2-Bromo-4,6-dinitrophenyl)azo]-5-[(2-cyanoethyl)(2-hydroxyethyl)amino]-4-methoxyphenyl]acetamide 219.
 4-[(2-Bromo-4,6-dinitrophenyl)azo]-*N,N*-dimethylbenzenamine 220.
 4-Bromoisoquinoline-*N*-oxide 221.
 Bromo(methanol)(phthalocyaninato)rhodium(III) 222.
 1-(Bromomethyl)naphthalene 223.
 1-Bromonaphthalene 224.
 2-Bromonaphthalene 225.
 4-[(2-Bromo-4-nitrophenyl)azo]-*N,N*-dimethylbenzenamine 226.
 9-Bromophenanthrene 227.
 4-Bromophenol/Eosin 558.
 (*E*)-1-Bromo-4-(2-phenylethenyl)benzene 229.
 1-Bromopyrene 228.
trans-4-Bromostilbene 229.
 2-Bromotriphenylene 230.
 1,3-Butadiene, 1,4-diphenyl- 535.
 1,3-Butadiene, 1,1,4,4-tetraphenyl- 1052.
 Butadiyne, 1,4-diphenyl- 536.
 1,3-Butanedione, 1-phenyl- 146.
 1,3-Butanedione, 4,4,4-trifluoro-1-phenyl- 158.
 2,3-Butanedione 164.
 1,2,3-Butanetrione, 1-(4-nitrophenyl)-2-[(4-methoxyphenyl)hydrazon]- 680.
 2-Butanone 231.
 1-Butene-3-one 761.
 3-Buten-2-one, 4-(2,6,6-trimethyl-1-cyclohexen-1-yl)- 638.
 Butyl benzoylformate 232.
 5-Butyl-2-[5-butyl-3-oxobenzo[*b*]thien-2(3*H*)-ylidene]-benzo[*b*]thiophen-3(2*H*)-one 364.
 4-*tert*-Butyl-3,5-dinitroanisole 233.
 2-[1-(Butylimino)ethyl]-5-methylphenol 234.
 2-*tert*-Butyl-4-methylindazole 235.
N-Butyl-5-nitro-2-furamide 236.
 1-Butynylbenzene 237.
 Butyraldehyde 238.
 9-Butyrylanthracene 239.
 Cadmium(II) tetrabenzoporphyrin 240.
 Cadmium(II) tetraphenylporphyrin 241.
 Caffeine 242.
 11-*cis*-C₂₂ aldehyde 250.
 7,9-*dicis*-C₁₅ aldehyde 758.
 7-*cis*-C₁₅ aldehyde 756.
 9-*cis*-C₁₅ aldehyde 757.
all-trans-C₂₂ aldehyde 251.
all-trans-C₂₄ aldehyde 501.
all-trans-C₁₇ aldehyde 750.
all-trans-C₁₅ aldehyde 755.
 Camphoroquinone 243.
 Canthaxanthin 244.
 Carbazole 245.
 Carbazole, 4*a*,5*a*-dihydro- 410.
 Carbazole, *N,N'*-dimethyl-3-anilino- 471.
 Carbazole, *N*-ethyl- 570.
 Carbazole, 9-[2-(4-methoxycarbonylbenzoxy)ethyl]- 665.
 Carbazole, 9-[3-(4-methoxycarbonylbenzoxy)propyl]- 666.
 Carbazole-(CH₂)₂-(terephthalic acid methyl ester) 665.
 Carbazole-(CH₂)₃-(terephthalic acid methyl ester) 666.
 3-Carbethoxypsoralen 246.
 2,2'-Carbocyanine, 1,1'-diethyl-, chloride, 392.
 4,4'-Carbocyanine, 1,1'-diethyl-, iodide 393.
 Carbostyryl 247.
 Carbostyryl, 7-amino-4-methyl- 49.
 Carbostyryl, negative ion 248.
 4-Carboxybenzophenone 249.
N-[9-(2-Carboxyphenyl)-6-(diethylamino)-3*H*-xanthen-3-ylidene]-*N*-ethylethanaminium chloride 983.
 11-*cis*-β-*apo*-14'-Carotenal 250.
 β-*apo*-14'-Carotenal 251.
 β-*apo*-8'-Carotenal 252.
 ψ,ψ-Carotene, *all-trans*-3,4-didehydro-1,2,7',8'-tetrahydro-1-methoxy- 996.
 ψ,ψ-Carotene, *all-trans*-3,4-didehydro-1,2,7',8'-tetrahydro-1-methoxy-2-oxo- 997.
 β-Carotene, 7,7'-dihydro- 412.
 ψ,ψ-Carotene, *all-trans*-7,8-dihydro- 808.
 β,ε-Carotene, 3,3'-dihydroxy-, (3*R*,3'*R*,6'*R*)- 648.
 β,β-Carotene, 3,3'-dihydroxy-, (3*R*,3'*R*)- 1123.
 ψ-Carotene, 15-*cis*-7,7',8,8',11,11',12,12'-octahydro- 908.
 ψ,ψ-Carotene, 7,7',8,8',11,11',12,12'-octahydro- 909.
 ψ,ψ-Carotene, *all-trans*-3,3',4,4'-tetrahydro-1,1',2,2'-tetrahydro-1,1'-dimethoxy- 998.
 ψ-Carotene, 7,8,11,12-tetrahydro- 255.
 15,15'-*cis*-β-Carotene 253.
 β-Carotene 254.
all-trans-ψ-Carotene 649.
 (3*R*,3'*R*)-β,β-Carotene-3,3'-diol 1123.
 (3*R*,3'*R*,6'*R*)-β,ε-Carotene-3,3'-diol 648.
 β-Carotene-4,4'-dione, 3,3'-dihydroxy- 92.
 β,β-Carotene-4,4'-dione 244.
all-trans-β,β-Carotene-3,3',4,4'-tetrone 91.
 ζ-Carotene 255.
 β-*apo*-8'-Carotenoic acid, methyl ester 256.
 β-*apo*-6'-Carotenoic acid, 4-[tri(4-methylphenyl)porphyrin]-3-phenoxypropyl ester 1090.
 4-(β-*apo*-7'-Carotenyl)benzyl pyropheophorbide 257.

- Chalcone 258.
Chalcone, 4'-cyano- 310.
Chalcone, 4-(dimethylamino)- 455.
Chalcone, 4-methoxy- 667.
Chloranil 259.
Chloranil/Benzo[*b*]triphenylene 144.
Chloranil/Triphenylamine 260.
Chloroaluminum phthalocyanine 261.
1-Chloroanthracene 262.
1-Chloroanthraquinone 263.
2-Chloroanthraquinone 264.
10-Chloro-9-anthryl phenyl ketone 150.
Chlorobenzene 265.
4-Chlorobenzophenone 266.
(2'-Chlorobenzoyl)amino-2- Δ^2 -thiazoline 267.
(4'-Chlorobenzoyl)amino-2- Δ^2 -thiazoline 268.
4-Chloro-6-[4-(diethylamino)phenyl]-*N,N*-dimethyl-1,3,5-triazin-2-amine 269.
2-Chloro-10-dimethylaminopropylphenothiazine 292.
2-Chloro-*N,N*-dimethyl-10*H*-phenothiazine-10-propanamine 292.
N-[2-[(2-Chloro-4,6-dinitrophenyl)azo]-5-[(2-cyanoethyl)(2-hydroxyethyl)amino]-4-methoxyphenyl]acetamide 270.
3-Chloro-2,6-diphenyl-1,4-benzoquinone/4-Methoxy-*N,N*-dimethylaniline 271.
3-Chloro-2,6-diphenyl-1,4-benzoquinone/Naphthalene 272.
3-Chloro-2,6-diphenyl-1,4-benzoquinone/*N,N,N',N'*-Tetramethyl-*p*-phenylenediamine 273.
3-Chloro-2,6-diphenyl-1,4-benzoquinone/1,2,3-Trimethoxybenzene 274.
3-Chloro-2,6-diphenyl-1,4-benzoquinone/Triphenylamine 275.
Chloroindium(III) phthalocyanine 276.
Chloro(methanol)(phthalocyaninato)rhodium(III) 277.
1-(Chloromethyl)naphthalene 278.
1-Chloronaphthalene 279.
2-Chloronaphthalene 280.
3-Chloro-1-(2-naphthyl)-1-propanone 281.
(*Z*)-1-Chloro-4-(2-phenylethenyl)benzene 288.
(*E*)-1-Chloro-4-(2-phenylethenyl)benzene 289.
1-(2-Chlorophenyl)naphthalene 282.
1-(4-Chlorophenyl)naphthalene 283.
Chlorophthalocyanatoindium(III) 276.
Chlorophyll *a* 284.
Chlorophyll *b* 285.
Chlorophyll *a* dimer 286.
Chlorophylline 287.
cis-4-Chlorostilbene 288.
trans-4-Chlorostilbene 289.
2-Chlorothioxanthene-9-one 290.
6-Chloro-1,3,3-trimethyl-6'-nitro-spiro-(2,2'-indolin[2*H*-1]benzopyran) 291.
Chlorpromazine 292.
4-Chromanone 293.
2*H*-Chromene, 8-methoxy-2,2,3-trimethyl-6-nitro- 686.
Chromone 294.
Chromone, 3-hydroxy- 616.
Chrysene 295.
9-Cinnamoylanthracene 296.
Citropten 449.
7,9-*dicis*-C₁₈ ketone 754.
7-*cis*-C₁₈ ketone 752.
9-*cis*-C₁₈ ketone 753.
all-trans-C₁₈ ketone 751.
Copper(II) phthalocyanine 297.
Coproporphyrin I, tetramethyl ester 300.
Coproporphyrin III 298.
Coproporphyrin III, tetramethyl ester 299.
Coronene 301.
Coronene-*d*₁₂ 302.
Coumarin 303.
Coumarin, 7-diethylamino-4-methyl- 390.
Coumarin, 5,7-dimethoxy- 449.
Coumarin, 4-hydroxy- 617.
Coumarin, 4-hydroxy-6-methyl- 623.
Coumarin, 7-hydroxy-4-methyl- 624.
p-Cresol 304.
Crocetin 305.
 α -Crocetin 305.
Croconate Blue dianion 306.
Cryptocyanine 393.
Crystal Violet 307.
2,2'-Cyanine, 1,1'-diethyl-, iodide 394.
2,2'-Cyanine, 1,1'-diethyl-6-bromo-, iodide 391.
2,2'-Cyanine, 1,1'-diethyl-6-iodo-, iodide 396.
9-Cyanoanthracene 308.
4-Cyanobiphenyl 309.
4'-Cyanochalcone 310.
trans-4-Cyano-4'-dimethylaminostilbene 311.
N-[5-[(2-Cyanoethyl)(2-hydroxyethyl)amino]-2-[(2,4-dinitrophenyl)azo]-4-methoxyphenyl]-acetamide 312.
1-Cyano-4-methoxynaphthalene 674.
trans-4-Cyano-4'-methoxystilbene 313.
1-Cyanonaphthalene 314.
9-Cyano-10-nitroanthracene 315.
trans-4-Cyanostilbene 316.
Cycl[3.3.3]azine 951.
Cyclobuta[1,2:3,4]dicycloheptene, 1,2,3,4,5,6,7,8,9,10-decahydro-1,1,5,5,6,6,10,10-octamethyl- 330.
1,2-Cyclobutanedicarboxylic acid, 3,4-di-1-naphthyl-, dimethyl ester, (1 α ,2 α ,3 β ,4 β)- 506.
trans-9,9'-(1,2-Cyclobutanediyl)bis-9*H*-carbazole 366.
Cyclobutene, 1,2-diphenyl- 537.
1,3-Cycloheptadiene 317.
2-Cyclohepten-1-one 318.
1-(1-Cyclohepten-1-yl)naphthalene 319.
2-(1-Cyclohepten-1-yl)naphthalene 320.
1,3-Cyclohexadiene 321.
2,5-Cyclohexadiene-1,4-dione, 2,3,5,6-tetramethyl-, conjugate monoacid 554.
Cyclohexanone, 2,6-bis[5-(dimethylamino)-2,4-pentadienylidene]- 197.

- Cyclohexanone, 2,6-bis[3-(dimethylamino)-2-propenylidene]- 199.
- Cyclohexanone, 2-[7-(dimethylamino)-2,4,6-heptatrienylidene]-6-[5-(dimethylamino)-2,4-pentadienylidene]- 458.
- Cyclohexene, 1-phenyl- 881.
- Cyclohexene-1-one, 3,5,5-trimethyl-2-phenoxy-2- 1089.
- 2-Cyclohexen-1-one, 2-(2-naphthyloxy)-3,5,5-trimethyl- 799.
- 2-Cyclohexen-1-one 322.
- 1-(1-Cyclohexen-1-yl)naphthalene 323.
- 1,3-Cyclooctadiene 324.
- 1-(1-Cycloocten-1-yl)naphthalene 325.
- Cyclopentadiene 326.
- Cyclopentanone, 2,5-bis[7-(dimethylamino)-2,4,6-heptatrienylidene]- 193.
- Cyclopentanone, 2,5-bis[5-(dimethylamino)-2,4-pentadienylidene]- 198.
- 2-Cyclopentenone 327.
- 1-(1-Cyclopenten-1-yl)naphthalene 328.
- 2-(1-Cyclopenten-1-yl)naphthalene 329.
- 9*H*-Cyclopropa[*e*]pyrene, 8*b*,9*a*-dihydro- 413.
- DNA/Benzo[*a*]pyrene 135.
- DNA/Benzo[*e*]pyrene 136.
- 1,2,3,4,5,6,7,8,9,10-Decahydro-1,1,5,5,6,6,10,10-octamethylcyclobuta[1,2:3,4]dicycloheptene 330.
- (*all-E*)-1,3,5,7,9-Decapentaene, 3,8-dimethyl-1,10-bis(2,6,6-trimethyl-1-cyclohexen-1-yl)- 477.
- Decapreno- β -carotene 331.
- 3,5,7,9-Decatetraen-2-one, 10-(dimethylamino)- 456.
- Deoxybenzoin 332.
- Deuteroporphyrin, dimethyl ester 333.
- Diacetyl 164.
- trans-N,N'*-Diacetylidido 334.
- 3,6-Diaminoacridine, conjugate diacid 922.
- 3,6-Diaminoacridine, conjugate monoacid 923.
- 3,6-Diaminoacridine, hydrochloride 921.
- 1,4-Diaminoanthraquinone 335.
- 4,4'-Diaminobiphenyl 111.
- 3,6-Diamino-2,7-dimethylacridine, conjugate monoacid 27.
- 3,6-Diamino-2,7-dimethyl-9-phenylacridine, monohydrochloride 119.
- 3,7-Diamino-2,8-dimethyl-5-phenylphenazinium 990.
- 3,7-Diamino-2,8-dimethyl-5-phenylphenazinium, conjugate diacid 991.
- 3,7-Diamino-2,8-dimethyl-5-phenylphenazinium, conjugate monoacid 992.
- 4,7-Diamino-1*H*-isoindole-1,3(2*H*)-dione 336.
- 3,6-Diamino-10-methylacridinium 30.
- 3,7-Diamino-5-phenoselenazinium, conjugate monoacid 994.
- 3,7-Diamino-5-phenoselenazinium 993.
- 3,7-Diamino-5-phenothiazinium, conjugate monoacid 1064.
- 3,7-Diamino-5-phenothiazinium 1063.
- 3,7-Diamino-5-phenoxazinium, conjugate monoacid 836.
- 3,7-Diamino-5-phenoxazinium 835.
- 3,6-Diamino-9-phenylacridine, conjugate monoacid 898.
- 3,6-Diaminophthalimide 336.
- p,p'*-Diaminoquaterphenyl 959.
- p,p'*-Diaminoterphenyl 1011.
- 3,6-Diamino-2,7,9-trimethylacridine, conjugate monoacid 580.
- 5,5'-Di-*tert*-amylthioindigo 337.
- 8,8'-Diapocarotenedioic acid 305.
- 1,5-Diazabicyclo[3,3,0]octa-3,7-diene-2,6-dione, 3,4,7,8-tetramethyl- 1047.
- 1,5-Diazaphenanthrene 862.
- 1,8-Diazaphenanthrene 865.
- 2,10-Diazaphenanthrene 127.
- 2,5-Diazaphenanthrene 863.
- 3,10-Diazaphenanthrene 126.
- 3,5-Diazaphenanthrene 864.
- 4,10-Diazaphenanthrene 125.
- 4,5-Diazaphenanthrene 861.
- 1,2,5,6-Dibenzacridine 339.
- Dibenz[*a,j*]acridine/PrOH 338.
- Dibenz[*a,h*]acridine 339.
- 1,2,7,8-Dibenzanthanthrene 937.
- 1,2,3,4-Dibenzanthracene 145.
- 1,2,5,6-Dibenzanthracene 341.
- Dibenz[*a,h*]anthracene/Chloranil 340.
- Dibenz[*a,h*]anthracene 341.
- 5*H*-Dibenzobismole, 5-phenyl- 880.
- 3,4,5,6-Dibenzocarbazole 342.
- 7*H*-Dibenzo[*c,g*]carbazole 342.
- Dibenzo[*b,def*]chrysene 343.
- Dibenzo[*def,mno*]chrysene 344.
- Dibenzo[*def,p*]chrysene 345.
- Dibenzo[*def,mno*]chrysene-6,12-dione 346.
- Dibenzo[*g,p*]chrysene 347.
- 5*H*-Dibenzo[*a,d*]cycloheptene 348.
- 5*H*-Dibenzogermole, 5,5-diphenyl- 540.
- Dibenzo[*c,g*]phenanthrene 847.
- Dibenzophenazine 349.
- 5*H*-Dibenzophosphole, 5-phenyl- 897.
- 6*H*-Dibenzo[*b,d*]pyran-6-one 350.
- 1,2,3,4-Dibenzopyrene 345.
- 1,2,4,5-Dibenzopyrene 783.
- 3,4,8,9-Dibenzopyrene 343.
- 3,4,9,10-Dibenzopyrene 129.
- Dibenzo[*a,e*]pyrene 783.
- Dibenzo[*a,h*]pyrene 343.
- Dibenzo[*a,i*]pyrene 129.
- Dibenzo[*a,l*]pyrene 345.
- 5,6,7,8-Dibenzoquinoxaline 351.
- Dibenzo[*f,h*]quinoxaline 351.
- 5*H*-Dibenzosilole, 5,5-diphenyl- 549.
- 5*H*-Dibenzostannole, 5,5-diethyl- 402.
- 5*H*-Dibenzostibole, 5-phenyl- 899.
- Dibenzo[*c,g*]triphenylene 352.
- 2,3-Dibenzoylbicyclo[2.2.1]hepta-2,5-diene 353.
- 2,3-Dibenzoylbicyclo[2.2.2]octa-2,5-diene 354.
- Dibenzoylmethane 355.
- 2,3-Dibenzoyl-2,5-norbornadiene 353.

- 2,4-Dibenzoylpentane 482.
 2,3:6,7-Dibenzphenanthrene 848.
N-(Dibenzylamino)methylphthalimide 356.
 2,5-Di(4-biphenyl)oxazole 357.
 9,10-Dibromoanthracene 358.
 4,4'-Dibromobiphenyl 359.
 Dibromofluorescein dianion 360.
 1,4-Dibromonaphthalene 361.
 (Z)-*N,N*-Dibutyl-2,3-dihydro-3-oxo-2-(3-oxonaphtho[1,2-*b*]thien-2(3*H*)-ylidene)-naphtho[1,8-*bc*]thiopyran-6-sulfonamide 362.
 (E)-*N,N*-Dibutyl-2,3-dihydro-3-oxo-2-(3-oxonaphtho[1,2-*b*]thien-2(3*H*)-ylidene)-naphtho[1,8-*bc*]thiopyran-6-sulfonamide 363.
 5,5'-Dibutylthioindigo 364.
 5,5'-Di-*tert*-butylthioindigo 365.
trans-1,2-Di(*N*-carbazolyl)cyclobutane 366.
meso (DL)-2,4-Di(*N*-carbazolyl)pentane 367.
rac (DD,LL)-2,4-Di(*N*-carbazolyl)pentane 368.
 2,2'-Dicarbocyanine iodide, 1,1'-diethyl- 395.
 1,5-Dichloroanthracene 369.
 9,10-Dichloroanthracene 370.
 1,8-Dichloroanthraquinone 371.
 4,4'-Dichlorobenzophenone 372.
 Dichlorobis(4,7-dimethyl-1,10-phenanthroline)iridium(III) ion 373.
 Dichlorobis(5,6-dimethyl-1,10-phenanthroline)iridium(III) ion 374.
 Dichlorobis(1,10-phenanthroline)iridium(III) ion 375.
 3,5-Dichloro-2,6-diphenyl-1,4-benzoquinone/Triphenylamine 376.
 1,4-Dichloronaphthalene 377.
 4,4'-Dichlorostilbene 378.
 4-(4,6-Dichloro-1,3,5-triazin-2-yl)-*N,N*-diethylaniline 379.
 Dicumarol 380.
 9,10-Dicyanoanthracene 381.
 1,2-Dicyanobenzene 382.
 1,3-Dicyanobenzene 383.
 1,4-Dicyanobenzene 384.
 1,4-Dicyanonaphthalene 385.
trans-4,4'-Dicyanostilbene 386.
all-trans-3',4'-Didehydro- β,ψ -16'-carotenal 387.
 5,7:14,16-Dietheno-8*H*,13*H*-diindeno[2,1-*h*:1',2'-*i*]-[1,4]dioxacyclotridecin, 10,11-dihydro- 414.
 5,7:20,22-Dietheno-8*H*,19*H*-diindeno[2,1-*n*:1',2'-*o*]-[1,4,7,10]tetraoxacyclononadecin, 10,11,13,14,16,17-hexahydro- 609.
 5,7:15,17-Dietheno-2,20-heptano-*H*-cyclotetradeca[1,2-*a*:1,14-*a'*]diindene, 9,10,11,12,13,14-hexahydro- 610.
 5,7:16,18-Dietheno-2,21-octanocyclopentadeca[1,2-*a*:1,15-*a'*]diindene, 8,9,10,11,12,13,14,15-octahydro- 829.
 2,17-Diethenyl-1,10,19,22,23,24-hexahydro-3,7,13,17-tetramethyl-1,19-dioxo-21*H*-bilin-8,12-dipropanoic acid 168.
 3,18-Diethenyl-1,19,22,24-tetrahydro-2,7,13,17-tetramethyl-1,19-dioxo-21*H*-bilin-8,12-dipropanoic acid 169.
 6,6'-Diethoxythioindigo 388.
N-[2-(Diethylamino)ethyl]phthalimide 389.
 7-Diethylamino-4-methylcoumarin 390.
 Diethylaniline/Pyrene 942.
 1,1'-Diethyl-6-bromo-2,2'-cyanine iodide 391.
 1,1'-Diethyl-2,2'-carbocyanine chloride 392.
 1,1'-Diethyl-4,4'-carbocyanine iodide 393.
 1,1'-Diethyl-2,2'-cyanine iodide 394.
 5,5-Diethyl-5*H*-dibenzostannole 402.
 1,1'-Diethyl-2,2'-dicarbocyanine iodide 395.
 Diethyl ether/Pyromellitic dianhydride 954.
 1,1'-Diethyl-6-iodo-2,2'-cyanine iodide 396.
 3,3'-Diethyl-9-methoxy-2,2'-thiacarbocyanine iodide 397.
 3,3'-Diethyl-2,2'-oxacarbocyanine iodide 398.
 3,3'-Diethyl-2,2'-oxadiazocarbocyanine iodide 399.
N,N'-Diethylrhodamine 400.
 5,5'-Diethylselenoindigo 401.
 9,9-Diethyl-9*H*-9-stannafuorene 402.
 3,3'-Diethyl-2,2'-thiacarbocyanine iodide 403.
 3,3'-Diethyl-2,2'-thiacyanine iodide 404.
 3,3'-Diethyl-2,2'-thiadiazocarbocyanine iodide 405.
 3,3'-Diethyl-2,2'-thiatricarbocyanine iodide 406.
 Difurobenzene, 2-formyl-5'-methyl- 600.
 6,6'-Dihexyloxythioindigo 407.
 12*a*,12*b*-Dihydrobenzo[ghi]perylene 408.
 3,4-Dihydro-1,1'-binaphthyl 409.
 4*a*,5*a*-Dihydrocarbazole 410.
 4',5'-Dihydro-3-carbethoxypsoralen 411.
 7,7'-Dihydro- β -carotene 412.
 8*b*,9*a*-Dihydro-9*H*-cyclopropa[*e*]pyrene 413.
 10,11-Dihydro-5,7:14,16-dietheno-8*H*,13*H*-diindeno[2,1-*h*:1',2'-*i*]-[1,4]dioxacyclotridecin 414.
 6,12-Dihydro-3,9-dimethyl(indeno[1,2-*b*]fluorene) 484.
 12*a*,12*b*-Dihydro-5,8-dimethyl(indolino[2,3-*c*]carbazole) 415.
 5,10-Dihydro-5,10-dimethylphenazine 416.
 (R)-4,5-Dihydrodinaphtho[2,1-*e*:1',2'-*g*][1,4]dioxocin 417.
 (R)-5,6-Dihydro-4*H*-dinaphtho[2,1-*f*:1',2'-*h*][1,5]dioxonin 418.
 5,10-Dihydro-5,10-diphenylphenazine 419.
 7,10-Dihydro-7,10-ethenocyclohepta[*de*]naphthalen-8-yl)phenyl]methanone 153.
 2,3-Dihydro-7*H*-furobenzopyran-7-one 430.
 5,10-Dihydroindeno[2,1-*a*]indene 420.
 3,4-Dihydro-2'-methyl-1,1'-binaphthyl 421.
 11,12-Dihydro-5-methylindeno[2,1-*a*]fluorene 725.
 2(1,3-Dihydro-1-methyl-3-oxo-2*H*-indol-2-ylidene)-1,2-dihydro-1-methyl-3*H*-indol-3-one 487.
 5,10-Dihydro-5-methyl-10-phenylphenazine 422.
 5,12-Dihydro-5-methylquino[2,3-*b*]acridine-7,14-dione 423.
 9,10-Dihydro-9-oxo-2-acridinesulfonate ion 424.
 9,10-Dihydro-9-oxo-2-acridinesulfonic acid 425.
 2,3-Dihydro-7-oxo-7*H*-furo[3,2-*g*][1]benzopyran-6-carboxylic acid, ethyl ester 411.

- 2,3-Dihydrophenalene 426.
 3,4-Dihydrophenanthrene 427.
 9,10-Dihydrophenanthrene 428.
 1-(1a,9b-Dihydrophenanthro[9,10-b]oxiren-4-yl)ethanone 15.
 9,10-Dihydro-9-phenylacridine 874.
 9,10-Dihydro-N-phenylacridine 875.
 1,2-Dihydro-3-phenylnaphthalene 429.
 4',5'-Dihydropsoralen 430.
 5,12-Dihydroquino[2,3-b]acridine-7,14-dione 431.
 1,2-Dihydro-2,2,4,6-tetramethylquinoline 432.
 1,4-Dihydro-5,6,11,12-tetraphenyl-1,4-epidioxytetracene 433.
 5,12-Dihydro-5,6,11,12-tetraphenyl-5,12-epidioxytetracene 434.
 (E)-5-[3,4-Dihydro[1,4]thiazino[3,4-b]benzothiazol-1-yl)methylene]-3-ethyl-2-thioxo-4-thiazolidinone 435.
 10H,10'H-10,10'-Dihydroxybianthrylidene 436.
 4,4'-Dihydroxybiphenyl 437.
 3,3'-Dihydroxy- β -carotene-4,4'-dione 92.
 4,7-Dihydroxy-1H-isoindole-1,3(2H)-dione 444.
 Dihydroxylcopene 438.
 2,4-Dihydroxy-5-methylpyrimidine 1072.
 2,3-Dihydroxynaphthalene, conjugate base 443.
 1,5-Dihydroxynaphthalene 439.
 1,8-Dihydroxynaphthalene 440.
 2,3-Dihydroxynaphthalene 441.
 2,7-Dihydroxynaphthalene 442.
 3,6-Dihydroxyphtalimide 444.
 4,4'-Dimethoxybenzophenone 445.
 2,3-Dimethoxy-1,4-benzoquinone 446.
 2,5-Dimethoxy-1,4-benzoquinone 447.
 4,4'-Dimethoxybiphenyl 448.
 5,7-Dimethoxycoumarin 449.
 2,3-Dimethoxy-5-methyl-1,4-benzoquinone 450.
 1,4-Dimethoxynaphthalene 451.
 1,2-Dimethoxy-4-nitrobenzene 452.
 (E)-1,4-Dimethoxy-2-[2-(4-nitrophenyl)ethenyl]benzene 453.
 trans-2,5-Dimethoxy-4'-nitrostilbene 453.
 4,4'-Dimethoxythiobenzophenone 454.
 7,8-Dimethylalloxazine 644.
 4-(Dimethylamino)benzoic acid, methyl ester 717.
 4-(Dimethylamino)chalcone 455.
 10-(Dimethylamino)-3,5,7,9-decatetraen-2-one 456.
 2-(Dimethylamino)ethyl benzoate 457.
 2-[7-(Dimethylamino)-2,4,6-heptatrienylidene]-6-[5-(dimethylamino)-2,4-pentadienylidene]-cyclohexanone 458.
 N-[(Dimethylamino)methyl]phthalimide 459.
 4-(Dimethylamino)nitrobenzene 497.
 1-Dimethylamino-4-nitronaphthalene 460.
 trans-4-Dimethylamino-4'-nitrostilbene 461.
 trans-4-Dimethylamino- β -nitrostyrene 465.
 8-(Dimethylamino)-3,5,7-octatrien-2-one 462.
 9-[4-(Dimethylamino)phenyl]anthracene 463.
 (E)-4-[2-[4-(Dimethylamino)phenyl]ethenyl]benzonitrile 311.
 4-[2-[4-(Dimethylamino)phenyl]ethyl]benzophenone 464.
 trans-1-(4-Dimethylaminophenyl)-2-nitroethylene 465.
 3-[4-(Dimethylamino)phenyl]-1-phenyl-2-propen-1-one 455.
 2-(Dimethylamino)purine 466.
 2-(Dimethylamino)pyridine 467.
 4-(Dimethylamino)pyridine 468.
 2-(Dimethylamino)pyridine/4-Nitroaniline 469.
 N,N-Dimethylaniline 470.
 N,N'-Dimethyl-3-anilino-carbazole 471.
 1,3-Dimethylantracene 472.
 9,10-Dimethylantracene 473.
 2,5-Dimethyl-1,4-benzoquinone 474.
 2,6-Dimethyl-1,4-benzoquinone 475.
 3,3'-Dimethylbiphenyl 476.
 (all-E)-3,8-Dimethyl-1,10-bis(2,6,6-trimethyl-1-cyclohexen-1-yl)-1,3,5,7,9-decapentaene 477.
 1-(3,3-Dimethyl-1-buten-2-yl)naphthalene 478.
 (Z)-2-(3,3-Dimethyl-1-butenyl)naphthalene 494.
 (E)-2-(3,3-Dimethyl-1-butenyl)naphthalene 495.
 2,7-Dimethyl-1,8-diazaphenanthrene 499.
 N,N'-Dimethyl-5,11-dihydroindolo[3,2-b]carbazole 479.
 cis-2,3-Dimethyl-2,3-di-(2-naphthyl)oxirane 480.
 trans-2,3-Dimethyl-2,3-di-(2-naphthyl)oxirane 481.
 2,4-Dimethyl-1,5-diphenyl-1,5-pentanedione 482.
 N,N'-Dimethyl-N,N'-diphenyl-1,4-phenylenediamine 483.
 2-(1,1-Dimethylethyl)-5-methoxy-1,3-dinitrobenzene 233.
 3,9-Dimethyl-trans-fluorenone 484.
 2,5-Dimethyl-2,4-hexadiene 485.
 1,3-Dimethylindazole 486.
 N,N'-Dimethylindigo 487.
 1,4-Dimethyl-7-isopropylazulene 488.
 1-(2,2-Dimethyl-1-methylenepropyl)naphthalene 478.
 1,2-Dimethylnaphthalene 489.
 2,3-Dimethylnaphthalene 490.
 2,7-Dimethylnaphthalene 491.
 N,N-Dimethyl-2-naphthylamine, conjugate acid 493.
 N,N-Dimethyl-2-naphthylamine 492.
 cis-3,3-Dimethyl-1-(2-naphthyl)-1-butene 494.
 trans-3,3-Dimethyl-1-(2-naphthyl)-1-butene 495.
 2,2-Dimethyl-1-(1-naphthyl)-1-phenylethylene 496.
 N,N-Dimethyl-4-nitroaniline 497.
 N,N-Dimethyl-4-[(4-nitrophenyl)azo]benzenamine 498.
 (E,Z)-2,6-Dimethyl-2,4,6-octatriene 806.
 (E)-10-(1,3-Dimethyl-10-oxo-9-anthracenylidene)-2,4-dimethyl-9-anthracenone 1046.
 3,8-Dimethyl-4,7-phenanthroline 499.
 N,N-Dimethyl-10H-phenothiazine-10-propanamine 924.
 N,N-Dimethyl-p-phenylenediamine 500.
 1,3-Dimethyl-2,4-pyrimidinedione 502.
 (E,E,E,E,E)-7,11-Dimethyl-7-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2,4,6,8,10,12-tridecahexaenol 501.
 1,3-Dimethyluracil 502.
 Dinaphtho[2,1-d:1',2'-f][1,3,2]dioxaphosphepin, 4-methoxy-, 4-oxide, (R)- 668.
 Dinaphtho[2,1-d:1',2'-f][1,3,2]dioxaphosphepin, 4-methoxy-, 4-oxide 669.
 Dinaphtho[2,1-b:1',2'-d][1,6]dioxecin, 4,5,6,7-tetrahydro, (R)- 1029.
 (S)-Dinaphtho[2,1-d:1',2'-f][1,3]dioxepin 503.
 Dinaphtho[2,1-e:1',2'-g][1,4]dioxocin, 4,5-dihydro-, (R)- 417.

- 4*H*-Dinaphtho[2,1-*f*:1',2'-*h*] [1,5]dioxonin, 5,6-dihydro-, (*R*)- 418.
 Di-2-naphthylamine 504.
 1,4-Di-1-naphthylbutane 505.
t-3,4-Di-(1-naphthyl)-*r*-1,4-dimethoxycarbonyl-cyclobutane 506.
 1,2-Di-1-naphthylethane 507.
 Di-1-naphthylmethane 508.
 2,5-Di(1-naphthyl)-1,3,4-oxadiazole 509.
cis-2,3-Di-(2-naphthyl)oxirane 510.
trans-2,3-Di-(2-naphthyl)oxirane 511.
 1,3-Di-1-naphthylpropane 512.
 1,3-Di-2-naphthylpropane 513.
 5,5'-Dineopentylthioindigo 514.
 3,5-Dinitroanisole 515.
 3,5-Dinitroanisole/Water 516.
 1,4-Dinitrobenzene/Bis(dimethylsulfoxide) phthalocyaninatoruthenium(II) 206.
 1,2-Dinitronaphthalene 517.
 1,4-Dinitronaphthalene 518.
 1,8-Dinitronaphthalene 519.
 4-[(2,4-Dinitrophenyl)azo]-*N,N*-dimethylbenzenamine 520.
 3-[[4-[(2,4-Dinitrophenyl)azo]phenyl](2-hydroxyethyl)amino]-4-propanenitrile 521.
 (*E*)-2,4-Dinitro-1-(2-phenylethenyl)benzene 522.
trans-2,4-Dinitrostilbene 522.
cis-4,4'-Dinitrostilbene 523.
trans-4,4'-Dinitrostilbene 524.
 1-(2,6-Dioxacyclohexyl)naphthalene 779.
 2-(2,6-Dioxacyclohexyl)naphthalene 780.
 1,3-Dioxane, 2-(1-naphthalenyl)- 779.
 1,3-Dioxane, 2-(2-naphthalenyl)- 780.
 1,4-Dioxin, 2,3,5,6-tetraphenyl- 1054.
 2,6-Dioxo-1,2,3,6-tetrahydro-4-pyrimidinecarboxylic acid 832.
 2,6-Dipentylbenzo[1,2-*c*:4,5-*c'*]dipyrrole-1,3,5,7(2*H*,6*H*)-tetrone 525.
N,N'-Dipentylpyromellitic diimide 525.
 1,3-Di(9-phenanthryl)propane 526.
 Diphenylacetylene 527.
 Diphenylamine 528.
 Diphenylamine/2,6-Diphenyl-1,4-benzoquinone 532.
 9,10-Diphenylanthracene 529.
 4,4'-Diphenylbenzophenone 530.
 2,6-Diphenyl-1,4-benzoquinone 531.
 2,6-Diphenyl-1,4-benzoquinone/Diphenylamine 532.
 2,6-Diphenyl-1,4-benzoquinone/TMPD 533.
 2,6-Diphenyl-1,4-benzoquinone/*N,N,N',N'*-Tetramethyl-*p*-phenylenediamine 533.
 2,6-Diphenyl-1,4-benzoquinone/Triphenylamine 534.
 1,4-Diphenyl-1,3-butadiene 535.
 1,4-Diphenylbutadiyne 536.
 1,2-Diphenylcyclobutene 537.
 5,5-Diphenyl-5*H*-dibenzogermole 540.
 5,5-Diphenyl-5*H*-dibenzosilole 549.
 1,4-Diphenyl-2,3-dibenzoyl-1,4-epoxy-1,4-dihydronaphthalene 538.
 1,1-Diphenylethylene 539.
 9,9-Diphenyl-9*H*-9-germafluorene 540.
 1,6-Diphenyl-1,3,5-hexatriene 541.
 1,4-Diphenylnaphthalene 542.
 1,5-Diphenylnaphthalene 543.
 1,8-Diphenyl-1,3,5,7-octatetraene 544.
 2,5-Diphenyl-1,3,4-oxadiazole 545.
 2,5-Diphenyloxazole 546.
N,N'-Diphenyl-*p*-phenylenediamine, conjugate acid 548.
N,N'-Diphenyl-*p*-phenylenediamine 547.
 1,3-Diphenyl-1,3-propanedione 355.
 1,3-Diphenyl-2-propen-1-one 258.
 9,9-Diphenyl-9*H*-9-silafluorene 549.
N-[(Dipropylamino)methyl]phthalimide 550.
 Disperse Blue 14 207.
 Dithiazanine iodide 405.
 Durene/Tetracyanobenzene 551.
 Duroquinone 552.
 Duroquinone, conjugate monoacid 554.
 Duroquinone/Triphenylamine 553.
 Echinenone 555.
 Eosin 556.
 Eosin/Aniline 557.
 Eosin/4-Bromophenol 558.
 Eosin/2-Naphthol 559.
 Eosin/Resorcinol 560.
 Eosin/Sulfanilic acid 561.
 Eosin Y 556.
 Eosin Yellowish 556.
 Eosin dianion 562.
 5-(1,4-Epidioxyphenyl)-6,11,12-triphenyltetracene 563.
 1,4-Epidioxytetracene, 1,4-dihydro-5,6,11,12-tetraphenyl- 433.
 5,12-Epidioxytetracene, 5,12-dihydro-5,6,11,12-tetraphenyl- 434.
 Ergosterol 564.
 Erythrosin 565.
 Erythrosin dianion 566.
 Ethanone, 1,2,2-triphenyl 882.
 (*E*)-4,4'-(1,2-Ethenediyl)bisbenzonitrile 386.
 2-Ethoxynaphthalene 567.
 Ethyl Violet 568.
 10-Ethyl-9(10*H*)-acridinone 569.
N-Ethylacridinone 569.
 2-[6-(Ethylamino)-3-(ethylimino)-3*H*-xanthen-9-yl]benzoic acid, monohydrochloride 400.
N-Ethylcarbazole 570.
 Ethylene, 1-acenaphthyl-1-phenyl- 4.
 Ethylene, 1-(2-anthryl)-2-(2-naphthyl)- 88.
 Ethylene, 1-(2-anthryl)-2-phenyl- 89.
 Ethylene, 1-(2-anthryl)-2-(2-thienyl)- 90.
 Ethylene, 1,1-bis(1-naphthyl)- 208.
 Ethylene, *trans*-1,2-bis(1-naphthyl)- 209.
 Ethylene, *trans*-1,2-bis(2-naphthyl)- 210.
 Ethylene, 2,2-dimethyl-1-(1-naphthyl)-1-phenyl- 496.
 Ethylene, 1,1-diphenyl- 539.
 Ethylene, 1-methyl-1-(1-naphthyl)- 738.
 Ethylene, 1-methyl-1-(2-naphthyl)- 739.
 Ethylene, *trans*-1-(1-naphthyl)-2-(2-naphthyl)- 796.

- Ethylene, 1-(1-naphthyl)-1-phenyl- 800.
 Ethylene, *trans*-1-(1-naphthyl)-2-phenyl- 801.
 Ethylene, *trans*-1-(2-naphthyl)-2-phenyl- 802.
 Ethylene, 1,1,2-trimethyl-2-(2-naphthyl)- 1087.
 Ethylene, triphenyl- 1098.
 3-Ethyl-2-[3-(3-ethyl-2(3*H*)-benzothiazolylidene)-2-methoxy-1-propenyl]benzothiazolium iodide 397.
 3-Ethyl-2-[3-(3-ethyl-2(3*H*)-benzothiazolylidene)-1-propenyl]benzothiazolium iodide 403.
 3-Ethyl-2-[(3-ethyl-2(3*H*)-benzothiazolylidene)methyl]-benzothiazolium iodide 404.
 3-Ethyl-2-[5-(3-ethyl-2(3*H*)-benzothiazolylidene)-1,3-pentadienyl]benzothiazolium iodide 405.
 3-Ethyl-2-[7-(3-ethyl-2(3*H*)-benzothiazolylidene)-1,3,5-heptatrienyl]benzothiazolium iodide 406.
 (*E,Z*)-3-Ethyl-5[(3-ethyl-2(3*H*)-benzothiazolylidene)-ethylidene]-2-thioxo-4-thiazolidinone 571.
 3-Ethyl-2-[3-(3-ethyl-2(3*H*)-benzoxazolylidene)-1-propenyl]benzoxazolium iodide 398.
 3-Ethyl-2-[5-(3-ethyl-2(3*H*)-benzoxazolylidene)-1,3-pentadienyl]benzoxazolium iodide 399.
 1-Ethyl-2-[5-(1-ethyl-1,3-dihydro-3,3-dimethyl-2*H*-indol-2-ylidene)-3-fluoro-1,3-pentadienyl]-3,3-dimethyl-3*H*-indolium 572.
 1-Ethyl-2-[5-(1-ethyl-1,3-dihydro-3,3-dimethyl-2*H*-indol-2-ylidene)-1,3-pentadienyl]-3,3-dimethyl-3*H*-indolium 573.
 3-Ethyl-2-[3-(3-ethyl-6-dimethylamino-2(3*H*)-benzothiazolylidene)-1-propenyl]-6-(dimethylamino)benzothiazolium iodide 191.
 1-Ethyl-naphthalene 574.
 2-Ethyl-naphthalene 575.
 9-Ethylphenanthrene 576.
 1-Ethyl-2-phenylacetylene 237.
N-Ethylphthalimide 577.
 β -Ethylstyrene 578.
O-Ethyl-1-thionaphthoate 774.
 Ethynylbenzene 873.
 Etiochlorin II 579.
 Euchrysine, conjugate monoacid 580.
 Flavine mononucleotide 581.
 Flavine mononucleotide, conjugate monoacid 582.
 Flavone 583.
 Flavone, 3-hydroxy- 619.
 Flavone, 7-hydroxy- 620.
 9*H*-Fluorene-1-amine, 7-nitro- 52.
 Fluoranthene 584.
 2-Fluorenamine 44.
 Fluorene 585.
 Fluorene, 2-amino- 44.
 9*H*-Fluorene, 9-(9*H*-fluoren-9-ylidene)- 167.
 Fluorene, 1-hydroxy- 621.
 Fluorene, 2-hydroxy- 622.
 Fluoren-1-ol 621.
 Fluoren-2-ol 622.
 Fluoren-9-one 586.
 9-Fluorenone 586.
 1-Fluorenyl diisopropyl borate 587.
 2-Fluorenyl diisopropyl borate 588.
 Fluoren-9-ylidene 589.
 9-(9*H*-Fluoren-9-ylidene)-9*H*-fluorene 167.
 2-Fluorenyl phenyl ketone 590.
 Fluorescein 591.
 Fluorescein, conjugate monoacid 592.
 Fluorescein, 2',4',5',7'-tetrabromo-, dianion 562.
 Fluorescein, tetraiodo-, dianion 566.
 Fluorescein dianion 593.
 Fluorobenzene 594.
 4-Fluorobenzophenone 595.
 (*E*)-1-Fluoro-4-(2-phenylethenyl)benzene 596.
trans-4-Fluorostilbene 596.
 2-(*N*-Formylamino)acetophenone 17.
 2-(*N*-Formylamino)-3'-aminopropiophenone 55.
 γ -[2-(Formylamino)phenyl]- γ -oxo- α -aminobutyryl-L-lysine 597.
 3-Formylfurochromene 598.
N-Formylkynurenine 599.
 2-(*N*-Formyl-*N*-methylamino)acetophenone 19.
 2-Formyl-5'-methyldifurobenzene 600.
 3-Formyl-2-methylfurochromene 601.
 3-Formyl-8-methylfurochromene 602.
 2-Furamide, *N*-butyl-5-nitro- 236.
 2(3*H*)-Furanone, 3-benzyl-3,4,5-triphenyl- 163.
 2(3*H*)-Furanone, 3,3,4,5-tetraphenyl- 1055.
 2(3*H*)-Furanone, 3,3,5-triphenyl- 1099.
 7*H*-Furo[3,2-*g*][1]benzopyran-6-carboxaldehyde 598.
 2*H*-Furo[2,3-*h*]-1-benzopyran-2-one 640.
 7*H*-Furo[3,2-*g*][1]benzopyran-7-one 934.
 Furochromene, 3-formyl- 598.
 Furochromene, 3-formyl-2-methyl- 601.
 Furochromene, 3-formyl-8-methyl- 602.
 2-Furoic acid, 5-nitro- 818.
 Gentian Violet 307.
 9*H*-9-Germafluorene, 9,9-diphenyl- 540.
 Germane, methyl-1-naphthylphenyl- 740.
 D-Glucose phenylosazone 603.
 Guaiazulene 488.
 Guanine, 7-methyl- 726.
 Hematoporphyrin IX 604.
 2,4,6-Heptatrienal, 5-methyl-7-(2,6,6-trimethyl-1-cyclohexen-1-yl)-, (*E,E,E*)- 750.
all-trans-1,3,5-Heptatriene 605.
 1,4,6-Heptatrien-3-one, 1,7-bis(dimethylamino)- 192.
 Hexacene 606.
 Hexachloroethane/Zinc(II) etioporphyrin I 1127.
 1,3,5,7,9,11,13,15,17,19,21,23,25-Hexacosatridecaene, 3,7,11,16,20,24-hexamethyl-1,26-bis(2,6,6-trimethyl-1-cyclohex-1-yl)-, (*all-E*)- 331.
 2,4-Hexadiene, 2,5-dimethyl- 485.
 1,1,1,5,5,5-Hexafluoroacetylacetone 607.
 Hexahelicene 608.
 10,11,13,14,16,17-Hexahydro-5,7:20,22-dietheno-8*H*,19*H*-diindeno[2,1-*n*:1',2'-*o*]-[1,4,7,10]tetraoxacyclononadecin 609.
 9,10,11,12,13,14-Hexahydro-5,7:15,17-dietheno-2,20-heptano-*H*-cyclotetradeca[1,2-*a*:1,14-*a'*]diindene 610.
 Hexamethylbenzene/Tetracyanobenzene 611.

- 1,3,5-Hexatriene, 1,6-diphenyl- 541.
D-arabino-Hexos-2-ulose bis(phenylhydrazine) 603.
D-arabino-Hexulose phenylosazone 603.
 6-Hydroxy-7*H*-benz[*de*]anthracen-7-one 612.
 6-Hydroxybenzanthrone 612.
 2-Hydroxybenzophenone 613.
 4-Hydroxybenzophenone 614.
 7-Hydroxy-4*H*-1-benzopyran 620.
 4-Hydroxybiphenyl 615.
 4'-Hydroxy-4-biphenylamine 46.
 3-Hydroxychromone 616.
 4-Hydroxycoumarin 617.
 1-(2-Hydroxyethyl)-3,3-dimethyl-6'-nitro-spiro-(2,2'-indolin[2*H*-1]benzopyran) 618.
 3-Hydroxyflavone 619.
 7-Hydroxyflavone 620.
 1-Hydroxyfluorene 621.
 2-Hydroxyfluorene 622.
 4-Hydroxy-6-methylcoumarin 623.
 7-Hydroxy-4-methylcoumarin 624.
 2-Hydroxy-2-methyl-1-phenylpropan-1-one 625.
 2-Hydroxy-2-methyl-1-[4-(2-propyl)phenyl]propan-1-one 626.
 9-Hydroxyphenanthrene 627.
 α -Hydroxy- α -phenylacetophenone 124.
 4'-Hydroxypropiophenone 628.
 4-Hydroxypyrene 629.
 4-Hydroxyquinoline-2-carboxylic acid 643.
 4,4-(Imidocarbonyl)bis(*N,N*-dimethylaniline) monohydrochloride 93.
 Indazole, 2-*tert*-butyl-4-methyl- 235.
 Indazole, 1,3-dimethyl- 486.
 Indazole, 1-methyl- 727.
 Indazole, 2-methyl- 728.
 Indene, 2-phenyl- 890.
 Indeno[2,1-*a*]fluorene, 11,12-Dihydro-5-methyl- 725.
 Indeno[1,2-*b*]fluorene, 6,12-dihydro-3,9-dimethyl- 484.
 Indeno[2,1- α]indene, 5,10-dihydro- 420.
 1-(1*H*-Inden-3-yl)naphthalene 630.
 Indigo, *trans-N,N'*-diacetyl- 334.
 Indigo, *N,N'*-dimethyl- 487.
 Iridium(III) (chloro)phthalocyanine 276.
 Indole 631.
 Indole, 1-methyl- 729.
 Indole, 2-methyl- 730.
 Indole, 3-methyl- 731.
 Indolino[2,3-*c*]carbazole, 12*a*,12*b*-dihydro-5,8-dimethyl- 415.
 Indolo[3,2-*b*]carbazole], 5,11-dihydro-*N,N'*-dimethyl- 479.
 2-Iodoanthracene 632.
 9-Iodoanthracene 633.
 4-Iodobiphenyl 634.
 Iodo(methanol)(phthalocyaninato)rhodium(III) 635.
 1-Iodonaphthalene 636.
 2-Iodonaphthalene 637.
 β -Ionone 638.
 β -Ionylidene acetaldehyde 755.
 Iridium(III) dichlorobis(2,2'-bipyridine) 188.
 Iridium(III) dichlorobis(4,7-dimethyl-1,10-phenanthroline) 373.
 Iridium(III) dichlorobis(5,6-dimethyl-1,10-phenanthroline) 374.
 Iridium(III) dichlorobis(1,10-phenanthroline) 375.
 Iridium(III) tris(2,2'-bipyridine) 1103.
 1*H*-Isoindole-1,3(2*H*)-dione, 2-[2-[bis(phenylmethyl)amino]ethyl]- 356.
 1*H*-Isoindole-1,3(2*H*)-dione, 2-[2-(diethylamino)ethyl]- 389.
 1*H*-Isoindole-1,3(2*H*)-dione, 2-[(dimethylamino)methyl]- 459.
 1*H*-Isoindole-1,3(2*H*)-dione, 2-[(dipropylamino)methyl]- 550.
 1*H*-Isoindole-1,3(2*H*)-dione, *N*-ethyl- 577.
 1*H*-Isoindole-1,3(2*H*)-dione, 2-methyl- 745.
 1*H*-Isoindole-1,3(2*H*)-dione, 2-propyl- 927.
 Isophthalonitrile 383.
 2-Isopropylthioxanthene-9-one 639.
 Isopsoralen 640.
 Isoquinoline 641.
 Isoquinoline, 1-benzyl-, *N*-oxide 160.
 Isoquinoline, 4-bromo-, *N*-oxide 221.
 Isoquinoline, 4-bromo-1-cyano-, *N*-oxide 217.
 Isoquinoline, 1-methyl-, *N*-oxide 732.
 Isoquinoline-*N*-oxide 642.
 Isostilbene 1005.
 Kryptocyanine 393.
 Kynurenic acid 643.
 Lanthanum(III) tris(4,4,4-trifluoro-1-phenyl-1,3-butanedione) 1107.
 Limettin 449.
 Lumichrome 644.
 Lumiflavine 645.
 Lumiflavine, conjugate monoacid 646.
 Lumiflavine, negative ion 647.
all-trans-Lutein 648.
 Lutetium(III) tris(4,4,4-trifluoro-1-phenyl-1,3-butanedione) 1108.
 Lycopene, dihydroxy- 438.
all-trans-Lycopene 649.
 Lysyltryptophanylylsine 650.
 Magnesium(II) mesoporphyrin 651.
 Magnesium(II) octaethylporphyrin 652.
 Magnesium(II) phthalocyanine 653.
 Magnesium(II) tetraphenylporphyrin 654.
 Malachite Green 655.
 Menadione 737.
 Menaquinone 737.
 Mercury(II) tetraphenylporphyrin 656.
 Mesoporphyrin 657.
 Mesoporphyrin, dimethyl ester 658.
 1-(β -Methacryloxyethyl)-3,3-dimethyl-6'-nitro-spiro(indoline-2,2'-[2*H*]benzopyran) 659.
 Methane, triphenyl- 1100.
 8,11*b*-Methanocyclodeca[*cde*]naphthalene 660.
 Methoxsalen 684.
 3-Methoxyacetophenone 661.
 4-Methoxyacetophenone 662.
 3-Methoxy-7*H*-benz[*de*]anthracen-7-one 663.
 3-Methoxybenzanthrone 663.

- Methoxybenzene 63.
 4-Methoxybenzophenone 664.
 9-[2-(4-Methoxycarbonylbenzoxy)ethyl]carbazole 665.
 9-[3-(4-Methoxycarbonylbenzoxy)propyl]carbazole 666.
 4-Methoxychalcone 667.
 4-Methoxy-*N,N*-dimethylaniline/3-Chloro-2,6-diphenyl-1,4-benzoquinone 271.
 (*R*)-4-Methoxydinaphtho[2,1-*d*:1',2'-*f*]-[1,3,2]dioxaphosphepin 4-oxide 668.
 4-Methoxydinaphtho[2,1-*d*:1',2'-*f*]-[1,3,2]dioxaphosphepin 4-oxide 669.
 1-Methoxy-3,5-dinitrobenzene 515.
 9-Methoxy-7*H*-furo[3,2-*g*]1]benzopyran-7-one 684.
 8'-Methoxy-5-methyl-6'-nitrospiro[(5,6-dihydro-phenanthridine-6,2)-(2*H*)chromene] 670.
 2-Methoxy-2-methyl-1-phenyl-1-propanone 671.
 1-Methoxynaphthalene 672.
 2-Methoxynaphthalene 673.
 4-Methoxy-1-naphthonitrile 674.
 1-Methoxy-3-nitrobenzene 813.
cis,cis-1-(3'-Methoxy-5'-nitro-2'-oxo-3',5'-cyclohexadienyl)-3,4-dimethyl-1,3-pentadiene 675.
 (*E*)-1-Methoxy-3-[2-(4-nitrophenyl)ethenyl]benzene 676.
trans-3-Methoxy-4'-nitrostilbene 676.
cis-4-Methoxy-4'-nitrostilbene 677.
trans-4-Methoxy-4'-nitrostilbene 678.
trans-1-Methoxy- β -nitrostyrene 681.
 4-Methoxyphenol 679.
 (*E*)-4-[2-(4-Methoxyphenyl)ethenyl]benzonitrile 313.
 2-[1-(4-Methoxyphenyl)-hydrazinyl-2-ylidene]-1-(4-nitrophenyl)-1,3-diketobutane 680.
trans-1-(4-Methoxyphenyl)-2-nitroethylene 681.
 1-(4-Methoxyphenyl)-3-phenyl-1-propanone 891.
 3-(4-Methoxyphenyl)-1-phenyl-2-propen-1-one 667.
 4'-Methoxypropiophenone 682.
 5-Methoxypsoralen 683.
 8-Methoxypsoralen 684.
trans-4-Methoxystilbene 685.
 8-Methoxy-2,2,3-trimethyl-6-nitro-2*H*-1-benzopyran 686.
 8-Methoxy-2,2,3-trimethyl-6-nitro-2*H*-chromene 686.
 6-Methoxy-1,3,3-trimethyl-6'-nitrospiro-(2,2'-indolin[2*H*-1]benzopyran) 687.
 7'-Methoxy-1,3,3-trimethyl-6'-nitrospiro-(2,2'-indolin[2*H*-1]benzopyran) 688.
 Methyl 9-anthrylacacetate 67.
 Methyl Violet 689.
 Methyl Violet 2B 689.
 2-Methylaceanthren-1-one 690.
 3-Methylacetophenone 691.
 4-Methylacetophenone 692.
 9-Methylacridine 693.
 10-Methyl-9(10*H*)-acridinethione 694.
 4-(Methylamino)benzoic acid 695.
N-Methyl-1-anilinonaphthalene 696.
N-Methyl-2-anilinonaphthalene 697.
N-Methyl-2-anilino-6-naphthalenesulfonate ion 698.
 2-Methylanthracene 699.
 9-Methylanthracene 700.
 9-Methylanthracene-*d*₁₂ 701.
 Methyl benzoate 702.
 6-Methylbenzo[1,2-*b*:5,4-*b'*]difuran-2-carboxaldehyde 600.
 2-Methylbenzoic acid 703.
 3-Methylbenzoic acid 704.
 4-Methylbenzoic acid 705.
 2-Methylbenzophenone 706.
 4-Methylbenzophenone 707.
 Methyl-1,4-benzoquinone 708.
 (2'-Methylbenzoyl)amino-2- Δ^2 -thiazoline 709.
 (4'-Methylbenzoyl)amino-2- Δ^2 -thiazoline 710.
 Methyl benzoylformate 711.
 Methylchlorophyllide *a* 712.
 1-Methyl-4-[4-cyanostyryl]pyridinium 713.
 10-Methyl-5-deazaalloxazine-3-propanesulfonate ion 714.
 3-Methyl-5-deazalumiflavine 715.
 1-Methyldeoxybenzoin 716.
 Methyl 4-dimethylaminobenzoate 717.
N-Methyldiphenylamine 718.
 Methylene Blue, dimer 721.
 Methylene Blue cation 719.
 Methylene Blue cation, conjugate monoacid 720.
 Methylene Green cation 722.
 4,4'-Methylenebis(1,1'-biphenyl) 723.
 3,3'-Methylenebis(4-hydroxycoumarin) 380.
 2-(1-Methylethenyl)naphthalene 739.
 Methyl ethyl ketone 231.
 α -Methyl- β -ethylstyrene 724.
 5-Methyl-*endo-cis*-fluorene 725.
 7-Methyl-7*H*-furo[3,2-*g*]1]benzopyran-6-carboxaldehyde 601.
 9-Methyl-7*H*-furo[3,2-*g*]1]benzopyran-6-carboxaldehyde 602.
 7-Methylguanine 726.
 1-Methylindazole 727.
 2-Methylindazole 728.
 1-Methylindole 729.
 2-Methylindole 730.
 3-Methylindole 731.
 1-Methylisoquinoline-*N*-oxide 732.
 Methylum, triphenyl- 1101.
 1-Methylnaphthalene 733.
 2-Methylnaphthalene 734.
 1-Methylnaphthalene/Pyromellitic dianhydride 735.
 Methyl 2-naphthoate 736.
 2-Methyl-1,4-naphthoquinone 737.
 1-Methyl-1-(1-naphthyl)ethylene 738.
 1-Methyl-1-(2-naphthyl)ethylene 739.
 Methyl 1-naphthyl ketone 5.
 Methyl 2-naphthyl ketone 6.
 Methyl-1-naphthylphenylgermane 740.
 5-Methyl-6'-nitrospiro[(5,6-dihydrophenanthridine-6,2)-(2*H*)chromene] 741.
 1-Methyl-4-[4-nitrostyryl]pyridinium 742.
 5-Methylphenazinium methyl sulfate 743.
N-Methylphenazonium methosulfate 743.
 4-Methylphenol 304.
 10-Methylphenothiazine 744.

- N*-Methylphenothiazine 744.
 1-Methyl-2-phenylacetylene 928.
 Methyl phenylglyoxylate 711.
N-Methylphthalimide 745.
 4-(2-Methylpropionyloxy)benzophenone 746.
 Methyl pyropheophorbide *a* 747.
all-trans-Methyl retinoate 748.
 2-Methylthioxanthene-9-one 749.
 (*E,E,E*)-5-Methyl-7-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2,4,6-heptatrienal 750.
 (*E,E,E*)-6-Methyl-8-(2,6,6-trimethyl-1-cyclohexen-1-yl)-3,5,7-octatrien-2-one 751.
 (*E,E,Z*)-6-Methyl-8-(2,6,6-trimethyl-1-cyclohexen-1-yl)-3,5,7-octatrien-2-one 752.
 (*E,Z,E*)-6-Methyl-8-(2,6,6-trimethyl-1-cyclohexen-1-yl)-3,5,7-octatrien-2-one 753.
 (*E,Z,Z*)-6-Methyl-8-(2,6,6-trimethyl-1-cyclohexen-1-yl)-3,5,7-octatrien-2-one 754.
 (*E,E*)-3-Methyl-5-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2,4-pentadienal 755.
 (*E,Z*)-3-Methyl-5-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2,4-pentadienal 756.
 (*Z,E*)-3-Methyl-5-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2,4-pentadienal 757.
 (*Z,Z*)-3-Methyl-5-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2,4-pentadienal 758.
 3-Methyl-3,4,5-triphenyl-2(3*H*)-furanone 759.
 1-Methyl-L-tryptophan 760.
 β-Methylumbelliferone 624.
 5-Methyluracil 1072.
 Methyl vinyl ketone 761.
 1-Methyl-2-vinylnaphthalene 762.
 2-Methyl-3-vinylnaphthalene 763.
 Michler's ketone 190.
 α-NPO 805.
 Naphthacene 1021.
 1-Naphthaldehyde 764.
 2-Naphthaldehyde 765.
 2-Naphthalenamine, conjugate acid 767.
 2-Naphthalenamine 766.
 Naphthalene 768.
 Naphthalene, 2-amino- 766.
 Naphthalene, 1-amino-4-nitro- 53.
 Naphthalene, 1-anilino- 62.
 Naphthalene, 1-bromo- 224.
 Naphthalene, 2-bromo- 225.
 Naphthalene, 1-(bromomethyl)- 223.
 Naphthalene, 1,1'-(1,4-butanediyl)bis- 505.
 Naphthalene, 1-chloro- 279.
 Naphthalene, 2-chloro- 280.
 Naphthalene, 1-(chloromethyl)- 278.
 Naphthalene, 1-(2-chlorophenyl)- 282.
 Naphthalene, 1-(4-chlorophenyl)- 283.
 Naphthalene, 1-cyano- 314.
 Naphthalene, 1-cyano-4-methoxy- 674.
 Naphthalene, 1-(1-cyclohepten-1-yl)- 319.
 Naphthalene, 2-(1-cyclohepten-1-yl)- 320.
 Naphthalene, 1-(1-cyclohexen-1-yl)- 323.
 Naphthalene, 1-(1-cycloocten-1-yl)- 325.
 Naphthalene, 1-(1-cyclopenten-1-yl)- 328.
 Naphthalene, 2-(1-cyclopenten-1-yl)- 329.
 Naphthalene, 1,4-dibromo- 361.
 Naphthalene, 1,4-dichloro- 377.
 Naphthalene, 1,4-dicyano- 385.
 Naphthalene, 1,2-dihydro-3-phenyl- 429.
 Naphthalene, 1,2-dihydro-4-(phenylthio)- 900.
 Naphthalene, 2,3-dihydroxy-, conjugate base 443.
 Naphthalene, 1,5-dihydroxy- 439.
 Naphthalene, 1,8-dihydroxy- 440.
 Naphthalene, 2,3-dihydroxy- 441.
 Naphthalene, 2,7-dihydroxy- 442.
 Naphthalene, 1,4-dimethoxy- 451.
 Naphthalene, 1,2-dimethyl- 489.
 Naphthalene, 2,3-dimethyl- 490.
 Naphthalene, 2,7-dimethyl- 491.
 Naphthalene, 1-dimethylamino-4-nitro- 460.
 Naphthalene, 1-(3,3-dimethyl-1-buten-2-yl)- 478.
 Naphthalene, 2-(3,3-dimethyl-1-butenyl)-, (*Z*)- 494.
 Naphthalene, 2-(3,3-dimethyl-1-butenyl)-, (*E*)- 495.
 Naphthalene, 1,2-dinitro- 517.
 Naphthalene, 1,4-dinitro- 518.
 Naphthalene, 1,8-dinitro- 519.
 Naphthalene, 2-(2,6-dioxacyclohexyl)- 780.
 Naphthalene, 1,4-diphenyl- 542.
 Naphthalene, 1,5-diphenyl- 543.
 Naphthalene, 1,4-diphenyl-2,3-dibenzoyl-1,4-epoxy-1,4-dihydro- 538.
 Naphthalene, 1,1'-(1,2-ethanediyl)bis- 507.
 Naphthalene, 2-ethoxy- 567.
 Naphthalene, 1-ethyl- 574.
 Naphthalene, 2-ethyl- 575.
 Naphthalene, 1-hydroxy- 786.
 Naphthalene, 2-hydroxy- 787.
 Naphthalene, 1-(1*H*-inden-3-yl)- 630.
 Naphthalene, 1-iodo- 636.
 Naphthalene, 2-iodo- 637.
 Naphthalene, 1-methoxy- 672.
 Naphthalene, 2-methoxy- 673.
 Naphthalene, 1-methyl- 733.
 Naphthalene, 2-methyl- 734.
 Naphthalene, 1,1'-methylenebis- 508.
 Naphthalene, 2-(1-methylethenyl)- 739.
 Naphthalene, 1-methyl-2-vinyl- 762.
 Naphthalene, 2-methyl-3-vinyl- 763.
 Naphthalene, 1-nitro- 819.
 Naphthalene, 2-nitro- 820.
 Naphthalene, 1-phenyl- 893.
 Naphthalene, 2-phenyl- 894.
 Naphthalene, 1-(1-phenylethenyl)- 800.
 Naphthalene, 1-[(phenylmethyl)sulfonyl]methyl]- 892.
 Naphthalene, 1,1'-(1,3-propanediyl)bis- 512.
 Naphthalene, 2,2'-(1,3-propanediyl)bis- 513.
 Naphthalene, 2-vinyl- 1119.
 Naphthalene-*d*₈ 769.
 Naphthalene/Chloranil 770.
 Naphthalene/3-Chloro-2,6-diphenyl-1,4-benzoquinone 272.
 Naphthalene/Pyromellitic dianhydride 771.

- Naphthalene/Tetrachlorophthalic anhydride 772.
 Naphthalene-*d*₈/Tetrachlorophthalic anhydride 773.
 1-Naphthalenecarbothioic acid, *O*-ethyl ester 774.
 1-Naphthalenecarbothioic acid, *O*-(2-phenylethyl) ester 888.
 2-Naphthalenecarbothioic acid, *O*-(2-phenylethyl) ester 889.
 1-Naphthalenecarboxylate ion 781.
 2-Naphthalenecarboxylate ion 782.
 2-Naphthalenecarboxylic acid, methyl ester 736.
 1-Naphthalenecarboxylic acid 784.
 2-Naphthalenecarboxylic acid 785.
 1,4-Naphthalenedicarbonitrile 385.
 1,5-Naphthalenediol 439.
 1,8-Naphthalenediol 440.
 2,3-Naphthalenediol 441.
 2,7-Naphthalenediol 442.
 2-Naphthalenesulfonate ion, 6-(methylphenylamino)- 698.
 2-Naphthalenesulfonic acid, sodium salt 775.
 1-Naphthalenesulfonic acid 776.
anti-[2.2](1,4)-Naphthalenophane 777.
syn-[2.2](1,4)-Naphthalenophane 778.
 2-(1-Naphthalenyl)-1,3-dioxane 779.
 2-(2-Naphthalenyl)-1,3-dioxane 780.
 β-Naphthiazoline, 2-benzoyl-*N*-methyl- 152.
 1-Naphthoate ion 781.
 2-Naphthoate ion 782.
 Naphtho[*de*-2.3.4]bicyclo[3.2.2]nona-2,6,8-triene, 8-benzoyl- 153.
 Naphtho[1,2,3,4-*def*]chrysene 783.
 Naphtho[1,2-*g*]chrysene 352.
 1-Naphthoic acid 784.
 2-Naphthoic acid 785.
 1-Naphthol 786.
 2-Naphthol 787.
 2-Naphthol/Eosin 559.
 1-Naphthol/Triethylamine 788.
 2-Naphthol/Triethylamine 789.
 1-Naphtholate ion 797.
 2-Naphtholate ion 798.
 1-Naphthonitrile 314.
 1,4-Naphthoquinone, 2-methyl- 737.
 Naphtho[1,8-*bc*]thiopyran-6-sulfonamide, *N,N*-dibutyl-2,3-dihydro-3-oxo-2-(3-oxonaphtho[1,2-*b*]thien-2(3*H*)-ylidene), (*E*)- 362.
 Naphtho[1,8-*bc*]thiopyran-6-sulfonamide, *N,N*-dibutyl-2,3-dihydro-3-oxo-2-(3-oxonaphtho[1,2-*b*]thien-2(3*H*)-ylidene), (*Z*)- 363.
 9-Naphthoylanthracene 790.
 1-Naphthyl acetate 791.
 2-Naphthylamine, conjugate acid 767.
 2-Naphthylamine, *N,N*-dimethyl-, conjugate acid 493.
 2-Naphthylamine, *N,N*-dimethyl- 492.
 1-Naphthylamine, *N*-methyl-*N*-phenyl- 696.
 2-Naphthylamine, *N*-methyl-*N*-phenyl- 697.
 2-Naphthylamine, *N*-phenyl- 895.
 2-Naphthylamine 766.
N-1-Naphthylaniline 62.
N-2-Naphthylaniline 895.
 2-(1-Naphthyl)benzoxazole 792.
 8-(1-Naphthyl)-6,7-dihydro-5*H*-benzocycloheptene 793.
 1-Naphthyl diisopropyl borate 794.
 2-Naphthyl diisopropyl borate 795.
trans-1-(1-Naphthyl)-2-(2-naphthyl)ethylene 796.
 1-Naphthyloxide ion 797.
 2-Naphthyloxide ion 798.
 2-(2-Naphthyloxy)-3,5,5-trimethyl-2-cyclohexen-1-one 799.
 1-(1-Naphthyl)-1-phenylethylene 800.
trans-1-(1-Naphthyl)-2-phenylethylene 801.
trans-1-(2-Naphthyl)-2-phenylethylene 802.
 2-(1-Naphthyl)-5-phenyl-1,3,4-oxadiazole 803.
 2-(2-Naphthyl)-5-phenyl-1,3,4-oxadiazole 804.
 2-(1-Naphthyl)-5-phenyloxazole 805.
 Neo-alloocimene 806.
 Neomethylene Blue cation 807.
all-trans-Neurosporene 808.
 Neutral Red cation 809.
 Neutral Red cation, conjugate diacid 810.
 5-Nitroacenaphthene 811.
 4-Nitroaniline 812.
 4-Nitroaniline/2-(Dimethylamino)pyridine 469.
 3-Nitroanisole 813.
 4-Nitroanisole 814.
 9-Nitroanthracene 815.
 10-Nitro-9-anthryl phenyl ketone 154.
 4-Nitrobiphenyl 816.
 2-[2-(5-Nitro-2-furanyl)]ethenylquinoline 817.
 5-Nitro-2-furoic acid 818.
 1-Nitronaphthalene 819.
 2-Nitronaphthalene 820.
 4-Nitro-1-naphthylamine 53.
 (*E*)-4-[2-(4-Nitrophenyl)ethenyl]benzenamine 54.
 (*E*)-1-Nitro-3-(2-phenylethenyl)benzene 823.
 1-(4-Nitrophenyl)-2-[(4-methoxyphenyl)hydrazono]-1,2,3-butanetrione 680.
 4-Nitropyridine-*N*-oxide 821.
 4-Nitroquinoline-*N*-oxide 822.
 5-Nitro-2-(2-quinolyloethenyl)furan 817.
trans-3-Nitrostilbene 823.
trans-4-Nitrostilbene 824.
trans-4-Nitrostilbene/Zinc(II) etioporphyrin I 1128.
 4-Nitro-*p*-terphenyl 825.
 2-Nitrothiophene 826.
 4-Nitrotoluene/Zinc(II) etioporphyrin I 1129.
 4-Nitroveratrole 452.
 1,3,6,8-Nonatetraen-5-one, 1,9-bis(dimethylamino)- 194.
 1,4,6,8-Nonatetraen-3-one, 1,9-bis(dimethylamino)- 195.
 2,5-Norbornadiene, 2,3-dibenzoyl- 353.
 1,3,5,7,9,11,13,15,17-Octadecanonaene, 3,7,12,16-tetramethyl-1,18-bis-(2,6,6-trimethyl-1-cyclohexen-1-yl)-, (*all-E*)- 254.
 Octaethylporphinatopalladium(II) 827.
 Octaethylporphinatotin(IV) dichloride 828.
 8,9,10,11,12,13,14,15-Octahydro-5,7:16,18-dietheno-2,21-octanocyclopentadeca[1,2-*a*:1,15-*a'*]diindene 829.
 (*all-E*)-3,7,11,15,20,24,28,32-Octamethyl-1,34-bis-(2,6,6-trimethyl-1-cyclohexen-1-yl)-tritriacontahaptadecaene 830.

- 1,3,5,7-Octatetraene, 1,8-diphenyl- 544.
 2,4,6-Octatriene, 2,6-dimethyl-, (*E,Z*)- 806.
 3,5,7-Octatrien-2-one, 8-(dimethylamino)- 462.
 3,5,7-Octatrien-2-one, 6-methyl-8-(2,6,6-trimethyl-1-cyclohexen-1-yl)-, (*E,E,E*)- 751.
 3,5,7-Octatrien-2-one, 6-methyl-8-(2,6,6-trimethyl-1-cyclohexen-1-yl)-, (*E,E,Z*)- 752.
 3,5,7-Octatrien-2-one, 6-methyl-8-(2,6,6-trimethyl-1-cyclohexen-1-yl)-, (*E,Z,E*)- 753.
 3,5,7-Octatrien-2-one, 6-methyl-8-(2,6,6-trimethyl-1-cyclohexen-1-yl)-, (*E,Z,Z*)- 754.
 Orotate ion 831.
 Orotic acid 832.
 Osmium(II) tris(2,2'-bipyridine) 1104.
 2,2'-Oxacarboxyanine iodide, 3,3'-diethyl- 398.
 1,3,4-Oxadiazole, 2-(4-biphenyl)-5-phenyl- 181.
 1,3,4-Oxadiazole, 2,5-di(1-naphthyl)- 509.
 1,3,4-Oxadiazole, 2,5-diphenyl- 545.
 1,3,4-Oxadiazole, 2-(1-naphthyl)-5-phenyl- 803.
 1,3,4-Oxadiazole, 2-(2-naphthyl)-5-phenyl- 804.
 7-Oxa-2,3-dibenzoylbicyclo[2.2.1]hepta-2,5-diene 833.
 7-Oxa-2,3-dibenzoylbicyclo[2.2.1]hept-2-ene 834.
 2,2'-Oxadibenzocyanine iodide, 3,3'-diethyl- 399.
 Oxazole, 2-(4-biphenyl)-5-phenyl- 182.
 Oxazole, 2,5-di(4-biphenyl)- 357.
 Oxazole, 2,5-diphenyl- 546.
 Oxazole, 2-(1-naphthyl)-5-phenyl- 805.
 Oxazole, 2,2'-(1,4-phenylene)bis[5-(4-butoxyphenyl)- 884.
 Oxazole, 2,2'-(1,4-phenylene)bis[5-phenyl- 885.
 Oxirane, *cis*-2,3-dimethyl-2,3-di-(2-naphthyl)- 480.
 Oxirane, *trans*-2,3-dimethyl-2,3-di-(2-naphthyl)- 481.
 Oxirane, *cis*-2,3-di-(2-naphthyl)- 510.
 Oxirane, *trans*-2,3-di-(2-naphthyl)- 511.
 2-(3-Oxobenzo[*b*]selenophen-2(3*H*)-ylidene)benzo[*b*]selenophen-3(2*H*)-one 401.
 2-(3-Oxobenzo[*b*]thien-2(3*H*)-ylidene)benzo[*b*]thiophen-3(2*H*)-one 1062.
 7-Oxo-7*H*-furo[3,2-*g*]1]benzopyran-6-carboxylic acid, ethyl ester 246.
cis-2-(1-Oxonaphtho[2,1-*b*]thiophen-2(1*H*)-ylidene)naphtho[1,8-*bc*]thiopyran-3(2*H*)-one 849.
trans-2-(1-Oxonaphtho[2,1-*b*]thiophen-2(1*H*)-ylidene)naphtho[1,8-*bc*]thiopyran-3(2*H*)-one 850.
 Oxonine cation 835.
 Oxonine cation, conjugate monoacid 836.
 4-(1-Oxo-3-phenyl-2-propenyl)benzonitrile 310.
 Oxotitanium(IV) tetraphenylporphyrin 837.
 PMDA 953.
 POPOP 885.
 PPO 546.
 Palladium(II) etioporphyrin I 838.
 Palladium(II) octaethylporphyrin 827.
 Palladium(II) tetrabenzoporphyrin 839.
 Palladium(II) tetrakis(carboxyphenyl)porphyrin 840.
 Palladium(II) tetrakis(4-*N*-methylpyridyl)porphyrin 841.
 Palladium(II) tetrakis(*p*-sulfonatophenyl)porphyrin 842.
 Palladium(II) tetraphenylporphyrin 843.
 [2.2]Paracyclophane 844.
 Pentacene 845.
 Pentacyclo[18.2.2.2^{9,12}.0^{4,15}.0^{6,17}]hexacos-4,6(17),9,11,15,20,22,23,25-nonane-10,25-dicarbonitrile 846.
 1,3,6,8,10,12,14-Pentadecaheptaen-5-one, 1,15-bis(dimethylamino)-, *all-trans*- 196.
 2,4-Pentadienal, 3-methyl-5-(2,6,6-trimethyl-1-cyclohexen-1-yl)-, (*E,E*)- 755.
 2,4-Pentadienal, 3-methyl-5-(2,6,6-trimethyl-1-cyclohexen-1-yl)-, (*E,Z*)- 756.
 2,4-Pentadienal, 3-methyl-5-(2,6,6-trimethyl-1-cyclohexen-1-yl)-, (*Z,E*)- 757.
 2,4-Pentadienal, 3-methyl-5-(2,6,6-trimethyl-1-cyclohexen-1-yl)-, (*Z,Z*)- 758.
 Pentahelicene 847.
 1,5-Pentanedione, 2,4-dimethyl-1,5-diphenyl- 482.
 2,4-Pentanedione, 3-phenyl- 872.
 2,4-Pentanedione, 1,1,1-trifluoro- 1080.
 2,4-Pentanedione 11.
 Pentaphene 848.
cis-Perinaphthothioindigo 849.
trans-Perinaphthothioindigo 850.
 1,4-Perinaphthyldiyl 851.
 Perylene 852.
 Perylene/Chloranil 853.
 1*H*-Phenylene, 2,3-dihydro- 426.
 Phenanthrazine 1019.
 9-Phenanthrenamine 56.
 Phenanthrene 854.
 Phenanthrene, 3-acetyl-9,10-epoxy-9,10-dihydro- 15.
 Phenanthrene, 9-amino- 56.
 Phenanthrene, 9-bromo- 227.
 Phenanthrene, 3,4-dihydro- 427.
 Phenanthrene, 9,10-dihydro- 428.
 Phenanthrene, 9-ethyl- 576.
 Phenanthrene, 9-hydroxy- 627.
 Phenanthrene-*d*₁₀ 855.
 Phenanthrene/Pyromellitic dianhydride 856.
 Phenanthrene/Tetrachlorophthalic anhydride 857.
 Phenanthrene-*d*₁₀/Tetrachlorophthalic anhydride 858.
 Phenanthridine 859.
 6(5*H*)-Phenanthridinone 860.
 Phenanthro[9,10-*b*]furan-2(3*H*)-one, 3-benzyl-3-phenyl- 162.
 9-Phenanthrol 627.
 4,7-Phenanthroline, 3,8-dimethyl- 499.
 1,10-Phenanthroline 861.
 1,7-Phenanthroline 862.
 1,8-Phenanthroline 863.
 1,9-Phenanthroline 864.
 4,7-Phenanthroline 865.
 Phenanthro[3,4-*c*]phenanthrene 608.
 Phenazine 866.
 Phenazine, conjugate monoacid 867.
 Phenazine, 5,10-dihydro-5,10-dimethyl- 416.
 Phenazine, 5,10-dihydro-5,10-diphenyl- 419.
 Phenazine, 5,10-dihydro-5-methyl-10-phenyl- 422.
 Phenazine methosulfate 743.
 Phenazinium, 3,7-diamino-2,8-dimethyl-5-phenyl- 990.
 Phenazinium, 3,7-diamino-2,8-dimethyl-5-phenyl-, conjugate diacid 991.

- Phenazinium, 3,7-diamino-2,8-dimethyl-5-phenyl-, conjugate monoacid 992.
- Phenol 868.
- Phenol, 2-[1-(butylimino)ethyl]-5-methyl- 234.
- Phenol, 4-methoxy- 679.
- Phenol, 4-methyl- 304.
- Phenol, 4-phenyl- 615.
- Phenosafuranin 869.
- 5-Phenoselenazinium, 3,7-diamino- 993.
- 5-Phenoselenazinium, 3,7-diamino-, conjugate monoacid 994.
- Phenothiazine 870.
- Phenothiazine, 2-chloro-10-dimethylaminopropyl- 292.
- 10*H*-Phenothiazine, 10-methyl- 744.
- 10*H*-Phenothiazine-10-propanamine, 2-chloro-*N,N*-dimethyl- 292.
- 10*H*-Phenothiazine-10-propanamine, *N,N*-dimethyl- 924.
- Phenothiazin-5-ium, 3,7-bis(dimethylamino)- 719.
- Phenothiazin-5-ium, 3,7-bis(dimethylamino)-, conjugate monoacid 720.
- Phenothiazin-5-ium, 3,7-bis(dimethylamino)-4-nitro- 722.
- Phenothiazin-5-ium, 3,7-diamino- 1063.
- Phenoxazine 871.
- Phenoxazin-5-ium, 3,7-bis(diethylamino)- 1024.
- Phenoxazin-5-ium, 3,7-diamino-, conjugate monoacid 836.
- Phenoxazin-5-ium, 3,7-diamino- 835.
- α -Phenylacetophenone 332.
- 3-Phenylacetylacetone 872.
- Phenylacetylene 873.
- 9-Phenylacridan 874.
- N*-Phenylacridan 875.
- Phenylalanine 876.
- 4-(Phenylamino)-7*H*-benz[*de*]anthracen-7-one 877.
- 9-Phenylanthracene 878.
- 4-Phenylbenzaldehyde 175.
- 4-Phenylbenzophenone 879.
- 9-Phenyl-9*H*-9-bisfluorene 880.
- 1-Phenyl-1,3-butanedione 146.
- 1-Phenyl-1-butene 578.
- 1-Phenylcyclohexene 881.
- 1-Phenyldeoxybenzoin 882.
- 5-Phenyl-5*H*-dibenzobismole 880.
- 5-Phenyl-5*H*-dibenzophosphole 897.
- 5-Phenyl-5*H*-dibenzostibole 899.
- 2,2'-(1,4-Phenylene)bisbenzoxazole 883.
- 2,2'-(1,4-Phenylene)bis[5-(4-butoxyphenyl)oxazole] 884.
- 2,2'-(1,4-Phenylene)bis[5-phenyloxazole] 885.
- 1,4-Phenylenediamine, *N,N'*-dimethyl-*N,N'*-diphenyl- 483.
- p*-Phenylenediamine, *N,N*-dimethyl- 500.
- p*-Phenylenediamine, *N,N'*-diphenyl- 547.
- p*-Phenylenediamine, *N,N'*-diphenyl-, conjugate acid 548.
- p*-Phenylenediamine, *N,N,N',N'*-tetramethyl- 1049.
- (*E*)-4-(2-Phenylethenyl)benzonitrile 316.
- 2-(1-Phenylethenyl)biphenyl 180.
- (*E*)-2-(2-Phenylethenyl)naphthalene 802.
- O*-(2-Phenylethyl) 4-(dimethylamino)benzenecarbothioate 886.
- O*-(2-Phenylethyl) 4-methoxybenzenecarbothioate 887.
- O*-(2-Phenylethyl) 1-naphthalenecarbothioate 888.
- O*-(2-Phenylethyl) 2-naphthalenecarbothioate 889.
- 2-Phenylindene 890.
- β -Phenyl-4'-methoxypropiofenone 891.
- 1-[(Phenylmethyl)sulfonyl]methyl]naphthalene 892.
- 1-Phenylnaphthalene 893.
- 2-Phenylnaphthalene 894.
- N*-Phenyl-1-naphthylamine 62.
- N*-Phenyl-2-naphthylamine 895.
- 2-Phenyl-2-norbornene 896.
- 2-Phenyl-2-pentene 724.
- 4-Phenylphenol 615.
- 9-Phenyl-9*H*-9-phosphafluorene 897.
- 9-Phenylproflavine, conjugate monoacid 898.
- 1-Phenyl-1-propyne 928.
- Phenyl-4-pyridyl ketone 157.
- 9-Phenyl-9*H*-9-stibafluorene 899.
- 1-Phenylthio-3,4-dihydronaphthalene 900.
- Phenyl[2,4,6-tris(1-methylethyl)phenyl]methanone 1084.
- (1-Phenylvinyl)acenaphthene 4.
- Pheophytin *a* 901.
- Pheophytin *b* 902.
- 3-Phorbinepropanoic acid, 9-ethenyl-14-ethyl-4,8,13,18-tetramethyl-20-oxo-, methyl ester, (3*SS*-trans)- 747.
- 3-Phorbinepropanoic acid, 9-ethenyl-14-ethyl-21-(methoxycarbonyl)- 901.
- 3-Phorbinepropanoic acid, 9-ethenyl-14-ethyl-13-formyl-21-(methoxycarbonyl)- 902.
- 9*H*-9-Phosphafluorene, 9-phenyl- 897.
- Photoprotoporphyrin isomer "A", dimethyl ester 903.
- Photoprotoporphyrin isomer "B", dimethyl ester 904.
- Phthalazine 905.
- Phthalimide, 3,6-Dihydroxy- 444.
- Phthalimide, 3-amino-*N*-methyl- 50.
- Phthalimide, 3,6-diamino- 336.
- Phthalimide, *N*-[(dipropylamino)methyl]- 550.
- Phthalimide, *N*-methyl- 745.
- Phthalimide, *N*-propyl- 927.
- Phthalocyanatocopper(II) 297.
- Phthalocyaninatobis(pyridine)ruthenium(II) 906.
- Phthalocyaninatomagnesium(II) 653.
- Phthalocyaninatozinc(II) 1130.
- Phthalocyanine 907.
- Phthalonitrile 382.
- 15-*cis*-Phytoene 908.
- all-trans*-Phytoene 909.
- Picene 910.
- Pinacyanol chloride 392.
- 2-Piperidinoanthraquinone 911.
- Pivalophenone 912.
- Poly[oxy[2-(1-pyrenylmethyl)-1,3-propanediyl]oxy(1,4-dioxo-1,4-butanediyl)] 913.
- Poly[oxy[2-(1-pyrenylmethyl)-1,3-propanediyl]oxy(1,4-dioxo-1,4-butanediyl)]/1,4-Dicyanobenzene 914.
- Poly[oxy[2-(1-pyrenylmethyl)-1,3-propanediyl]oxy(1,4-dioxo-1,6-hexanediyl)] 915.
- Poly[oxy[2-(1-pyrenylmethyl)-1,3-propanediyl]oxy(1,4-dioxo-1,6-hexanediyl)]/1,4-Dicyanobenzene 916.

- Poly(phenylisopropenyl ketone) 917.
Poly(4-vinylbenzophenone) 918.
Poly(2-vinylnaphthalene) 919.
Porphinatozinc(II) 1131.
21*H*,23*H*-Porphine, 2,3-dihydro-5,10,15,20-tetraphenyl- 1053.
21*H*,23*H*-Porphine, 5,10,15,20-tetraethyl- 1026.
21*H*,23*H*-Porphine, 2,8,12,18-tetraethyl-2,3-dihydro-3,7,13,17-tetramethyl- 579.
Porphine, tetrakis(2,6-dimethyl-4-sulfonatophenyl)- 1032.
Porphine, tetrakis(2-*N*-methylpyridyl)- 1033.
Porphine, tetrakis(3-*N*-methylpyridyl)- 1034.
Porphine, tetrakis(4-*N*-methylpyridyl)- 1035.
Porphine, tetrakis(2-pyridyl)-, conjugate diacid 1039.
Porphine, tetrakis(3-pyridyl)-, conjugate diacid 1040.
Porphine, tetrakis(4-pyridyl)-, conjugate diacid 1041.
Porphine, tetrakis(2-pyridyl)- 1036.
Porphine, tetrakis(3-pyridyl)- 1037.
Porphine, tetrakis(4-pyridyl)- 1038.
Porphine, tetrakis(*p*-sulfonatophenyl)- 1043.
Porphine, tetrakis(trimethylaminophenyl)- 1044.
Porphine, *meso*-tetraphenyl- 1057.
21*H*,23*H*-Porphine 920.
21*H*,23*H*-Porphine-2,18-dipropionic acid, 2,3-dihydro-3,3,7,12,17-pentamethyl- 213.
Porphine-6,7-dipropionic acid, 1,3,5,8-tetramethyl-2,4-divinyl- 930.
Porphyrin, tetraphenyl-, dianion 1058.
PrOH/Dibenz[*a,j*]acridine 338.
Proflavine 921.
Proflavine, conjugate diacid 922.
Proflavine, conjugate monoacid 923.
Promazine 924.
1,3-Propandiol diacetate, 2-(1-pyrenylmethyl)- 949.
Propane, 1,3-di(9-phenanthryl)- 526.
Propanedinitrile, 2,2',2''-(4,5-dihydroxy-2-oxo-4-cyclopentene-1,3-diylidene)bis-, dianion 306.
1,3-Propanedione, 1,3-diphenyl- 355.
4-Propanenitrile, 3-[[4-[(2,4-dinitrophenyl)azo]phenyl]-(2-hydroxyethyl)amino]- 521.
Propanoic acid, 2-methyl-4-benzoylphenyl ester 746.
1-Propanone, 2-acetoxy-2-methyl-1-phenyl- 10.
1-Propanone, 1-(4-biphenyl)-3-chloro- 179.
1-Propanone, 3-chloro-1-(2-naphthyl)- 281.
1-Propanone, 2,2-dimethyl-1-phenyl- 912.
1-Propanone, 1,2-diphenyl 716.
1-Propanone, 2-hydroxy-2-methyl-1-phenyl- 625.
1-Propanone, 2-hydroxy-2-methyl-1-[4-(2-propyl)phenyl]- 626.
1-Propanone, 1-(4-hydroxyphenyl)- 628.
1-Propanone, 2-methoxy-2-methyl-1-phenyl- 671.
1-Propanone, 1-(4-methoxyphenyl)- 682.
1-Propanone, 1-(4-methoxyphenyl)-3-phenyl- 891.
2-Propen-1-one, 1-(9-anthracenyl)- 87.
2-Propen-1-one, 3-[4-(dimethylamino)phenyl]-1-phenyl- 455.
2-Propen-1-one, 1,3-diphenyl- 258.
2-Propen-1-one, 3-(4-methoxyphenyl)-1-phenyl- 667.
Propionate ion, 2-amino-3-(4-methoxy-6-benzothiazolyl)- 47.
9-Propionylantracene 925.
Propiophenone 926.
Propiophenone, 2-(*N*-formylamino)-3'-amino- 55.
Propiophenone, 4'-hydroxy- 628.
N-Propylphthalimide 927.
1-Propynylbenzene 928.
Protoporphyrin 929.
Protoporphyrin IX 930.
Protoporphyrin IX, dimethyl ester 931.
Protoporphyrin IX, dimethyl ester, conjugate diacid 932.
Protoporphyrin IX, dimethyl ester, conjugate monoacid 933.
Pseudoisocyanine 394.
Psoralen 934.
Psoralen, 4'-aminomethyl-4,5',8-trimethyl- 51.
Psoralen, 3-carbethoxy- 246.
Psoralen, 4',5'-dihydro- 430.
Psoralen, 4',5'-dihydro-3-carbethoxy- 411.
Psoralen, 5-methoxy- 683.
Psoralen, 8-methoxy- 684.
Psoralen, 4,5',8-trimethyl- 1091.
4',5'-Psoralen-thymine photo adduct 935.
4-[3*H*]Pteridinone, 2-amino- 57.
Pterin 57.
1*H*-Purin-2-amine 58.
Purine 936.
Purine, 2-amino- 58.
Purine, 2-(dimethylamino)- 466.
Pyranthrene 937.
Pyrazine 938.
1*H*,5*H*-Pyrazolo[1,2-*a*]pyrazole-1,5-dione, 2,3,6,7-tetramethyl- 1047.
Pyrene 939.
Pyrene, 1-bromo- 228.
Pyrene, 4-hydroxy- 629.
Pyrene, 1,3,6,8-tetraphenyl- 1059.
Pyrene-*d*₁₀ 940.
Pyrene/Chloranil 941.
Pyrene/Diethylaniline 942.
Pyrene/Tetracyanobenzene, 1:1 943.
Pyrene/Tetracyanobenzene, 2:1 944.
1-Pyrenecarboxaldehyde 945.
Pyrene-1-sulfonate 946.
Pyrene-3-sulfonate 947.
4-Pyrenol 629.
[2,2](2,7)-Pyrenophane 948.
2-(1-Pyrenylmethyl)-1,3-propandiol diacetate 949.
2-(1-Pyrenylmethyl)-1,3-propandiol diacetate/1,4-Dicyanobenzene 950.
Pyridine, 2-amino- 59.
Pyridine, 4-amino- 60.
Pyridine, 2-benzoyl- 155.
Pyridine, 3-benzoyl- 156.
Pyridine, 4-benzoyl- 157.
Pyridine, 2-(dimethylamino)- 467.
Pyridine, 4-(dimethylamino)- 468.

- J. Phys. Chem. Ref. Data, Vol. 15, No. 1, 1986

- Spiro-(2,2'-indolin[2*H*-1]benzopyran),
1,3,3-trimethyl-6'-nitro- 1088.
- Spiro(indoline-2,2'-[2*H*]benzopyran),
1-(β -methacryloxyethyl)-3,3-dimethyl-6'-nitro- 659.
- Spiro[isobenzofuran-1(3*H*),9'-[9*H*]xanthen]-3-one,
2',4',5',7'-tetrabromo-3',6'-dihydroxy-, disodium salt 556.
- Spiroisobenzofuran-1(3*H*),9'[9*H*]xanthen-3-one,
2',4',5',7'-tetrabromo-3',6'-dihydroxy- 562.
- Spiroisobenzofuran-1(3*H*),9'[9*H*]xanthen-3-one,
3',6'-dihydroxy-2',4',5',7'-tetraiodo-, ion(2-) 566.
- Spiro[isobenzofuran-1(3*H*),9'-[9*H*]xanthen-3-one,
3',7'-dihydroxy- 591.
- Spiro[isobenzofuran-1(3*H*),9'-[9*H*]xanthen-3-one,
3',7'-dihydroxy-, conjugate monoacid 592.
- Spiro[isobenzofuran-1(3*H*),9'-[9*H*]xanthen-3-one,
3',7'-dihydroxy-, ion(1-) 593.
- Spiro[isobenzofuran-1(3*H*),9'-[9*H*]xanthen]-3-one,
4,7-dichloro-3',6'-dihydroxy-2',4',5',7'-tetraiodo, di-
potassium salt 988.
- 9*H*-9-Stannafluorene, 9,9-diethyl- 402.
- 9*H*-9-Stibafluorene, 9-phenyl- 899.
- Stilbene 1000.
- Stilbene, *trans*-4-amino-4'-nitro- 54.
- Stilbene, *trans*-4-bromo- 229.
- Stilbene, *cis*-4-chloro- 288.
- Stilbene, *trans*-4-chloro- 289.
- Stilbene, *trans*-4-cyano- 316.
- Stilbene, *trans*-4-cyano-4'-dimethylamino- 311.
- Stilbene, *trans*-4-cyano-4'-methoxy- 313.
- Stilbene, 4,4'-dichloro- 378.
- Stilbene, *trans*-4,4'-dicyano- 386.
- Stilbene, *trans*-2,5-dimethoxy-4'-nitro- 453.
- Stilbene, *trans*-4-dimethylamino-4'-nitro- 461.
- Stilbene, *trans*-2,4-dinitro- 522.
- Stilbene, *cis*-4,4'-dinitro- 523.
- Stilbene, *trans*-4,4'-dinitro- 524.
- Stilbene, *trans*-4-fluoro- 596.
- Stilbene, 4-methoxy-, *trans*- 685.
- Stilbene, *trans*-3-methoxy-4'-nitro- 676.
- Stilbene, *cis*-4-methoxy-4'-nitro- 677.
- Stilbene, *trans*-4-methoxy-4'-nitro- 678.
- Stilbene, *trans*-3-nitro- 823.
- Stilbene, *trans*-4-nitro- 824.
- trans*-Stilbene-2,2'-*d*₂ 1001.
- trans*-Stilbene-2,3,4,5,6-*d*₅ 1002.
- trans*-Stilbene-4,4'-*d*₂ 1003.
- trans*-Stilbene-7,7'-*d*₂ 1004.
- cis*-Stilbene 1005.
- trans*-Stilbene 1006.
- trans*-Stilbene-*d*₁₂ 1007.
- Styrene 1008.
- Styrene, *trans*-1-methoxy- β -nitro- 681.
- Styrene, β -(2-anthryl)- 89.
- Styrene, *trans*-4-dimethylamino- β -nitro- 465.
- Styrene, β,β -dimethyl- α -naphthyl- 496.
- Styrene, β -ethyl- 578.
- Styrene, α -methyl- β -ethyl- 724.
- Styrene, α -naphthyl- 800.
- Sulfacetamide 1009.
- Sulfanilamide, *N*'-acetyl- 1009.
- Sulfanilic acid/Eosin 561.
- 4-Sulfomethylbenzophenone 1010.
- (4-Sulfomethylphenyl)phenylethanedione 958.
- TMPD 1049.
- TMPD/2,6-Diphenyl-1,4-benzoquinone 533.
- Terephthalonitrile 384.
- Terphenyl, 4,4'-bis(diethylamino)- 1028.
- p*-Terphenyl, 4-nitro- 825.
- [1,1':4',1''-Terphenyl]-4,4''-diamine 1011.
- m*-Terphenyl 1012.
- o*-Terphenyl 1013.
- p*-Terphenyl 1014.
- p*-Terphenyl-*d*₁₄ 1015.
- 4-(4-*p*-Terphenyl)methylbenzophenone 1016.
- 2-([1,1':4',1''-Terphenyl]-4-ylmethyl)triphenylene 1017.
- Testosterone 1018.
- 1,2,3,4,5,6,7,8-Tetrabenzonaphthalene 347.
- Tetrabenzofulvalene 167.
- Tetrabenzophenazine 1019.
- Tetrabenzoporphinatocadmium(II) 240.
- Tetrabenzoporphinatozinc(II) 1133.
- Tetrabenzoporphine 1020.
- 2',4',5',7'-Tetrabromofluorescein dianion 562.
- Tetrabromofluorescein disodium salt 556.
- Tetracene 1021.
- Tetracene, 5-(1,4-epidioxyphenyl)-6,11,12-triphenyl- 563.
- Tetracene, 5,6,11,12-tetraphenyl- 989.
- 1,5,9,10-Tetrachloroanthracene 1022.
- Tetrachlorophthalic anhydride/Naphthalene-*d*₈ 773.
- Tetrachlorophthalic anhydride/Phenanthrene-*d*₁₀ 858.
- 1,2,4,5-Tetracyanobenzene 1023.
- 1,3,5,7,9,11,13-Tetradecaheptaene, 3,7,12-trimethyl-
1,14-bis(2,6,6-trimethyl-1-cyclohexen-1-yl)-, (*all-E*)- 1086.
- p,p'*-Tetraethyldiaminoquaterphenyl 1027.
- p,p'*-Tetraethyldiaminoterphenyl 1028.
- N,N,N',N'*-Tetraethyloxonine cation, conjugate
monoacid 1025.
- N,N,N',N'*-Tetraethyloxonine cation 1024.
- Tetraethylporphinatozinc(II) 1134.
- Tetraethylporphine 1026.
- N,N,N',N'*-Tetraethyl-[1,1':4',1'':4'',1'''-quaterphenyl]-
4,4'''-diamine 1027.
- N,N,N',N'*-Tetraethyl-[1,1':4',1''-terphenyl]-4,4''-
diamine 1028.
- 7,8,9,10-Tetrahydrobenzo[*a*]pyrene-7,8,9,10-tetrol 1031.
- 6*ba*,6*ca*,12*ba*,12*ca*-Tetrahydrocyclobuta[1,2-*a*:3,4-*a'*]-
diacenaphthylene 2.
- 6*ba*,6*c* β ,12*ba*,12*c* β -Tetrahydrocyclobuta[1,2-*a*:3,4-*a'*]-
diacenaphthylene 3.
- 7,8,11,12-Tetrahydro- ψ -carotene 255.
- 6,7,12,13-Tetrahydro-6,13:7,12-di-*o*-benzenobenzo[5,6]-
cycloocta[1,2-*b*]naphthalene 72.
- (*R*)-4,5,6,7-Tetrahydrodinaphtho[2,1-*b*:1',2'-*d'*][1,6]-
dioxecin 1029.
- 6,7,8,9-Tetrahydro-4-hydroxythiazolo[4,5-*h*]isoquinoline-
7-carboxylate ion 1030.
- 7,8,9,10-Tetrahydrotetrahydroxybenzo[*a*]pyrene 1031.

- Tetraiodofluorescein dianion **566**.
 Tetrakis(carboxyphenyl)porphinatopalladate(II) ion **840**.
 Tetrakis(carboxyphenyl)porphinatozincate(II) ion **1135**.
 Tetrakis(2,6-dimethyl-4-sulfonatophenyl)porphine **1032**.
 Tetrakis(2,6-dimethyl-4-sulfonatophenyl)porphinatozincate(II) ion **1136**.
 Tetrakis(4-*N*-methylpyridyl)porphinatopalladium(II) ion **841**.
 Tetrakis(2-*N*-methylpyridyl)porphine **1033**.
 Tetrakis(3-*N*-methylpyridyl)porphine **1034**.
 Tetrakis(4-*N*-methylpyridyl)porphine **1035**.
 Tetrakis(2-*N*-methylpyridyl)porphinatozinc(II) ion **1137**.
 Tetrakis(3-*N*-methylpyridyl)porphinatozinc(II) ion **1138**.
 Tetrakis(4-*N*-methylpyridyl)porphinatozinc(II) ion **1139**.
 Tetrakis(2-pyridyl)porphine, conjugate diacid **1039**.
 Tetrakis(3-pyridyl)porphine, conjugate diacid **1040**.
 Tetrakis(4-pyridyl)porphine, conjugate diacid **1041**.
 Tetrakis(2-pyridyl)porphine **1036**.
 Tetrakis(3-pyridyl)porphine **1037**.
 Tetrakis(4-pyridyl)porphine **1038**.
 Tetrakis(*p*-sulfonatophenyl)porphinatopalladate(II) ion **842**.
 Tetrakis(*p*-sulfonatophenyl)porphinatorhodate(III) ion **1042**.
 Tetrakis(*p*-sulfonatophenyl)porphine **1043**.
 Tetrakis(*p*-sulfonatophenyl)porphinatozincate(II) ion **1140**.
 Tetrakis(trimethylaminophenyl)porphine **1044**.
 Tetrakis-4-(*N,N,N*-trimethylammonio)phenylporphine zinc(II) ion **1141**.
N,N,N,N'-Tetramethylbenzidine **1045**.
 2,3,5,6-Tetramethylbenzoquinone **552**.
 Tetramethyl-*p*-benzoquinone **552**.
 1,1',3,3'-Tetramethyl- $[\Delta^{9,9'}\text{-bianthracene}]$ -10,10'-dione **1046**.
 1,1',3,3'-Tetramethyldianthrone **1046**.
 3,4,7,8-Tetramethyl-1,5-diazabicyclo[3,3,0]octa-3,7-diene-2,6-dione **1047**.
 1,3,5,8-Tetramethyl-2,4-divinylporphine-6,7-dipropionic acid **930**.
 1,3,3,6-Tetramethyl-6'-nitro-spiro-(2,2'-indolin[2*H*-1]benzopyran) **1048**.
N,N,N,N'-Tetramethyl-*p*-phenylenediamine/3-Chloro-2,6-diphenyl-1,4-benzoquinone **273**.
N,N,N,N'-Tetramethyl-*p*-phenylenediamine **1049**.
 2,3,6,7-Tetramethyl-1*H*,5*H*-pyrazolo[1,2-*a*]pyrazole-1,5-dione **1047**.
 Tetramethylthiuram disulfide **1050**.
 Tetramethylthiuram monosulfide **1051**.
 Tetraphene **106**.
 1,1,4,4-Tetraphenyl-1,3-butadiene **1052**.
 Tetraphenylchlorin **1053**.
 Tetraphenyl-*p*-dioxin **1054**.
 3,3,4,5-Tetraphenyl-2(3*H*)-furanone **1055**.
 5,6,11,12-Tetraphenylnaphthacene **989**.
 Tetraphenylporphinatocadmium(II) **241**.
 Tetraphenylporphinatomanganese(II) **654**.
 Tetraphenylporphinatomercury(II) **656**.
 Tetraphenylporphinatopalladium(II) **843**.
 Tetraphenylporphinatorrhodium(III) chloride **1056**.
 Tetraphenylporphinatozinc(II) **1143**.
meso-Tetraphenylporphine **1057**.
 Tetraphenylporphyrin dianion **1058**.
 1,3,6,8-Tetraphenylpyrene **1059**.
 5,6,11,12-Tetraphenyltetracene **989**.
 2,2'-Thiacarbocyanine, 6,6'-bis(dimethylamino)-3,3'-diethyl-, iodide **191**.
 2,2'-Thiacarbocyanine, 3,3'-diethyl-, iodide **403**.
 2,2'-Thiacarbocyanine, 3,3'-diethyl-9-methoxy-, iodide **397**.
 2,2'-Thiacyanine iodide, 3,3'-diethyl- **404**.
 2,2'-Thiadicarbocyanine iodide, 3,3'-diethyl- **405**.
 2,2'-Thiatricarbocyanine iodide, 3,3'-diethyl- **406**.
 4-Thiazolidinone, 5[(3,4-dihydro[1,4]thiazino[3,4-*b*]-benzothiazol-1-yl)methylene]-3-ethyl-2-thio-, (*E*)- **435**.
 4-Thiazolidinone, 3-ethyl-5[(3-ethyl-2(3*H*)-benzothiazolylidene)ethylidene]-2-thio-, (*E,Z*)- **571**.
 2- Δ^2 -Thiazoline, benzoylamino- **147**.
 2- Δ^2 -Thiazoline, (2'-chlorobenzoyl)amino- **267**.
 2- Δ^2 -Thiazoline, (4'-chlorobenzoyl)amino- **268**.
 2- Δ^2 -Thiazoline, (2'-methylbenzoyl)amino- **709**.
 2- Δ^2 -Thiazoline, (4'-methylbenzoyl)amino- **710**.
 Thiazolo[4,5-*h*]isoquinoline-7-carboxylate, 6,7,8,9-tetrahydro-4-hydroxy-, ion **1030**.
 Thiobenzophenone **1060**.
 Thiobenzophenone, 4,4'-bis(dimethylamino)- **200**.
 Thiobenzophenone, 4,4'-dimethoxy- **454**.
 Thiodicarbonic diamide, tetramethyl- **1051**.
 Thioflavine **1061**.
 Thioflavine T **1061**.
 Thioflavine TCN **1061**.
 Thioindigo **1062**.
 Thioindigo, 5,5'-di-*tert*-amyl- **337**.
 Thioindigo, 5,5'-dibutyl- **364**.
 Thioindigo, 6,6'-diethoxy- **388**.
 Thioindigo, 6,6'-dihexyloxy- **407**.
 Thioindigo, 5,5'-dineopentyl- **514**.
 Thionine cation **1063**.
 Thionine cation, conjugate monoacid **1064**.
 Thiophene, 2,5-bis(5-*tert*-butyl-2-benzoxazolyl)- **189**.
 Thiophene, 2-nitro- **826**.
 Thiopyronine **1065**.
 Thiopyronine cation **1066**.
 4-Thiouridine **1067**.
 9*H*-Thioxanthene-9-thione **1068**.
 Thioxanthen-9-one, 2-chloro- **290**.
 Thioxanthen-9-one, 2-isopropyl- **639**.
 Thioxanthen-9-one, 2-methyl- **749**.
 Thioxanthen-9-one **1069**.
 9*H*-Thioxanthen-9-one **1069**.
 Thioxanthone **1069**.
 Thymidine **1070**.
 Thymidine 5'-monophosphate **1071**.
 5'-Thymidylic acid **1071**.
 Thymine **1072**.
 Thymine, negative ion **1073**.
 Tin(IV) octaethylporphyrin dichloride **828**.
 Tolan **527**.
 Toluene/Tetracyanobenzene **1074**.
m-Toluic acid **704**.
o-Toluic acid **703**.
p-Toluic acid **705**.
p-Toluquinone **708**.

- Torularhodinaldehyde 387.
 1,3,5-Triazin-2-amine,
 4-chloro-6-[4-(diethylamino)phenyl]-*N,N*-dimethyl- 269.
 1,3,5-Triazine, 4,6-dichloro-2-[4-(diethylamino)phenyl]- 379.
s-Triazine 1075.
 1,2,3,4,6,7-Tribenzophenazine 1076.
 Tribenzo[*a,c,h*]phenazine 1076.
 1,3,5-Tri(biphenyl-4-yl)benzene 183.
 2,9,10-Trichloroanthracene 1077.
 1,3,5-Tricyanobenzene 1078.
 Tricyclo[8.2.2.2^{4,7}]hexadeca-4,6,10,12,13,15-hexaene 844.
 Tricyclo[8.2.2.2^{4,7}]hexadeca-4,6,10,12,13,15-hexaene-
 5,15-dicarbonitrile 1079.
 2,4,6,8,10,12-Tridecahexaenal, 7,11-dimethyl-
 7-(2,6,6-trimethyl-
 1-cyclohexen-1-yl)-, (*E,E,E,E,E,E*)- 501.
 1,3,5,8,10,12-Tridecahexaen-7-one,
 1,13-bis(dimethylamino)-, *all-trans*- 201.
 1,3,6,8,10,12-Tridecahexaen-5-one,
 1,13-bis(dimethylamino)-, *all-trans*- 202.
 Triethylamine/Benzil 113.
 1,1,1-Trifluoroacetylacetone 1080.
 4-(Trifluoromethyl)acetophenone 1081.
 4-(Trifluoromethyl)benzophenone 1082.
 1,1,1-Trifluoro-2,4-pentanedione 1080.
 4,4,4-Trifluoro-1-phenyl-1,3-butanedione 158.
 4,4,4-Trifluoro-1-phenyl-1,3-butanedionatosodium 1083.
 2,4,6-Triisopropylbenzophenone 1084.
 1,2,3-Trimethoxybenzene/3-Chloro-2,6-diphenyl-
 1,4-benzoquinone 274.
 7,8,10-Trimethylbenzo[*g*]pteridine-2,4-(3*H*,10*H*)-dione 645.
 2,3,5-Trimethyl-1,4-benzoquinone 1085.
 (*all-E*)-3,7,12-Trimethyl-
 1,14-bis(2,6,6-trimethyl-1-cyclohexen-1-yl)-
 1,3,5,7,9,11,13-tetradecaheptaene 1086.
 4-(2,6,6-Trimethyl-1-cyclohexen-1-yl)-3-buten-2-one 638.
 1,1,2-Trimethyl-2-(2-naphthyl)ethylene 1087.
 1,3,3-Trimethyl-6'-nitro-
 spiro-(2,2'-indolin[2*H*-1]benzopyran) 1088.
 3,5,5-Trimethyl-2-phenoxy-2-cyclohexene-1-one 1089.
 4-[Tri(4-methylphenyl)porphyrin]-3-phenoxypropyl
 β-*apo*-6'-carotenate 1090.
 4,5',8-Trimethylpsoralen 1091.
 Trioxsalen 1091.
 Triphenylamine 1092.
 Triphenylamine, conjugate acid 1093.
 Triphenylamine/3-Chloro-2,6-diphenyl-1,4-
 benzoquinone 275.
 Triphenylamine/3,5-Dichloro-2,6-diphenyl-1,4-
 benzoquinone 376.
 1,3,5-Triphenylbenzene 1094.
 Triphenylene 1095.
 Triphenylene, 2-bromo- 230.
 Triphenylene-*d*₁₂ 1096.
 Triphenylene/Chloranil 1097.
 Triphenylethylene 1098.
 3,3,5-Triphenyl-2(3*H*)-furanone 1099.
 Triphenylmethane 1100.
 Triphenylmethyl cation 1101.
 Triphenylmethyl cation 1101.
 1,3,3-Triphenylprop-2-en-1-one 1102.
 Triptycene, 3-acetyl- 20.
 Tris(2,2'-bipyridine)iridium(III) ion 1103.
 Tris(2,2'-bipyridine)osmium(II) ion 1104.
 Tris(2,2'-bipyridine)ruthenium(II) ion 1105.
 Tris(1,10-phenanthroline)rhodium(III) ion 1106.
 Tris(4,4,4-trifluoro-1-phenyl-1,3-butanedionato)-
 lanthanum(III) 1107.
 Tris(4,4,4-trifluoro-1-phenyl-1,3-butanedionato)-
 lutetium(III) 1108.
 1,3,5,7,9,11,13,15,17,19,21,23,25,27,29,31,33-
 Tritriacontahaptadecaene, 3,7,11,15,20,24,28,32-
 octamethyl-1,34-bis(2,6,6-trimethyl-1-
 cyclohexen-1-yl)-, (*all-E*)- 830.
 Trypaflavine cation 30.
 Tryptophan 1109.
 L-Tryptophan, 1-methyl- 760.
 DL-Tryptophan 1110.
 L-Tryptophan 1111.
 Tyrosine 1112.
 L-Tyrosylglycine 1113.
 Ubiquinone 30 1114.
 1,3,6,8,10-Undecapentaen-5-one,
 1,11-bis(dimethylamino)-, *all-trans*- 203.
 Uracil 1115.
 Uracil, 1,3-dimethyl- 502.
 Uridine 1116.
 Uridine monophosphate 1117.
 Uridine 5'-monophosphate 1117.
 Uridylic acid 1117.
 Uroporphyrin I, octamethyl ester 1118.
 2-Vinylnaphthalene 1119.
 Vitamin A 974.
 Vitamin A acid 973.
 Water/3,5-Dinitroanisole 516.
 9*H*-Xanthene-9-thione 1120.
 9-Xanthenone 1121.
 Xanthone 1121.
 Xanthotoxin 684.
 Xanthylium, 9-(2-carboxyphenyl)-3,6-bis(diethylamino)-,
 chloride 983.
 Xanthylium, 3,6-diamino-9-(2-carboxyphenyl)-, chloride
 982.
 Xanthylium, 9-[2-(ethoxycarbonyl)phenyl]-
 3,6-bis(ethylamino)-2,7-dimethyl-, chloride 984.
o-Xylene 1122.
 Zeaxanthin 1123.
 Zinc(II) chlorophyll *a* 1124.
 Zinc(II) cytochrome *c* 1125.
 Zinc(II) etioporphyrin I 1126.
 Zinc(II) etioporphyrin I/Hexachloroethane 1127.
 Zinc(II) etioporphyrin I/*trans*-4-Nitrostilbene 1128.
 Zinc(II) etioporphyrin I/4-Nitrotoluene 1129.
 Zinc(II) phthalocyanine 1130.
 Zinc(II) porphyrin 1131.
 Zinc(II) protoporphyrin 1132.
 Zinc(II) tetrabenzoporphyrin 1133.
 Zinc(II) tetraethylporphyrin 1134.

Zinc(II) tetrakis(carboxyphenyl)porphyrin 1135.
 Zinc(II) tetrakis(2,6-dimethyl-4-sulfonatophenyl)porphyrin 1136.
 Zinc(II) tetrakis(2-*N*-methylpyridyl)porphyrin 1137.
 Zinc(II) tetrakis(3-*N*-methylpyridyl)porphyrin 1138.
 Zinc(II) tetrakis(4-*N*-methylpyridyl)porphyrin 1139.
 Zinc(II) tetrakis(*p*-sulfonatophenyl)porphyrin 1140.
 Zinc(II) tetrakis(trimethylaminophenyl)porphyrin 1141.
 Zinc(II) tetraphenylchlorin 1142.
 Zinc(II) tetraphenylporphyrin 1143.

10.2. Formula Index

$C_3H_3N_3$ *s*-Triazine 1075.
 $C_3H_4O_3$ Pyruvic acid 957.
 C_3H_6O Acetone 7.
 $C_4H_3NO_2S$ 2-Nitrothiophene 826.
 $C_4H_4N_2$ Pyrazine 938.
 Pyrimidine 952.
 $C_4H_4N_2O_2$ Uracil 1115.
 C_4H_6O 1-Butene-3-one 761.
 $C_4H_6O_2$ 2,3-Butanedione 164.
 C_4H_8O 2-Butanone 231.
 Butyraldehyde 238.
 $C_5H_2F_6O_2$ 1,1,1,5,5,5-Hexafluoroacetylacetone 607.
 $C_5H_3NO_5$ 5-Nitro-2-furoic acid 818.
 $C_5H_3N_2O_4^-$ Orotate ion 831.
 $C_5H_4N_2O_3$ 4-Nitropyridine-*N*-oxide 821.
 $C_5H_4N_2O_4$ Orotic acid 832.
 $C_5H_4N_4$ Purine 936.
 $C_5H_3F_3O_2$ 1,1,1-Trifluoro-2,4-pentanedione 1080.
 $C_5H_3N_2O_2^-$ Thymine, negative ion 1073.
 $C_5H_5N_5$ 2-Aminopurine 58.
 C_5H_6 Cyclopentadiene 326.
 $C_5H_6N_2$ 2-Aminopyridine 59.
 4-Aminopyridine 60.
 $C_5H_6N_2O_2$ Thymine 1072.
 C_5H_6O 2-Cyclopentenone 327.
 $C_5H_8O_2$ 2,4-Pentanedione 11.
 $C_6Cl_4O_2$ Chloranil 259.
 C_6D_6 Benzene-*d*₆ 110.
 $C_6H_4O_2$ 1,4-Benzoquinone 143.
 C_6H_5Cl Chlorobenzene 265.
 C_6H_5F Fluorobenzene 594.
 $C_6H_5N_2O$ Pterin 57.
 C_6H_6 Benzene 109.
 $C_6H_6N_2O_2$ 4-Nitroaniline 812.
 C_6H_6O Phenol 868.
 C_6H_7N Aniline 61.
 $C_6H_7N_3O$ 7-Methylguanine 726.
 C_6H_8 1,3-Cyclohexadiene 321.
 $C_6H_8ClN_7O$ Amiloride 32.
 $C_6H_8N_2O_2$ 1,3-Dimethyluracil 502.
 C_6H_8O 2-Cyclohexen-1-one 322.
 $C_6H_{12}N_2S_3$ Tetramethylthiuram monosulfide 1051.
 $C_6H_{12}N_2S_4$ Tetramethylthiuram disulfide 1050.
 C_7H_5N Benzonitrile 128.
 $C_7H_6N_2O_5$ 3,5-Dinitroanisole 515.
 C_7H_6O Benzaldehyde 104.

$C_7H_6O_2$ Benzoic acid 123.
 C_7H_7NO Methyl-1,4-benzoquinone 708.
 $C_7H_7NO_2$ Benzamide 105.
 2-Aminobenzoic acid 39.
 4-Aminobenzoic acid 40.
 $C_7H_7NO_3$ 3-Nitroanisole 813.
 4-Nitroanisole 814.
 C_7H_8O Anisole 63.
 C_7H_8O *p*-Cresol 304.
 $C_7H_8O_2$ 4-Methoxyphenol 679.
 $C_7H_8N_5$ 2-(Dimethylamino)purine 466.
 C_7H_{10} 1,3-Cycloheptadiene 317.
 $C_7H_{10}N_2$ *all-trans*-1,3,5-Heptatriene 605.
 $C_7H_{10}N_2$ 2-(Dimethylamino)pyridine 467.
 $C_7H_{10}O$ 4-(Dimethylamino)pyridine 468.
 $C_8H_4N_2$ 2-Cyclohepten-1-one 318.
 $C_8H_4N_2$ 1,2-Dicyanobenzene 382.
 $C_8H_4N_2$ 1,3-Dicyanobenzene 383.
 $C_8H_4N_2$ 1,4-Dicyanobenzene 384.
 $C_8H_5NO_4$ 3,6-Dihydroxyphthalimide 444.
 C_8H_6 Phenylacetylene 873.
 $C_8H_6N_2$ Phthalazine 905.
 $C_8H_6N_2O_2$ Quinoxaline 963.
 C_8H_7N Quinoxaline-1,4-dioxide 964.
 C_8H_7N Indole 631.
 $C_8H_7N_3O_2$ 3,6-Diaminophthalimide 336.
 C_8H_8 Styrene 1008.
 $C_8H_8N_2$ 1-Methylindazole 727.
 C_8H_8O 2-Methylindazole 728.
 C_8H_8O Acetophenone 8.
 $C_8H_8O_2$ 2,5-Dimethyl-1,4-benzoquinone 474.
 $C_8H_8O_2$ 2,6-Dimethyl-1,4-benzoquinone 475.
 $C_8H_8O_2$ Methyl benzoate 702.
 $C_8H_8O_2$ *o*-Toluic acid 703.
 $C_8H_8O_2$ *m*-Toluic acid 704.
 $C_8H_8O_2$ *p*-Toluic acid 705.
 $C_8H_8O_4$ 2,3-Dimethoxy-1,4-benzoquinone 446.
 $C_8H_8O_4$ 2,5-Dimethoxy-1,4-benzoquinone 447.
 $C_8H_9NO_2$ 4-(Methylamino)benzoic acid 695.
 $C_8H_9NO_4$ 1,2-Dimethoxy-4-nitrobenzene 452.
 $C_8H_9O^+$ Acetophenone, conjugate acid 9.
 C_8H_{10} *o*-Xylene 1122.
 $C_8H_{10}N_2O_2$ *N,N*-Dimethyl-4-nitroaniline 497.
 $C_8H_{10}N_2O_3S$ Sulfacetamide 1009.
 $C_8H_{10}N_4O_2$ Caffeine 242.
 $C_8H_{11}N$ *N,N*-Dimethylaniline 470.
 C_8H_{12} 1,3-Cyclooctadiene 324.
 $C_8H_{12}N_2$ *N,N*-Dimethyl-*p*-phenylenediamine 500.
 C_8H_{14} 2,5-Dimethyl-2,4-hexadiene 485.
 $C_9F_3H_7O$ 4-(Trifluoromethyl)acetophenone 1081.
 $C_9H_3N_3$ 1,3,5-Tricyanobenzene 1078.
 C_9H_6BrNO 4-Bromoisoquinoline-*N*-oxide 221.
 $C_9H_6NO^-$ Carbostyryl, negative ion 248.
 $C_9H_6N_2O_3$ 4-Nitroquinoline-*N*-oxide 822.
 $C_9H_6O_2$ Chromone 294.
 $C_9H_6O_3$ Coumarin 303.
 $C_9H_6O_3$ 3-Hydroxycoumarin 616.
 $C_9H_6O_3$ 4-Hydroxycoumarin 617.

C_9H_7N	Isoquinoline 641. Quinoline 961.	$C_{10}H_7O^-$	1-Naphthyloxy ion 797. 2-Naphthyloxy ion 798.
C_9H_7NO	Carbostyryl 247. Isoquinoline- <i>N</i> -oxide 642.	$C_{10}H_7O_2^-$	2,3-Dihydroxynaphthalene, conjugate base 443.
$C_9H_7NO_2$	<i>N</i> -Methylphthalimide 745.	$C_{10}H_8$	Azulene 95. Naphthalene 768.
C_9H_8	1-Propynylbenzene 928.	$C_{10}H_8N_2O_2$	1-Amino-4-nitronaphthalene 53.
$C_9H_8N^+$	Quinoline, conjugate acid 962.	$C_{10}H_8N_2O_4$	1,4-Dinitronaphthalene 518.
$C_9H_8N_2O_2$	3-Amino- <i>N</i> -methylphthalimide 50.	$C_{10}H_8O$	1-Naphthol 786. 2-Naphthol 787.
$C_9H_8O_2$	4-Chromanone 293.	$C_{10}H_8O_2$	1,5-Dihydroxynaphthalene 439. 1,8-Dihydroxynaphthalene 440. 2,3-Dihydroxynaphthalene 441. 2,7-Dihydroxynaphthalene 442.
$C_9H_8O_3$	Methyl benzoylformate 711.	$C_{10}H_8O_3$	4-Hydroxy-6-methylcoumarin 623. 7-Hydroxy-4-methylcoumarin 624.
C_9H_9N	1-Methylindole 729. 2-Methylindole 730. 3-Methylindole 731.	$C_{10}H_8O_3S$	1-Naphthalenesulfonic acid 776.
$C_9H_9NO_2$	2-(<i>N</i> -Formylamino)acetophenone 17.	$C_{10}H_9ClN_2OS$	(2'-Chlorobenzoyl)amino-2- Δ^2 -thiazoline 267. (4'-Chlorobenzoyl)amino-2- Δ^2 -thiazoline 268.
$C_9H_9NO_3$	<i>trans</i> -1-Methoxy- β -nitrostyrene 681.	$C_{10}H_9N$	2-Naphthylamine 766.
$C_9H_{10}N_2$	1,3-Dimethylindazole 486.	$C_{10}H_9NO$	1-Methylisoquinoline- <i>N</i> -oxide 732.
$C_9H_{10}O$	3-Methylacetophenone 691. 4-Methylacetophenone 692. Propiophenone 926.	$C_{10}H_9NO_2$	<i>N</i> -Ethylphthalimide 577.
$C_9H_{10}O_2$	4'-Hydroxypropiophenone 628. 3-Methoxyacetophenone 661. 4-Methoxyacetophenone 662.	$C_{10}H_{10}$	1-Butynylbenzene 237.
$C_9H_{10}O_4$	2,3,5-Trimethyl-1,4-benzoquinone 1085. 2,3-Dimethoxy-5-methyl-1,4-benzoquinone 450.	$C_{10}H_{10}N^+$	2-Naphthylamine, conjugate acid 767.
$C_9H_{11}NO_2$	Phenylalanine 876.	$C_{10}H_{10}N_2O$	7-Amino-4-methylcarbostyryl 49.
$C_9H_{11}NO_3$	Tyrosine 1112.	$C_{10}H_{10}N_2O_5$	Benzoylamino-2- Δ^2 -thiazoline 147.
$C_9H_{12}N_2$	4-Pyrrolidinopyridine 956.	$C_{10}H_{10}O_2$	1-Benzoylacetone 146.
$C_9H_{12}N_2O_4$	<i>N</i> -Butyl-5-nitro-2-furamide 236.	$C_{10}H_{11}NO_2$	2-(<i>N</i> -Acetylamino)acetophenone 16. 2-(<i>N</i> -Formyl- <i>N</i> -methylamino)-acetophenone 19.
$C_9H_{12}N_2O_5S$	4-Thiouridine 1067.	$C_{10}H_{12}$	1-Phenyl-1-butene 578.
$C_9H_{12}N_2O_6$	Uridine 1116.	$C_{10}H_{12}N_2O_2$	2-(<i>N</i> -Formylamino)-3'-aminopropiophenone 55. <i>trans</i> -4-Dimethylamino- β -nitrostyrene 465.
$C_9H_{13}N_2O_9P$	Uridine monophosphate 1117.		3,4,7,8-Tetramethyl-1,5-diazabicyclo-[3,3,0]octa-3,7-diene-2,6-dione 1047.
$C_{10}D_8$	Naphthalene- <i>d</i> ₈ 769.	$C_{10}H_{12}O_2$	Duroquinone 552. 2-Hydroxy-2-methyl-1-phenylpropan-1-one 625.
$C_{10}H_2N_4$	1,2,4,5-Tetracyanobenzene 1023.		4'-Methoxypropiophenone 682.
$C_{10}H_2O_6$	1,2,4,5-Benzenetetracarboxylic anhydride 953.	$C_{10}H_{13}NO_2$	Methyl 4-dimethylaminobenzoate 717.
$C_{10}H_5BrN_2O$	4-Bromo-1-cyanoisoquinoline- <i>N</i> -oxide 217.	$C_{10}H_{13}O_2^+$	Duroquinone, conjugate monoacid 554.
$C_{10}H_6Br_2$	1,4-Dibromonaphthalene 361.	$C_{10}H_{14}N_2O_5$	Thymidine 1070.
$C_{10}H_6Cl_2$	1,4-Dichloronaphthalene 377.	$C_{10}H_{14}O_2$	Camphoroquinone 243.
$C_{10}H_6F_3NaO_2$	4,4,4-Trifluoro-1-phenyl-1,3-butanedionatosodium 1083.	$C_{10}H_{15}NO$	8-(Dimethylamino)-3,5,7-octatrien-2-one 462.
$C_{10}H_6N_2O_4$	1,2-Dinitronaphthalene 517. 1,8-Dinitronaphthalene 519.	$C_{10}H_{15}N_2O_8P$	Thymidine 5'-monophosphate 1071.
$C_{10}H_6N_4O_2$	Alloxazine 31.	$C_{10}H_{16}$	Neo-alloocimene 806.
$C_{10}H_7Br$	1-Bromonaphthalene 224. 2-Bromonaphthalene 225.	$C_{10}H_{16}N_2$	<i>N,N,N',N'</i> -Tetramethyl- <i>p</i> -phenylenediamine 1049.
$C_{10}H_7Cl$	1-Chloronaphthalene 279. 2-Chloronaphthalene 280.	$C_{11}H_6O_3$	Angelicin 640. Psoralen 934.
$C_{10}H_7F_3O_2$	4,4,4-Trifluoro-1-phenyl-1,3-butanedione 158.	$C_{11}H_7N$	1-Naphthonitrile 314.
$C_{10}H_7I$	1-Iodonaphthalene 636. 2-Iodonaphthalene 637.	$C_{11}H_7O_2^-$	1-Naphthoate ion 781. 2-Naphthoate ion 782.
$C_{10}H_7NO_2$	1-Nitronaphthalene 819. 2-Nitronaphthalene 820.		
$C_{10}H_7NO_3$	4-Hydroxyquinoline-2-carboxylic acid 643.		
$C_{10}H_7NaO_3S$	2-Naphthalenesulfonic acid, sodium salt 775.		

$C_{11}H_8O$	1-Naphthaldehyde 764. 2-Naphthaldehyde 765.	$C_{12}H_9N$	Carbazole 245.
$C_{11}H_8O_2$	2-Methyl-1,4-naphthoquinone 737. 1-Naphthoic acid 784. 2-Naphthoic acid 785.	$C_{12}H_9NO$	Pyrido[2,1,6- <i>de</i>]quinolizine 951. 2-Benzoylpyridine 155. 3-Benzoylpyridine 156. 4-Benzoylpyridine 157. 4-Methoxy-1-naphthonitrile 674. Phenoxazine 871.
$C_{11}H_8O_3$	4',5'-Dihydropsoralen 430.	$C_{12}H_9NO_2$	5-Nitroacenaphthene 811. 4-Nitrobiphenyl 816.
$C_{11}H_8Br$	1-(Bromomethyl)naphthalene 223.	$C_{12}H_9NS$	Phenothiazine 870.
$C_{11}H_9Cl$	1-(Chloromethyl)naphthalene 278.	$C_{12}H_9N_2^+$	Phenazine, conjugate monoacid 867.
$C_{11}H_{10}$	1-Methylnaphthalene 733. 2-Methylnaphthalene 734.	$C_{12}H_9N_2O_3S^-$	6,7,8,9-Tetrahydro-4-hydroxythiazolo-[4,5- <i>h</i>]isoquinoline-7-carboxylate ion 1030.
$C_{11}H_{10}O$	1-Methoxynaphthalene 672. 2-Methoxynaphthalene 673.	$C_{12}H_{10}$	Acenaphthene 1. Biphenyl 173. 2-Vinylnaphthalene 1119.
$C_{11}H_{10}O_4$	5,7-Dimethoxycoumarin 449.	$C_{12}H_{10}N_3O^+$	Oxonine cation 835.
$C_{11}H_{11}NO_2$	<i>N</i> -Propylphthalimide 927.	$C_{12}H_{10}N_3S^+$	Thionine cation 1063.
$C_{11}H_{11}NO_3S^-$	2-Amino-3-(4-methoxy-6-benzothiazolyl)-propionate ion 47.	$C_{12}H_{10}N_3Se^+$	Selenine cation 993.
$C_{11}H_{12}N_2OS$	(2'-Methylbenzoyl)amino-2- Δ^2 -thiazoline 709. (4'-Methylbenzoyl)amino-2- Δ^2 -thiazoline 710.	$C_{12}H_{10}N_4O_2$	Lumichrome 644.
$C_{11}H_{12}N_2O_2$	<i>N</i> -[(Dimethylamino)methyl]phthalimide 459. Tryptophan 1109. DL-Tryptophan 1110. L-Tryptophan 1111.	$C_{12}H_{10}O$	1'-Acetonaphthone 5. 2'-Acetonaphthone 6. 4-Phenylphenol 615.
$C_{11}H_{12}N_2O_4$	<i>N</i> -Formylkynurenine 599.	$C_{12}H_{10}O_2$	<i>p,p'</i> -Biphenol 437. Methyl 2-naphthoate 736.
$C_{11}H_{12}O_2$	3-Phenylacetylacetone 872.	$C_{12}H_{11}N$	1-Naphthyl acetate 791. 4-Aminobiphenyl 43. 4 <i>a</i> ,5 <i>a</i> -Dihydrocarbazole 410.
$C_{11}H_{13}NO_2$	2 (<i>N</i> -Acetyl- <i>N</i> -methylamino)acetophenone 18.	$C_{12}H_{11}NO$	Diphenylamine 528.
$C_{11}H_{14}$	2-Phenyl-2-pentene 724.	$C_{12}H_{11}N_3O^{2+}$	4-Amino-4'-hydroxybiphenyl 46. Oxonine cation, conjugate monoacid 836.
$C_{11}H_{14}N_2O_4$	L-Tyrosylglycine 1113.	$C_{12}H_{11}N_3S^{2+}$	Thionine cation, conjugate monoacid 1064.
$C_{11}H_{14}N_2O_5$	4- <i>tert</i> -Butyl-3,5-dinitroanisole 233.	$C_{12}H_{11}N_3Se^{2+}$	Selenine cation, conjugate monoacid 994.
$C_{11}H_{14}O$	Pivalophenone 912.	$C_{12}H_{12}$	1,2-Dimethylnaphthalene 489. 2,3-Dimethylnaphthalene 490. 2,7-Dimethylnaphthalene 491.
$C_{11}H_{14}O_2$	2-Methoxy-2-methyl-1-phenyl-1-propanone 671.	$C_{12}H_{12}O$	1-Ethylnaphthalene 574.
$C_{11}H_{15}NO_2$	2-(Dimethylamino)ethyl benzoate 457.	$C_{12}H_{12}N_2$	2-Ethylnaphthalene 575.
$C_{11}H_{18}N_2O$	1,7-Bis(dimethylamino)-1,4,6-heptatrien-3-one 192.	$C_{12}H_{12}N_2O_2$	Benidine 111. 1-Dimethylamino-4-nitronaphthalene 460.
$C_{11}N_4O_3^{2-}$	Croconate Blue dianion 306.	$C_{12}H_{12}O$	2-Ethoxynaphthalene 567.
$C_{12}D_{10}$	Biphenyl- <i>d</i> ₁₀ 174.	$C_{12}H_{12}O_2$	1,4-Dimethoxynaphthalene 451.
$C_{12}H_6N_2$	1,4-Dicyanonaphthalene 385.	$C_{12}H_{13}N$	<i>N,N</i> -Dimethyl-2-naphthylamine 492.
$C_{12}H_8$	Diphenylene 176.	$C_{12}H_{14}$	1-Phenylcyclohexene 881.
$C_{12}H_8Br_2$	4,4'-Dibromobiphenyl 359.	$C_{12}H_{14}N^+$	<i>N,N</i> -Dimethyl-2-naphthylamine, conjugate acid 493.
$C_{12}H_8N_2$	Benzo[<i>c</i>][1,5]naphthyridine 125. Benzo[<i>c</i>][1,6]naphthyridine 126. Benzo[<i>c</i>][1,7]naphthyridine 127. 1,10-Phenanthroline 861. 1,7-Phenanthroline 862. 1,8-Phenanthroline 863. 1,9-Phenanthroline 864. 4,7-Phenanthroline 865. Phenazine 866.	$C_{12}H_{14}N_2O_2$	1-Methyl-L-tryptophan 760.
$C_{12}H_8O_3$	3-Formylfurochromene 598.	$C_{12}H_{14}O_3$	2-Acetoxy-2-methyl-1-phenyl-1-propanone 10. Butyl benzoylformate 232.
$C_{12}H_8O_4$	2-Formyl-5'-methylidifurobenzene 600.	$C_{12}H_{16}N_2$	2- <i>tert</i> -Butyl-4-methylindazole 235.
$C_{12}H_8Br$	5-Methoxypsoralen 683.	$C_{12}H_{17}NO$	10-(Dimethylamino)-3,5,7,9-decatetraen-2-one 456.
$C_{12}H_9Br$	8-Methoxypsoralen 684.		
$C_{12}H_9I$	4-Bromobiphenyl 216. 4-Iodobiphenyl 634.		

$C_{13}D_9N$	Acridine- <i>d</i> ₉ 24.	$C_{13}H_{13}N$	<i>N</i> -Methyldiphenylamine 718.
$C_{13}H_7ClOS$	2-Chlorothioxanthen-9-one 290.	$C_{13}H_{13}N_3^{2+}$	Proflavine, conjugate diacid 922.
$C_{13}H_8$	Fluoren-9-ylidene 589.	$C_{13}H_{13}N_4O_2^+$	Lumiflavine, conjugate monoacid 646.
$C_{13}H_8Cl_2O$	4,4'-Dichlorobenzophenone 372.	$C_{13}H_{14}$	2-Phenyl-2-norbornene 896.
$C_{13}H_8NO_4S^-$	9,10-Dihydro-9-oxo-2-acridinesulfonate ion 424.	$C_{13}H_{14}Cl_2N_4$	4-(4,6-Dichloro-1,3,5-triazin-2-yl)- <i>N,N</i> -diethylaniline 379.
$C_{13}H_8O$	Fluoren-9-one 586.	$C_{13}H_{15}NO_4$	8-Methoxy-2,2,3-trimethyl-6-nitro-2 <i>H</i> -chromene 686.
$C_{13}H_8OS$	Thioxanthen-9-one 1069.	$C_{13}H_{15}N_3O_2$	<i>N</i> -Acetyl-L-tryptophanamide 21.
$C_{13}H_8O_2$	9 <i>H</i> -Xanthene-9-thione 1120.	$C_{13}H_{17}N$	1,2-Dihydro-2,2,4,6-tetramethylquinoline 432.
	6 <i>H</i> -Dibenzo[<i>b,d</i>]pyran-6-one 350.	$C_{13}H_{18}O_2$	2-Hydroxy-2-methyl-1-[4-(2-propyl)phenyl]propan-1-one 626.
	Xanthone 1121.	$C_{13}H_{19}NO$	2-[1-(Butylimino)ethyl]-5-methylphenol 234.
$C_{13}H_8S_2$	9 <i>H</i> -Thioxanthene-9-thione 1068.	$C_{13}H_{20}N_2O$	1,9-Bis(dimethylamino)-1,3,6,8-nonatetraen-5-one 194.
$C_{13}H_9ClO$	4-Chlorobenzophenone 266.		1,9-Bis(dimethylamino)-1,4,6,8-nonatetraen-3-one 195.
$C_{13}H_9FO$	4-Fluorobenzophenone 595.		β -Ionone 638.
$C_{13}H_9N$	Acridine 23.		Anthracene- <i>d</i> ₁₀ 65.
	Benzo[<i>f</i>]quinoline 139.	$C_{13}H_{20}O$	Phenanthrene- <i>d</i> ₁₀ 855.
	Benzo[<i>h</i>]quinoline 140.	$C_{14}D_{10}$	<i>trans</i> -Stilbene- <i>d</i> ₁₂ 1007.
	4-Cyanobiphenyl 309.	$C_{14}D_{12}$	1,8-Dichloroanthraquinone 371.
	Phenanthridine 859.	$C_{14}H_8Cl_2O_2$	1,5,9,10-Tetrachloroanthracene 1022.
$C_{13}H_9NO$	9(10 <i>H</i>)-Acridinone 29.	$C_{14}H_6Cl_4$	Anthraquinone-2,6-disulfonate ion 77.
	6(5 <i>H</i>)-Phenanthridinone 860.	$C_{14}H_6O_8S_2^-$	1-Chloroanthraquinone 263.
$C_{13}H_9NO_4S$	9,10-Dihydro-9-oxo-2-acridinesulfonic acid 425.	$C_{14}H_7ClO_2$	2-Chloroanthraquinone 264.
$C_{13}H_{10}$	Fluorene 585.		2,9,10-Trichloroanthracene 1077.
	1,4-Perinaphthylidyl 851.	$C_{14}H_7Cl_3$	<i>trans</i> -Stilbene-2,3,4,5,6- <i>d</i> ₅ 1002.
$C_{13}H_{10}N^+$	Acridinium ion 28.	$C_{14}H_7D_5$	9,10-Anthraquinone-2-sulfonate ion 78.
	Benzo[<i>f</i>]quinolizinium 141.	$C_{14}H_7O_5S^-$	9,10-Dibromoanthracene 358.
	Benzo[<i>h</i>]quinolizinium 142.	$C_{14}H_8Br_2$	1,5-Dichloroanthracene 369.
$C_{13}H_{10}N_2$	9-Aminoacridine 33.	$C_{14}H_8Cl_2$	9,10-Dichloroanthracene 370.
$C_{13}H_{10}N_2O_2$	1-Amino-7-nitrofluorene 52.		Anthraquinone 76.
$C_{13}H_{10}O$	Benzophenone 133.	$C_{14}H_8O_2$	9-Bromoanthracene 215.
	4-Biphenylcarboxaldehyde 175.	$C_{14}H_9Br$	9-Bromophenanthrene 227.
	1-Hydroxyfluorene 621.		1-Chloroanthracene 262.
	2-Hydroxyfluorene 622.	$C_{14}H_9Cl$	4-(Trifluoromethyl)benzophenone 1082.
$C_{13}H_{10}O_2$	2-Hydroxybenzophenone 613.	$C_{14}H_9F_3O$	2-Iodoanthracene 632.
	4-Hydroxybenzophenone 614.	$C_{14}H_9I$	9-Iodoanthracene 633.
$C_{13}H_{10}O_3$	3-Formyl-2-methylfurochromene 601.		1-Aminoanthraquinone 36.
	3-Formyl-8-methylfurochromene 602.	$C_{14}H_9NO_2$	2-Aminoanthraquinone 37.
$C_{13}H_{10}S$	Thiobenzophenone 1060.		9-Nitroanthracene 815.
$C_{13}H_{11}ClO$	3-Chloro-1-(2-naphthyl)-1-propanone 281.	$C_{14}H_9NO_3$	1-Amino-4-hydroxyanthraquinone 45.
$C_{13}H_{11}N$	Acridan 22.	$C_{14}H_{10}$	Anthracene 64.
	2-Aminofluorene 44.		Benz[<i>a</i>]azulene 108.
$C_{13}H_{11}NO$	3-Aminobenzophenone 41.		Diphenylacetylene 527.
	4-Aminobenzophenone 42.		Phenanthrene 854.
$C_{13}H_{11}NS$	10-Methylphenothiazine 744.	$C_{14}H_{10}Cl_2$	4,4'-Dichlorostilbene 378.
$C_{13}H_{11}N_3$	Proflavine 921.	$C_{14}H_{10}D_2$	<i>trans</i> -Stilbene-2,2'- <i>d</i> ₂ 1001.
$C_{13}H_{11}N_4O_2^-$	Lumiflavine, negative ion 647.		<i>trans</i> -Stilbene-4,4'- <i>d</i> ₂ 1003.
$C_{13}H_{11}O^+$	Benzophenone, conjugate acid 134.		<i>trans</i> -Stilbene-7,7'- <i>d</i> ₂ 1004.
$C_{13}H_{12}$	2,3-Dihydrophenalene 426.	$C_{14}H_{10}N_2O$	2,5-Diphenyl-1,3,4-oxadiazole 545.
	1-Methyl-1-(1-naphthyl)ethylene 738.	$C_{14}H_{10}N_2O_2$	1,4-Diaminoanthraquinone 335.
	2-(1-Methylethenyl)naphthalene 739.	$C_{14}H_{10}N_2O_4$	<i>trans</i> -2,4-Dinitrostilbene 522.
	1-Methyl-2-vinylnaphthalene 762.		<i>cis</i> -4,4'-Dinitrostilbene 523.
	2-Methyl-3-vinylnaphthalene 763.		<i>trans</i> -4,4'-Dinitrostilbene 524.
$C_{13}H_{12}ClN_3S$	Azure C 99.		
$C_{13}H_{12}N_3^+$	Proflavine, conjugate monoacid 923.		
$C_{13}H_{12}N_4O_2$	Lumiflavine 645.		
$C_{13}H_{12}OS$	1-Naphthalenecarbothioic acid, <i>O</i> -ethyl ester 774.		

$C_{14}H_{10}O$	1-Anthrol 80. 2-Anthrol 81. 9-Anthrol 82. Anthrone 83. 9-Phenanthrol 627.	$C_{14}H_{17}NO_2$ $C_{14}H_{17}NO_4$	7-Diethylamino-4-methylcoumarin 390. <i>cis,cis</i> -1-(3'-Methoxy-5'-nitro-2'-oxo-3',5'-cyclohexadienyl)-3,4-dimethyl-1,3-pentadiene 675.
$C_{14}H_{10}OS$	2-Methylthioxanthen-9-one 749.	$C_{14}H_{18}N_2O_2$	<i>N</i> -[2-(Diethylamino)ethyl]phthalimide 389.
$C_{14}H_{10}O_2$	Benzil 112.	$C_{15}D_{12}$	9-Methylanthracene- <i>d</i> ₁₂ 701.
$C_{14}H_{10}O_3$	4-Carboxybenzophenone 249.	$C_{15}H_8N_2O_2$	9-Cyano-10-nitroanthracene 315.
$C_{14}H_{10}O_5$	3-Carboxyypsoralen 246.	$C_{15}H_9N$	9-Cyanoanthracene 308.
$C_{14}H_{11}Br$	<i>trans</i> -4-Bromostilbene 229.	$C_{15}H_9O_2^-$	2-Anthroate ion 79.
$C_{14}H_{11}Cl$	<i>cis</i> -4-Chlorostilbene 288. <i>trans</i> -4-Chlorostilbene 289.	$C_{15}H_{10}N_2O_3$	5-Nitro-2-(2-quinolylolethenyl)furan 817.
$C_{14}H_{11}F$	<i>trans</i> -4-Fluorostilbene 596.	$C_{15}H_{10}O$	Anthracene-9-carboxaldehyde 68.
$C_{14}H_{11}N$	9-Aminophenanthrene 56. 9-Methylacridine 693.	$C_{15}H_{10}O_2$	Anthracene-9-carboxylic acid 70. Flavone 583.
$C_{14}H_{11}NO_2$	<i>trans</i> -3-Nitrostilbene 823. <i>trans</i> -4-Nitrostilbene 824.	$C_{15}H_{10}O_3$	3-Hydroxyflavone 619. 7-Hydroxyflavone 620.
$C_{14}H_{11}NS$	10-Methyl-9(10 <i>H</i>)-acridinethione 694.	$C_{15}H_{11}N$	<i>trans</i> -4-Cyanostilbene 316.
$C_{14}H_{11}O_4S^-$	4-Sulfomethylbenzophenone anion 1010.	$C_{15}H_{11}NO$	Anthracene-9-carboxamide 69. 2,5-Diphenyloxazole 546.
$C_{14}H_{12}$	3,4-Dihydrophenanthrene 427. 9,10-Dihydrophenanthrene 428. 1,1-Diphenylethylene 539. Stilbene 1000. <i>cis</i> -Stilbene 1005. <i>trans</i> -Stilbene 1006.	$C_{15}H_{11}O_5S^-$ $C_{15}H_{12}$	Quantacure SKS anion 958. 5 <i>H</i> -Dibenzo[<i>a,d</i>]cycloheptene 348. 2-Methylanthracene 699. 9-Methylanthracene 700. 2-Phenylindene 890.
$C_{14}H_{12}N^+$	2-Aminoanthracene, conjugate acid 35.	$C_{15}H_{12}NO_3$	4'-Aminomethyl-4,5',8-trimethylpsoralen 51.
$C_{14}H_{12}N_2$	3,8-Dimethyl-4,7-phenanthroline 499.	$C_{15}H_{12}N_2O_2$	1-Amino-4-(<i>N</i> -methylamino)-anthraquinone 48.
$C_{14}H_{12}N_2O_2$	<i>trans</i> -4-Amino-4'-nitrostilbene 54.	$C_{15}H_{12}O$	Chalcone 258.
$C_{14}H_{12}N_5O_2Br$	4-[(2-Bromo-4,6-dinitrophenyl)azo]- <i>N,N</i> -dimethylbenzenamine 220.	$C_{15}H_{12}O_2$	Dibenzoylmethane 355.
$C_{14}H_{12}O$	4-Acetylbiphenyl 14. Deoxybenzoin 332. 2-Methylbenzophenone 706. 4-Methylbenzophenone 707.	$C_{15}H_{13}ClO$	1-(4-Biphenyl)-3-chloro-1-propanone 179.
$C_{14}H_{12}O_2$	Benzoin 124. 4-Methoxybenzophenone 664.	$C_{15}H_{13}NO$ $C_{15}H_{13}NO_3$	10-Ethyl-9(10 <i>H</i>)-acridinone 569. <i>trans</i> -3-Methoxy-4'-nitrostilbene 676. <i>trans</i> -4-Methoxy-4'-nitrostilbene 678.
$C_{14}H_{12}O_3$	4,5',8-Trimethylpsoralen 1091.	$C_{15}H_{13}N_2^+$	1-Methyl-4-[4-cyanostyryl]pyridinium 713.
$C_{14}H_{12}O_5$	4',5'-Dihydro-3-carboxyypsoralen 411.	$C_{15}H_{13}NO_3$	<i>cis</i> -4-Methoxy-4'-nitrostilbene 677.
$C_{14}H_{13}N_2O_2^+$	1-Methyl-4-[4-nitrostyryl]pyridinium 742.	$C_{15}H_{14}$	1-(1-Cyclopenten-1-yl)naphthalene 328.
$C_{14}H_{13}N_4O_2Br$	4-[(2-Bromo-4-nitrophenyl)azo]- <i>N,N</i> -dimethylbenzenamine 226.	$C_{15}H_{14}O$	2-(1-Cyclopenten-1-yl)naphthalene 329.
$C_{14}H_{13}N_5O_4$	4-[(2,4-Dinitrophenyl)azo]- <i>N,N</i> -dimethylbenzenamine 520.	$C_{15}H_{14}O_2S$	<i>trans</i> -4-Methoxystilbene 685.
$C_{14}H_{13}O^+$	Benzyl phenyl ketone, conjugate acid 161.	$C_{15}H_{14}O_3$	1-Methyldeoxybenzoin 716.
$C_{14}H_{14}$	3,3'-Dimethylbiphenyl 476.	$C_{15}H_{15}N_3O_2$	4,4'-Dimethoxythiobenzophenone 454.
$C_{14}H_{14}ClN_3S$	Azure A 96.	$C_{15}H_{16}$	4,4'-Dimethoxybenzophenone 445.
$C_{14}H_{14}N_2$	5,10-Dihydro-5,10-dimethylphenazine 416.	$C_{15}H_{16}BrN_3S$ $C_{15}H_{16}N_3^+$	3-Methyl-5-deazaluminiflavine 715. 1,1,2-Trimethyl-2-(2-naphthyl)ethylene 1087.
$C_{14}H_{14}N_2O_4S$	Phenazine methosulfate 743.	$C_{15}H_{16}N_4^+$	Azure B 98.
$C_{14}H_{14}N_3^+$	Acriflavine cation 30.	$C_{15}H_{18}$	Acridine Yellow, conjugate monoacid 27.
$C_{14}H_{14}N_3S^+$	Azure A cation 97.	$C_{15}H_{18}N_4^{3+}$	Auorophosphine, conjugate monoacid 94.
$C_{14}H_{14}N_4O_2$	<i>N,N</i> -Dimethyl-4-[(4-nitrophenyl)azo]-benzenamine 498.	$C_{15}H_{18}O_2$	Neutral Red cation 809.
$C_{14}H_{14}O_2$	4,4'-Dimethoxybiphenyl 448. 2-(1-Naphthalenyl)-1,3-dioxane 779. 2-(2-Naphthalenyl)-1,3-dioxane 780.	$C_{15}H_{20}ClN_5$	Guaiazulene 488. Neutral Red cation, conjugate diacid 810.

$C_{15}H_{20}N_2O_2$	<i>N</i> -[(Dipropylamino)methyl]phthalimide 550.	$C_{16}H_{16}$	1-(1-Cyclohexen-1-yl)naphthalene 323.
$C_{15}H_{22}N_2O$	<i>all-trans</i> -1,11-Bis(dimethylamino)-1,3,6,8,10-undecapentaen-5-one 203.	$C_{16}H_{16}N_2OS_3$	[2.2]Paracyclophane 844.
$C_{15}H_{22}O$	(<i>E,E</i>)-3-Methyl-5-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2,4-pentadienal 755.	$C_{16}H_{16}N_2O_2$	(<i>E,Z</i>)-3-Ethyl-5[(3-ethyl-2(3 <i>H</i>)-benzothiazolylidene)ethylidene]-2-thioxo-4-thiazolidinone 571.
	(<i>E,Z</i>)-3-Methyl-5-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2,4-pentadienal 756.		<i>trans</i> -4-Dimethylamino-4'-nitrostilbene 461.
	(<i>Z,E</i>)-3-Methyl-5-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2,4-pentadienal 757.	$C_{16}H_{16}O_2$	1-(4-Methoxyphenyl)-3-phenyl-1-propanone 891.
	(<i>Z,Z</i>)-3-Methyl-5-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2,4-pentadienal 758.	$C_{16}H_{16}O_2S$	<i>O</i> -(2-Phenylethyl) 4-methoxybenzenecarbothioate 887.
$C_{16}D_{10}$	Pyrene- <i>d</i> ₁₀ 940.	$C_{16}H_{17}N_4O_2S^+$	Methylene Green cation 722.
$C_{16}H_8N_2$	9,10-Dicyanoanthracene 381.	$C_{16}H_{18}$	1-(3,3-Dimethyl-1-buten-2-yl)naphthalene 478.
$C_{16}H_8O_2S_2$	Thioindigo 1062.		<i>cis</i> -3,3-Dimethyl-1-(2-naphthyl)-1-butene 494.
$C_{16}H_6Br$	1-Bromopyrene 228.		<i>trans</i> -3,3-Dimethyl-1-(2-naphthyl)-1-butene 495.
$C_{16}H_9O_3S^-$	Pyrene-1-sulfonate 946.		Euchrysine, conjugate monoacid 580.
	Pyrene-3-sulfonate 947.	$C_{16}H_{18}N_3^+$	Methylene Blue cation 719.
$C_{16}H_{10}$	1,4-Diphenylbutadiyne 536.	$C_{16}H_{18}N_3S^+$	9,9-Diethyl-9 <i>H</i> -9-stannafuorene 402.
	Fluoranthene 584.	$C_{16}H_{18}Sn$	Methylene Blue cation, conjugate monoacid 720.
	Pyrene 939.	$C_{16}H_{19}N_3S^{2+}$	<i>N,N,N',N'</i> -Tetramethylbenzidine 1045.
$C_{16}H_{10}N_2$	Benzophenazine 132.	$C_{16}H_{20}N_2$	1-Naphthyl diisopropyl borate 794.
	Dibenzo[<i>f,h</i>]quinoxaline 351.	$C_{16}H_{21}BO_3$	2-Naphthyl diisopropyl borate 795.
	<i>trans</i> -4,4'-Dicyanostilbene 386.	$C_{16}H_{24}N_2O$	2,6-Bis[3-(dimethylamino)-2-propenylidene]cyclohexanone 199.
$C_{16}H_{10}O$	4-Hydroxypyrene 629.		Benzanthrone 107.
$C_{16}H_{11}Cl$	1-(2-Chlorophenyl)naphthalene 282.	$C_{17}H_{10}O$	1-Pyrenecarboxaldehyde 945.
	1-(4-Chlorophenyl)naphthalene 283.	$C_{17}H_{10}O_2$	6-Hydroxybenzanthrone 612.
$C_{16}H_{11}N$	Benzo[<i>a</i>]carbazole 114.	$C_{17}H_{11}N$	Benz[<i>a</i>]acridine 101.
	Benzo[<i>b</i>]carbazole 115.		Benz[<i>c</i>]acridine 102.
	Benzo[<i>c</i>]carbazole 116.	$C_{17}H_{11}NO$	6-Aminobenzanthrone 38.
$C_{16}H_{11}NO$	4'-Cyanochalcone 310.		Benz[<i>b</i>]acridin-12(5 <i>H</i>)-one 103.
$C_{16}H_{12}$	5,10-Dihydroindeno[2,1- <i>a</i>]indene 420.		2-(1-Naphthyl)benzoxazole 792.
	1-Phenyl-naphthalene 893.	$C_{17}H_{12}$	Benzo[<i>a</i>]fluorene 120.
	2-Phenyl-naphthalene 894.		Benzo[<i>b</i>]fluorene 121.
$C_{16}H_{12}N_2O_3$	4',5'-Psoralen-thymine photo adduct 935.		Benzo[<i>c</i>]fluorene 122.
$C_{16}H_{12}O$	1-Acetylanthracene 12.		8 <i>b</i> ,9 <i>a</i> -Dihydro-9 <i>H</i> -cyclopropa[<i>e</i>]pyrene 413.
	9-Acetylanthracene 13.		8,11 <i>b</i> -Methanocyclodeca[<i>cde</i>]naphthalene 660.
$C_{16}H_{12}O_2$	3-Acetyl-9,10-epoxy-9,10-dihydro-phenanthrene 15.	$C_{17}H_{12}O$	9-Anthryl ethenyl ketone 87.
$C_{16}H_{13}N$	1-Anilinonaphthalene 62.		2-Methylaceanthren-1-one 690.
	<i>N</i> -Phenyl-2-naphthylamine 895.	$C_{17}H_{13}N_2O_5S^-$	Brilliant Sulfaflavine anion 214.
$C_{16}H_{13}NO$	1-Benzylisoquinoline- <i>N</i> -oxide 160.	$C_{17}H_{14}NO_3S^-$	<i>N</i> -Methyl-2-anilino-6-naphthalenesulfonate ion 698.
	<i>trans</i> -4-Cyano-4'-methoxystilbene 313.		10-Methyl-5-deazaisoalloxazine-3-propanesulfonate ion 714.
$C_{16}H_{14}$	1,2-Dihydro-3-phenyl-naphthalene 429.	$C_{17}H_{14}N_3O_5S^-$	9-Propionylantracene 925.
	1,3-Dimethylantracene 472.		9-Anthraceneacetic acid, methyl ester 67.
	9,10-Dimethylantracene 473.	$C_{17}H_{14}O_2$	<i>N</i> -Methyl-1-anilinonaphthalene 696.
	1,4-Diphenyl-1,3-butadiene 535.	$C_{17}H_{15}N$	<i>N</i> -Methyl-2-anilinonaphthalene 697.
	1,2-Diphenylcyclobutene 537.		2-[1-(4-Methoxyphenyl)-hydrazinyl-2-ylidene]-1-(4-nitrophenyl)-1,3-diketobutane 680.
$C_{16}H_{14}N_2OS_3$	9-Ethylphenanthrene 576.		Methyl-1-naphthylphenylgermane 740.
	(<i>E</i>)-5[(3,4-Dihydro[1,4]thiazino-[3,4- <i>b</i>]benzothiazol-1-yl)methylene]-3-ethyl-2-thioxo-4-thiazolidinone 435.	$C_{17}H_{15}N_3O_5$	
$C_{16}H_{14}N_2O_2$	1,4-Bis(methylamino)anthraquinone 207.		
$C_{16}H_{14}OS$	2-Isopropylthioxanthen-9-one 639.		
$C_{16}H_{14}O_2$	4-Methoxychalcone 667.		
$C_{16}H_{14}S$	1-Phenylthio-3,4-dihydronaphthalene 900.		
$C_{16}H_{15}NO_4$	<i>trans</i> -2,5-Dimethoxy-4'-nitrostilbene 453.		

$C_{17}H_{16}N_6O_5$	3-[[4-[(2,4-Dinitrophenyl)azo]phenyl]- (2-hydroxyethyl)amino]- 4-propanenitrile 521 .	$C_{18}H_{14}$	1-(1-Naphthyl)-1-phenylethylene 800 . <i>trans</i> -1-(1-Naphthyl)-2-phenylethylene 801 . <i>trans</i> -1-(2-Naphthyl)-2-phenylethylene 802 . <i>m</i> -Terphenyl 1012 . <i>o</i> -Terphenyl 1013 . <i>p</i> -Terphenyl 1014 . Tricyclo[8.2.2.2 ^{4,7}]hexadeca- 4,6,10,12,13,15-hexaene- 5,15-dicarbonitrile 1079 . <i>N,N'</i> -Dimethylindigo 487 . Phenosafranin 869 . Triphenylamine 1092 . 1,6-Diphenyl-1,3,5-hexatriene 541 . Triphenylamine, conjugate acid 1093 . <i>N,N'</i> -Diphenyl- <i>p</i> -phenylenediamine 547 . [1,1':4',1''-Terphenyl]-4,4''-diamine 1011 . 9-Butyrylanthracene 239 . 1-[[[(Phenylmethyl)sulfonyl]methyl]- naphthalene 892 . <i>N,N'</i> -Diphenyl- <i>p</i> -phenylenediamine, conjugate acid 548 . 1-(1-Cycloocten-1-yl)naphthalene 325 . 6,6'-Biquinoline, conjugate monoacid 187 . D-Glucose phenylosazone 603 . 6,6'-Biquinoline, conjugate diacid 185 . (<i>E,E,E</i>)-6-Methyl-8-(2,6,6-trimethyl- 1-cyclohexen-1-yl)- 3,5,7-octatrien-2-one 751 . (<i>E,E,Z</i>)-6-Methyl-8-(2,6,6-trimethyl- 1-cyclohexen-1-yl)- 3,5,7-octatrien-2-one 752 . (<i>E,Z,E</i>)-6-Methyl-8-(2,6,6-trimethyl- 1-cyclohexen-1-yl)- 3,5,7-octatrien-2-one 753 . (<i>E,Z,Z</i>)-6-Methyl-8-(2,6,6-trimethyl- 1-cyclohexen-1-yl)- 3,5,7-octatrien-2-one 754 . Dicumarol 380 . 2-(4-Biphenyl)benzoxazole 178 . 2-(1-Naphthyl)-5-phenyloxazole 805 . 1-(1 <i>H</i> -Inden-3-yl)naphthalene 630 . 4-Benzoylbiphenyl 879 . Triphenylmethyl cation 1101 . 9-Phenylacridan 874 . <i>N</i> -Phenylacridan 875 . Triphenylmethane 1100 . 5,10-Dihydro-5-methyl-10-phenyl- phenazine 422 . 9-Phenylproflavine, conjugate mono- acid 898 . <i>O</i> -(2-Phenylethyl) 1-naphthalenecarbothioate 888 . <i>O</i> -(2-Phenylethyl) 2-naphthalenecarbothioate 889 . 6-Chloro-1,3,3-trimethyl-6'-nitrospiro- (2,2'-indolin[2 <i>H</i> -1]benzopyran) 291 .
$C_{17}H_{16}O_3$	4-(2-Methylpropionyloxy)benzophenone 746 .	$C_{18}H_{14}N_2$	
$C_{17}H_{17}NO$	4-(Dimethylamino)chalcone 455 .	$C_{18}H_{14}N_2O_2$	
$C_{17}H_{18}$	1-(1-Cyclohepten-1-yl)naphthalene 319 . 2-(1-Cyclohepten-1-yl)naphthalene 320 . <i>trans</i> -4-Cyano-4'-dimethylaminostilbene 311 .	$C_{18}H_{15}ClN_4$	
$C_{17}H_{18}N_2$		$C_{18}H_{15}N$	
$C_{17}H_{19}ClN_2O$	Pyronine 955 .	$C_{18}H_{16}$	
$C_{17}H_{19}ClN_2S$	Chlorpromazine 292 . Thioflavine 1061 . Thiopyronine 1065 . <i>O</i> -(2-Phenylethyl) 4-(dimethylamino)- benzenecarbothioate 886 .	$C_{18}H_{16}N^+$	
$C_{17}H_{19}NOS$		$C_{18}H_{16}N_2$	
$C_{17}H_{19}N_2S^+$	Thiopyronine cation 1066 .		
$C_{17}H_{19}N_2Se^+$	Selenopyronine 995 .	$C_{18}H_{16}O$	
$C_{17}H_{19}N_3$	Acridine Orange, free base 26 .	$C_{18}H_{16}O_2S$	
$C_{17}H_{20}N_2O$	4,4'-Bis(dimethylamino)benzophenone 190 .	$C_{18}H_{17}N_2^+$	
$C_{17}H_{20}N_2S$	4,4'-Bis(dimethylamino)thiobenzophenone 200 . Promazine 924 . Acridine Orange, conjugate monoacid 25 .	$C_{18}H_{20}$	
$C_{17}H_{20}N_3^+$		$C_{18}H_{22}N_3^+$	
$C_{17}H_{20}N_3S^+$	Neomethylene Blue cation 807 .	$C_{18}H_{22}N_4O_4$	
$C_{17}H_{21}N_3$	Auramine O 93 .	$C_{18}H_{23}N_3^+$	
$C_{17}H_{21}N_4O_6^+$	Riboflavin, conjugate monoacid 987 . <i>all-trans</i> -1,13-Bis(dimethylamino)- 1,3,5,8,10,12-tridecahexaen-7-one 201 . <i>all-trans</i> -1,13-Bis(dimethylamino)- 1,3,6,8,10,12-tridecahexaen-5-one 202 . γ -[2-(Formylamino)phenyl]- γ -oxo- α -aminobutyl-L-lysine 597 . (<i>E,E,E</i>)-5-Methyl-7-(2,6,6-trimethyl- 1-cyclohexen-1-yl)- 2,4,6-heptatrienal 750 . Triphenylene- <i>d</i> ₁₂ 1096 . <i>p</i> -Terphenyl- <i>d</i> ₁₄ 1015 . 2-Bromotriphenylene 230 . Benz[<i>a</i>]anthracene 106 . Benzo[<i>c</i>]phenanthrene 131 . Chrysene 295 . Benz[<i>b</i>]anthracene 1021 . Triphenylene 1095 . 2,2'-Biquinoline 184 . 2-(1-Naphthyl)-5-phenyl-1,3,4-oxadiazole 803 . 2-(2-Naphthyl)-5-phenyl-1,3,4-oxadiazole 804 . 2,6-Diphenyl-1,4-benzoquinone 531 . 3-Methoxybenzanthrone 663 . 9-Phenyl-9 <i>H</i> -9-bisfluorene 880 . 4-Nitro- <i>p</i> -terphenyl 825 . 2,2'-Biquinoline, conjugate monoacid 186 . 9-Phenyl-9 <i>H</i> -9-phosphafluorene 897 . 9-Phenyl-9 <i>H</i> -9-stibafluorene 899 .	$C_{18}H_{26}O$	
$C_{17}H_{24}N_4O_5$		$C_{19}H_{12}O_6$	
$C_{17}H_{24}O$		$C_{19}H_{13}NO$	
$C_{18}D_{12}$		$C_{19}H_{14}$	
$C_{18}D_{14}$		$C_{19}H_{14}O$	
$C_{18}H_{11}Br$		$C_{19}H_{15}^+$	
$C_{18}H_{12}$		$C_{19}H_{15}N$	
$C_{18}H_{12}N_2$		$C_{19}H_{16}$	
$C_{18}H_{12}N_2O$		$C_{19}H_{16}N_2$	
$C_{18}H_{17}O_7$		$C_{19}H_{16}N_3^+$	
$C_{18}H_{13}Bi$		$C_{19}H_{16}OS$	
$C_{18}H_{13}NO_2$			
$C_{18}H_{13}N_2^+$			
$C_{18}H_{13}P$			
$C_{18}H_{13}Sb$			

$C_{19}H_{17}NO_2$	2-Piperidinoanthraquinone 911.	$C_{20}H_{16}O$	1-Phenyldeoxybenzoin 882.
$C_{19}H_{18}N_2O_3$	1,3,3-Trimethyl-6'-nitrospiro-(2,2'-indolin[2 <i>H</i> -1]benzopyran) 1088.	$C_{20}H_{16}O_2Se_2$	5,5'-Diethylselenoindigo 401.
$C_{19}H_{19}IN_2S_2$	3,3'-Diethyl-2,2'-thiacyanine iodide 404.	$C_{20}H_{16}O_3$	7-Oxa-2,3-dibenzoyl-bicyclo[2.2.1]hept-2-ene 834.
$C_{19}H_{20}O_2$	2,4-Dimethyl-1,5-diphenyl-1,5-pentanedione 482.	$C_{20}H_{16}O_4$	7,8,9,10-Tetrahydrotetrahydroxybenzo[<i>a</i>]pyrene 1031.
	2-(2-Naphthyl-3,5,5-trimethyl-2-cyclohexen-1-one 799.	$C_{20}H_{16}O_4S_2$	6,6'-Diethoxythioindigo 388.
$C_{19}H_{23}BO_3$	1-Fluorenyl diisopropyl borate 587.	$C_{20}H_{18}$	2,2-Dimethyl-1-(1-naphthyl)-1-phenylethylene 496.
$C_{19}H_{26}N_2O$	2-Fluorenyl diisopropyl borate 588.		1,8-Diphenyl-1,3,5,7-octatetraene 544.
	<i>all-trans</i> -1,15-Bis(dimethylamino)-1,3,6,8,10,12,14-pentadecaheptaen-5-one 196.	$C_{20}H_{18}N_2$	12 <i>a</i> ,12 <i>b</i> -Dihydro-5,8-dimethyl-(indolino[2,3- <i>c</i>]carbazole) 415.
	2,5-Bis[5-(dimethylamino)-2,4-pentadienylidene]cyclopentanone 198.	$C_{20}H_{19}N_4^+$	<i>N,N'</i> -Dimethyl-3-anilinocarbazole 471.
$C_{19}H_{28}O_2$	Testosterone 1018.	$C_{20}H_{20}N_2$	Safranine cation 990.
$C_{20}H_4Cl_4I_4O_5$	Rose Bengal 988.		<i>N,N'</i> -Dimethyl- <i>N,N'</i> -diphenyl-1,4-phenylenediamine 483.
$C_{20}H_6Br_4Na_2O_5$	Eosin 556.	$C_{20}H_{20}N_2O_3$	1,3,3,6-Tetramethyl-6'-nitrospiro-(2,2'-indolin[2 <i>H</i> -1]benzopyran) 1048.
$C_{20}H_6Br_4O_5^{2-}$	Eosin dianion 562.	$C_{20}H_{20}N_2O_4$	1-(2-Hydroxyethyl)-3,3-dimethyl-6'-nitrospiro-(2,2'-indolin[2 <i>H</i> -1]benzopyran) 618.
$C_{20}H_6I_4O_5^{2-}$	Erythrosin dianion 566.		6-Methoxy-1,3,3-trimethyl-6'-nitrospiro-(2,2'-indolin[2 <i>H</i> -1]benzopyran) 687.
$C_{20}H_8Br_2O_5^{2-}$	Dibromofluorescein dianion 360.		7'-Methoxy-1,3,3-trimethyl-6'-nitrospiro-(2,2'-indolin[2 <i>H</i> -1]benzopyran) 688.
$C_{20}H_8I_4O_5$	Erythrosin 565.	$C_{20}H_{20}N_4^{2+}$	Safranine cation, conjugate monoacid 992.
$C_{20}H_{10}O_5^{2-}$	Fluorescein dianion 593.	$C_{20}H_{20}N_7O_7Br$	<i>N</i> -[2-[(2-Bromo-4,6-dinitrophenyl)azo]-5-[(2-cyanoethyl)(2-hydroxyethyl)-amino]-4-methoxyphenyl]acetamide 219.
$C_{20}H_{12}$	Benzo[<i>a</i>]pyrene 137.		<i>N</i> -[2-[(2-Chloro-4,6-dinitrophenyl)azo]-5-[(2-cyanoethyl)(2-hydroxyethyl)-amino]-4-methoxyphenyl]acetamide 270.
	Benzo[<i>e</i>]pyrene 138.	$C_{20}H_{21}N_4^{3+}$	Safranine cation, conjugate diacid 991.
	Perylene 852.	$C_{20}H_{21}N_7O_7$	<i>N</i> -[5-[(2-Cyanoethyl)(2-hydroxyethyl)-amino]-2-[(2,4-dinitrophenyl)azo]-4-methoxyphenyl]acetamide 312.
$C_{20}H_{12}N_2$	Dibenzophenazine 349.		1-Anthryl diisopropyl borate 84.
$C_{20}H_{12}N_2O_2$	5,12-Dihydroquino[2,3- <i>b</i>]acridine-7,14-dione 431.	$C_{20}H_{23}BO_3$	2-Anthryl diisopropyl borate 85.
	2,2'-(1,4-Phenylene)bisbenzoxazole 883.		9-Anthryl diisopropyl borate 86.
$C_{20}H_{12}N_4Zn$	Zinc(II) porphyrin 1131.	$C_{20}H_{24}N_2O_4$	<i>N,N'</i> -Dipentylpyromellitic diimide 525.
$C_{20}H_{12}O_5$	Fluorescein 591.	$C_{20}H_{24}O_4$	α -Crocin 305.
$C_{20}H_{13}N$	7 <i>H</i> -Dibenzo[<i>c,g</i>]carbazole 342.	$C_{20}H_{26}N_3O^+$	<i>N,N,N',N'</i> -Tetraethyloxonine cation 1024.
$C_{20}H_{13}O_5^+$	Fluorescein, conjugate monoacid 592.		<i>N,N,N',N'</i> -Tetraethyloxonine cation, conjugate monoacid 1025.
$C_{20}H_{14}$	1,1'-Binaphthyl 171.	$C_{20}H_{27}N_3O^{2+}$	2,6-Bis[5-(dimethylamino)-2,4-pentadienylidene]cyclohexanone 197.
	2,2'-Binaphthyl 172.	$C_{20}H_{28}N_2O$	Retinal 965.
	9-Phenylanthracene 878.		11- <i>cis</i> -Retinal 966.
$C_{20}H_{14}N_2O$	2-(4-Biphenyl)-5-phenyl-1,3,4-oxadiazole 181.	$C_{20}H_{28}O_2$	13- <i>cis</i> -Retinal 967.
$C_{20}H_{14}N_2O_2$	1-Amino-4-anilinoanthraquinone 34.	$C_{20}H_{30}O$	7,9- <i>cis</i> -Retinal 968.
$C_{20}H_{14}N_2O_4$	<i>trans-N,N'</i> -Diacetylindigo 334.	$C_{21}H_{13}BrO$	7- <i>cis</i> -Retinal 969.
$C_{20}H_{14}N_4$	21 <i>H</i> ,23 <i>H</i> -Porphine 920.	$C_{21}H_{13}ClO$	9- <i>cis</i> -Retinal 970.
$C_{20}H_{14}O$	2-Fluorenyl phenyl ketone 590.		<i>all-trans</i> -Retinal 972.
$C_{20}H_{14}O_3$	7-Oxa-2,3-dibenzoylbicyclo[2.2.1]-hepta-2,5-diene 833.		<i>all-trans</i> -Retinoic acid 973.
$C_{20}H_{14}S$	1-(2-Anthryl)-2-(2-thienyl)ethylene 90.		<i>all-trans</i> -Retinol 974.
$C_{20}H_{15}ClN_2O_3$	Rhodamine 110 982.		9-Benzoyl-10-bromoanthracene 149.
$C_{20}H_{15}N$	Di-2-naphthylamine 504.		9-Benzoyl-10-chloroanthracene 150.
$C_{20}H_{15}NOS$	2-Benzoyl- <i>N</i> -methyl- β -naphthiazoline 152.		
$C_{20}H_{16}$	1-Acenaphthyl-1-phenylethylene 4.		
	2-(1-Phenylethenyl)biphenyl 180.		
	3,4-Dihydro-1,1'-binaphthyl 409.		
	Triphenylethylene 1098.		
$C_{20}H_{16}Cl_2IrN_4^+$	Bis(2,2'-bipyridine)dichloroiridium(III) ion 188.		
$C_{20}H_{16}N_2$	<i>N,N'</i> -Dimethyl-5,11-dihydro-indolo[3,2- <i>b</i>]carbazole 479.		

$C_{21}H_{13}N$	Dibenz[<i>a,h</i>]acridine 339.	$C_{22}H_{16}O_2$	Benzyl 9-anthroate 159.
$C_{21}H_{13}NO_3$	9-Benzoyl-10-nitroanthracene 154.		(<i>R</i>)-4,5-Dihydrodinaphtho[2,1- <i>e</i> :1',2'- <i>g</i>] [1,4]dioxocin 417.
$C_{21}H_{14}N_2O_2$	5,12-Dihydro-5-methyl- quino[2,3- <i>b</i>]acridine-7,14-dione 423.		3,3,5-Triphenyl-2(3 <i>H</i>)-furanone 1099.
$C_{21}H_{14}O$	9-Benzoylanthracene 148.	$C_{22}H_{18}$	3,9-Dimethyl- <i>trans</i> -fluorenacene 484.
$C_{21}H_{14}O_2$	(<i>S</i>)-Dinaphtho[2,1- <i>d</i> :1',2'- <i>f</i>][1,3]dioxepin 503.	$C_{22}H_{18}O_2$	1,2-Di-1-naphthylethane 507.
$C_{21}H_{15}NO$	2-(4-Biphenyl)-5-phenyloxazole 182.		2,3-Dibenzoylbicyclo[2.2.2]octa-2,5- diene 354.
$C_{21}H_{15}O_4P$	(<i>R</i>)-4-Methoxydinaphtho[2,1- <i>d</i> :1',2'- <i>f</i>] [1,3,2]dioxaphosphopin 4-oxide 668.	$C_{22}H_{19}N$	9-[4-(Dimethylamino)phenyl]anthracene 463.
	4-Methoxydinaphtho[2,1- <i>d</i> :1',2'- <i>f</i>] dioxaphosphopin 4-oxide 669.	$C_{22}H_{23}IN_2OS_2$	3,3'-Diethyl-9-methoxy- 2,2'-thiacarbocyanine iodide 397.
$C_{21}H_{16}$	Di-1-naphthylmethane 508.	$C_{22}H_{28}O$	2,4,6-Triisopropylbenzophenone 1084.
$C_{21}H_{16}O$	5-Methyl- <i>endo-cis</i> -fluorenacene 725.	$C_{22}H_{30}N_2O$	2-[7-(Dimethylamino)- 2,4,6-heptatrienyldiene]-
$C_{21}H_{16}O_2$	1,3,3-Triphenylprop-2-en-1-one 1102.		6-[5-(dimethylamino)- 2,4-pentadienyldiene]cyclohexanone 458.
	2,3-Dibenzoylbicyclo[2.2.1]hepta-2,5-diene 353.		11- <i>cis</i> - β - <i>apo</i> -14'-Carotenal 250.
$C_{21}H_{18}$	3,4-Dihydro-2'-methyl-1,1'-binaphthyl 421.	$C_{22}H_{30}O$	β - <i>apo</i> -14'-Carotenal 251.
	8-(1-Naphthyl)-6,7-dihydro- 5 <i>H</i> -benzocycloheptene 793.	$C_{22}H_{32}O_2$	Retinyl acetate 975.
$C_{21}H_{20}ClN_3$	Benzoflavine 119.	$C_{22}H_{36}$	1,2,3,4,5,6,7,8,9,10-Decahydro- 1,1,5,5,6,6,10,10-octamethyl- cyclobuta[1,2:3,4]dicycloheptene 330.
$C_{21}H_{20}N_7O_5Br$	<i>N</i> -[2-[(2-Bromo-6-cyano-4-nitrophenyl) azo]-5-[(2-cyanoethyl)(2-hydroxyethyl)- amino]-4-methoxyphenyl]acetamide 218.	$C_{23}H_{15}NO$	4-Anilinobenzanthrone 877.
$C_{21}H_{21}IN_2O_2$	3,3'-Diethyl-2,2'-oxacarbocyanine iodide 398.	$C_{23}H_{16}O$	8-Benzoylnaphtho[<i>de</i> -2.3.4]bicyclo[3.2.2]- nona-2,6,8-triene 153.
$C_{21}H_{21}IN_2S_2$	3,3'-Diethyl-2,2'-thiacarbocyanine iodide 403.		9-Cinnamoylanthracene 296.
$C_{21}H_{28}N_2O$	2,5-Bis[7-(dimethylamino)-2,4,6- heptatrienyldiene]cyclopentanone 193.	$C_{23}H_{18}N_2O_4$	8'-Methoxy-5-methyl-6'-nitro- spiro[(5,6-dihydrophenanthridine-6,2)- (2 <i>H</i>)chromene] 670.
$C_{21}H_{30}O_2$	<i>all-trans</i> -Methyl retinoate 748.	$C_{23}H_{18}O_2$	(<i>R</i>)-5,6-Dihydro-4 <i>H</i> -dinaphtho- [2,1- <i>f</i> :1',2'- <i>h</i>][1,5]dioxonin 418.
$C_{22}H_{10}O_2$	Dibenzo[<i>def,mno</i>]chrysene-6,12-dione 346.		3-Methyl-3,4,5-triphenyl-2(3 <i>H</i>)-furanone 759.
$C_{22}H_{12}$	Benzo[<i>ghi</i>]perylene 130.	$C_{23}H_{19}NO_4$	9-[2-(4-Methoxycarbonylbenzoxy)ethyl]- carbazole 665.
	Dibenzo[<i>def,mno</i>]chrysene 344.	$C_{23}H_{20}$	1,3-Di-1-naphthylpropane 512.
$C_{22}H_{13}NO$	9-Benzoyl-10-cyanoanthracene 151.		1,3-Di-2-naphthylpropane 513.
$C_{22}H_{14}$	Benzo[<i>b</i>]chrysene 117.	$C_{23}H_{22}BrIN_2$	1,1'-Diethyl-6-bromo-2,2'-cyanine iodide 391.
	Benzo[<i>b</i>]triphenylene 145.	$C_{23}H_{22}I_2N_2$	1,1'-Diethyl-6-iodo-2,2'-cyanine iodide 396.
	Dibenz[<i>a,h</i>]anthracene 341.	$C_{23}H_{23}IN_2$	1,1'-Diethyl-2,2'-cyanine iodide 394.
	12 <i>a</i> ,12 <i>b</i> -Dihydrobenzo[<i>ghi</i>]perylene 408.	$C_{23}H_{23}IN_2S_2$	3,3'-Diethyl-2,2'-thiadcarbocyanine iodide 405.
	Pentacene 845.	$C_{23}H_{23}NO$	4-[2-[4-(Dimethylamino)phenyl]ethyl]- benzophenone 464.
	Pentahelicene 847.	$C_{23}H_{23}N_2O_2I$	3,3'-Diethyl-2,2'-oxadcarbocyanine iodide 399.
	Pentaphene 848.	$C_{23}H_{36}N_6O_4^{2+}$	Lysyltryptophanyllysine 650.
	Picene 910.	$C_{24}D_{12}$	Coronene- <i>d</i> ₁₂ 302.
$C_{22}H_{14}N_2O$	2,5-Di(1-naphthyl)-1,3,4-oxadiazole 509.	$C_{24}H_{12}$	Coronene 301.
$C_{22}H_{16}$	1-(2-Anthryl)-2-phenylethylene 89.	$C_{24}H_{12}O_2S_2$	<i>cis</i> -Perinaphthothioindigo 849.
	1,1-Bis(1-naphthyl)ethylene 208.		<i>trans</i> -Perinaphthothioindigo 850.
	<i>trans</i> -1,2-Bis(1-naphthyl)ethylene 209.		Benzo[<i>rst</i>]pentaphene 129.
	<i>trans</i> -1,2-Bis(2-naphthyl)ethylene 210.		Dibenzo[<i>b,def</i>]chrysene 343.
	1,4-Diphenylnaphthalene 542.		Dibenzo[<i>def,p</i>]chrysene 345.
	1,5-Diphenylnaphthalene 543.		Naphtho[1,2,3,4- <i>def</i>]chrysene 783.
	<i>trans</i> -1-(1-Naphthyl)-2-(2-naphthyl)- ethylene 796.	$C_{24}H_{14}$	
$C_{22}H_{16}N_2O_3$	5-Methyl-6'-nitrospiro[(5,6-dihydro- phenanthridine-6,2)-(2 <i>H</i>)chromene] 741.		
$C_{22}H_{16}O$	3-Acetyltriptycene 20.		
	<i>cis</i> -2,3-Di-(2-naphthyl)oxirane 510.		
	<i>trans</i> -2,3-Di-(2-naphthyl)oxirane 511.		

$C_{24}H_{14}N_2$	Tribenzo[<i>a,c,h</i>]phenazine 1076.	$C_{25}H_{25}ClN_2$	1,1'-Diethyl-2,2'-carbocyanine chloride 392.
$C_{24}H_{16}$	Acenaphthylene, <i>cis</i> -photodimer 2.	$C_{25}H_{25}IN_2$	1,1'-Diethyl-4,4'-carbocyanine iodide 393.
$C_{24}H_{16}Cl_2IrN_4^+$	Dichlorobis(1,10-phenanthroline)-iridium(III) ion 375.	$C_{25}H_{25}IN_2S_2$	3,3'-Diethyl-2,2'-thiatricbocyanine iodide 406.
$C_{24}H_{16}N_2$	9,9'-Bicarbazole 166.	$C_{25}H_{30}ClN_3$	Crystal Violet 307.
$C_{24}H_{16}N_2O_2$	2,2'-(1,4-Phenylene)bis[5-phenyloxazole] 885.	$C_{26}H_{16}$	Bifluorenylidene 167.
$C_{24}H_{18}$	<i>p</i> -Quaterphenyl 960.		Dibenzo[<i>g,p</i>]chrysene 347.
$C_{24}H_{18}Ge$	1,3,5-Triphenylbenzene 1094.		Dibenzo[<i>c,g</i>]triphenylene 352.
$C_{24}H_{18}N_2$	9,9-Diphenyl-9 <i>H</i> -9-germafluorene 540.		Hexacene 606.
$C_{24}H_{18}N_2$	5,10-Dihydro-5,10-diphenylphenazine 419.		Hexahelicene 608.
$C_{24}H_{18}Si$	9,9-Diphenyl-9 <i>H</i> -9-silafluorene 549.	$C_{26}H_{18}$	1-(2-Anthryl)-2-(2-naphthyl)ethylene 88.
$C_{24}H_{20}$	<i>anti</i> -[2.2](1,4)-Naphthalenophane 777.	$C_{26}H_{18}O$	9,10-Diphenylanthracene 529.
$C_{24}H_{20}N_2$	<i>syn</i> -[2.2](1,4)-Naphthalenophane 778.	$(C_{26}H_{24}O_4)_{32}$	Spiro[9,10-dihydro-9-oxoanthracene-10,2'-5',6'-benzindan] 999.
	[1,1':4',1'':4'',1''':4''']-Quaterphenyl]-4,4'''-diamine 959.		Poly[oxy[2-(1-pyrenylmethyl)-1,3-propanediyl]oxy(1,4-dioxo-1,6-hexanediyl)] 915.
$C_{24}H_{20}O$	<i>cis</i> -2,3-Dimethyl-2,3-di-(2-naphthyl)-oxirane 480.	$C_{26}H_{26}N_2O_2S$	2,5-Bis(5- <i>tert</i> -butyl-2-benzoxazolyl)-thiophene 189.
	<i>trans</i> -2,3-Dimethyl-2,3-di-(2-naphthyl)-oxirane 481.	$C_{26}H_{28}O_2S_2$	5,5'-Di- <i>tert</i> -amylthioindigo 337.
$C_{24}H_{20}O_2$	(<i>R</i>)-4,5,6,7-Tetrahydro-dinaphtho[2,1- <i>b</i> :1',2'- <i>d</i>][1,6]dioxecin 1029.	$C_{26}H_{32}N_2$	5,5'-Dineopentylthioindigo 514.
$(C_{24}H_{20}O_4)_{13}$	Poly[oxy[2-(1-pyrenylmethyl)-1,3-propanediyl]oxy(1,4-dioxo-1,4-butanediyl)] 913.	$C_{27}H_{19}NO$	<i>N,N,N',N'</i> -Tetraethyl-[1,1':4',1''-terphenyl]-4,4''-diamine 1028.
$C_{24}H_{21}NO_4$	9-[3-(4-Methoxycarbonylbenzoxy)propyl]-carbazole 666.	$C_{27}H_{27}N_2I$	2,5-Di(4-biphenyl)oxazole 357.
$C_{24}H_{22}$	1,4-Di-1-naphthylbutane 505.		1,1'-Diethyl-2,2'-dicarbocyanine iodide 395.
$C_{24}H_{22}N_2O_2$	<i>N</i> -[(Dibenzylamino)methyl]phthalimide 356.	$C_{27}H_{33}N_9O_5P_2$	Flavine mononucleotide 581.
$C_{24}H_{22}O_4$	2-(1-Pyrenylmethyl)-1,3-propandiol diacetate 949.	$C_{27}H_{34}N_9O_5P_2^+$	Flavine mononucleotide, conjugate monoacid 582.
$C_{24}H_{23}ClN_2O_3$	<i>N,N'</i> -Diethylrhodamine 400.	$C_{27}H_{39}IN_4S_2$	6,6'-Bis(dimethylamino)-3,3'-diethyl-2,2'-thiacarbocyanine iodide 191.
$C_{24}H_{24}N_2O_5$	1-(β-Methacryloxyethyl)-3,3-dimethyl-6'-nitro-spiro(indoline-2,2'-[2 <i>H</i>]benzopyran) 659.	$C_{28}H_{14}$	Benzo[<i>a</i>]coronene 118.
$C_{24}H_{24}O_2S_2$	5,5'-Dibutylthioindigo 364.	$C_{28}H_{16}N_2$	Tetrabenzophenazine 1019.
$C_{24}H_{28}ClN_3$	5,5'-Di- <i>tert</i> -butylthioindigo 365.	$C_{28}H_{18}$	9,9'-Bianthryl 165.
$C_{24}H_{32}O$	Methyl Violet 689.	$C_{28}H_{20}$	Anthracene photodimer 71.
	(<i>E,E,E,E,E</i>)-7,11-Dimethyl-7-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2,4,6,8,10,12-tridecahexaenal 501.	$C_{28}H_{20}O_2$	10 <i>H</i> ,10' <i>H</i> -10,10'-Dihydroxybianthrylidene 436.
$C_{24}H_{37}N$	<i>N</i> -11- <i>cis</i> -Retinylidene- <i>n</i> -butylamine 976.		Tetraphenyl- <i>p</i> -dioxin 1054.
	<i>N</i> -13- <i>cis</i> -Retinylidene- <i>n</i> -butylamine 977.	$C_{28}H_{22}$	3,3,4,5-Tetraphenyl-2(3 <i>H</i>)-furanone 1055.
	<i>N</i> -9- <i>cis</i> -Retinylidene- <i>n</i> -butylamine 978.	$C_{28}H_{22}N_2$	1,1,4,4-Tetraphenyl-1,3-butadiene 1052.
	<i>N</i> -all- <i>trans</i> -Retinylidene- <i>n</i> -butylamine 979.		<i>trans</i> -1,2-Di(<i>N</i> -carbazolyl)cyclobutane 366.
$C_{24}H_{38}N^+$	<i>N</i> -11- <i>cis</i> -Retinylidene- <i>n</i> -butylamine, conjugate acid 980.	$C_{28}H_{24}Cl_2IrN_4^+$	Dichlorobis(4,7-dimethyl-1,10-phenanthroline)iridium(III) ion 373.
	<i>N</i> -all- <i>trans</i> -Retinylidene- <i>n</i> -butylamine, conjugate acid 981.		Dichlorobis(5,6-dimethyl-1,10-phenanthroline)iridium(III) ion 374.
$C_{25}H_{16}O$	9-Naphthoanthracene 790.	$C_{28}H_{24}N_2$	Pentacyclo[18.2.2.2 ^{9,12} .0 ^{4,15} .0 ^{6,17}]-hexacos-4,6(17),9,11,15,20,22,23,25-nonane-10,25-dicarbonitrile 846.
$C_{25}H_{17}NO$	2-Biphenylphenylbenzoxazole 177.	$C_{28}H_{24}O_4$	<i>t</i> -3,4-Di-(1-naphthyl)- <i>r</i> -1,4-2-dimethoxycarbonylcyclobutane 506.
$C_{25}H_{18}O$	4,4'-Diphenylbenzophenone 530.	$C_{28}H_{28}N_4Zn$	Zinc(II) tetraethylporphyrin 1134.
$C_{25}H_{20}$	4,4'-Methylenebis(1,1'-biphenyl) 723.	$C_{28}H_{30}N_4$	Tetraethylporphine 1026.
		$C_{28}H_{30}O_4S_2$	6,6'-Dihexyloxythioindigo 407.

- $C_{28}H_{31}ClN_2O_3$ Rhodamine B **983**.
Rhodamine 6G **984**.
 $C_{28}H_{32}N_2O_3^{2+}$ Rhodamine 6G cation, conjugate monoacid **985**.
 $C_{28}H_{44}O$ Ergosterol **564**.
 $C_{29}H_{16}$ Pyranthrene **937**.
 $C_{29}H_{20}O$ 3-Benzyl-3-phenylphenanthro-[9,10-*b*]furan-2(3*H*)-one **162**.
 $C_{29}H_{22}O_2$ 3-Benzyl-3,4,5-triphenyl-2(3*H*)-furanone **163**.
10,11-Dihydro-5,7:14,16-dietheno-8*H*,13*H*-diindeno[2,1-*h*:1',2'*i*]-[1,4]dioxacyclotridecin **414**.
 $C_{29}H_{26}N_2$ *meso* (DL)-2,4-Di(*N*-carbazolyl)pentane **367**.
rac (DD,LL)-2,4-Di(*N*-carbazolyl)pentane **368**.
 $C_{29}H_{34}FN_2^+$ 1-Ethyl-2-[5-(1-ethyl-1,3-dihydro-3,3-dimethyl-2*H*-indol-2-ylidene)-3-fluoro-1,3-pentadienyl]-3,3-dimethyl-3*H*-indolium **572**.
 $C_{29}H_{35}N_2^+$ 1-Ethyl-2-[5-(1-ethyl-1,3-dihydro-3,3-dimethyl-2*H*-indol-2-ylidene)-1,3-pentadienyl]-3,3-dimethyl-3*H*-indolium **573**.
 $C_{30}H_{18}$ 9,10-Bis(phenylethynyl)anthracene **211**.
 $C_{30}H_{18}F_9LaO_6$ Tris(4,4,4-trifluoro-1-phenyl-1,3-butanedionato)lanthanum(III) **1107**.
 $C_{30}H_{18}F_9LuO_6$ Tris(4,4,4-trifluoro-1-phenyl-1,3-butanedionato)lutetium(III) **1108**.
 $C_{30}H_{24}IrN_6^{3+}$ Tris(2,2'-bipyridine)iridium(III) ion **1103**.
 $C_{30}H_{24}N_6Os^{2+}$ Tris(2,2'-bipyridine)osmium(II) ion **1104**.
 $C_{30}H_{24}N_6Ru^{2+}$ Tris(2,2'-bipyridine)ruthenium(II) ion **1105**.
 $C_{30}H_{40}O$ β -*apo*-8'-Carotenal **252**.
 $C_{30}H_{44}$ (*all-E*)-3,8-Dimethyl-1,10-bis(2,6,6-trimethyl-1-cyclohexen-1-yl)-1,3,5,7,9-decapentaene **477**.
 $C_{31}H_{24}$ 1,3-Di(9-phenanthryl)propane **526**.
 $C_{31}H_{34}N_4O_4$ Bonellin **213**.
 $C_{31}H_{42}ClN_3$ Ethyl Violet **568**.
 $C_{31}H_{42}O_2$ β -*apo*-8'-Carotenoic acid, methyl ester **256**.
 $C_{32}H_{16}AlClN_8$ Chloroaluminum phthalocyanine **261**.
 $C_{32}H_{16}ClInN_8$ Chloroindium(III) phthalocyanine **276**.
 $C_{32}H_{16}CuN_8$ Copper(II) phthalocyanine **297**.
 $C_{32}H_{16}MgN_8$ Magnesium(II) phthalocyanine **653**.
 $C_{32}H_{16}N_8Zn$ Zinc(II) phthalocyanine **1130**.
 $C_{32}H_{18}N_8$ Phthalocyanine **907**.
 $C_{32}H_{22}$ Anthracene-tetracene photodimer **72**.
 $C_{32}H_{24}$ [2.2](1,4)(9,10)-Anthracenophane **73**.
anti-[2.2](1,4)-Anthracenophane **74**.
syn-[2.2](1,4)-Anthracenophane **75**.
 $C_{32}H_{24}O$ 4-(4-*p*-Terphenylmethyl)benzophenone **1016**.
 $C_{32}H_{24}O_2$ (*E*)-10-(1,3-Dimethyl-10-oxo-9-anthracenylidene)-2,4-dimethyl-9-anthracenone **1046**.
 $C_{32}H_{29}NO_4S_3$ (*Z*)-*N,N*-Dibutyl-2,3-dihydro-3-oxo-2-(3-oxonaphtho[1,2-*b*]-thien-2(3*H*)-ylidene)naphtho[1,8-*bc*]-thiopyran-6-sulfonamide **362**.
(*E*)-*N,N*-Dibutyl-2,3-dihydro-3-oxo-2-(3-oxonaphtho[1,2-*b*]-thien-2(3*H*)-ylidene)naphtho[1,8-*bc*]-thiopyran-6-sulfonamide **363**.
 $C_{32}H_{32}N_2O_4$ 2,2'-(1,4-Phenylene)bis[5-(4-butoxyphenyl)oxazole] **884**.
 $C_{32}H_{34}N_4O_4$ Deuteroporphyrin, dimethyl ester **333**.
 $C_{32}H_{36}N_2$ *N,N,N',N'*-Tetraethyl-[1,1':4',1'':4'',1''':4''']-quaterphenyl]-4,4'''-diamine **1027**.
 $C_{32}H_{36}N_4Pd$ Palladium(II) etioporphyrin I **838**.
 $C_{32}H_{36}N_4Zn$ Zinc(II) etioporphyrin I **1126**.
 $C_{32}H_{40}N_4$ Etiochlorin II **579**.
 $C_{33}H_{20}BrN_8ORh$ Bromo(methanol)(phthalocyaninato)-rhodium(III) **222**.
 $C_{33}H_{20}ClN_8ORh$ Chloro(methanol)(phthalocyaninato)-rhodium(III) **277**.
 $C_{33}H_{20}IN_8ORh$ Iodo(methanol)(phthalocyaninato)-rhodium(III) **635**.
 $C_{33}H_{30}O_4$ 10,11,13,14,16,17-Hexahydro-5,7:20,22-dietheno-8*H*,19*H*-diindeno[2,1-*n*:1',2'-*o*]-[1,4,7,10]tetraoxacyclononadecin **609**.
 $C_{33}H_{34}N_4O_6$ Biliverdin **169**.
 $C_{33}H_{36}N_4O_6$ Bilirubin **168**.
 $C_{34}H_{24}O_3$ 1,4-Diphenyl-2,3-dibenzoyl-1,4-epoxy-1,4-dihydronaphthalene **538**.
 $C_{34}H_{32}N_4O_4Zn$ Zinc(II) protoporphyrin **1132**.
 $C_{34}H_{34}N_4O_4$ Protoporphyrin IX **930**.
 $C_{34}H_{36}MgN_4O_4$ Magnesium(II) mesoporphyrin **651**.
 $C_{34}H_{36}N_4O_3$ Methyl pyrophosphoribide *a* **747**.
 $C_{34}H_{38}N_4O_4$ Mesoporphyrin **657**.
 $C_{34}H_{38}N_4O_6$ Hematoporphyrin IX **604**.
 $C_{35}H_{38}N_4O_6$ Biliverdin, dimethyl ester **170**.
 $C_{35}H_{50}$ (*all-E*)-3,7,12-Trimethyl-1,14-bis(2,6,6-trimethyl-1-cyclohexen-1-yl)-1,3,5,7,9,11,13-tetradecaheptacene **1086**.
 $C_{36}H_{20}CdN_4$ Cadmium(II) tetrabenzoporphyrin **240**.
 $C_{36}H_{20}N_4Pd$ Palladium(II) tetrabenzoporphyrin **839**.
 $C_{36}H_{20}N_4Zn$ Zinc(II) tetrabenzoporphyrin **1133**.
 $C_{36}H_{22}N_4$ Tetrabenzoporphine **1020**.
 $C_{36}H_{24}$ [2.2](2,7)-Pyrenophane **948**.
 $C_{36}H_{24}N_6Rh^{3+}$ Tris(1,10-phenanthroline)rhodium(III) ion **1106**.
 $C_{36}H_{26}O$ *rac*-Bis[1-(1-pyrenyl)ethyl]ether **212**.
 $C_{36}H_{28}N_8O_2RuS_2$ Bis(dimethylsulfoxide)phthalocyanatoruthenium(II) **205**.
 $C_{36}H_{36}MgN_4O_5$ Methylchlorophyllide *a* **712**.
 $C_{36}H_{36}N_4O_6$ Photoporphyrin isomer "A", dimethyl ester **903**.
Photoporphyrin isomer "B", dimethyl ester **904**.
 $C_{36}H_{38}N_4O_4$ Protoporphyrin IX, dimethyl ester **931**.
 $C_{36}H_{38}N_4O_8$ Coproporphyrin III **298**.

$C_{36}H_{39}N_4O_4^+$	Protoporphyrin IX, dimethyl ester, conjugate monoacid 933 .	$C_{42}H_{28}$	Rubrene 989 .
$C_{36}H_{40}N_4O_4^{2+}$	Protoporphyrin IX, dimethyl ester, conjugate diacid 932 .	$C_{42}H_{30}$	5''-[1,1'-Biphenyl]-4-yl-1,1':4',1'':3'',1''':4''',1''''-quinquephenyl 183 .
$C_{36}H_{42}N_4O_4$	Mesoporphyrin, dimethyl ester 658 .	$C_{42}H_{60}O_2$	<i>all-trans</i> -Spirilloxanthin 998 .
$C_{36}H_{44}Cl_2N_4Sn$	Octaethylporphinatotin(IV) dichloride 828 .	$C_{44}H_{24}N_4O_{12}PdS_4^{4-}$	Tetrakis(<i>p</i> -sulfonatophenyl)-porphinatopalladate(II) ion 842 .
$C_{36}H_{44}MgN_4$	Magnesium(II) octaethylporphyrin 652 .	$C_{44}H_{24}N_4O_{12}RhS_4^{3-}$	Tetrakis(<i>p</i> -sulfonatophenyl)-porphinatorhodate(III) ion 1042 .
$C_{36}H_{44}N_4Pd$	Octaethylporphinatopalladium(II) 827 .	$C_{44}H_{24}N_4O_{12}S_4Zn^{4-}$	Tetrakis(<i>p</i> -sulfonatophenyl)-porphinatozincate(II) ion 1140 .
$C_{37}H_{26}$	2-([1,1':4',1'':Terphenyl]-4-ylmethyl)-triphenylene 1017 .	$C_{44}H_{26}N_4O_{12}S_4^{4-}$	Tetrakis(<i>p</i> -sulfonatophenyl)porphine 1043 .
$C_{38}H_{30}N_{10}O_2Ru$	Bis(dimethylformamide)-phthalocyaninatoruthenium(II) 204 .	$C_{44}H_{28}CdN_4$	Tetraphenylporphinatocadmium(II) 241 .
$C_{39}H_{40}$	9,10,11,12,13,14-Hexahydro-5,7:15,17-ietheeno-2,20-heptano- <i>H</i> -cyclotetradeca[1,2- <i>a</i> :1,14- <i>a'</i>]-diindene 610 .	$C_{44}H_{28}ClN_4Rh$	Tetraphenylporphinatorrhodium(III) chloride 1056 .
$C_{39}H_{58}O_4$	Ubiquinone 30 1114 .	$C_{44}H_{28}HgN_4$	Tetraphenylporphinatomercury(II) 656 .
$C_{40}H_{26}$	1,3,6,8-Tetraphenylpyrene 1059 .	$C_{44}H_{28}MgN_4$	Tetraphenylporphinato-magnesium(II) 654 .
$C_{40}H_{26}N_8$	Tetrakis(2-pyridyl)porphine 1036 .	$C_{44}H_{28}N_4^{2-}$	Tetraphenylporphyrin dianion 1058 .
	Tetrakis(3-pyridyl)porphine 1037 .	$C_{44}H_{28}N_4OTi$	Oxotitanium(IV) tetraphenylporphyrin 837 .
	Tetrakis(4-pyridyl)porphine 1038 .	$C_{44}H_{28}N_4Pd$	Tetraphenylporphinatopalladium(II) 843 .
$C_{40}H_{28}N_8^{2+}$	Tetrakis(2-pyridyl)porphine, conjugate diacid 1039 .	$C_{44}H_{28}N_4Zn$	Tetraphenylporphinatozinc(II) 1143 .
	Tetrakis(3-pyridyl)porphine, conjugate diacid 1040 .	$C_{44}H_{30}N_4Zn$	<i>meso</i> -Tetraphenylporphine 1057 .
	Tetrakis(4-pyridyl)porphine, conjugate diacid 1041 .	$C_{44}H_{32}N_4$	Zinc(II) tetraphenylchlorin 1142 .
$C_{40}H_{46}N_4O_8$	Coproporphyrin III, tetramethyl ester 299 .	$C_{44}H_{36}N_8Pd^{4+}$	Tetraphenylchlorin 1053 .
	Coproporphyrin I, tetramethyl ester 300 .	$C_{44}H_{36}N_8Zn^{4+}$	Palladium(II) tetrakis-(4- <i>N</i> -methylpyridyl)porphyrin 841 .
$C_{40}H_{48}O_4$	Astacin 91 .		Tetrakis(2- <i>N</i> -methylpyridyl)-porphinatozinc(II) ion 1137 .
$C_{40}H_{52}O$	<i>all-trans</i> -3',4'-Didehydro- β,ψ -16'-carotenal 387 .	$C_{44}H_{38}N_8^{4+}$	Tetrakis(3- <i>N</i> -methylpyridyl)-porphinatozinc(II) ion 1138 .
$C_{40}H_{52}O_2$	Canthaxanthin 244 .		Tetrakis(4- <i>N</i> -methylpyridyl)-porphinatozinc(II) ion 1139 .
$C_{40}H_{52}O_4$	Astaxanthin 92 .		Tetrakis(2- <i>N</i> -methylpyridyl)-porphine 1033 .
$C_{40}H_{54}O$	Echinenone 555 .	$C_{48}H_{24}N_4O_8Pd^{4-}$	Tetrakis(3- <i>N</i> -methylpyridyl)-porphine 1034 .
$C_{40}H_{56}$	15,15'- <i>cis</i> - β -Carotene 253 .	$C_{48}H_{24}N_4O_8Zn$	Tetrakis(4- <i>N</i> -methylpyridyl)-porphine 1035 .
	β -Carotene 254 .		Palladium(II) tetrakis-(carboxyphenyl)porphyrin 840 .
$C_{40}H_{56}O_2$	<i>all-trans</i> -Lycopene 649 .	$C_{48}H_{28}O_2$	Tetrakis(carboxyphenyl)porphinatozincate(II) ion 1135 .
	Dihydroxylycopene 438 .		1,4-Dihydro-5,6,11,12-tetraphenyl-1,4-epidioxytetracene 433 .
	<i>all-trans</i> -Lutein 648 .		5,12-Dihydro-5,6,11,12-tetraphenyl-5,12-epidioxytetracene 434 .
	Zeaxanthin 1123 .		5-(1,4-Epidioxyphenyl)-6,11,12-triphenyltetracene 563 .
$C_{40}H_{58}$	<i>all-trans</i> -Neurosporene 808 .	$C_{48}H_{54}N_4O_{16}$	Uroporphyrin I, octamethyl ester 1118 .
$C_{40}H_{60}$	ζ -Carotene 255 .	$C_{50}H_{68}$	Decapreno- β -carotene 331 .
$C_{40}H_{64}$	7,7'-Dihydro- β -carotene 412 .	$C_{52}H_{40}N_4O_{12}S_4Zn^{4-}$	Tetrakis(2,6-dimethyl-4-sulfonatophenyl)porphinatozincate(II) ion 1136 .
	15- <i>cis</i> -Phytoene 908 .		Tetrakis(2,6-dimethyl-4-sulfonatophenyl)porphine 1032 .
	<i>all-trans</i> -Phytoene 909 .		
$C_{41}H_{44}$	8,9,10,11,12,13,14,15-Octahydro-5,7:16,18-dietheno-2,21-octanocyclopentadeca-[1,2- <i>a</i> :1,15- <i>a'</i>]diindene 829 .		
	<i>all-trans</i> -Spheroidenone 997 .		
$C_{41}H_{58}O_2$	<i>all-trans</i> -Spheroidene 996 .		
$C_{41}H_{60}O$	Phthalocyaninobis(pyridine)-ruthenium(II) 906 .		
$C_{42}H_{26}N_{10}Ru$			

- $C_{52}H_{54}N_4O_{12}$ Malachite Green 655.
 $C_{55}H_{70}MgN_4O_6$ Chlorophyll *b* 285.
 $C_{55}H_{72}MgN_4O_5$ Chlorophyll *a* 284.
 $C_{55}H_{72}N_4O_5Zn$ Zinc(II) chlorophyll *a* 1124.
 $C_{55}H_{72}N_4O_6$ Pheophytin *b* 902.
 $C_{55}H_{74}N_4O_5$ Pheophytin *a* 901.
 $C_{56}H_{60}N_8Zn^{4+}$ Tetrakis-4-(*N,N,N*-trimethylammonio)-phenylporphine zinc(II) ion 1141.
 $C_{56}H_{62}N_8^{4+}$ Tetrakis(trimethylaminophenyl)porphine 1044.
 $C_{60}H_{80}$ (*all-E*)-3,7,11,15,20,24,28,32-Octamethyl-1,34-bis(2,6,6-trimethyl-1-cyclohexen-1-yl)-1,3,5,7,9,11,13,15,17,19,21,23,25,27,29,31,33-tritriacontaheptadecaene 830.
 $C_{71}H_{80}N_4O_3$ 4-(β -*apo*-7'-Carotenyl)benzyl pyro-pheophorbide 257.
 $C_{82}H_{82}N_4O_3$ 4-[Tri(4-methylphenyl)porphynyl]-3-phenoxypropyl β -*apo*-6'-carotenate 1090.

10.3. Author Index

- Abrahamson, E.W. 285, 866, 972
 Abuin, E.B. 1121
 Achiba, Y. 128, 735, 771, 953
 Acuna, A.U. 123, 703
 Adachi, T. 620
 Adamczyk, A. 234
 Adams, R.G. 972
 Aikawa, M. 133
 Aikens, D.A. 809, 990
 Al'-Ainen, S.A. 141, 1101
 Al'perovich, M.A. 435
 Al-Chalabi, A.O. 144
 Al-Saigh, H.Y. 989
 Albrecht, A.C. 61, 699
 Alchalel, A. 966, 974, 976, 980
 Alfimov, M.V. 8, 43, 64, 173, 362, 768, 1062, 1100
 Alkaitis, S.A. 870, 1045
 Allen, J.C. 684
 Almgren, M. 254
 Alpert, B. 930
 Alvarez, V.L. 905
 Amand, B. 279
 Amano, M. 154
 Amirzadeh, G. 290, 332
 Amouyal, E. 143, 446, 552
 Amrein, W. 153
 Anderson, R.W., Jr. 963
 Andreev, O.M. 173
 Andrews, L.J. 586, 720
 Andrievskii, A.M. 350
 Angliker, H. 330, 606
 Anitoff, O. 1138
 Arai, S. 7, 173, 822
 Arai, T. 427
 Arakawa, M. 1083
 Arce, R. 936
 Arimitsu, S. 470
 Arvis, M. 392, 394
 Asano, M. 284
 Asher, M. 463
 Astier, R. 11, 64, 109, 173, 733
 Aubailly, M. 57
 Averbek, D. 246, 598
 Azerad, R. 966
 Baba, H. 439, 672
 Babichev, F.S. 435
 Bagdasar'yan, Kh.S. 1, 23, 133, 166, 570, 719, 1095
 Baiardo, J. 854
 Bailey, D.N. 867
 Balzani, V. 1105
 Bandyopadhyay, B.N. 861
 Bar-Eli, K.H. 719
 Baral-Tosh, S. 133
 Barigelletti, F. 746
 Barthels, M.R. 308
 Barwise, A.J.G. 173
 Batekha, I.G. 43, 125
 Bates, A.L. 91
 Bauer, H. 137
 Baugher, J.F. 284, 505
 Baumgartner, C.E. 990
 Daxendale, J.II. 64
 Bayrakceken, F. 348, 433
 Bazin, M. 57, 631, 1109
 Beaumont, P.C. 684
 Bebelaar, D. 64
 Beck, G. 870
 Becker, R.S. 250, 251, 502, 748, 1070
 Bedwell, E. 729
 Bell, I.P. 939
 Bell, J.A. 133
 Bender, C.O. 153
 Bennett, R.G. 369
 Bensasson, R. 6, 51, 133, 143, 246, 251, 253, 255, 257, 279, 446, 502, 541, 548, 552, 598, 640, 751, 919, 935, 966, 972, 1067, 1070, 1072, 1090, 1105, 1114, 1140
 Bent, D.V. 63, 461, 631, 876, 938, 1075
 Bentley, P. 38
 Bercovici, T. 1046
 Berens, K. 58
 Berg, R.A. 392
 Bergwerf, Ph. 233, 525
 Bir, E.Sh. 423
 Bircher, P. 235
 Bisagni, E. 246
 Bianchi, J.-P. 133
 Blume, H. 535, 672
 Bocherkov, A.I. 125
 Bokobza, A. 64
 Boldridge, D.W. 764
 Bolot'ko, L.M. 50, 178
 Bolotnikova, T.N. 64
 Bolton, J.R. 743
 Bonamy, A. 958
 Bonneau, R. 4, 96, 180, 318, 578, 768, 1063

- Bonnet, R. 1035
Bonnett, R. 299
Borisevich, N.A. 50
Borisevich, Yu.E. 192
Borovkova, V.A. 23, 166, 570
Bortolus, P. 746
Bowers, P.G. 64, 284
Braslavsky, S.E. 153, 213
Braterman, P.S. 1105
Braun, A.M. 133, 479, 907, 1143
Brede, O. 133, 861
Breheret, E. 8, 245, 691
Bridge, N.K. 552
Briegleb, G. 768
Brinen, J.S. 65, 106, 173, 295, 769, 1016
Brown, R.G. 190, 743
Bruehlmann, U. 416
Brunschwig, B.S. 652
Buben, N.Ya. 8
Buettner, A.V. 64, 133, 391
Bulska, H. 184
Buono-Core, G.E. 82
Burger, U. 153
Burland, D.M. 245
Burnett, M.N. 851
Busse, R. 526
Byers, G.W. 939
Cabaret, D. 414
Cadogan, K.D. 61
Caldwell, R.A. 258
Camilleri, P. 213
Canbaeck, G. 152
Cannistraro, S. 254
Capellos, C. 53, 460, 517, 518, 733, 819, 820
Carapellucci, P.A. 586, 866
Carioli, A. 1106
Carlini, C. 746
Carroll, F.A. 1126
Cauzzo, G. 604
Ceballos, A. 123, 703
Cessna, A.J. 7
Chachaty, C. 1114
Chahidi, C. 57
Chakachery, E. 353
Chalvet, O. 246
Chantrell, S.J. 931
Chaquin, P. 238
Charalambides, A.A. 299
Charlier, M. 133
Chattopadhyay, S.K. 133, 385, 529, 535, 662, 945
Chauhan, S.M.S. 353
Chedekel, M.R. 47
Chene, G. 284
Chessin, M. 254
Chevalier, S. 654
Chew, V.S.F. 743
Chibisov, A.K. 97, 191, 397, 400, 722, 869, 984, 986, 991
Chihara, K. 748
Chou, M. 1104
Chow, Y.L. 82
Chrysochoos, J. 556
Claesson, S. 285
Cogdell, R.J. 808
Cohen, S.G. 249
Collart, P. 212
Colson, E.C. 1059
Compton, R.H. 64
Connolly, J.S. 100
Cook, M.G. 768
Cooke, J. 64
Cooper, M. 939
Cooper, M.B. 966
Cornelisse, J. 515
Corval, A. 147
Costa, S.M. de B. 159
Cowley, D.J. 52, 269
Coyle, J.D. 356, 457
Craig, B.B. 1023
Craig, D.P. 64, 279, 768
Craw, M. 246, 305
Creutz, C. 1104
Cundall, R.B. 164
Currie, W.J.C. 32
Dabral, V. 353, 538
Dainton, F.S. 64, 133, 1005, 1006
Dallinger, R.F. 92
Damschen, D.E. 963
Danilova, V.I. 178
Danziger, R.M. 719
Darmanyan, A.P. 64, 358, 531, 572, 989
Das, P.K. 88, 133, 162, 200, 209, 250, 251, 385, 480, 529, 535, 538, 662, 694, 748, 945, 971, 1045, 1105
Davidson, R.S. 957
Davies, A.K. 292
Davis, G.A. 586, 866
Davis, H.F. 385, 694
Davis, S.G. 966
Davydov, S.N. 402
Dawe, E.A. 253, 966
Dawson, W.R. 65, 106, 118, 130, 940, 972
Dedinas, J. 133
de Groot, R.L. 80
de Gunst, G.P. 515
Dekker, R.H. 31
Delestinne, A. 308
Delouis, J.F. 556
Dempster, D.N. 49, 393, 399, 984
Demuth, M. 153
Deroulede, A. 586
De Sa E Melo, T.M. 51
De Schryver, F.C. 212, 366
Dilung, I.I. 132
Dixit, S.N. 1125
Dixon, R.S. 369
Doan, T.L. 21
Dodiuk, H. 1047
Dung, D. 923
Dunne, A. 982

- Dunston, I.M. 7
Dupuy, F. 1095
Eaker, D.W. 802
Ebbesen, T.W. 265, 1140
Efros, L.S. 670
Ege, S.N. 385
Eichler, J. 10, 625
Eisenthal, K.B. 773
El'tsov, A.V. 125
El-Sayed, M.A. 295, 608
Encinas, M.V. 112, 133, 232
Euing, W. 769
Evans, G.B. 164
Evans, R.F. 1109
Faidysh, A.N. 768
Fang, T.-S. 211
Fanghaenel, E. 995
Farmilo, A. 254
Farquharson, S. 92
Faulkner, L.R. 989
Faure, J. 96, 177, 332, 482, 556, 768
Favre, A. 1067
Feldberg, S.W. 652
Fendler, J.H. 714
Ferraudi, G. 204, 222, 906
Ferreira, M.I.C. 1063
Ferris, J.P. 727
Fielden, E.M. 64
Fischer, E. 352, 436, 1046
Fischer, G. 279, 768
Fischer, H. 1063
Fisher, G.J. 737
Fisher, M.M. 972
Flamigni, L. 746
Fleming, G.R. 62
Foerster, E.W. 410, 718
Fojtik, A. 861
Folcher, G. 1138
Foote, C.S. 308
Formosinho, S.J. 23
Fornier de Violet, Ph. 180, 957
Fouassier, J.-P. 177, 332, 482, 958
Fox, M.A. 306
Frank, A.J. 744
Fratev, F. 114
Freilich, S.C. 133
Fujara, F. 24
Fujimori, E. 284
Fujiwara, Y. 616
Fukushima, M. 767
Fuller, J. 173
Fushimi, K. 29
Gaevskii, A.S. 768
Gallivan, J.B. 295
Garcia Dominguez, J.A. 123
Garner, A. 631, 1121
Gautron, R. 147
Geacintov, N.E. 135, 1031
Gegiou, D. 871
Geissler, U. 768
Gennari, G. 604
George, M.V. 153, 162, 353, 538, 1054
Geraghty, N. 414
Gerhardt, G.E. 1016
Gerko, V.I. 43, 64, 173
Ghetti, F. 1140
Ghiggino, K.P. 1109, 1110
Ghiron, C.A. 729, 1109
Giraud, M. 643
Gisin, M. 774, 851
Givens, R.S. 892
Glagolev, V.L. 8, 173
Godfrey, T.S. 41
Goerner, H. 54, 95, 107, 218, 229, 311, 334, 420, 46
535, 678, 713, 802
Goldberg, M. 715
Goldschmidt, Ch.R. 64, 787
Goodwin, D. 957
Gorman, A.A. 95, 173, 317, 806
Gorman, D.S. 100
Goujon, P. 394
Gould, I.R. 133, 317, 892, 1045
Gouterman, M. 827
Goyal, G.C. 684
Grabowska, A. 963
Grabowski, Z.R. 463
Graetzel, C.K. 1140
Graetzel, M. 224, 284, 479, 552, 744, 870, 922, 1032,
1045, 1088, 1105, 1131, 1135, 1140, 1143
Graf, G.A. 133
Grajcar, L. 556
Gramain, J.-C. 133
Grattan, K.T.V. 64
Green, P.N. 958
Greene, B.I. 23
Grellmann, K.-H. 64, 71, 352, 388, 410, 415, 696, 718,
965
Gresser, J.D. 586, 866
Grieser, F. 216, 1043
Griffin, G.W. 385, 480, 1045
Griffiths, P.A. 164
Grodowski, M. 410, 966
Grodowski, M.S. 31
Gross, S. 939
Grossweiner, L.I. 562, 599, 684, 760
Gruen, H. 218
Gruzinskii, V.V. 178
Gschwind, R. 259
Gu, C. 308
Guenthard, Hs.H. 64, 1046
Guesten, H. 64, 420
Guglielmetti, R. 675
Guillory, W.A. 23
Gupta, A. 613
Gurinovich, G.P. 284
Gust, D. 257, 1090
Gusten, H. 529
Guyot, G. 179

- Gyoda, H. 259
Hadley, S.G. 64, 641, 768, 905
Haertel, H. 768
Hafner, M. 1063
Hama, Y. 837
Hamada, K. 946
Hamanoue, K. 13, 76, 148, 154, 263, 293, 706, 815
Hamblett, I. 95, 317, 806
Hammond, P.R. 357
Hamoudi, H.I. 32
Hanst, P.L. 133
Hara, K. 439, 672
Hardwick, R. 64
Harriman, A. 76, 241, 356, 861, 966, 1050, 1063, 1105
Harrison, R.J. 95
Haselbach, E. 259
Hashimoto, S. 1045
Hata, N. 247, 821
Hatano, Y. 570
Hatchard, C.G. 1064
Havinga, E. 515
Hayashi, H. 20, 133, 427, 577, 740, 1084
Hayashi, K. 171
Hayon, E. 5, 63, 631, 876, 938, 1075
Head, D.A. 768
Heath, G.A. 1105
Heelis, P.F. 32, 581
Heinrich, G. 420, 529, 535, 672
Heinzelmann, W. 106
Helene, A. 1109
Helene, C. 21, 133, 1109
Hellner, C. 845
Helmstreit, W. 133
Hemmerich, P. 715
Henrichs, P.M. 939
Henry, B.R. 245, 303
Hentzschel, P. 352, 388
Herbert, M.A. 831, 1115
Herbich, J. 963
Hercules, D.M. 867
Herkstroeter, W.G. 348, 966, 989
Hermann, H. 314
Herre, W. 768
Herrmann, H. 337
Herz, C.P. 10, 625
Hidaka, T. 13, 148, 706
Higuchi, M. 768
Hilinski, E.F. 83
Hilpern, J.W. 41
Hindman, J.C. 284
Hinzmann, S. 995
Hirase, S. 76
Hirata, Y. 23, 866
Hirayama, S. 13, 154, 815
Ho, S.K. 64, 173
Hochstrasser, R.M. 23, 64
Hodgkinson, K.A. 106, 939
Hodgson, W.G. 769
Hoffman, M.Z. 64, 720
Holmstroem, B. 285, 645
Holroyd, R.A. 64, 733
Holtén, D. 827
Holzer, G. 535
Honig, B. 980
Horrocks, A.R. 1, 878
Hoshi, T. 173, 437, 527
Hoshino, M. 71, 837
Hrdlovic, P. 179
Huber, J.R. 31, 416, 871, 1046
Hug, G.L. 251, 535, 748, 971
Huizer, A.H. 233
Hulme, B.E. 36
Hummel, A. 470
Hunt, J.W. 768, 1115
Hunt, R.V. 303
Hunter, T.F. 947
Huppert, D. 224, 1047
Hurley, J.K. 133
Ikawa, H. 1
Ikeda, N. 342
Imamura, M. 71, 173, 837, 939
Inbar, S. 249
Indelli, M.T. 1106
Infelta, P.P. 224, 1105
Inoue, A. 938
Inoue, H. 527
Inoue, T. 366
Irie, M. 171
Ishii, Y. 165
Ishikawa, M. 1091
Ishikawa, S. 73, 844
Itaya, A. 570
Ito, Y. 1084
Itoh, M. 15, 474, 552, 616, 619, 620, 1121
Itoh, N. 768
Ivanoff, N. 556
Ivanov, V.B. 67
Ivanov, V.L. 67, 141, 370, 673, 1101
Iwamura, H. 20
Iwanaga, C. 719
Iwaoka, T. 866
Iwata, S. 73, 551
Jackson, G. 23, 64
Jacques, P. 907
Jansen, K. 23
Japar, S.M. 866
Jardon, P. 147
Jensen, N.-H. 605, 972
Jimenez, L.A. 936
Johns, H.E. 831, 1072, 1073, 1115
Jones, R.A. 32
Jori, G. 298, 604
Journeaux, R. 284
Joussot-Dubien, J. 96, 768
Joy, A. 1090
Joy, A.M. 257
Justus, B.L. 764
Kaenzig, H. 484, 768

- Kagiya, V.T. 152
Kajiwara, Y. 76, 154, 263
Kalisky, Y. 284, 659
Kalyanasundaram, K. 840, 923, 1032, 1033, 1042, 1044, 1135
Kamachi, M. 133
Kamat, P.V. 306, 719, 1024, 1064
Kanamaru, N. 905, 1049
Kanety, H. 1047
Karstens, T. 769, 1062
Karyakin, A.V. 97, 397, 722, 869, 991
Kasama, K. 7, 23, 822
Kasha, M. 245
Kassem, A. 995
Kato, S. 23, 261, 276, 279, 562, 719, 1065, 1066, 1130
Katsuda, Y. 474
Katz, J.J. 284
Kawai, K. 259
Kawata, H. 964
Kayser, R.H. 719
Kazan, J. 1016
Keene, J.P. 64, 133, 164
Keller, R.A. 64, 768
Kellmann, A. 23, 25, 26, 675
Kellogg, R.E. 64
Kemp, D.R. 259, 552
Kemp, T.J. 64, 133, 236, 818, 826, 989
Ketsle, G.A. 984, 986, 988
Khelladi, F.Z. 768
Khosroffian, J.M. 135
Khudyakov, I.V. 441, 817
Kikuchi, K. 23, 29, 64, 80, 166, 245, 546, 570, 719, 787, 923, 964, 1063, 1091
Kikuchi, M. 719
Kikuta, Y. 133
Kilp, T. 682, 733
Kimoto, M. 148, 154
Kimura, E. 9
Kimura, K. 128, 470, 735, 771, 905, 953
Kingston, D.H. 457
Kira, A. 23, 171, 173, 939
Kirkor-Kaminska, E. 963
Kiryukhin, Yu.I. 1, 23, 133, 166, 719, 1095
Kissinger, P.T. 989
Kitamura, N. 366
Kiwi, J. 284, 1088
Klaening, U.K. 787
Klages, C.P. 849
Klasinc, L. 64
Klein, R. 631
Kleo, J. 254
Kliger, D.S. 699
Klinakova, A. 370
Knowles, A. 645
Kobashi, H. 1, 259
Kobayashi, S. 64
Kobayashi, T. 13, 83, 188, 528
Kobs, K. 849, 1062
Koizumi, M. 23, 492, 562, 719, 787, 866, 923
Kokrashvili, T.A. 346
Kokubun, H. 23, 29, 64, 80, 166, 245, 492, 546, 570, 719, 964, 1063, 1091
Kolobkov, V.P. 12
Kolosov, V.A. 166
Koltzenburg, G. 314
Komuro, M. 437
Kondo, R. 1
Koningstein, J.A. 284
Kopylova, T.N. 178
Korenstein, R. 436, 1046
Korobov, V.E. 397, 400
Koshihara, S. 528
Kosower, E.M. 1047
Kossanyi, J. 238
Kostin, A.K. 64
Kotlicka, J. 184
Koussini, R. 180
Kovalenko, N.P. 173
Kraljic, I. 64
Kramer, H.F.A. 25, 645, 715, 835, 1063
Krasnaya, Zh.A. 192
Krasnova, T.L. 402
Krebs, A. 330
Kreibich, U.T. 641
Krieg, M. 133
Krongauz, V. 1088
Kropp, J.L. 118, 130, 338, 940
Krowczynski, A. 463
Krueger, U. 565
Krysanov, S.A. 362, 1062
Kryukov, A.Yu. 618
Kuball, H.-G. 769
Kubo, Y. 577
Kuehnle, W. 526
Kuhlmann, R. 332, 912
Kumar, C.V. 162, 200, 353, 529, 535, 538, 662, 694, 945, 1054
Kuntz, R.R. 729, 1109
Kusabayashi, S. 570
Kuyumdzh, E.S. 8
Kuz'min, M.G. 67, 141, 381, 673, 1101
Kuz'min, V.A. 64, 97, 192, 245, 346, 435, 441, 531, 552, 572, 670, 989, 1031
Kuznetsov, V.A. 984
Kuznetsov, V.S. 125
Labhart, H. 106, 235
Lachish, U. 1105
Lacourbas, B. 245
Lafferty, J. 251, 541, 930
Laffitte, E. 26
Lahiri, S. 353
Lambert, C. 47, 305
Lamotte, M. 1095
Land, E.J. 6, 16, 23, 34, 36, 47, 51, 168, 169, 176, 244, 246, 251, 253, 254, 255, 299, 303, 446, 470, 541, 552, 598, 640, 684, 737, 751, 768, 808, 919, 931, 935, 966, 972, 973, 1009, 1035, 1090, 1114

- Lang, F. 53, 460
Langelaar, J. 118, 939
Langkilde, F.W. 605
Lapouyade, R. 4, 180, 320
Latowski, T. 410
Lavalette, D. 64, 118, 173, 176, 414, 669, 769
Lazare, S. 4, 320
Leclercq, J.M. 64, 173
le Cornelisse, J. 516
Ledger, M.B. 8, 64
Lee, C.W.B. 82
Lee, C.Y. 1058
Lee, E.D. 809
Lee, P.C.-C. 133
Lee, W.A. 1032
Lehman, W.R. 257
Lemaire, J. 179
Lemmer, D. 153
Le Roux, D. 1138
Leupin, W. 951
Levin, G. 301, 1021, 1058
Levin, P.P. 260, 346, 531, 552
Levkoev, I.I. 97, 435
Levshin, L.V. 984, 986, 988
Lewis, G.N. 528, 591
Lichtin, N.N. 719, 720, 1024, 1064, 1130
Liddell, P.A. 257
Lifanov, Yu.I. 397
Lim, B.T. 505
Lim, E.C. 30, 505, 938
Lin, J. 211
Lindqvist, L. 8, 23, 25, 64, 265, 285, 592, 593, 631, 675, 691, 845, 930
Linschitz, H. 78, 100, 133, 249, 284, 285, 548, 586, 718, 720
Lipkin, D. 528, 591
Lissi, E.A. 232
Liu, C.-T. 47
Liu, D.K.K. 989
Liu, R.S.H. 751, 966
Livingston, R. 64, 254, 284, 965
Lo, K.K.N. 751, 973
Lochet, R. 26, 768
Loeff, I. 78, 265
Lohray, B.B. 162
Lohse, C. 160
Long, D.A. 253
Lou, J.J. 338
Longnot, D.-I. 177, 332, 482, 958, 1010
Lower, S.K. 64
Lukac, I. 179
Lutz, H. 8, 265, 691
Luzhkov, V.B. 817
MacLachlan, A. 1006
Maciejewski, A. 164, 603
Maeda, Y. 464
Magde, D. 827
Magel, T.T. 591
Maki, A.H. 83
Makin, S.M. 64
Malkin, Ya.N. 245, 670
Manring, L.E. 308
Marchetti, A.P. 64
Mark, F. 420
Marks, G.T. 809
Marowsky, G. 885
Martens, F.M. 525
Martin, M. 25, 245
Martin, M.M. 342
Martins, L.J.A. 236, 818, 826
Martynov, I.Yu. 673
Martynova, V.P. 670
Maruyama, K. 577
Masuhara, H. 212, 366, 464, 526, 570, 846, 913, 946
Mataga, N. 64, 212, 342, 366, 464, 526, 570, 846, 913, 946
Mathis, P. 254, 284
Matsumoto, S. 383, 551, 719
Matsuura, T. 1084
Matthews, J.I. 213
Matuszewski, B. 892
Maudinas, B. 255
McAlpine, E. 34
McAuliffe, C.A. 931
McCallum, K.J. 23
McCarthy, R.L. 1006
McCartin, P.J. 369
McClure, D.S. 64, 133, 348
McKellar, J.F. 38, 581
McVie, J. 297, 680
Medinger, T. 529, 878
Mehnert, R. 133, 861
Melhuish, W.H. 64, 133, 279
Melo, E.C.C. 159
Memming, R. 565, 849, 965, 1062
Menzel, R. 848
Merkel, P.B. 989
Merlin, A. 332, 482
Merritt, C. 613
Metras, J.-C. 291
Meyer, Y.H. 11, 64, 109, 173, 733
Mialocq, J.-C. 392, 394, 1138
Miedlar, K. 1105
Miethke, E. 25
Mikawa, H. 259
Miller, J.R. 173
Mills, A. 76
Milton, S.V. 83
Mintas, M. 64
Misumi, S. 73, 846
Mitchell, A.D. 680
Mitchell, M.B. 23
Miyake, T. 76, 154, 293
Miyashita, Y. 23
Miyata, N. 15
Moan, J. 1111
Mochida, K. 740
Moiseeva, Z.Z. 423

- Molera, M.J. 123, 703
Monti, S. 675
Monzikoff, A. 57
Moore, A.L. 257, 1090
Moore, T.A. 257, 1090
Moorthy, P.N. 938
Morgante, C.G. 632
Morita, H. 383
Morita, M. 719, 1065, 1066
Morita, T. 1, 165, 259
Moroi, Y. 479
Morozov, Yu.V. 25
Morris, J.M. 112
Morrow, T. 49, 64, 393, 399, 984
Mosse, M. 291
Mueller, A. 106
Mukherjee, R. 854
Muller, A.J. 385, 1045
Munn, R.W. 931
Munro, I.H. 106, 173, 939
Muralidharan, S. 222
Murata, I. 413
Murata, K. 15
Murgia, S.M. 254, 604, 719
Murofushi, K. 173, 437, 527
Murty, B.A.R.C. 538
Muszkat, K.A. 436, 1046
Mutai, K. 452
Nafisi-Movaghar, J. 552
Nagakura, S. 6, 73, 83, 383, 452, 551, 577, 802, 844, 905, 1084
Nagumo, T. 259
Nahor, G.S. 1043
Naito, I. 7, 625, 912
Nakajima, A. 586
Nakajima, K. 13, 263
Nakamura, H. 62, 1049
Nakamura, J. 73, 427, 452, 551, 844
Nakamura, S. 1049
Nakamura, T. 939
Nakashima, N. 20, 62, 64, 109, 679, 1049
Nakasui, K. 413
Nakato, Y. 1049
Nakayama, T. 13, 76, 148, 154, 263, 293, 706, 768, 815
Naumova, T.M. 64
Navaratnam, S. 32, 292, 1009
Nelipovich, K.I. 768
Nemeth, G.A. 257
Neta, P. 552
Netzel, T.L. 1104
Neumann-Spallart, M. 840
Newport, G.L. 356
Nicholls, C.H. 1109, 1110
Nicol, M. 65
Niizuma, S. 866
Nikoforov, V.I. 64
Nishida, Y. 23
Nishijima, Y. 570
Nishimura, T. 64
Nohara, S. 1049
Nosaka, Y. 939
Nouchi, G. 25, 26, 64, 106, 307, 768, 1095
Novak, J.R. 64
Noyes, W.A., Jr. 575
Nozakura, S. 133
Nurmukhametov, R.N. 984
O'Dowd, R.F. 64
O'Hare, A. 64
O'Sullivan, M. 956
Obuchi, H. 133
Ohashi, Y. 188, 374
Ohmine, I. 109
Ohno, T. 261, 276, 279, 562, 719, 1130
Ohwada, S. 570, 913
Oka, M. 437
Okada, T. 342
Okamoto, K. 570
Okamoto, M. 154, 706
Okamoto, T. 15
Okumura, M. 1104
Olbrich, G. 420
Ono, I. 821
Orloff, M.K. 173
Orlowski, T.E. 659
Ortmann, W. 995
Osaka, K. 821
Osif, T.L. 719
Osipov, V.V. 132
Ostertag, R. 768
Ota, K. 173, 437, 527
Ottolenghi, M. 64, 463, 548, 787, 939, 966, 974, 976, 980
Pagni, R.M. 851
Pailthorpe, M.T. 1109, 1110
Pakula, B. 963
Pantke, E.R. 235
Parker, C.A. 1064
Parsons, B.J. 32, 581, 684, 1009
Pasha, I. 269
Pasqua, A. 604
Pasternak, C. 536
Pavlopoulos, T.G. 138, 214, 295, 303, 357, 768, 855, 939, 1021
Pecht, I. 715
Pekkarinen, L. 100
Peng, C.T. 1005
Pepmiller, C. 729
Peradejordi, F. 176
Pereyre, J. 1063
Periasamy, N. 23
Pernot, C. 631
Persy, G. 413
Peteleski, N. 173
Peters, K.S. 133
Peterson, F.C. 64
Petrov, S.N. 125
Phillips, D. 106
Phillips, G.O. 32, 36, 38, 292, 581, 684, 1009
Pierce, R.A. 392

- Pikel'ni, V.F. 166
Pileni, M.-P. 16, 643, 654, 922, 1131, 1143
Pineault, R.L. 632
Pirogov, N.O. 245
Plantenga, F.L. 233, 516
Plotnikov, V.G. 1100
Polansky, O.E. 114
Poletti, A. 254, 604, 719
Pomazan, Yu.V. 8, 173
Poplavskii, A.N. 350
Popov, L.S. 64, 173
Poppe, W. 684
Porter, G. 7, 8, 23, 35, 41, 62, 64, 106, 190, 259, 284, 552, 743, 819
Poskocil, J. 719
Potashnik, R. 64, 939
Poulos, A.T. 1031
Prabhu, K.V. 727
Pradevan, G.-O. 1143
Prasad, D.R. 204, 906
Prass, B. 24
Pratt, A.C. 931
Pratte, J.F. 513, 575
Prell, G. 693, 901
Previtali, C.M. 265
Priestley, E.B. 768
Prusik, T. 135
Ptak, M. 1109
Pugh, A.C. 64
Qin, L. 200
Quinn, M.F. 49, 982, 984
Quinn, M.J. 768
Rabani, J. 1043
Rabek, J.F. 152
Raemme, G. 152
Ramanan, G. 64
Ramsay, I.A. 173
Ranald, U.R. 484, 768
Ranby, B. 152
Rand, S.D. 224
Rankin, R. 393, 399
Ranogajec, F. 64
Rapp, W. 848
Rauscher, W. 25
Rayez, J.-C. 509, 803
Razumov, V.F. 64
Reddi, E. 298
Rehak, V. 719
Reichel, L.W. 385
Reilley, C.N. 989
Rentzepis, P.M. 83, 224
Reske, G. 137
Reynolds, A.H. 224
Rhodes, W. 608
Richards, J.T. 470, 939, 1049
Richtol, H.H. 809, 990
Ridge, R.J. 1035
Rivera, V. 936
Robbins, R.J. 62
Roberge, P.C. 509, 545, 803, 845, 1059
Robert, M.-P. 320
Roberts, J.P. 64, 369
Robinson, E.A. 1006
Robinson, G.W. 768
Rodgers, M.A.J. 91, 92, 173, 298, 939, 1023
Rodionov, A.N. 402
Roe, D.K. 867
Roe, E.M.F. 645
Roitman, G.P. 97, 722
Romanov, N.N. 64, 435
Romashov, L.V. 23, 719
Rommel, E. 413, 606, 851
Ronfard-Haret, J.-C. 143, 238, 246, 919
Rosenfeld, T. 966, 974, 976, 980
Ross, I.G. 64
Rothenberger, G. 224
Rougee, M. 1140
Rousset, A. 1095
Roy, J.K. 1126, 1128
Rudolph, E. 901
Rulliere, C. 509, 545, 803, 1059
Ruppel, D. 173
Rushton, F.A.P. 768, 935
Ruziewicz, Z. 1095
Rygalov, L.N. 869
Sa E Melo, M.T. 246, 598
Sabbah, S. 238
Sadovskii, N.A. 381
Sagun, E.I. 284
Saito, T. 464, 946
Saito, Y. 1049
Sakaguchi, Y. 427
Sakata, Y. 73, 846
Sakhno, T.V. 350
Sakuragi, H. 427
Salet, C. 51, 502, 598, 640, 935, 1067, 1070, 1072, 1105, 1114
Salmon, G.A. 64, 133, 1005, 1006
Saltiel, J. 802
Salvin, R. 482
Santos, O. 956
Santus, R. 16, 21, 57, 599, 631, 643, 760, 1109
Sapunov, V.V. 240, 579
Sarkanen, K. 284
Sato, K. 913
Savenko, A.K. 25
Scaiano, J.C. 82, 112, 133, 232, 353, 552, 613, 854, 1010, 1054, 1121
Scandola, F. 1106
Schaeffer, C.G. 133
Schaffner, K. 153
Scheerer, R. 552
Schmalstieg, H. 330
Schmitt, U. 696
Schnabel, W. 7, 10, 290, 332, 506, 625, 912
Schneider, B. 22
Schneider, R. 211

- Schomburg, H. 885
Schoof, S. 529
Schrader, U. 526
Schreiner, S. 645, 1063
Schulte-Frohlinde, D. 54, 95, 218, 229, 314, 334, 337, 420, 461, 535, 672, 678, 713
Schulz, R.C. 768
Schuster, G.B. 223, 589, 1069
Schuster, H. 768
Schwarz, H.A. 64
Scott, A.R. 164
Scott, G. 1050
Scott, G.W. 613, 764, 963
Seely, G.R. 100
Seki, H. 71
Seki, Y. 913
Selvarajan, N. 251
Selwyn, J.C. 854
Setif, P. 284
Shakhverdov, P.A. 287, 653
Shamraev, V.N. 984
Shamshev, V.N. 8, 173
Shatrov, V.D. 125
Shaw, R.W. 65
Shcherbo, S.N. 29, 423, 424
Sheck, Yu.B. 43
Sheena, H.H. 1050
Shekk, Yu.B. 64
Shepanski, J.F. 284
Shibuya, E. 173
Shigorin, D.N. 29, 350, 402, 423, 424
Shimizu, K. 552
Shinitzky, M. 536
Shioyama, H. 366, 946
Shirota, Y. 259
Shizuka, H. 9, 133, 165, 767
Shubin, V.V. 400
Shudo, K. 15
Shvedova, L.A. 64, 192
Shvindt, N.N. 991
Siegel, S. 64, 173, 768
Siemiarczuk, A. 463
Silvie, C. 307
Sime, M.E. 106
Sinai, N. 133
Sinclair, R.S. 34, 47, 297, 299, 541, 680, 931, 1035
Singer, L.A. 211
Singh, A. 164, 768
Singh, M. 258
Sinitsyna, Z.A. 1, 133, 1095
Skvortsov, B.V. 869, 991
Skvortsov, V.I. 768
Slavnova, T.D. 984, 986
Slifkin, M.A. 66, 144, 343, 536
Slocum, G.H. 223
Sloper, R.W. 684
Small, R.D. 133
Smalley, J.F. 652
Smirnov, V.A. 173, 1100
Smith, G.R. 23
Soboleva, I.V. 381
Soep, B. 25
Sokolova, L.K. 988
Solov'ev, K.N. 240, 579
Soloveichik, O.M. 370
Sommer, U. 106
Sopchyshyn, F. 164
Spikes, J.D. 298, 604
Srinivasan, B.N. 31
Staneva, T.G. 50
Stehlik, D. 24
Stein, G. 787
Steiner, U. 645, 1063, 1066
Stewart, L.C. 232
Stewart, T. 768
Strong, R.L. 727
Struve, W.S. 632
Subudhi, P.C. 505
Sugamori, S.E. 7
Sugawara, T. 20
Sumitani, M. 6, 73, 109, 802, 1049
Suryanarayanan, K. 517, 518, 820
Sutin, N. 1104
Swallow, A.J. 768
Swenson, G.W. 30
Sykes, A. 244, 254, 649, 974
Synowiec, J.A. 62
Syutkina, O.P. 402
Szabo, A.G. 58, 1069
Szoc, K. 586
Szwarc, M. 1021
Szymanski, M. 164
Taaffe, J.K. 64
Tai, S. 148
Tait, D. 299, 931, 1035
Takahashi, T. 546
Takamuku, S. 506
Takashima, M. 1121
Takematsu, A. 7, 822
Takemura, T. 439, 672, 748
Talley, L.D. 963
Tamai, N. 366, 526
Tammimga, J.J. 515, 516
Tamura, S.-I. 719, 1063
Tanabe, T. 366
Tanaka, I. 23, 247, 866, 1083
Tanaka, J. 62, 1049
Tanaka, J.A. 212
Tanaka, S. 133
Tanimoto, Y. 474, 552, 619, 1121
Tanner, D.W. 64
Tanno, T. 261
Tatemitsu, H. 846
Tatikolov, A.S. 64, 192, 441, 552
Tatischeff, I. 631
Tazuke, S. 366, 913
Tegner, L. 645
Teitelbaum, Z. 939

- ten Brink, H.T. 939
Teply, J. 861
Teranishi, H. 13, 76, 148, 154, 263, 293, 706, 815
Terenin, A.N. 653
Testa, A.C. 59, 467, 468, 812, 956
Tetreau, C. 173, 176, 414, 669
Tfibel, F. 23, 64, 245, 675
Theard, L.M. 64
Thomas, J.K. 171, 216, 254, 470, 768, 939, 1045, 1049
Thompson, G.F. 64, 393, 399
Tobita, S. 1083
Tokumaru, K. 427
Tokumura, K. 15, 619
Tolg, P.K. 1069
Tom, R. 1090
Tomioka, Y. 259
Tomlinson, M. 173
Topp, M.R. 106, 133
Torres, C. 936
Toth, M. 133
Toulme, J.J. 21
Traber, R. 715
Treinin, A. 5, 78
Truscott, T.G. 34, 47, 244, 246, 251, 254, 297, 299, 303, 541, 649, 680, 684, 751, 808, 930, 931, 972, 973, 974, 1035
Tsubomura, H. 133, 259, 470, 679, 1049
Tsvirko, M.P. 240, 579
Tung, C. 892
Tung, C.-H. 1045
Turro, N.J. 133, 892, 1045
Udagawa, H. 474, 552
Uji-ie, K. 23
Umehara, Y. 1084
Urruti, E.H. 733
Usacheva, M.N. 132
Usui, Y. 719, 1063
Uzhinov, B.M. 673
Val'kova, G.A. 29, 350, 423, 424
Vala, M. 854
Vandenburg, M.J. 32
Vandendriessche, J. 366
van den Ende, C.A.M. 516
Vander Donckt, E. 35, 64, 167, 308
Vanderkooi, J.M. 1125
van der Ploeg, P.M. 233
Vannikov, A.V. 64, 618
van Sinoy, A. 167
van Vooren, C. 64
van Voorst, J.D.W. 118, 939
van Zeyl, P.H.M. 516
Varma, C.A.G.O. 233, 515, 516, 525
Verhoeven, J.W. 525
Veyret, B. 31, 966
Vinogradov, A.M. 435
Viovy, R. 284
Visser, A.J.W.G. 714
Vogelmann, E. 25, 835
Volkert, W.A. 1109
Volkov, S.V. 12
Walmsley, R.H. 66, 343
Walrant, P. 599
Wardman, P. 64
Ware, W.R. 180
Waring, A.J. 1125
Warman, J.M. 470
Warwick, P. 768
Watanabe, S. 837
Watarai, H. 787
Watkins, A.R. 133, 348
Webber, S.E. 513, 575, 919
Webster, D. 505, 938
Weers, J.G. 83
Wegdam-van Beek, J. 118, 939
Weisman, R.B. 23
Weiss, K. 31, 410, 719, 871, 966, 972
Weller, H. 415
Welvert, Z. 414
Werner, T. 1069
West, G. 939
West, M.A. 23
Whillans, D.W. 1072, 1073, 1115
Whitten, D.G. 1126, 1128
Wieder, F. 177
Wierzchowski, K.L. 58
Wilbrandt, R. 605, 972
Wild, U.P. 64, 484, 641, 725, 768, 1046
Wildes, P.D. 720
Wilkinson, F. 1, 64, 133, 234, 254, 529, 552, 631, 768, 878, 1121
Williams, D.J. 284, 659
Willsher, C.J. 133, 768
Windsor, M.W. 7, 64, 65, 827, 940
Winter, G. 1063, 1066
Wirz, J. 330, 413, 606, 774, 851, 886, 951
Wismontski-Knittel, T. 88, 209, 352, 682
Wojtczak, J. 164
Wolff, T. 799, 900
Wolleben, J. 59, 467, 812
Wood, B. 1023
Woodruff, W.H. 92
Woods, R.J. 23
Wulff, V.J. 972
Wyman, G.M. 337
Wyrzykowska, K. 410
Yamada, A. 261
Yamada, H. 679
Yamada, T. 211
Yamamoto, M. 570
Yamamoto, N. 133, 470, 1049
Yamamoto, S. 23, 64, 71, 80, 245, 570, 822
Yamamoto, S.-A. 23, 80, 166
Yamamuro, T. 247
Yamashita, H. 492
Yasoshima, S. 946
Yates, S.F. 1069
Yavrouian, A. 613
Yellowlees, L.J. 1105

- | | | | |
|-------------------|--------------------------------|-------------------|------------------|
| Yildiz, A. | 989 | Zakhs, E.R. | 670 |
| Yip, R.W. | 7, 1069 | Zander, M. | 114 |
| Yokoyama, K. | 263, 452, 1049 | Zanker, V. | 22, 25, 693, 901 |
| Yorozu, T. | 171 | Zanocco, A. | 232 |
| Yoshida, K. | 171 | Zavyalov, Yu.A. | 1100 |
| Yoshida, M. | 846, 966 | Zen'kevich, E.I. | 284 |
| Yoshihara, K. | 6, 20, 62, 109, 112, 802, 1049 | Zhmyreva, I.A. | 12 |
| Yoshino, J. | 173, 437 | Zhuravlyeva, T.S. | 237 |
| Young, R.H. | 719 | Zupancic, J.J. | 589 |
| Younis, A.I. | 947 | Zwart, J.P. | 233 |
| Zachariasse, K.A. | 526 | Zwicker, E.F. | 562 |
| Zador, E. | 470 | | |