

Thermodynamic Properties of Twenty-One Monocyclic Hydrocarbons

Cite as: Journal of Physical and Chemical Reference Data **15**, 437 (1986); <https://doi.org/10.1063/1.555773>
Published Online: 15 October 2009

O. V. Dorofeeva, L. V. Gurvich, and V. S. Jorish



View Online



Export Citation

ARTICLES YOU MAY BE INTERESTED IN

NIST-JANAF Thermochemical Tables. II. Three Molecules Related to Atmospheric Chemistry: HNO_3 , H_2SO_4 , and H_2O_2

Journal of Physical and Chemical Reference Data **32**, 879 (2003); <https://doi.org/10.1063/1.1547435>

NIST-JANAF Thermochemical Tables. I. Ten Organic Molecules Related to Atmospheric Chemistry

Journal of Physical and Chemical Reference Data **30**, 475 (2001); <https://doi.org/10.1063/1.1364518>

Ideal Gas Thermodynamic Properties of Ethane and Propane

Journal of Physical and Chemical Reference Data **2**, 427 (1973); <https://doi.org/10.1063/1.3253123>

Where in the **world** is AIP Publishing?
Find out where we are exhibiting next

AIP
Publishing

Thermodynamic Properties of Twenty-One Monocyclic Hydrocarbons

O. V. Dorofeeva, L. V. Gurvich, and V. S. Jorish

Institute for High Temperatures, Izhorskaya ul. 13/19, Moscow 127412 USSR

The available structural parameters, fundamental frequencies, and relative energies of different stable conformers, if any, for cyclopropane, cyclopropene, cyclobutane, cyclobutene, 1,3-cyclobutadiene, cyclopentane, cyclopentene, 1,3-cyclopentadiene, cyclohexane, cyclohexene, 1,3-cyclohexadiene, 1,4-cyclohexadiene, cycloheptane, cycloheptene, 1,3-cycloheptadiene, 1,3,5-cycloheptatriene, cyclooctane, cyclooctene, 1,3-cyclooctadiene, 1,5-cyclooctadiene, and 1,3,5,7-cyclooctatetraene were critically evaluated and the recommended values selected. Molecular constants for some molecules were estimated as the experimental values for these compounds are not available. This information was utilized to calculate the ideal gas thermodynamic properties C_p° , S° , $-(G^\circ - H_0^\circ)/T$, $H^\circ - H_0^\circ$, and $\log K_r$ from 100 to 1500 K. The thermal functions were obtained using the rigid-rotor harmonic-oscillator approximation. The contributions derived for the inversion motion of cyclobutane and cyclopentene were obtained from energy levels calculated with the potential functions. For cyclopentane the pseudorotational contributions to thermal functions were calculated by assuming the pseudorotation as the free rotation of the molecule. The calculated values of the thermal functions are compared with those reported in other work. Agreement with experimental data, where such are available, is satisfactory within the experimental uncertainties.

Key words: ideal gas thermodynamic properties; molecular structure; monocyclic hydrocarbons with carbon atom numbers from 3 to 8; vibrational assignments.

Contents

1. Introduction	438
2. Cyclopropane	445
3. Cyclopropene	446
4. Cyclobutane	446
5. Cyclobutene	447
6. 1,3-Cyclobutadiene	448
7. Cyclopentane	448
8. Cyclopentene	449
9. 1,3-Cyclopentadiene	450
10. Cyclohexane	451
11. Cyclohexene	451
12. 1,3-Cyclohexadiene	452
13. 1,4-Cyclohexadiene	452
14. Cycloheptane	453
15. Cycloheptene	454
16. 1,3-Cycloheptadiene	454
17. 1,3,5-Cycloheptatriene	455
18. Cyclooctane	455
19. <i>trans</i> -Cyclooctene	456
20. <i>cis,cis</i> -1,3-Cyclooctadiene	457
21. <i>cis,cis</i> -1,5-Cyclooctadiene	458
22. 1,3,5,7-Cyclooctatetraene	458
23. References	462

List of Tables

1. Molecular and thermochemical constants for the three-membered rings	438
2. Molecular and thermochemical constants for the four-membered rings	439
3. Molecular and thermochemical constants for the five-membered rings	439
4. Molecular and thermochemical constants for the six-membered rings	440
5. Vibrational frequencies (cm^{-1}) for the six-membered rings	441
6. Molecular and thermochemical constants for the seven-membered rings	442
7. Vibrational frequencies (cm^{-1}) for the seven-membered rings	443
8. Molecular and thermochemical constants for the eight-membered rings	444
9. Vibrational frequencies (cm^{-1}) for the eight-membered rings	445
10. Ideal gas thermodynamic properties for cyclopropane	446
11. Ideal gas thermodynamic properties for cyclopropene	446
12. Ideal gas thermodynamic properties for cyclobutane	447
13. Ideal gas thermodynamic properties for cyclobutene	447

©1986 by the U. S. Secretary of Commerce on behalf of the United States.
This copyright is assigned to the American Institute of Physics and the American Chemical Society.
Reprints available from ACS; see Reprints List at back of issue.

14. Ideal gas thermodynamic properties for 1,3-cyclobutadiene	448	24. Ideal gas thermodynamic properties for 1,3-cycloheptadiene	455
15. Ideal gas thermodynamic properties for cyclopentane	449	25. Ideal gas thermodynamic properties for 1,3,5-cycloheptatriene	455
16. Ideal gas thermodynamic properties for cyclopentene	450	26. Ideal gas thermodynamic properties for cyclooctane	456
17. Ideal gas thermodynamic properties for 1,3-cyclopentadiene	450	27. Ideal gas thermodynamic properties for <i>trans</i> -cyclooctene	457
18. Ideal gas thermodynamic properties for cyclohexane	451	28. Ideal gas thermodynamic properties for <i>cis,cis</i> -1,3-cyclooctadiene	457
19. Ideal gas thermodynamic properties for cyclohexene	452	29. Ideal gas thermodynamic properties for <i>cis, cis</i> -1,5-cyclooctadiene	458
20. Ideal gas thermodynamic properties for 1,3-cyclohexadiene	452	30. Ideal gas thermodynamic properties for 1,3,5,7-cyclooctatetraene	458
21. Ideal gas thermodynamic properties for 1,4-cyclohexadiene	453	31. The uncertainties of the calculated thermal functions and adopted enthalpies of formation..	459
22. Ideal gas thermodynamic properties for cycloheptane	454	32. Comparison of experimental entropy with calculated values (in $\text{J K}^{-1} \text{mol}^{-1}$)	460
23. Ideal gas thermodynamic properties for cycloheptene	454	33. Comparison of experimental heat capacities with calculated values (in $\text{J K}^{-1} \text{mol}^{-1}$)	461

1. Introduction

This report involves the calculation of the ideal gas thermodynamic properties of 21 monocyclic hydrocarbons with carbon atom numbers from 3 to 8 (cycloalkanes and their unsaturated analogs). For a few of these molecules, thermodynamic properties have been reported. Recently, more complete and reliable information has become available on the structure and vibrational assignments of monocyclic hydrocarbons. This information permits us to make more precise calculations of thermal functions of some molecules and to calculate the thermal functions of others for the first time. A number of substances, viz., 1,4-cycloheptadiene, 1,4-cyclooctadiene, 1,3,5-, and 1,3,6-cyclooctatriene were not considered in this work due to the unreliability of structural and conformational data and due to the lack of vibrational assignments.

The available data on vibrational frequencies, structural parameters, and energies of different stable conformations have been critically examined and the most reliable values have been selected. Molecular constants for some compounds were estimated in the present work as the experimental values for these molecules are not available, incomplete, or unreliable. The selected molecular constants are given in Tables 1-9.

Based on the selected values of molecular constants, the ideal gas thermal functions {heat capacity (C_p°), entropy (S°), Gibbs energy function [$-(G^\circ - H_0^\circ)/T$], and enthalpy ($H^\circ - H_0^\circ$)} were calculated by the standard statistical mechanical method using rigid-rotor harmonic-oscillator approximations. The enthalpy of formation ($\Delta_f H^\circ$) and the calculated thermal functions have been used to calculate the logarithm of the equilibrium constant of formation ($\log K_f$) by the usual thermodynamic formulas [subscript f denotes formation by the reaction $n\text{C}(\text{c, graphite}) + m\text{H}_2(\text{g}) = \text{C}_n\text{H}_{2m}(\text{g})$]. The procedures for calculation of thermodynamic properties are similar to

those used in *Thermodynamic Properties of Individual Substances* by Gurvich *et al.*¹ The fundamental physical constants and thermodynamic properties of the elements in their reference states used in the calculations are also taken from Ref. 1. The chemical thermodynamic property values

Table 1. Molecular and thermochemical constants for the three-membered rings






	Cyclopropane  C_3H_6	Cyclopropene  C_3H_4
Molecular weight	42.0804	40.0646
Point group	D_{3h}	C_{2v}
Symmetry number	6	2
Ground state statistical weight	1	1
Product of the three principal moments of inertia, $\text{g}^2 \text{cm}^6 \times 10^{11}$	117	65.3
Vibrational frequencies, cm^{-1} :		
ν_1	A_1' 3038	A_1 3152
ν_2	1479	2909
ν_3	1188	1653
ν_4	A_1'' 1126	1483
ν_5	A_2' 1070	1105
ν_6	A_2'' 3101.7	905
ν_7	854	A_2 996
ν_8	E' 3024.4	815
ν_9	1457.7	B_1 3110
ν_{10}	1028.4	1043
ν_{11}	868.5	1011
ν_{12}	E'' 3082	769
ν_{13}	1188	B_2 2995
ν_{14}	739	1088
ν_{15}		569
Enthalpy of formation at 298.15 K, kJ mol^{-1}	53.3	277.1




Table 2. Molecular and thermochemical constants for the four-membered rings

	Cyclobutane  C ₄ H ₈	Cyclobutene  C ₄ H ₆	1,3-Cyclobutadiene  C ₄ H ₄
Molecular weight	56.1072	54.0914	52.0756
Point group	D _{2d}	C _{2v}	D _{2h}
Symmetry number	8 ^a	2	4
Ground state statistical weight	1	1	1
Product of the three principal moments of inertia, g ² cm ⁶ ×10 ¹¹⁶	62	55	36
Vibrational frequencies, cm ⁻¹ :			
ν ₁	A ₁ 2974	A ₁ 3063	A _g 3050
ν ₂			1510
ν ₃			1120
ν ₄			900
ν ₅			A _u 1100
ν ₆			570
ν ₇	A ₂ 1300		B _{1g} 3030
ν ₈			1235
ν ₉	B ₁ 1234	A ₂ 2955	770
ν ₁₀			B _{2g} 900
ν ₁₁			B _{2g} 900
ν ₁₂	B ₂ 2987		B _{1u} 570
ν ₁₃			B _{2u} 3030
ν ₁₄		B ₁ 3056	1240
ν ₁₅			720
ν ₁₆			B _{3u} 3040
ν ₁₇	E 2965		1520
ν ₁₈			990
ν ₁₉			
ν ₂₀			
ν ₂₁		B ₂ 2961	
ν ₂₂			
ν ₂₃			
ν ₂₄			
Enthalpy of formation at 298.15 K, kJ mol ⁻¹	28.4	156.7	385

^a Although cyclobutane has the nonplanar conformation (D_{2d} symmetry), the symmetry number 8 is given in the table due to the fact that the molecule is undergoing inversion through its planar configuration (D_{4h} symmetry).

^b The ring-puckering frequency ν₆ = 199 cm⁻¹ is not given in table as the contribution derived for the inversion motion of cyclobutane has been obtained by the direct summation over the energy levels.

Table 3. Molecular and thermochemical constants for the five-membered rings

	Cyclopentane  C ₅ H ₁₀	Cyclopentene  C ₅ H ₈	1,3-Cyclopentadiene  C ₅ H ₆
Molecular weight	70.134	68.1182	66.1024
Point group	C ₁	C _s	C _{2v}
Symmetry number	1	2 ^c	2
Ground state statistical weight	1	1	1
Product of the three principal moments of inertia, g ² cm ⁶ ×10 ¹¹⁵	38.75	28.39	19.96
Vibrational frequencies, cm ⁻¹ :			
ν ₁	2960(5) ^a	A ₁ 3070	A ₁ 3091
ν ₂	2880(5)		3075
ν ₃	1480(3)		2886
ν ₄	1455(2)		1500
ν ₅	1310(2)		1378
ν ₆	1207(2)		1365
ν ₇	1250(2)		1106
ν ₈	1210(2)		994
ν ₉	1160(2)		915
ν ₁₀	1027(2)		802
ν ₁₁	1022		A ₂ 1100
ν ₁₂	985	A ₂ 2938	941
ν ₁₃	949		700
ν ₁₄	896		516
ν ₁₅	886		B ₁ 3105
ν ₁₆	858		3043
ν ₁₇	827		1580
ν ₁₈	770	B ₁ 3068	1292
ν ₁₉	617		1239
ν ₂₀	545		1090
ν ₂₁	283		959
ν ₂₂	- ^b		805
ν ₂₃			B ₂ 2900
ν ₂₄			925
ν ₂₅			891
ν ₂₆			664
ν ₂₇		B ₂ 2963	350
ν ₂₈			2933
ν ₂₉			1207
ν ₃₀			1047
ν ₃₁			695
ν ₃₂			593
ν ₃₃			- ^d
Enthalpy of formation at 298.15 K, kJ mol ⁻¹	-78.4	32.7	130.8





^a Numbers in parentheses represent the accidental multiplicities since some of the close-lying frequencies were averaged due to their large uncertainty.

^b The pseudorotational contribution was calculated by free rotation formulae¹.

^c Although cyclopentene has the nonplanar structure (C_s symmetry), the symmetry number 2 is given in the table due to the fact that the molecule is undergoing inversion through its planar configuration (C_{2v} symmetry).

^d The ring-puckering frequency ν₃₃ = 127 cm⁻¹ is not given in table as the contribution due to inversion of cyclopentene was obtained by direct summation over the energy levels.

Table 4. Molecular and thermochemical constants for the six-membered rings





	Cyclohexane ^a  C ₆ H ₁₂	Cyclohexene  C ₆ H ₁₀	1,3-Cyclohexadiene  C ₆ H ₈	1,4-Cyclohexadiene  C ₆ H ₈
Molecular weight	84.1608	82.145	80.1292	80.1292
Point group	D _{3d}	C ₂	C ₂	D _{2h}
Symmetry number	6	2	2	4
Number of optical isomers	1	2	2	1
Ground state statistical weight	1	1	1	1
Product of the three principal moments of inertia, g ³ cm ⁶ ×10 ¹¹⁴	13.35	10.71	8.517	9.26
Vibrational frequencies	see Table 5	see Table 5	see Table 5	see Table 5
Enthalpy of formation at 298.15 K, kJ mol ⁻¹	-123.3	-4.6	106.3	109.0

^a Another stable conformer was also taken into account for cyclohexane: point group D₂, symmetry number 4, number of optical isomers 2, statistical weight 1, relative energy 1925 cm⁻¹.

Table 5. Vibrational frequencies (cm^{-1}) for the six-membered rings

	Cyclohexane		Cyclohexene		1,3-Cyclohexadiene		1,4-Cyclohexadiene	
ν_1	A_{1g}	2936	A	3040	A	3050	A_g	3032
ν_2		2853		2940		3050		2822
ν_3		1465		2916		2939		1680
ν_4		1158		2865		2838		1426
ν_5		802		2839		1577		1197
ν_6		384		1660		1444		854
ν_7	A_{1u}	1380		1460		1330		530
ν_8		1150		1445		1243	A_u	1250
ν_9		1100		1353		1223		970
ν_{10}		1350		1343		1178		370
ν_{11}		1100		1240		1150	B_{1g}	1240
ν_{12}	A_{2g}	2914		1222		1059		706
ν_{13}		2863		1140		994	B_{2g}	3032
ν_{14}	A_{2u}	1457		1095		945		1377
ν_{15}		1039		1068		850		1280
ν_{16}		522		966		753		1035
ν_{17}	E_g	2924		905		559		574
ν_{18}		2095		612		506	B_{3g}	2875
ν_{19}		1445		657		201		1010
ν_{20}		1347		520	B	3050		985
ν_{21}		1268		392		3050		403
ν_{22}		1029		270		2004	B_{1u}	3042
ν_{23}		785	B	3078		2838		2840
ν_{24}		427		2960		1602		1439
ν_{25}	E_u	2934		2890		1435		1405
ν_{26}		2863		2878		1377		962
ν_{27}		1457		2858		1178		888
ν_{28}		1346		1455		1165	B_{2u}	2889
ν_{29}		1260		1450		1100		962
ν_{30}		906		1343		1040		625
ν_{31}		862		1325		1016		108
ν_{32}		241		1269		927	B_{3u}	3042
ν_{33}				1215		745		1642
ν_{34}				1140		658		1362
ν_{35}				1039		468		1159
ν_{36}				1009		298		887
ν_{37}				919				
ν_{38}				877				
ν_{39}				719				
ν_{40}				638				
ν_{41}				450				
ν_{42}				165				

Table 6. Molecular and thermochemical constants for the seven-membered rings

	Cycloheptane ^a  C ₇ H ₁₄	Cycloheptene ^b  C ₇ H ₁₂	1,3-Cycloheptadiene ^c  C ₇ H ₁₀	1,3,5-Cycloheptatriene  C ₇ H ₈
Molecular weight	98.1876	96.1718	94.156	92.1402
Point group	C ₂	C _{2v}	C _{2v}	C _{2v}
Symmetry number	2	1	1	1
Number of optical isomers	2	1	1	1
Ground state statistical weight	1	1	1	1
Product of the three principal moments of inertia, g ³ cm ⁶ × 10 ¹¹⁴	39.1	32.1	29.12	21.43
Vibrational frequencies	see Table 7	see Table 7	see Table 7	see Table 7
Enthalpy of formation at 298.15 K, kJ mol ⁻¹	-116.2	-9.4	94.2	182.8

Other stable conformers were also taken into account for cycloheptane, cycloheptene and 1,3-cycloheptadiene (point group, symmetry number (σ), number of optical isomers (n), statistical weight (p) and relative energy (T) are listed for each conformer):

^a C₂, $\sigma = 2$, $n = 2$, $p = 1$, $T = 1200$ cm⁻¹;

^b C_{2v}, $\sigma = 2$, $n = 2$, $p = 1$, $T = 500$ cm⁻¹;

^c C₁, $\sigma = 1$, $n = 2$, $p = 1$, $T = 200$ cm⁻¹;






C₂, $\sigma = 2$, $n = 2$, $p = 1$, $T = 500$ cm⁻¹.

Table 7. Vibrational frequencies (cm^{-1}) for the seven-membered rings

Cycloheptane	Cycloheptene	1,3-Cycloheptadiene	1,3,5-Cycloheptatriene
2925(7) ^a	\underline{A}' 3024	\underline{A}' 3018	3050(6) ^a
2860(7)	2964	3018	2950
1467	2926	2927	2850
1450	2881	2885	1650(3)
1446	2852	2858	1450(3)
1440	2852	2830	1400(2)
1430(3)	2837	1613	1200(3)
1360	1656	1433	1100(2)
1350	1457	1430	1000(4)
1310(3)	1443	1372	950(2)
1285(2)	1434	1356	900(2)
1230(3)	1339	1257	800
1210	1332	1252	750
1200(2)	1252	1043	700
1125	1234	1004	650
1100	1200	955	450(3)
1040	1072	882	350
1020	1042	856	300
1005(2)	983	767	225
950	875	683	
915	824	585	
850	746	363	
830	691	347	
810	479	242	
800	417	\underline{A}'' 3062	
735	353	3062	
690	190	2963	
650	\underline{A}'' 3062	2905	
515	2963	1645	
490	2924	1448	
400	2854	1397	
335	2842	1341	
320	1447	1300	
273	1439	1223	
186	1391	1195	
123	1357	1153	
	1323	1077	
	1270	971	
	1234	895	
	1207	853	
	1144	835	
	1104	730	
	1024	577	
	985	377	
	960	165	
	889		
	832		
	585		
	469		
	312		
	209		

^a Numbers in parentheses are the accidental multiplicities (some of the close-lying frequencies were averaged due to their large uncertainty).

Table 8. Molecular and thermochemical constants for the eight-membered rings

	Cyclooctane ^a  C ₈ H ₁₆	Cyclooctene ^b  C ₈ H ₁₄	1,3-Cyclooctadiene ^c  C ₈ H ₁₂	1,5-Cyclooctadiene ^d  C ₈ H ₁₂	1,3,5,7-Cyclooctatetraene  C ₈ H ₈
Molecular weight	112.2144	110.1986	108.1828	108.1828	104.1512
Point group	C _{8h}	C ₂	C ₂	C ₂	D _{2d}
Symmetry number	1	2	2	2	4
Number of optical isomers	1	2	2	2	1
Ground state statistical weight	1	1	1	1	1
Product of the three principal moments of inertia, g ³ cm ⁶ × 10 ¹¹⁴	92	80	66	66	51.67
Vibrational frequencies	see Table 9	see Table 9	see Table 9	see Table 9	see Table 9
Enthalpy of formation at 298.15 K, kJ mol ⁻¹	-124.4	-27.0	80.5	58.6	297.6

Other stable conformers were also taken into account for cyclooctane, cyclooctene, 1,3-cyclooctadiene, and 1,5-cyclooctadiene (point group, symmetry number (σ), number of optical isomers (n), statistical weight (p), and relative energy (T) are listed for each conformer):

$$\begin{aligned}
 & \left. \begin{array}{l}
 {}^a \underline{D}_2, \sigma = 4, n = 2 \\
 \underline{C}_1, \sigma = 1, n = 2 \\
 \underline{C}_{2v}, \sigma = 2, n = 1 \\
 \underline{C}_2, \sigma = 2, n = 2 \\
 \underline{C}_1, \sigma = 1, n = 2
 \end{array} \right\} p = 1, T = 275 \text{ cm}^{-1} \\
 & {}^b \underline{C}_2, \sigma = 2, n = 2, p = 1, T = 1000 \text{ cm}^{-1}; \\
 & {}^c \underline{C}_1, \sigma = 1, n = 2, p = 1, T = 200 \text{ cm}^{-1}; \\
 & \left. \begin{array}{l}
 {}^d \underline{C}_{2h}, \sigma = 2, n = 1 \\
 \underline{C}_s, \sigma = 1, n = 1
 \end{array} \right\} p = 1, T = 1500 \text{ cm}^{-1}.
 \end{aligned}$$

Table 9. Vibrational frequencies (cm^{-1}) for the eight-membered rings

Cyclooctane	Cyclooctene	1,3-Cyclooctadiene	1,5-Cyclooctadiene	1,3,5,7-Cyclooctatetraene
2933	3040(2) ^a	3040(4) ^a	3040(4) ^a	A ₁ 3013
2921	2920(4)	2930(2)	2910(4)	1651
2915(5) ^a	2900(2)	2900(2)	2850(4)	1202
2908	2850(6)	2885(2)	1660(2)	873
2885(2)	1660	2860(2)	1450(4)	758
2880	1450(6)	1635(2)	1390(2)	194
2872	1390	1450(3)	1350	A ₂ 2950
2867(2)	1350(4)	1360(5)	1310(2)	1400
2859	1300(3)	1310(2)	1260(2)	1000
2853	1240(3)	1250(2)	1200(3)	900
1483	1140(3)	1200(2)	1160(2)	190
1472	1080(3)	1110(3)	1060(2)	E ₁ 2954
1455	1050(2)	1090(2)	1020	1459
1442	980(2)	970(2)	980(2)	994
1400(6)	900(2)	930(2)	940(2)	967
1380(3)	870(2)	870(2)	890	655
1364	800	830(2)	820(3)	249
1360	770(2)	800	760	B ₂ 3015
1350	680	740(2)	720	1608
1297	550	620(2)	660	1204
1291	460(2)	530	590(2)	669
1260	350(4)	410	490(2)	285
1250	300	370	380	E 3015
1230	200	310	310	2965
1217	150	260	270(2)	1640
1175		220	210	1400
1137		190	120	1223
1135		140		967
1110				945
1088(2)				800
1046				625
988				365
956				
930(2)				
862				
854				
799				
768				
725(2)				
691				
667				
540				
514				
478				
367				
326				
292				
255				
220				
125				

^a Numbers in parentheses are the accidental multiplicities (some of the close-lying frequencies were averaged due to their large uncertainty).

at temperatures from 100 to 1500 K for a pressure of 1 atm (101 325 Pa) are given in Tables 10–30.

The enthalpy of formation $\Delta_f H^\circ$ (298.15 K) for all substances but 1,3-cyclobutadiene, 1,4-cyclohexadiene, and 1,3-

cyclooctadiene was taken from the compilation of Pedley and Rylance.²

The inversion motion contributions to the thermodynamic properties for cyclobutane and cyclopentene were calculated by the use of the potential function of type $V(x) = \frac{1}{2}kx^2 + ax^4$. These potential functions are based on experimentally observed transitions and barrier heights of the inversion mode for the respective molecules. The contributions due to inversion were obtained by the summation over the energy levels calculated from the potential functions. The procedure for energy levels calculations was described previously.^{3,4}

The pseudorotation contributions to the thermodynamic properties of cyclopentane were calculated using the free rotation model.¹

The thermodynamic properties of molecules with several stable conformations were calculated by assuming the same fundamental frequencies and principal moments of inertia for all conformers (only the discrepancies in symmetry numbers and relative energies were taken into account).

It should be emphasized that the chiral conformations (C_1, C_2, D_2 symmetry) exist as an equimolar mixture of two enantiomeric forms. The contribution to the thermodynamic properties of two optical isomers is obtained by adding the entropy of mixing, $S^\circ_{\text{mix}} = R \ln 2$, to S° and $-(G^\circ - H^\circ_0)/T$, which is equivalent to assuming the effective symmetry number $\sigma_{\text{eff}} = \sigma/n$ (σ is the symmetry number, $n = 2$ is the number of optical isomers). For that reason Tables 4, 6, and 8 show the numbers of optical isomers together with other molecular constants.

Uncertainties in the calculated thermal functions (Table 31) were obtained by taking into account the inaccuracy of selected molecular structural and spectroscopic data and inaccuracy due to the deviation from the rigid-rotor harmonic-oscillator model. The procedure for the approximate evaluation of these uncertainties was described previously.¹ Uncertainties in adopted enthalpies of formation (Table 31) for most of the substances were taken from data of Pedley and Rylance.² For 1,3-cyclobutadiene, 1,4-cyclohexadiene, and 1,3-cyclooctadiene uncertainties in $\Delta_f H^\circ$ (298.15 K) were estimated in the present work. The selection of the enthalpy of formation values for these three substances will be described below.

Comparisons of the calculated and experimental entropy and heat capacity values are given in Tables 32 and 33, respectively.

2. Cyclopropane

The rotational constants B_0 and C_0 were obtained from high-resolution infrared spectra^{5–7} and from pure rotational Raman spectra,^{8–10} respectively. The structural parameters estimated from rotational constants^{9,10} are consistent with electron diffraction data.¹¹ The product of the principal moments of inertia of cyclopropane (Table 1) was calculated using the more precise rotational constants of McCubbin *et al.*⁷ and Butcher and Jones.¹⁰

Various workers have studied the infrared and Raman spectra of cyclopropane.^{5–8,10,12–15} Duncan and McKean¹⁴

Table 10. Ideal gas thermodynamic properties for cyclopropane

T	C _p ^o	-(G ^o -H ₀ ^o)/T	S ^o	H ^o -H ₀ ^o	log K _f
K	J K ⁻¹ mol ⁻¹			kJ mol ⁻¹	
100	33.321	161.608	194.871	3.326	-39.474
200	38.907	184.814	218.993	6.830	-23.159
298.15	55.571	199.109	237.378	11.410	-18.298
300	55.941	199.346	237.723	11.513	-18.241
400	76.052	211.291	256.589	18.119	-16.051
500	93.859	222.249	275.532	26.641	-14.832
600	108.542	232.677	293.986	36.786	-14.099
700	120.682	242.708	311.658	48.265	-13.612
800	130.910	252.385	328.459	60.859	-13.268
900	139.658	261.731	344.396	74.398	-13.014
1000	147.207	270.760	359.510	88.750	-12.817
1100	153.749	279.486	373.855	103.806	-12.660
1200	159.432	287.923	387.482	119.471	-12.532
1300	164.377	296.084	400.443	135.668	-12.425
1400	168.688	303.982	412.767	152.326	-12.332
1500	172.453	311.631	424.556	169.387	-12.252

Table 11. Ideal gas thermodynamic properties for cyclopropene

T	C _p ^o	-(G ^o -H ₀ ^o)/T	S ^o	H ^o -H ₀ ^o	log K _f
K	J K ⁻¹ mol ⁻¹			kJ mol ⁻¹	
100	33.444	167.767	200.987	3.328	-147.962
200	39.353	190.989	223.401	6.882	-74.339
298.15	52.881	205.344	243.441	11.373	-50.367
300	53.164	205.580	243.619	11.472	-50.068
400	67.459	217.324	261.176	17.540	-38.066
500	80.503	227.765	277.736	24.985	-30.935
600	90.659	237.408	293.343	33.561	-26.219
700	98.992	246.455	307.964	43.056	-22.873
800	105.964	255.008	321.652	53.315	-20.378
900	111.949	263.134	334.488	64.219	-18.446
1000	117.084	270.679	346.556	75.677	-16.907
1100	121.526	278.280	357.928	87.613	-15.651
1200	125.378	285.370	368.672	99.963	-14.608
1300	128.725	292.172	378.843	112.672	-13.727
1400	131.639	298.716	388.491	125.693	-12.973
1500	134.161	305.004	397.662	138.987	-12.320

reported the complete vibrational assignment from their infrared measurements for the gaseous and solid states of cyclopropane. We have adopted the vibrational fundamentals, as shown in Table 1, of Duncan and McKean,¹⁴ with the exception that the more precise values of ν_6 and ν_8 - ν_{11} were taken from the high-resolution infrared spectrum of Duncan.⁶ Duncan and Burns¹⁶ confirmed their assignment by carrying out a normal coordinate analysis. More recent experimental and theoretical studies¹⁷⁻²⁰ are consistent with the vibrational assignment of Duncan.

Ideal gas thermodynamic properties for cyclopropane are listed in Table 10. The calculated value of S° (298.15 K) is in good agreement with the calorimetric value of Ruehrwein and Powell²¹ and with the other statistically calculated entropy values²¹⁻²³ (Table 32). The calculated values of C_p° agree well with available experimental data^{254,255} (Table 33). The thermal functions reported by Chang *et al.*²⁴ were obtained with the molecular constants estimated from molecular mechanics (empirical force field) calculation; the agreement with their functions is worse than with other calculations (for S° see Table 32; the discrepancy in C_p° amounted to 2 J K⁻¹ mol⁻¹ at 298.15 K). Due to the discrepancy in the molecular constants used, the agreement with the data of Karuppannan *et al.*²³ increases as the temperature increases (for S° see Table 32; the discrepancy in C_p° amounted to 2 J K⁻¹ mol⁻¹ at 1000 K). As we have used more reliable input data, our calculated thermodynamic properties are believed to be the most reliable.

3. Cyclopropene

The molecular structure of cyclopropene was investigated by electron diffraction²⁵ and microwave^{26,27} techniques. The structural parameters obtained in these studies are mutually consistent. The rotational constants determined by Stigliani *et al.*,²⁷ which are believed to be more precise, were used to compute the product of the three principal moments of inertia (Table 1).

Vibrational spectra of cyclopropene were studied by various workers.²⁸⁻³⁰ The fundamental frequencies used in our calculations (Table 1) were taken from the more reliable work of Yum and Eggers.³⁰ These authors observed vibra-

tional frequencies from infrared and Raman spectra for the vapor and liquid states and carried out a normal coordinate analysis to confirm the vibrational assignment. The frequencies calculated by Wieberg *et al.*^{31,256} are consistent with vibrational assignment given by Yum and Eggers.³⁰

Ideal gas thermodynamic properties for cyclopropene are listed in Table 11. Unfortunately, no experimental data on gaseous C_p° and S° are available for comparison; however, the thermal functions computed in this work are practically identical with those reported earlier^{30,32} (the comparison for S° is given in Table 32).

4. Cyclobutane

Infrared,³³⁻³⁷ Raman,³⁴⁻³⁶ nuclear magnetic resonance,³⁸ and electron diffraction³⁹⁻⁴² studies have shown that cyclobutane undergoes inversion and the ground-state molecular structure is puckered (D_{2d} symmetry). *Ab initio*^{43,44} and semiempirical⁴⁵⁻⁴⁹ calculations confirm these experimental results. Different conclusions, however, have been reported with regard to the degree of ring puckering and the value of methylene group rocking. The product of the principal moments of inertia for cyclobutane (Table 2) was calculated from electron diffraction structural data obtained by Takabayashi *et al.*,⁴² except for the methylene group rocking angle. Takabayashi *et al.*⁴² estimated this angle, $\tau = 6^\circ$, assuming the linear dependence of this parameter from the ring puckering angle. We adopted $\tau = 4^\circ$ according to data^{38,44,48,49} as they appear more reliable. The ring puckering angle $\phi = 26^\circ$ determined by Takabayashi *et al.*⁴² agrees well with the values obtained from nuclear magnetic resonance data³⁸ and the high quality *ab initio* calculation.⁴⁴ This result significantly differs from the previous conclusions^{33-36,41} ascribing to the cyclobutane ring much higher nonplanarity of approximately 35° .

Infrared and Raman spectra for the vapor, liquid, and solid phases of cyclobutane were investigated.^{33,35,36,50-53} Until recently, the vibrational frequencies determined by Miller *et al.*⁵² were believed to be the most reliable. However, later Bánhegyi *et al.*⁵⁴ carried out the *ab initio* calculation and reassigned some experimental vibrational frequencies. Aleksanyan and Antipov⁵³ reinvestigated the infrared and

Table 12. Ideal gas thermodynamic properties for cyclobutane

T	C_p°	$-(G^\circ-H_0^\circ)/T$	S°	$H^\circ-H_0^\circ$	$\log K_f$
K	$J K^{-1} mol^{-1}$			$kJ mol^{-1}$	
100	38.169	177.164	211.958	3.479	-33.224
200	40.633	202.400	241.155	7.750	-22.473
298.15	70.564	219.003	264.396	13.534	-19.645
300	71.048	219.286	264.831	13.663	-19.614
400	97.065	233.667	288.960	22.101	-18.532
500	122.029	247.104	313.397	33.176	-18.062
600	142.030	260.221	337.522	46.381	-17.842
700	159.915	272.942	360.846	61.533	-17.738
800	174.524	285.338	383.180	78.274	-17.688
900	186.985	297.403	404.473	96.363	-17.667
1000	197.008	309.131	424.743	115.612	-17.659
1100	206.856	320.524	444.044	135.850	-17.657
1200	214.761	331.585	462.368	156.940	-17.658
1300	221.590	342.325	479.658	178.764	-17.659
1400	227.496	352.745	496.478	201.226	-17.660
1500	232.622	362.861	512.354	224.239	-17.659

Raman spectra of cyclobutane in the three aggregation states and confirmed the results of Bánhegyi *et al.*³⁴ The vibrational frequencies given in Table 2 are those obtained by Miller *et al.*⁵² from infrared and Raman spectral measurements of gaseous cyclobutane, with the exception of ν_4 , ν_7 , ν_8 , ν_{10} , and ν_{15} modes, for which the values were taken from Refs. 53 and 54.

Cyclobutane is a molecule which undergoes inversion. The thermodynamic-property contributions due to inversion of the cyclobutane ring were obtained by the direct summation over the energy levels calculated with the potential function. The procedure for energy levels calculation was described previously.^{3,4} Several investigations of the double minimum ring-puckering potential function of cyclobutane have been reported.³⁴⁻³⁶ The potential function given by Miller and Capwell³⁶ was used to calculate 60 energy levels. The potential function for inversion is $V(x) = 6.932 \times 10^5 x^4 - 3.79 \times 10^4 x^2$ (where x is the ring-puckering coordinate) with a barrier height of $(518 \pm 5) \text{ cm}^{-1}$. The partition function for inversion is based on these 60 levels with the first 7 levels being the experimental levels from which the potential function was derived.

Ideal gas thermodynamic properties for cyclobutane are given in Table 12. The calculated value of S° (298.15 K) is $1 \text{ J K}^{-1} \text{ mol}^{-1}$ less than that obtained from calorimetric measurements⁵⁵ (Table 32). This discrepancy is practically within uncertainties of the experimental and calculated values. Another two available calculations (Table 32) are not suitable for comparison with our result. Rathjens *et al.*⁵⁰ obtained the excellent agreement with experiment by actually fitting the vibrational assignment for cyclobutane. The discrepancy between data of Rathjens *et al.*⁵⁰ and thermal functions given in Table 12 increases significantly at higher temperatures [up to $4.2 \text{ J K}^{-1} \text{ mol}^{-1}$ for S° (1000 K) and $3 \text{ J K}^{-1} \text{ mol}^{-1}$ for C_p° (1000 K)]. As Rathjens *et al.*⁵⁰ used unreliable molecular constants and calculated inversion contributions to thermal functions by an approximate method we suppose that our data are more precise. Chang *et al.*²⁴ used the molecular constants evaluated from the molecular mechanics calculation. Therefore, the good agreement between their and our values of entropy (Table 32) is fortu-

tous (the discrepancy in C_p° amounted to $2.8 \text{ J K}^{-1} \text{ mol}^{-1}$ at 500 K).

5. Cyclobutene

The molecular structure of cyclobutene has been studied by the visual electron diffraction method⁵⁶ and by microwave techniques.^{57,58} Cyclobutene has a planar structure. The rotational constants determined in microwave studies^{57,58} agree well. Bak *et al.*⁵⁸ studied the microwave spectra of cyclobutene and four of its isotopic species. From these data they calculated the structural parameters of cyclobutene. These parameters are in good agreement with those obtained by Wieberg and Wendoloski⁵⁹ from *ab initio* calculation. The product of the principal moments of inertia, given in Table 2, was calculated using the rotational constants of Bak *et al.*⁵⁸

Lord and Rea⁶⁰ observed both the Raman spectrum of the liquid and the infrared spectrum of gas and liquid, and made a complete assignment of the vibrational frequencies for cyclobutene. Unobserved ν_9 - ν_{12} modes of A_2 symmetry were evaluated by comparison with suitable fundamentals of B_2 symmetry. Sverdlov and Krainov⁶¹ carried out the normal coordinate analysis and proposed new assignments for several frequencies. The reassignment of spectral data⁶⁰ was also made by Suzuki and Nibler⁶² from their investigation of vibrational spectra of *cis*-3,4-dichlorocyclobutene. The most reliable assignment seems to be that reported by Aleksanyan and Garkusha⁶³ from infrared and Raman spectra of gaseous, liquid, and solid states of cyclobutene. This assignment was adopted in the present work, except for two unobserved frequencies of A_2 symmetry, ν_9 and ν_{11} (Table 2). We have preferred the vapor phase values whenever available. For the CH_2 -stretching mode (ν_9) the value of 2955 cm^{-1} was chosen in accordance with the estimation made by Lord and Rea.⁶⁰ The assignment of the CH_2 -rocking fundamental (ν_{11}) has been less certain. Lord and Rea⁶⁰ have evaluated it as 640 cm^{-1} , whereas Sverdlov and Krainov,⁶¹ using a normal coordinate treatment, have assigned the band at 1276 cm^{-1} to a ν_{11} fundamental. From comparison with the CH_2 -rocking fundamentals of cyclopentene⁶⁴ we estimated this frequency as $1000 \pm 150 \text{ cm}^{-1}$.

Table 13. Ideal gas thermodynamic properties for cyclobutene

T	C_p°	$-(G^\circ-H_0^\circ)/T$	S°	$H^\circ-H_0^\circ$	$\log K_f$
K	$J K^{-1} mol^{-1}$			$kJ mol^{-1}$	
100	35.049	180.384	214.009	3.362	-92.470
200	44.897	216.308	240.897	7.266	-49.177
298.15	64.415	219.844	261.962	12.557	-35.429
300	64.841	220.105	262.362	12.677	-35.260
400	88.075	233.393	284.220	20.331	-28.549
500	108.624	245.756	306.149	30.197	-24.649
600	125.542	257.612	327.501	41.933	-22.115
700	139.443	269.068	347.931	55.204	-20.341
800	151.035	280.149	367.330	69.745	-19.032
900	160.833	290.866	385.701	85.352	-18.026
1000	169.184	301.227	403.090	101.864	-17.228
1100	176.339	311.242	419.559	119.149	-16.579
1200	182.493	320.925	435.173	137.098	-16.041
1300	187.801	330.288	449.995	155.619	-15.586
1400	192.393	339.346	464.085	174.634	-15.197
1500	196.378	348.113	477.498	194.078	-14.858

Ideal gas thermodynamic properties for cyclobutene are presented in Table 13. There are no experimental data for comparison. The thermal functions calculated by Danti⁶⁵ are different from ours by up to $3.4 \text{ J K}^{-1} \text{ mol}^{-1}$ for S° (1000 K) and $2.7 \text{ J K}^{-1} \text{ mol}^{-1}$ for C_p° (298.15 K) (see also Table 32). As we used the most recent and complete data, our thermal functions are believed to be more reliable.

6. 1,3-Cyclobutadiene

The question of the ground-state structure of cyclobutadiene has been of great interest to both experimental and theoretical chemists over the past 15 years. The observed infrared spectrum of matrix isolated cyclobutadiene originally has been interpreted as indicating a square-planar structure for the molecule because the number of bands agreed with the number expected,⁶⁶⁻⁶⁸ and because the frequencies agreed with those calculated theoretically for a square ground state.^{69,70} However, a number of semiempirical⁷¹⁻⁷³ and good *ab initio*⁷⁴⁻⁸⁰ calculations of the geometry all agree in predicting a rectangular ground state instead. Masamune *et al.*⁸¹ concluded from an improved experimental infrared spectrum that the cyclobutadiene ring is not square but most likely rectangular. This is supported by the similarity found between Masamune's and the computed infrared spectrum of rectangular cyclobutadiene.^{77,79,80,82} Thus the evidence, both theoretical and experimental, leads to the conclusion that the cyclobutadiene ring in the ground state is not square but rectangular (D_{2h} symmetry).

There are no experimental data on the molecular structure of cyclobutadiene. The product of the principal moments of inertia (Table 2) was calculated based on the estimated structural parameters for planar rectangular geometry:

$$r(\text{C}=\text{C}) = 1.34 \pm 0.03 \text{ \AA},$$

$$r(\text{C}-\text{C}) = 1.55 \pm 0.03 \text{ \AA},$$

$$r(\text{C}-\text{H}) = 1.08 \pm 0.02 \text{ \AA},$$

and

$$\angle \text{C}-\text{C}-\text{H} = 135 \pm 3^\circ.$$

These parameters were adopted by comparing the results of *ab initio* calculations for cyclobutadiene^{75,76,78,80} and semiempirical calculations for cyclobutadiene and cyclobutene.⁷² (The experimental molecular structure is known for cyclobutene and it helps to estimate the uncertainty of theoretical predictions for cyclobutadiene.)

Only five vibrational fundamentals were assigned from the infrared spectrum of matrix isolated cyclobutadiene^{66,81,82}: 570 cm^{-1} (ν_{12} , B_{1u} , CH wagging); 1240 cm^{-1} (ν_{14} , B_{2u} , CCH deformation); 720 cm^{-1} (ν_{15} , B_{2u} , C-C stretching); 3040 cm^{-1} (ν_{16} , B_{3u} , CH stretching); and 1520 cm^{-1} (ν_{17} , B_{3u} , C=C stretching). These values were adopted in the present work (Table 2). The vibrational spectrum of cyclobutadiene was calculated by the *ab initio* method.^{77,79,80,82} While the calculated intensities were found to be in good agreement with those observed experimentally, an appreciable discrepancy exists between calculated and experimental frequency values (up to $100\text{--}250 \text{ cm}^{-1}$). To obtain more reliable data for unobserved fundamentals, a nor-

Table 14. Ideal gas thermodynamic properties for 1,3-cyclobutadiene

T	C_p°	$-(S^\circ - H_0^\circ)/T$	S°	$H^\circ - H_0^\circ$	$\log K_f$
K	$\text{J K}^{-1} \text{ mol}^{-1}$		kJ mol^{-1}		
100	33.620	172.314	205.614	3.330	-204.267
200	42.863	195.772	230.970	7.040	-102.414
298.15	60.968	210.733	251.328	12.103	-69.161
300	61.332	210.965	251.706	12.216	-68.745
400	80.069	223.707	271.974	19.307	-52.047
500	95.643	235.336	291.578	28.121	-42.098
600	108.029	246.273	310.154	38.328	-35.503
700	117.950	256.056	327.591	49.046	-30.814
800	126.171	265.553	343.867	61.867	-27.310
900	133.025	274.003	359.155	74.836	-24.595
1000	138.834	281.042	373.479	88.437	-22.426
1100	143.795	293.701	386.950	102.575	-20.657
1200	148.055	302.006	399.650	117.173	-19.186
1300	151.725	309.982	411.649	132.166	-17.943
1400	154.900	317.654	423.072	147.501	-16.878
1500	157.655	325.040	433.795	163.132	-15.957

mal coordinate calculation was performed. For this purpose we used the NCA (normal coordinate analysis and force constants refinement) program by Novikov.⁸³ The initial force field (12 force constants) was transferred from 1,3-cyclopentadiene.⁸⁴ After force constants refinement, the five observed band positions were reproduced with an accuracy of 1 cm^{-1} . The uncertainties of calculated values for unobserved frequencies (except for torsion frequencies) are less than 50 cm^{-1} . The vibrations involving the torsion motion are not observed experimentally and their force constants therefore cannot be evaluated in the same way as the other ones. These fundamentals [$\nu_5(A_u)$, $\nu_6(A_u)$, $\nu_{10}(B_{2g})$, and $\nu_{11}(B_{3g})$] were obtained by averaging the values calculated with the torsional force constants of 1,3-cyclopentadiene,⁸⁴ 1,3-cyclohexadiene,⁸⁵ and 1,3-cycloheptadiene⁸⁶; their uncertainties were estimated to be $50\text{--}150 \text{ cm}^{-1}$.

No experimental data on the enthalpy of formation of cyclobutadiene are available, but its value was calculated by *ab initio*⁷⁸ and semiempirical^{73,87} methods. We adopted the MINDO/3 value of $\Delta_f H^\circ$ (298.15 K) obtained by Kollmar *et al.*,⁷³ since in their opinion the extrapolation of the errors in the MINDO/3 enthalpy of formation for cyclobutane and cyclobutene suggests that the error in the case of cyclobutadiene is probably quite small.

Ideal gas thermodynamic properties for 1,3-cyclobutadiene, given in Table 14, are reported for the first time. No experimental data are available for comparison.

7. Cyclopentane

Kilpatrick *et al.*⁸⁸ introduced the concept of pseudorotation to explain the high gas phase entropy value of cyclopentane. The cyclopentane equilibrium conformation has been shown by Pitzer *et al.*^{88,89} to be puckered in contrast with the previously assumed planar conformation (D_{5h} symmetry), and to vary continuously on an equipotential energy surface. Semiempirical and *ab initio* calculations of the relative energies for different conformations of cyclopentane^{90-93,257} confirmed that a pseudorotation of the ring puckering takes place. This implies a close similarity between the energies of the nonplanar half-chair (C_2 symmetry) and envelope (C_s symmetry) conformations. The

most direct evidence for pseudorotation in cyclopentane has come from spectroscopic studies.⁹⁴⁻⁹⁶ Electron diffraction investigation⁹⁷ also agrees with the pseudorotation of cyclopentane and shows the ring to be decidedly nonplanar.

Since cyclopentane undergoes an almost unhindered pseudorotation, that is, all conformations of puckered molecule are energetically equivalent, the effective nonplanar structure (C_1 symmetry) is adopted for cyclopentane. The product of the principal moments of inertia, given in Table 3, was calculated from electron diffraction structural parameters obtained by Adams *et al.*⁹⁷ for nonplanar (C_2 and C_s symmetry) conformations of cyclopentane.

The vibrational spectrum of cyclopentane based on infrared and Raman studies^{98,99} was partly reassigned according to force field calculations.^{92,100} The frequencies, given in Table 3, were based on the assignments proposed in Refs. 92 and 100.

The contributions of unhindered pseudorotation to the thermal functions were calculated by free rotation formulas¹ assuming the reduced moment of inertia, $I_r = 11 \cdot 10^{-40}$ g cm² (spectroscopic value⁹⁴), and symmetry number, $\sigma_{\text{pseudo}} = 10$ [cyclopentane ring undergoes pseudorotation through its planar configuration (D_{5h} symmetry)].

Ideal gas thermodynamic properties for cyclopentane are listed in Table 15. The calculated values of S° at 298.15, 310.16, and 322.41 K coincide with the calorimetric entropies^{88,101} [for S° (298.15 K) see Table 32] and the calculated values of C_p° agree with experimental ones^{88,101,258} within 0.1–1.1 J K⁻¹ mol⁻¹ (Table 33). The agreement of calculated values of S° at 230, 260, and 323.2 K (274.4, 282.4, and 299.9 J K⁻¹ mol⁻¹, respectively) with earlier calorimetric data of Aston *et al.*²⁵⁹ (273.0, 282.4, and 297.9 J K⁻¹ mol⁻¹, respectively) is not as good as with more recent experimental data.^{88,101} Our thermal functions are in good agreement with those calculated by Kilpatrick *et al.*⁸⁸ and McCullough *et al.*¹⁰¹ at low temperatures. The discrepancies, however, increase at higher temperatures up to 1.8 J K⁻¹ mol⁻¹ for S° (1500 K)¹⁰¹ and 2.7 J K⁻¹ mol⁻¹ for C_p° (1500 K).⁸⁸ These discrepancies, as well as the discrepancies with thermal functions calculated by Chang *et al.*²⁴ are due to the difference in molecular constants used in calculations. As we have used more recent and reliable data, we

feel our calculated values of thermodynamic properties should be more reliable.

8. Cyclopentene

Microwave studies,¹⁰²⁻¹⁰⁴ electron diffraction investigation,¹⁰⁵ and analysis of the ring-puckering potential function from infrared and Raman spectra,¹⁰⁶⁻¹⁰⁸ as well as *ab initio* calculation²⁵⁷ established that cyclopentene has a nonplanar ground-state conformation (envelope, C_s symmetry) and the molecule undergoes inversion through its planar configuration (C_{2v} symmetry).

The values of rotational constants obtained in microwave investigations^{102,103} are somewhat different. Scharpen¹⁰⁴ reanalyzed the data of Butcher and Costain¹⁰³ taking into account vibrational-rotational coupling and obtained good agreement with the rotational constants of Rathjens.¹⁰² The product of the principal moments of inertia, given in Table 3, was calculated using the rotational constants of Rathjens.¹⁰² Assuming the values of some structural parameters, Rathjens¹⁰² evaluated the ring-puckering angle as 22°. From analysis of the ring-puckering potential of cyclopentene the dihedral angle was deduced to be 22–26°. ^{106,108,260} The electron diffraction value is somewhat larger (29°).¹⁰⁵ The value obtained by *ab initio* computation²⁵⁷ is probably too small (13.6°).

Vibrational spectra of cyclopentene were studied by various workers.^{64,109-113} The fundamental frequencies for gaseous cyclopentene used in our calculations (Table 3) were taken from the more detailed investigation of Villarreal *et al.*⁶⁴ These authors have measured the infrared and Raman spectra of cyclopentene and its isotopic species in all three aggregation states. In conjunction with a normal coordinate analysis, vibrational assignments are proposed on the basis of isotopic shift ratios, group frequency considerations, relative band intensities, and shapes, as well as depolarization ratios.

The thermodynamic-property contributions due to inversion of the cyclopentene ring were obtained by direct summation over the energy levels calculated with a potential function. The procedure for energy level calculation was described previously.^{3,4} Several investigations of the double minimum ring-puckering potential function of cyclopentene have been reported.^{104,106-108,260} Laane and Lord¹⁰⁶ have reported the potential function derived from experimentally observed transition of the inversion modes for cyclopentene. This function is based on an estimated value of reduced mass. More recently, Villarreal *et al.*²⁶¹ have calculated the value of reduced mass and the potential function was recalculated. In this work the potential function given by Villarreal *et al.*²⁶¹ based on the study of Laane and Lord¹⁰⁶ was used to calculate 60 energy levels. The potential function for inversion is $V(x) = 7.11 \times 10^5 x^4 - 25.7 \times 10^3 x^2$ (where x is the ring-puckering coordinate) with a barrier height (232 ± 5) cm⁻¹. The partition function for inversion is based on these 60 levels with the first 12 levels being the experimental levels from which the potential function was derived.

Ideal gas thermodynamic properties for cyclopentene

Table 15. Ideal gas thermodynamic properties for cyclopentane

T	C _p ^o	-(G ^o -H ₀ ^o)/T	S ^o	H ^o -H ₀ ^o	log K _f
K	J K ⁻¹ mol ⁻¹			kJ mol ⁻¹	
100	40.182	196.937	234.979	3.804	15.693
200	54.185	226.295	266.284	8.398	-3.338
298.15	82.759	242.469	292.857	15.023	-6.567
300	83.388	242.782	293.371	15.177	-6.652
400	118.150	258.986	322.122	25.255	-10.268
500	150.053	274.593	352.001	38.704	-12.671
600	177.072	289.986	381.820	55.100	-14.396
700	199.703	305.192	410.866	73.972	-15.694
800	219.795	320.160	438.814	94.923	-16.704
900	235.021	334.841	465.546	117.635	-17.511
1000	248.883	349.196	491.045	141.848	-18.167
1100	260.761	363.205	515.337	167.346	-18.709
1200	270.964	376.854	538.475	193.945	-19.163
1300	279.750	390.141	560.519	221.491	-19.546
1400	287.341	403.068	581.536	249.855	-19.873
1500	293.915	415.639	601.590	278.926	-20.155

Table 16. Ideal gas thermodynamic properties for cyclopentene

T	C_p^0	$-(G^0-H_0^0)/T$	S^0	$H^0-H_0^0$	$\log K_f$
K	$J K^{-1} mol^{-1}$			$kJ mol^{-1}$	
100	40.365	197.018	233.171	3.615	-34.310
200	54.759	223.495	264.743	8.250	-22.484
298.15	81.275	241.438	291.267	14.857	-19.295
300	81.836	241.746	291.772	15.008	-19.259
400	112.081	257.701	319.489	24.715	-17.986
500	138.986	272.862	347.473	37.306	-17.394
600	161.381	287.591	374.858	52.360	-17.089
700	179.949	301.952	401.174	69.455	-16.921
800	195.517	315.935	426.248	88.250	-16.822
900	208.709	329.526	450.059	108.460	-16.761
1000	219.901	342.718	472.646	129.928	-16.721
1100	229.601	355.513	494.075	152.418	-16.693
1200	237.882	367.915	514.417	175.802	-16.672
1300	245.019	379.934	533.746	199.956	-16.654
1400	251.186	391.582	552.136	224.776	-16.639
1500	256.530	402.875	569.652	250.165	-16.624

are presented in Table 16. The calculated value of S^0 (298.15 K) is $1.5 J K^{-1} mol^{-1}$ larger than that obtained from calorimetric measurements,¹⁰⁹ whereas the entropy values calculated in previous works^{109,114,115} agree well with the experimental entropy (Table 32). Beckett *et al.*¹⁰⁹ and Furuyama *et al.*¹¹⁴ calculated the thermal functions based on a classical partition function for inversion and unreliable vibrational frequencies. Draeger *et al.*¹¹⁵ used the reduced potential function¹⁰⁴ based on the study of Laane and Lord,¹⁰⁶ and employed the same method for computing of inversion contributions as in the present work. However, the rather unreliable vibrational assignment of Furuyama *et al.*¹¹⁴ was adopted by Draeger *et al.*¹¹⁵ Unfortunately, Draeger *et al.*¹¹⁵ have not reported the values of entropy contributions due to inversion and thus the discrepancies between their data and thermal functions given in Table 16 cannot be analyzed in detail. Nevertheless, we state that the main discrepancies between the thermal functions obtained in this work and calculated by other authors^{109,114,115} [up to $6.5 J K^{-1} mol^{-1}$ for S^0 (1500 K)¹¹⁵ and $2.5 J K^{-1} mol^{-1}$ for C_p^0 (298.15 K)¹¹⁵] are due to the difference in the adopted vibrational frequencies. If we had adopted the same values of vibrational frequencies as Draeger *et al.*¹¹⁵ had, we would have obtained the value of S^0 (298.15 K) = $290.0 J K^{-1} mol^{-1}$, which is in good agreement with the experimental value of S^0 (298.15 K) = $289.7 J K^{-1} mol^{-1}$. However, the vibrational assignment of Villarreal *et al.*⁶⁴ seems to be the most reliable (see above) and we cannot find any reason for its modification. Thus, the use of the vibrational frequencies of Villarreal *et al.*⁶⁴ has not enabled us to obtain good agreement between the calculated and experimental values of S^0 (298.15 K). One can suggest that the discrepancy between the calculated and experimental entropy values is within the experimental uncertainty. Beckett *et al.*¹⁰⁹ do not indicate the experimental uncertainty, but, as is seen from Table 32, it might be expected to fall within the range $0.8-1.5 J K^{-1} mol^{-1}$.

9. 1,3-Cyclopentadiene

Scharpen and Laurie¹¹⁶ have made microwave measurements of cyclopentadiene and three of its isotopic derivatives containing ^{13}C from which the structural param-

eters for the planar carbon atom ring were found. Damiani *et al.*¹¹⁷ have studied the microwave spectra of five deuterated species of cyclopentadiene and have calculated the complete molecular structure using the data of Scharpen and Laurie.¹¹⁶ The rotational constants obtained in this way do not significantly differ from the values reported by Scharpen and Laurie. For computation of the product of the principal moments of inertia (Table 3) the rotational constants of Damiani *et al.*¹¹⁷ were used. Experimental structural parameters^{116,117} are consistent with those obtained from theoretical calculations.^{59,118,257}

Various workers have studied the infrared and Raman spectra of cyclopentadiene,^{84,119-123} but the available vibrational assignments have differed appreciably from one another. The more reliable interpretation of the vibrational spectrum of cyclopentadiene was proposed by Gallinella *et al.*^{84,122} These authors investigated the infrared spectra in the three aggregation states and the liquid- and solid-phase Raman spectra of cyclopentadiene and its deuterated species. In order to confirm the vibrational assignment they also presented the results of a normal coordinate treatment. The fundamentals, given in Table 3, are those for liquid cyclopentadiene reported by Gallinella *et al.*,⁸⁴ as the assignment for vapor phase is incomplete. The use of liquid state frequencies instead of those for the vapor state causes small errors in calculated thermal functions (the average vapor-liquid shift is found to be $5 cm^{-1}$).

Ideal gas thermodynamic properties for 1,3-cyclopentadiene are listed in Table 17. Grant and Walsh¹²⁴ reported a measurement of the entropy of cyclopentadiene from a study of the equilibrium between H_2 , condensed cyclopentene, and condensed cyclopentadiene (see Table 32). There are no calorimetric determinations of this entropy. The entropy values computed in this work are in good agreement with those obtained in other calculations^{114,125} at low temperatures. All calculated values of S^0 (298.15 K) are different from the entropy value obtained using equilibrium data¹²⁴ (Table 32), but the inaccuracy of the latter is rather large. The difference in molecular constants used in this work and other calculations^{114,125} is the reason for discrepancies in entropy

Table 17. Ideal gas thermodynamic properties for 1,3-cyclopentadiene

T	C_p^0	$-(G^0-H_0^0)/T$	S^0	$H^0-H_0^0$	$\log K_f$
K	$J K^{-1} mol^{-1}$			$kJ mol^{-1}$	
100	35.045	188.227	221.808	3.358	-78.587
200	46.526	212.372	249.654	7.456	-42.074
298.15	75.367	228.642	274.039	13.535	-30.560
300	75.892	228.923	274.507	13.675	-30.419
400	103.300	243.515	300.164	22.666	-24.830
500	126.449	257.417	325.792	34.188	-21.594
600	145.032	270.892	350.552	47.796	-19.498
700	160.045	283.968	374.075	63.075	-18.033
800	172.413	296.632	396.299	79.717	-16.953
900	182.774	308.876	417.202	97.491	-16.125
1000	191.553	320.706	436.926	116.219	-15.468
1100	199.043	332.126	455.543	135.759	-14.934
1200	205.467	343.151	473.145	155.993	-14.492
1300	210.997	353.797	489.815	176.823	-14.118
1400	215.775	364.083	505.630	198.167	-13.798
1500	219.919	374.024	520.662	219.956	-13.520

values at high temperatures [up to $2.6 \text{ J K}^{-1} \text{ mol}^{-1}$ for S° (1000 K)¹²⁵] and in heat capacity values [up to 3.0^{125} and $4.2 \text{ J K}^{-1} \text{ mol}^{-1}$ ¹¹⁴ for C_p° (500 K)].

10. Cyclohexane

Electron diffraction studies,^{126–130} analysis of vibrational spectra,^{92,131–137} dynamic nuclear magnetic resonance spectroscopy,¹³⁸ and theoretical calculations^{90,139–143} indicate that the chair conformation (D_{3d} symmetry) is the most stable form of cyclohexane. The strain-energy calculations^{90,139–143} show that there are also two conformations, twist-boat (D_2 symmetry) and boat (C_2 symmetry), with an energy of about $20\text{--}25 \text{ kJ mol}^{-1}$ above that of the chair form. The twist-boat and boat forms can interconvert by pseudorotation. Calculations suggest that the twist-boat forms are at the minima of the pseudorotation path and that the boats are the transition states about 4 kJ mol^{-1} above the twist boats. Direct experimental information on the twist-boat form was obtained from nuclear magnetic resonance spectroscopy¹³⁸ and matrix-isolated infrared spectra.^{136,144} In the present work the twist-boat conformation was considered together with the basic stable chair form. Relative energy of the twist-boat form, $1925 \pm 70 \text{ cm}^{-1}$ ($23 \pm 0.8 \text{ kJ mol}^{-1}$), given in Table 4, was obtained from experimental data.^{138,144} This value is in a good agreement with estimations.^{90,139–143} The product of the principal moments of inertia for the chair conformation of cyclohexane (Table 4) was calculated using the structural parameters determined by Bastiansen *et al.*¹²⁹ from electron diffraction investigation.

Vibrational spectra of cyclohexane have been studied extensively.^{92,131–137} Various vibrational assignments have differed slightly from one another. Fundamentals, presented in Table 5, are those (except for $\nu_7\text{--}\nu_{11}$) obtained by Wieberg and Shrake.¹³³ These authors observed in the infrared spectra of vapor and liquid and Raman spectrum of liquid cyclohexane and carried out a normal coordinate analysis. We have preferred the vapor-phase values whenever available. The frequencies $\nu_7\text{--}\nu_9$ of A_{1u} symmetry and ν_{10} , ν_{11} of A_{2g} symmetry, unobserved by Wieberg and Shrake,¹³³ were selected according to experimental data^{135,136} and calculated values.^{131–133}

Ideal gas thermodynamic properties for cyclohexane are given in Table 18. The calculated standard entropy at 298.15 K is 1.5 and $0.9 \text{ J K}^{-1} \text{ mol}^{-1}$ less than the values obtained from the calorimetric measurements of Aston *et al.*²⁶² and Beckett *et al.*,¹⁴⁵ respectively (Table 32) and the calculated C_p° values are $0.2\text{--}2.7 \text{ J K}^{-1} \text{ mol}^{-1}$ less than the experimental data of Spitzer and Pitzer²⁵⁸ (Table 33). We suppose these discrepancies are within experimental uncertainties. The molecular constants of cyclohexane are rather reliable and we cannot find any reason for their modification. (Some different vibrational assignments for cyclohexane were considered in this work and for all of them the thermal functions are less than experimental values.) Beckett *et al.*¹⁴⁵ have achieved the agreement with the experimental value of S° (298.15 K) by fitting the molecular constants unknown at the time. The largest discrepancies with thermal

Table 18. Ideal gas thermodynamic properties for cyclohexane

T	C_p°	$-(G^\circ - H_0^\circ)/T$	S°	$H^\circ - H_0^\circ$	$\log K_f$
K	$\text{J K}^{-1} \text{ mol}^{-1}$			kJ mol^{-1}	
100	42.590	190.546	226.103	3.556	31.451
200	69.053	217.933	263.276	9.069	4.423
298.15	105.341	238.431	297.276	17.544	-5.590
300	106.108	238.796	297.930	17.740	-5.724
400	148.638	258.084	334.266	30.473	-11.324
500	188.678	277.066	371.822	47.378	-14.943
600	223.376	295.995	409.374	68.027	-17.489
700	252.620	314.827	446.071	91.870	-19.373
800	277.052	333.459	481.448	118.391	-20.819
900	297.419	351.797	515.292	147.145	-21.958
1000	314.416	369.772	547.535	177.763	-22.873
1100	328.656	387.337	578.189	209.937	-23.620
1200	340.645	404.466	607.315	243.419	-24.239
1300	350.793	421.143	634.993	278.005	-24.757
1400	359.439	437.366	661.314	313.528	-25.195
1500	366.845	453.136	686.372	349.851	-25.568

functions calculated by Beckett *et al.*¹⁴⁵ amount to $4.4 \text{ J K}^{-1} \text{ mol}^{-1}$ for S° (1500 K) and $3.9 \text{ J K}^{-1} \text{ mol}^{-1}$ for C_p° (1500 K). The thermal functions of Chang *et al.*²⁴ are unreliable, because the molecular parameters were obtained from molecular mechanics calculations.

11. Cyclohexene

Half-chair form (C_2 symmetry) has been found to be the most stable conformation of cyclohexene by electron diffraction measurements.^{146–148} Similar conclusions were reached by microwave techniques,^{149,150} infrared and Raman studies,^{151–153} nuclear magnetic resonance spectroscopy,¹⁵⁴ as well as *ab initio*^{155,257} and force-field^{156–160} calculations. To explain the difference between calculated and experimental values of entropy and heat capacity, Beckett *et al.*¹⁰⁹ have suggested that cyclohexene exists as a mixture of half-chair and half-boat (C_s symmetry) conformations where the former is the predominant form and is more stable than half-boat by 11.3 kJ mol^{-1} . However, the more recent experimental^{152,154} and force-field^{156–160} results have shown that the half-boat form is approximately $20\text{--}30 \text{ kJ mol}^{-1}$ higher in energy than the half-chair. According to most of the force-field calculations^{157–160} the half-boat is a transition state (an energy maximum). A metastable half-boat form (an energy minimum which lies slightly lower than the potential maximum), on the other hand, is favored by analysis of the twisting mode region of the Raman spectrum.¹⁵² Since the energy of half-boat conformation is rather high in both cases, we have not considered this form in our calculations.

The product of the principal moments of inertia for the half-chair conformation of cyclohexene (Table 4) was calculated using the rotational constants of Scharpen *et al.*¹⁴⁹ The adopted rotational constants are in excellent agreement with those obtained in another microwave study.¹⁵⁰ From microwave data,^{149,150} the molecular structure of cyclohexene was estimated assuming a number of structural parameters. All in all this structure agrees with those obtained from electron diffraction studies^{147,148} and *ab initio* calculations.^{155,257}

Neto *et al.*¹⁵¹ reported vibrational assignment for cyclo-

Table 19. Ideal gas thermodynamic properties for cyclohexene

T	C_p°	$-(G^\circ-H_0^\circ)/T$	S°	$H^\circ-H_0^\circ$	$\log K_f$
K	$J K^{-1} mol^{-1}$			$kJ mol^{-1}$	
100	43.061	204.644	240.867	3.622	-21.844
200	67.349	232.282	277.521	9.048	-18.943
298.15	101.462	252.593	310.518	17.270	-18.874
300	102.161	252.952	311.148	17.459	-18.879
400	139.702	271.804	345.721	29.567	-19.262
500	173.269	290.080	380.600	45.260	-19.700
600	201.437	308.028	414.759	64.039	-20.099
700	224.915	325.641	447.628	85.391	-20.441
800	244.650	342.865	478.986	108.897	-20.728
900	261.377	359.660	508.794	134.221	-20.967
1000	275.634	376.000	537.091	161.090	-21.166
1100	287.829	391.876	563.948	189.279	-21.331
1200	298.291	407.288	589.453	218.598	-21.470
1300	307.292	422.241	613.693	248.889	-21.585
1400	315.058	436.746	636.757	280.016	-21.680
1500	321.782	450.818	658.729	311.865	-21.760

hexene from the the infrared measurements in the gaseous and liquid states and the Raman spectrum in the liquid state. They carried out the normal coordinate analysis to confirm their assignment. We have adopted their frequencies (with the exception of $\nu_{18}-\nu_{22}$ and $\nu_{39}-\nu_{42}$ modes) preferring the vapor-phase values whenever available. Low-frequency out-of-plane skeletal bending and twisting fundamentals $\nu_{18}-\nu_{22}$ and $\nu_{39}-\nu_{42}$ reported by Smithson and Wieser¹⁵³ from the infrared spectrum of cyclohexene in the vapor phase were used in this work.

Ideal gas thermodynamic properties for cyclohexene are given in Table 19. The calculated value of S° (298.15 K) coincides with the calorimetric entropy¹⁰⁹ (Table 32), but the calculated value of C_p° (400 K) is $5.3 J K^{-1} mol^{-1}$ less than the experimental value of Beckett *et al.*¹⁰⁹ (Table 33). In order to achieve the agreement between experimental and calculated values of C_p° , Beckett *et al.*¹⁰⁹ suggested the existence of the half-boat form with relative energy of $11.3 kJ mol^{-1}$. However, the more recent results have shown that this suggestion is not correct (see above). We think that the accuracy of the experimental value of C_p° is not sufficiently high to discuss the disagreement between the experimental and calculated C_p° values. The largest discrepancies between thermal functions calculated by Beckett *et al.*¹⁰⁹ and given in Table 19 are equal to $7 J K^{-1} mol^{-1}$ for S° (1500 K) and $5.7 J K^{-1} mol^{-1}$ for C_p° (500 K) and are due to the difference in adopted molecular constants.

12. 1,3-Cyclohexadiene

The nonplanar twisted conformation of C_2 symmetry was found for 1,3-cyclohexadiene from electron diffraction studies¹⁶¹⁻¹⁶³ and semiempirical^{164,165} and *ab initio*^{155,166} calculations. The rotational constants of 1,3-cyclohexadiene were obtained by microwave studies.^{167,168} Assuming the bond distances and bond angles from related compounds, Butcher¹⁶⁷ estimated the dihedral angle in agreement with electron diffraction values¹⁶¹⁻¹⁶³ and showed that the moments of inertia calculated from the proposed molecular structure of C_2 symmetry are close to those obtained from rotational constants. The product of the principal moments of inertia, given in Table 4, was calculated using the rota-

Table 20. Ideal gas thermodynamic properties for 1,3-cyclohexadiene

T	C_p°	$-(G^\circ-H_0^\circ)/T$	S°	$H^\circ-H_0^\circ$	$\log K_f$
K	$J K^{-1} mol^{-1}$			$kJ mol^{-1}$	
100	40.779	202.970	238.278	3.531	-72.277
200	62.742	229.570	272.610	8.608	-41.357
298.15	94.165	248.782	303.304	16.256	-31.859
300	94.802	249.120	303.889	16.431	-31.745
400	128.546	266.795	335.842	27.619	-27.258
500	157.995	283.808	367.789	41.991	-24.727
600	182.235	300.407	398.812	59.043	-23.122
700	202.143	316.600	428.447	78.293	-22.019
800	218.699	332.353	456.553	99.360	-21.216
900	232.630	347.644	483.139	121.946	-20.605
1000	244.450	362.462	508.278	145.815	-20.122
1100	254.531	376.809	532.062	170.778	-19.731
1200	263.164	390.694	554.589	196.674	-19.407
1300	270.585	404.130	575.953	223.370	-19.131
1400	276.984	417.134	596.246	250.757	-18.894
1500	282.523	429.724	615.549	278.738	-18.687

tional constants of Butcher.¹⁶⁷ These constants are consistent with those of Luss and Harmony.¹⁶⁸

Carreira *et al.*¹⁶⁹ investigated the Raman spectrum of gaseous 1,3-cyclohexadiene and obtained the ring-twisting potential function from which the barrier to inversion of the ring through its planar configuration was determined to be $1099 \pm 50 cm^{-1}$. We do not consider the inversion motion in 1,3-cyclohexadiene because the barrier to inversion is sufficiently high.

Di Lauro *et al.*⁸⁵ reported the complete vibrational assignment for 1,3-cyclohexadiene from the infrared measurements in gaseous and liquid states and the Raman study in the liquid state as well as from a normal coordinate analysis. Some of these frequencies were reinterpreted by Warshel and Karplus¹⁶⁴ based on the force-field calculation. Carreira *et al.*¹⁶⁹ investigated the Raman spectrum of gaseous 1,3-cyclohexadiene, but no detailed assignment was proposed. Fundamentals, presented in Table 5, are those obtained by Di Lauro *et al.*⁸⁵ We prefer the vapor state values whenever available.

Ideal gas thermodynamic properties for 1,3-cyclohexadiene, given in Table 20, are reported for the first time. No experimental data are available for comparison.

13. 1,4-Cyclohexadiene

Over the years, conclusions regarding the equilibrium configuration of 1,4-cyclohexadiene were contradictory. An early infrared and Raman investigation¹⁷⁰ was interpreted on the basis of D_{2h} (planar) symmetry. More recently, Stidham¹⁷¹ has reinvestigated the vibrational spectra of 1,4-cyclohexadiene and supported the planar configuration, although he could not rule out a boat form (C_{2v} symmetry) that inverts through the planar form. Although subsequent studies involving electron diffraction,¹⁷² nuclear magnetic resonance,¹⁷³ and semiempirical calculations^{157,174} also supported a planar conformation, there were conflicting reports involving electron diffraction,¹⁶³ nuclear magnetic resonance,¹⁷⁵ and semiempirical calculation,¹⁷⁶ which supported a boat conformation. The electron diffraction study of Oberhammer and Bauer¹⁶³ constituted the major evidence

against a planar structure, but their conclusion was based on an erroneous interpretation of the experimental data.¹⁵⁵ The planarity of the 1,4-cyclohexadiene ring was conclusively determined by the combination of the far-infrared and Raman data for the ring-puckering mode.^{169,177} The conformation of 1,4-cyclohexadiene must closely correspond to the parabolic potential energy curve involving wide amplitude vibrations around the planar form, as described from infrared and Raman spectra in the low-frequency region^{169,177} and supported by the semiempirical^{174,178,179} and *ab initio*^{155,166,178} calculations.

The molecular structure of 1,4-cyclohexadiene was determined by electron diffraction measurements.^{163,172} Although Dallinga and Toneman¹⁷² adopted the planar structure, for a variety of reasons^{162,163} their data seem to be unreliable. Oberhammer and Bauer¹⁶³ have interpreted their data for a bent structure with a dihedral angle of 159.3° because no large amplitude vibrations have been taken into account. These authors have also tested a planar structure which showed worse agreement with experimental data. For lack of more reliable data, the structural parameters of the planar model obtained by Oberhammer and Bauer¹⁶³ were used for calculation of the product of the principal moments of inertia given in Table 4.

Stidham¹⁷¹ reported vibrational assignment for 1,4-cyclohexadiene on the basis of infrared spectra of gaseous and liquid phases and Raman spectrum of liquid phase. Gebhardt and Cyvin¹⁸⁰ reproduced completely Stidham's frequency values by the force constants refinement. Ermer and Lifson¹¹³ reassigned the values of some frequencies arising from the force-field calculation, their results being consistent with unpublished data of Neto *et al.*, reported in Ref. 113. Most of the fundamentals, presented in Table 5, are those of Stidham.¹⁷¹ We preferred the vapor-phase values whenever available. Unobserved vibrational frequencies (ν_8 – ν_{11} , ν_{15} , ν_{19}) and the values of ν_{25} , ν_{34} , and ν_{36} were taken from the assignment of Ermer and Lifson.¹¹³ The ring-puckering frequency, ν_{31} (B_{2u}), is that observed by Laane and Lord¹⁷⁷ from the far-infrared spectrum.

No experimental data on the enthalpy of formation of 1,4-cyclohexadiene are available. The value presented in Ta-

ble 4 is based on molecular mechanics calculations^{143,178} and agrees well with the value recommended by Shaw *et al.*¹⁸¹

Ideal gas thermodynamic properties for 1,4-cyclohexadiene, given in Table 21, are reported for the first time. No experimental values of S° and C_p° are available for comparison.

14. Cycloheptane

Extensive conformational energy calculations^{90,182–184,263} have been done on cycloheptane and these have led to an identification of four basic conformations: chair (C_s symmetry), boat (C_s symmetry), twist-chair (C_2 symmetry), and twist-boat (C_2 symmetry). On the energy profile during pseudorotation, obtained in these studies, the twist-chair and twist-boat forms are located at the minima and the chair and boat forms at the maxima. The twist-boat form is approximately 10–15 kJ mol^{−1} higher in energy than the most stable twist-chair conformation. Vibrational spectra of cycloheptane¹⁸⁵ were interpreted on the basis of one rigid conformation, viz., the twist-chair form. However, the best agreement with experimental electron diffraction data¹⁸⁶ was obtained for a mixture of two minimum-energy conformers: the most stable twist-chair form and the chair form with energy higher by 3.8 kJ mol^{−1}. We have performed calculations of S° (298.15 K) for models of the different conformational composition and made comparison of the experimental entropy with the calculated values.¹⁸⁷ An inspection of these values shows that an excellent agreement exists for the twist-chair conformer alone or for the mixture of twist-chair and twist-boat, while the mixture of twist-chair and chair reported in electron diffraction study¹⁸⁶ has a noticeable discrepancy. From these considerations only two stable conformations of cycloheptane, twist-chair, and twist-boat, were taken into account in the present work. Relative energy of the twist-boat form, 1200 cm^{−1} (14 kJ mol^{−1}), given in Table 6, was obtained by Bocian and Strauss¹⁸⁴ from the strain-energy calculations.

The gaseous electron diffraction structural data of Dillen and Geise¹⁸⁶ for the twist-chair form were used to calculate the product of the principal moments of inertia (Table 6). These structural parameters are consistent with those obtained by the force-field calculations.^{90,182–184,263}

The vibrational frequencies of cycloheptane used in our calculations (Table 7) were taken from a study by Bocian and Strauss.¹⁸⁵ These authors have examined in detail the vapor- and liquid-phase infrared spectra, vapor-phase far-infrared spectrum, and liquid-phase Raman spectrum; special attention has been given to the low-frequency region. In addition, a normal coordinate analysis was performed to confirm the vibrational assignment and to estimate the unobserved frequencies.

Ideal gas thermodynamic properties for cycloheptane are listed in Table 22. The calculated value of S° (298.15 K), as shown in Table 32, coincides with the calorimetric entropy¹⁸⁸ and is close to that calculated by Chang *et al.*²⁴ However, the discrepancies between the thermal function reported by Chang *et al.*²⁴ and listed in Table 22 increase to 2.3 J K^{−1} mol^{−1} for S° and 7.3 J K^{−1} mol^{−1} for C_p° as the

Table 21. Ideal gas thermodynamic properties for 1,4-cyclohexadiene

T	C_p°	$-(S^\circ - H_0^\circ)/T$	S°	$H^\circ - H_0^\circ$	$\log K_f$
K	J K ^{−1} mol ^{−1}			kJ mol ^{−1}	
100	42.617	193.232	230.380	3.715	−74.038
200	65.252	220.991	265.437	8.889	−42.432
298.15	94.051	240.689	296.225	16.558	−32.702
300	94.675	241.033	296.609	16.733	−32.585
400	127.751	258.946	328.630	27.873	−27.982
500	156.822	276.065	360.352	42.144	−25.382
600	180.966	292.696	391.149	59.072	−23.733
700	200.940	308.881	420.593	78.198	−22.601
800	217.630	324.608	448.546	99.151	−21.777
900	231.710	339.863	475.015	121.637	−21.150
1000	243.669	354.662	500.064	145.422	−20.656
1100	253.871	368.950	523.779	170.312	−20.256
1200	262.606	382.797	546.253	196.147	−19.924
1300	270.111	396.198	567.577	222.702	−19.642
1400	276.580	409.169	587.837	250.135	−19.400
1500	282.177	421.728	607.114	278.080	−19.188

Table 22. Ideal gas thermodynamic properties for cycloheptane

T	C_p°	$-(G^\circ-H_0^\circ)/T$	S°	$H^\circ-H_0^\circ$	$\log K_f$
K	J K ⁻¹ mol ⁻¹			kJ mol ⁻¹	
100	52.126	213.735	253.285	3.955	23.380
200	86.837	245.427	299.578	10.830	-1.733
298.15	132.014	270.263	342.332	21.487	-11.250
300	132.944	270.710	343.151	21.732	-11.378
400	183.225	294.436	388.326	37.556	-16.774
500	228.473	317.804	434.204	58.200	-20.291
600	266.447	340.982	479.323	83.005	-22.775
700	298.025	363.875	522.841	111.274	-24.619
800	324.491	386.365	564.415	142.440	-26.037
900	346.862	408.364	603.962	176.038	-27.155
1000	365.887	429.818	641.519	211.701	-28.054
1100	382.126	450.698	677.174	249.123	-28.789
1200	396.036	470.994	711.035	288.049	-29.397
1300	407.986	490.707	743.219	328.265	-29.906
1400	418.284	509.846	773.840	369.591	-30.335
1500	427.191	528.426	803.010	411.875	-30.700

temperature increases to 500 K. These discrepancies are due to a difference in molecular constants used in calculations (Chang *et al.*²⁴ evaluated the molecular constants from the molecular mechanic calculations).

15. Cycloheptene

The conformational properties of cycloheptene have been studied experimentally by vibrational spectroscopy¹⁸⁹ and by nuclear magnetic resonance techniques.¹⁹⁰ Observed spectra were, in both cases, interpreted in terms of a C_s symmetric chair form being the most stable conformation. Predictions of the conformations of the molecule have been made from force-field calculations. All recent calculations^{143,157,191-193} imply that the chair conformation is only slightly more stable than the C_2 twist conformation (the C_2 twist form is 2-7 kJ mol⁻¹ higher in energy). Another local minimum for the boat form (C_s symmetry) was supported by some workers,^{157,191} but this proposal was not confirmed by more reliable calculations.^{143,193} Two stable conformations of cycloheptene were considered in this work: the most stable chair form and the C_2 twist form with the energy higher by 500 cm⁻¹ (~6 kJ mol⁻¹). The relative energy of the C_2 form was estimated by averaging the values of the above parameters obtained in different force-field calculations.^{143,157,191-193}

There are no direct experimental data on the molecular structure of cycloheptene. The product of the principal moments of inertia for the chair conformation of cycloheptene (Table 6) was calculated using the structural parameters obtained by Ermer and Lifson¹⁹³ from the force-field study. Adopted parameters are in good agreement with those calculated in other force-field studies^{143,192} and are close to structural parameters in related compounds for which experimental data are available.

Neto *et al.*¹⁸⁹ have investigated the infrared spectra of cycloheptene in all three physical states and the Raman spectrum in the liquid state and carried out a normal coordinate analysis to confirm their vibrational assignment. The fundamentals, presented in Table 7, are those for liquid cycloheptene reported by Neto *et al.*¹⁸⁹ Vapor-liquid frequen-

Table 23. Ideal gas thermodynamic properties for cycloheptene

T	C_p°	$-(G^\circ-H_0^\circ)/T$	S°	$H^\circ-H_0^\circ$	$\log K_f$
K	J K ⁻¹ mol ⁻¹			kJ mol ⁻¹	
100	47.403	211.327	248.470	3.714	-26.132
200	82.290	240.912	291.566	10.131	-23.762
298.15	124.071	264.230	332.010	20.209	-24.030
300	124.903	264.650	332.780	20.439	-24.041
400	169.272	286.927	374.849	35.169	-24.661
500	208.626	308.734	416.987	54.126	-25.272
600	242.095	330.221	458.092	76.722	-25.802
700	269.902	351.369	497.562	102.363	-26.243
800	293.326	371.979	535.176	130.557	-26.607
900	313.209	392.114	570.904	160.911	-26.907
1000	330.170	411.703	604.805	193.102	-27.154
1100	344.086	430.733	636.972	226.863	-27.358
1200	357.141	449.203	667.511	261.970	-27.526
1300	367.856	467.121	696.531	298.233	-27.665
1400	377.107	484.502	724.139	335.493	-27.780
1500	385.116	501.361	750.436	373.613	-27.874

cy shifts were taken into account in estimation of the inaccuracies in the vibrational frequencies.

Ideal gas thermodynamic properties for cycloheptene, given in Table 23, are reported for the first time. No experimental values of S° and C_p° are available for comparison.

16. 1,3-Cycloheptadiene

A number of experimental and theoretical studies have been reported on this molecule with contradictory conclusions about the most stable conformation. Electron diffraction data^{194,195} indicated that the carbon skeleton is planar except for the one carbon atom (C_s form). A nuclear magnetic resonance study,¹⁹⁶ on the other hand, was consistent with a C_2 twist form. Schrader and Ansmann,¹⁹⁷ from infrared data on stretching vibrations of the double bonds, inferred that 1,3-cycloheptadiene exists in a C_s form. The vibrational spectrum of 1,3-cycloheptadiene was also interpreted in terms of a C_s form.⁸⁶ Avirah *et al.*¹⁹⁸ investigated the microwave spectrum of 1,3-cycloheptadiene. In their opinion, the agreement between the observed and calculated rotational constants based on the electron diffraction studies argues strongly for the C_s form. In addition to experimental results, molecular mechanics calculations on 1,3-cycloheptadiene have also been reported. In early work, Favini *et al.*^{199,200} found for the most stable conformation a C_2 form, which was replaced by the C_s structure in more recent studies.^{201,264} The existence of three forms which are relatively close in energy was found in the later work of Favini *et al.*²⁶⁴ The C_s form was found to be 7.6 kJ mol⁻¹ more stable than the form with C_2 symmetry and 3.4 kJ mol⁻¹ more stable than the intermediate nonsymmetrical form (C_1). Allinger and Sprague²⁰² predicted that the molecule undergoes an almost unhindered pseudorotation between a C_s form and an alternate nonsymmetrical form (C_1). This pseudorotating structure is in equilibrium with the C_2 form separated by a barrier of about 4 kJ mol⁻¹. According to *ab initio* calculations of Saebø and Boggs,²⁰³ there are three energy-minimum conformations of 1,3-cycloheptadiene, viz., the most stable C_s form and two C_2 forms with energies higher than that of C_s by 10.5 and 60.0 kJ mol⁻¹, respective-

Table 24. Ideal gas thermodynamic properties for 1,3-cycloheptadiene

T	C_p°	$-(G^\circ-H_0^\circ)/T$	S°	$H^\circ-H_0^\circ$	$\log K_f$
K	$J K^{-1} mol^{-1}$			$kJ mol^{-1}$	
100	52.524	211.542	251.010	3.942	-72.100
200	80.878	242.988	295.802	10.503	-43.745
298.15	117.672	266.801	334.710	20.247	-35.219
300	118.417	267.222	335.440	20.466	-35.117
400	158.028	289.219	374.993	34.309	-31.185
500	192.987	310.302	414.121	51.909	-29.017
600	222.108	330.781	451.964	72.710	-27.069
700	246.262	350.694	488.073	96.166	-26.759
800	266.498	370.026	522.316	121.832	-26.105
900	283.613	386.764	554.721	149.361	-25.608
1000	298.102	400.907	585.377	178.470	-25.218
1100	310.636	424.462	614.590	208.927	-24.902
1200	321.317	441.446	641.894	240.538	-24.638
1300	330.507	457.977	667.980	273.141	-24.413
1400	338.439	473.778	692.776	306.596	-24.218
1500	345.305	489.170	716.366	340.793	-24.045

Table 25. Ideal gas thermodynamic properties for 1,3,5-cycloheptatriene

T	C_p°	$-(G^\circ-H_0^\circ)/T$	S°	$H^\circ-H_0^\circ$	$\log K_f$
K	$J K^{-1} mol^{-1}$			$kJ mol^{-1}$	
100	42.240	208.479	243.860	3.538	-111.711
200	71.041	235.824	281.457	9.126	-60.943
298.15	106.249	256.568	316.250	17.794	-44.885
300	106.936	256.938	316.909	17.991	-44.688
400	142.799	276.399	352.668	30.508	-36.865
500	173.733	295.198	387.962	46.382	-32.326
600	199.178	313.514	421.966	65.070	-29.380
700	220.092	331.337	454.889	86.067	-27.319
800	237.493	348.634	484.849	108.972	-25.797
900	252.138	365.387	513.691	133.473	-24.628
1000	264.562	381.591	540.917	159.325	-23.700
1100	275.163	397.256	566.642	186.325	-22.945
1200	284.244	412.395	590.984	214.307	-22.318
1300	292.052	427.027	614.052	243.131	-21.787
1400	298.790	441.175	635.948	272.682	-21.332
1500	304.625	454.860	656.766	302.859	-20.936

ly. Saebø and Boggs²⁰³ consider their results to be close to that reported by Allinger and Sprague,²⁰² and believe the existence of a mixture of C_s and C_2 forms to explain the contradiction in the interpretation of electron diffraction^{194,195} and nuclear magnetic resonance¹⁹⁶ data. Three stable conformations of 1,3-cycloheptadiene were taken into account in this work: the most stable C_s form, as well as C_1 and C_2 forms with energies higher than the C_s by 200 cm^{-1} ($\sim 2.4 kJ mol^{-1}$) and 500 cm^{-1} ($\sim 6 kJ mol^{-1}$), respectively. The adopted relative energies of the C_1 and C_2 forms are intermediate between the results of Allinger and Sprague²⁰² and of Favini *et al.*,²⁶⁴ and nearer to the latter.

The product of the principal moments of inertia, given in Table 6, was calculated using the rotational constants of Avirah *et al.*¹⁹⁸ This value agrees with those calculated from electron diffraction data.^{194,195}

Ansmann and Schrader⁸⁶ investigated the infrared and Raman spectra of liquid 1,3-cycloheptadiene and presented the vibrational assignment on the basis of normal coordinate analysis. Their frequencies are listed in Table 7. Vapor-liquid frequency shifts were taken into account in estimating the inaccuracies in the adopted fundamentals.

Ideal gas thermodynamic properties for 1,3-cycloheptadiene, given in Table 24, are reported for the first time. No experimental data are available for comparisons.

17. 1,3,5-Cycloheptatriene

Cycloheptatriene has a nonplanar equilibrium conformation (boat form of C_s symmetry), as shown by electron diffraction,²⁰⁴ microwave,²⁰⁵ infrared and Raman,²⁰⁶ and nuclear magnetic resonance^{207,208} measurements as well as by force-field^{165,209-211} and *ab initio*^{203,212} calculations. A planar structure (C_{2v} symmetry) was found to be about 25 $kJ mol^{-1}$ higher in energy and to be a transition state on the energy surface, probably for interconversion of boat forms.^{203,207,208,212}

Conflicting results concerning the degree of nonplanarity of the cycloheptatriene ring have been published. Traetteberg²⁰⁴ determined the total molecular structure by electron

diffraction and found the values of two dihedral angles to be $\alpha = 40.5^\circ$ and $\beta = 36.5^\circ$. Butcher²⁰⁵ determined the rotational constants from microwave study of normal isotopic species of cycloheptatriene and estimated $\alpha = 29.5^\circ$ and $\beta = 50^\circ$, assuming bond lengths and angles, the values of dihedral angles being somewhat insensitive to the choice of structural parameters. Butcher's estimates of dihedral angles agree pretty well with theoretical results.^{165,203,209-212} The product of the principal moments of inertia, given in Table 6, was calculated using the rotational constants of Butcher.²⁰⁵


Evans and Lord²¹³ interpreted the infrared and Raman spectra of cycloheptatriene assuming a planar structure of C_{2v} symmetry. Later, La Lau and De Ruyter²⁰⁶ investigated the Raman spectrum of liquid cycloheptatriene and proposed the reassignment of Evans and Lord's data²¹³ for C_s symmetry. Paulick *et al.*²¹⁰ presented a new interpretation of vibrational spectra^{206,213} on the basis of the force-field calculation. Vibrational frequencies, listed in Table 7, were obtained by means of critical comparison of experimental^{206,213} and calculated²¹⁰ frequencies of cycloheptatriene with those for related compounds. The adopted vibrational assignment gives a better agreement between experimental and calculated values of S° (298.15 K) than those of La Lau and Ruyter²⁰⁶ or Paulick *et al.*²¹⁰

Ideal gas thermodynamic properties for 1,3,5-cycloheptatriene are presented in Table 25. The agreement between the calorimetric entropy¹⁸⁸ and our calculated value is satisfactory within the experimental uncertainty (Table 32). The value of S° (298.15 K) calculated by Evans and Lord²¹³ for a planar structure is about 2 $J K^{-1} mol^{-1}$ larger than that calculated in the present work.

18. Cyclooctane

Some families of cyclooctane conformations were identified by conformational energy calculations.^{91,139,182,214-216,263} The most stable conformation of cyclooctane is the boat-chair (BC) of C_s symmetry and according to most of calculations two families of conforma-


forms are close in energy:



[TBC: twist-boat-chair (C_2), crown (D_{4d}), CC: chair-chair (C_{2v}), TCC: twist-chair-chair (D_2)]. The energy differences among these low-energy conformers are estimated to be from 0.8 to 10 kJ mol⁻¹. It should be emphasized that force-field calculations give inconsistent results on the relative energies of conformers (except that for the BC form) and on the location of conformers on the potential energy surface (local minimum or saddle point). The analysis of torsional energy surfaces presented by Ivanov and Ōsawa²⁶³ shows that even slightly different force fields exhibit significantly different pictures for the dynamic behavior of the cyclooctane molecule.

Experimental investigations have led to a variety of conclusions. Nuclear magnetic resonance experiments^{217,218} provide evidence for the boat-chair and indicate the presence of a small amount of another form, probably of the crown family. The gas-phase electron diffraction data of Almenningen *et al.*²¹⁹ were not compatible with the assumption of any single geometry, but rather with a mixture of several conformations. However, good agreement between experimental and theoretical electron diffraction data was obtained for the boat-chair conformer in a recent reinvestigation by Dorofeeva *et al.*²²⁰ An early vibrational study²²¹ arrived at the conclusion of a tub form for cyclooctane, which is now known to be incorrect. Pakes *et al.*²²² examined the vibrational spectra of cyclooctane, giving special attention to the low-frequency region. Agreement between calculated and observed vibrational frequencies was obtained only for the boat-chair conformer.

Although several low-energy conformers of cyclooctane were predicted from theoretical calculations, reliable experimental evidence for their existence has not been obtained. We have performed the calculations of the standard entropy of gaseous cyclooctane for models of different conformational composition¹⁸⁷ and have shown the value of S° (298.15 K) for the single boat-chair form to be 14.7 J K⁻¹ mol⁻¹ less than the experimental value. It should be noted that the set of conformers calculated by Pakes *et al.*²¹⁶ is not sufficient to reproduce the experimental entropy. A good agreement between experimental and calculated entropy values was achieved by assuming that in two above families of the cyclooctane conformers the intermediate non-symmetrical (C_1) energy-minimum forms exist:



(The existence of C_1 forms is assumed by analogy with strain-energy calculation results 1,4-cyclooctadiene.²²³ The crown form is excluded from consideration because according to the calculation of Pakes *et al.*,²¹⁶ it is a saddle point. See also Ref. 187.) To calculate the thermal functions of cyclooctane, the six conformers of this molecule were taken into account (Table 8): the most stable boat-chair conformer and the five forms (TCC, C_1 , CC, C_1 , and TBC) with energy higher by 275 cm⁻¹ (3.3 kJ mol⁻¹). Average relative

Table 26. Ideal gas thermodynamic properties for cyclooctane

T	C_p°	$-(S^\circ - H_0^\circ)/T$	S°	$H^\circ - H_0^\circ$	$\log K_f$
K	J K ⁻¹ mol ⁻¹			kJ mol ⁻¹	
100	64.769	219.700	262.528	4.283	19.843
200	99.987	255.803	318.710	12.581	-6.067
298.15	146.194	284.435	366.725	24.535	-16.010
300	147.189	284.945	367.633	24.806	-16.145
400	202.155	311.819	417.501	42.273	-21.857
500	253.069	338.025	468.208	65.092	-25.624
600	296.638	363.914	518.311	92.636	-28.307
700	335.376	389.461	566.877	124.191	-30.314
800	366.461	414.572	613.479	159.126	-31.865
900	390.901	439.165	657.974	196.929	-33.094
1000	413.473	463.182	700.359	237.177	-34.086
1100	432.791	486.592	740.697	279.515	-34.898
1200	449.361	509.381	779.084	323.644	-35.573
1300	463.605	531.544	815.629	369.310	-36.137
1400	475.896	553.089	850.446	416.300	-36.614
1500	486.229	574.020	883.651	464.434	-37.019

energy of the five conformers was estimated from the energies of the TCC, CC, and TBC forms calculated by Pakes *et al.*²¹⁶

The electron diffraction structural parameters obtained by Dorofeeva *et al.*²²⁰ were used to calculate the product of the principal moments of inertia (Table 8). These parameters are consistent with those obtained by force-field calculations.^{216,220}

The vibrational frequencies of cyclooctane used in our calculations (Table 9) were taken from the study of Pakes *et al.*²²² These authors have examined in detail the vapor and liquid-phase infrared spectra, vapor-phase far-infrared spectrum, and liquid-phase Raman spectrum, and carried out a normal coordinate analysis to confirm their vibrational assignment.

Ideal gas thermodynamic properties for cyclooctane are given in Table 26. The calculated value of S° (298.15 K), as shown in Table 32, coincides with the calorimetric entropy.¹⁸⁸ Chang *et al.*²⁴ have calculated the thermal functions of the D_2 cyclooctane conformation using the molecular constants obtained from molecular mechanics calculations. Their value of S° (298.15 K) is much less than the value obtained in this work (Table 32). However, this discrepancy will be markedly decreased if the C_s symmetry is employed. Unfortunately, Chang *et al.*²⁴ did not publish the molecular constants they used and the reliability of their functions cannot be estimated.

19. *trans*-Cyclooctene

Trans-cyclooctene, which is the most stable cyclooctene isomer, can exist as the twist-chair-chair (C_2 symmetry) or chair (C_2 symmetry) form. The electron diffraction study of Gavin and Wang²²⁴ and early force-field calculation of Buemi *et al.*²²⁵ favored a chair conformation. Traetteberg²²⁶ has interpreted the electron diffraction data of *trans*-cyclooctene as indicative of a twist-chair-chair conformation. The same form was also predicted by the force-field calculations.^{157,160,193,201} As the reliability of electron diffraction data of Gavin and Wang²²⁴ raise some doubts,^{193,226} the most stable conformation of *trans*-cyclooctene should be twist-chair-chair. A less stable chair conformation is also

Table 27. Ideal gas thermodynamic properties for *trans*-cyclooctene

T	C_p°	$-(G^\circ-H_0^\circ)/T$	S°	$H^\circ-H_0^\circ$	$\log K_f$
K	$J K^{-1} mol^{-1}$			$kJ mol^{-1}$	
100	52.282	217.505	256.296	3.879	-24.007
200	95.703	249.271	304.923	11.130	-25.376
298.15	142.912	275.180	351.225	22.673	-27.063
300	143.904	275.652	352.112	22.938	-27.093
400	196.926	300.831	400.841	40.004	-28.514
500	244.054	325.755	449.991	62.118	-29.642
600	285.393	350.491	496.077	88.552	-30.531
700	316.026	374.900	544.290	118.373	-31.236
800	343.353	398.849	588.326	151.581	-31.799
900	366.444	422.246	630.137	187.102	-32.254
1000	386.079	445.035	669.790	224.755	-32.622
1100	402.844	467.192	707.394	264.223	-32.924
1200	417.206	486.708	743.078	305.244	-33.172
1300	429.546	509.590	776.972	347.597	-33.377
1400	440.784	529.849	809.203	391.096	-33.545
1500	449.384	549.504	839.894	435.586	-33.685

considered in the present work (Table 8). The relative energy of this form, 1000 cm^{-1} (12 kJ mol^{-1}), was estimated from the force-field calculations.^{157,193,201}

The product of the principal moments of inertia for *trans*-cyclooctene, given in Table 8, was calculated from the electron diffraction structural parameters of Traetteberg.²²⁶ These parameters are consistent with those obtained by force-field calculations.^{160,193,226,227}

Until now no vibrational spectra of *trans*-cyclooctene have been reported. In the present work the fundamental vibrations were estimated by normal coordinate calculations. The similarity of force fields for cyclohexene,¹⁵¹ cycloheptene,¹⁸⁹ and related compounds allows a decision to be made concerning the transferability of the force fields for cycloalkenes. Thus, the force field used was transferred from cycloheptene.¹⁸⁹ For normal coordinate calculations we used the NCA program written by Novikov.⁸³ The fundamental frequencies, given in Table 9, were selected on the basis of our calculations and the comparison with fundamentals of cyclohexene and cycloheptene; their uncertainties are estimated to be $20\text{--}50 \text{ cm}^{-1}$.

Ideal gas thermodynamic properties for *trans*-cyclooctene, given in Table 27, are reported for the first time. No experimental data are available for comparisons.

20. *cis,cis*-1,3-Cyclooctadiene

Cis,cis-1,3-cyclooctadiene is the most stable isomer of 1,3-cyclooctadiene. Braude,²²⁸ from ultraviolet spectroscopy, and Schrader and Ansmann,¹⁹⁷ from infrared data on stretching vibrations of the double bonds, deduced that the diene system in *cis,cis*-1,3-cyclooctadiene must be nonplanar with a torsional angle of about $40^\circ\text{--}45^\circ$ and 38° , respectively. Traetteberg²²⁹ has investigated the structure of *cis,cis*-1,3-cyclooctadiene in the gas phase by electron diffraction and has reported that the conformation is irregular twist-boat (C_1 symmetry) with a diene twist angle of 38° . Allinger *et al.*²³⁰ studied *cis,cis*-1,3-cyclooctadiene by the iterative force-field method. The molecule is calculated to be a mixture of two conformations, one of which has a C_2 axis, and one is irregular. The energies of these two conformations are calculated to be very nearly the same. These results are con-

sistent with the noniterative force-field findings.^{199,200} Anet and Yavari²³¹ have investigated *cis,cis*-1,3-cyclooctadiene by dynamic nuclear magnetic resonance spectroscopy and have also reported iterative force-field calculations of the interconversion paths of *cis,cis*-1,3-cyclooctadiene conformations. According to their data, the compound exists as a nearly equal mixture of symmetrical (C_2) twist-boat-chair and unsymmetrical (C_1) twist-boat conformers. This is in agreement with previous calculations.^{199,200,230} The mixture of the two conformers of *cis,cis*-1,3-cyclooctadiene was considered in this work (Table 8), where the twist-boat-chair is the predominant conformation and is more stable than twist-boat by 200 cm^{-1} (2.4 kJ mol^{-1}). (Anet and Yavari²³¹ have estimated this energy as $\sim 2.1 \text{ kJ mol}^{-1}$.)

Electron diffraction structural parameters of the C_1 conformer determined by Traetteberg²²⁹ were used to calculate the product of the principal moments of inertia (Table 8). Allinger *et al.*²³⁰ suppose that this structure must be some kind of average over the two conformations of *cis,cis*-1,3-cyclooctadiene.

Only infrared data on the stretching vibrations of $C=C$ bonds have been reported for 1,3-cyclooctadiene.¹⁹⁷ The vibrational frequencies, given in Table 9, were estimated on the basis of normal coordinate calculations. The similarity of force fields for 1,3-cyclohexadiene,⁸⁵ 1,3-cycloheptadiene,⁸⁶ and related compounds allows a decision to be made concerning the transferability of the force fields for 1,3-cycloalkadienes. Thus, the force constants used were transferred from 1,3-cycloheptadiene.⁸⁶ For normal coordinate calculations we used the NCA program by Novikov.⁸³ The fundamental frequencies, given in Table 9, were selected on the basis of our calculations and the comparison with fundamentals of 1,3-cyclohexadiene and 1,3-cycloheptadiene. The uncertainties of calculated frequencies are estimated to be $20\text{--}50 \text{ cm}^{-1}$.

The enthalpy of formation, presented in Table 8, was estimated by Kozina *et al.*²³² from available experimental data.

Ideal gas thermodynamic properties for *cis,cis*-1,3-cyclooctadiene, given in Table 28, are reported for the first time. No experimental data are available for comparisons.

Table 28. Ideal gas thermodynamic properties for *cis,cis*-1,3-cyclooctadiene

T	C_p°	$-(G^\circ-H_0^\circ)/T$	S°	$H^\circ-H_0^\circ$	$\log K_f$
K	$J K^{-1} mol^{-1}$			$kJ mol^{-1}$	
100	61.276	217.979	260.574	4.263	-71.586
200	95.587	253.045	313.431	12.077	-66.081
298.15	138.825	280.512	359.358	23.508	-38.652
300	139.704	281.001	360.219	23.766	-38.565
400	186.556	306.638	406.895	40.163	-35.260
500	227.963	331.527	453.101	60.887	-33.502
600	262.458	355.375	497.812	85.463	-32.444
700	291.077	376.797	540.486	113.183	-31.747
800	315.664	401.561	580.966	143.524	-31.254
900	335.360	423.643	619.280	176.073	-30.886
1000	352.644	445.036	655.532	210.450	-30.598
1100	367.422	465.746	689.854	246.516	-30.363
1200	380.100	485.789	722.381	283.910	-30.167
1300	391.005	505.186	753.247	322.479	-29.997
1400	400.418	523.961	782.576	362.062	-29.846
1500	408.569	542.140	810.487	402.521	-29.712

21. *cis,cis*-1,5-Cyclooctadiene

It is known from electron diffraction study²³³ and from interpretation of the nuclear magnetic resonance spectra²³⁴ and molecular mechanics calculations,^{143,235,236} that the lowest energy form of the *cis,cis*-1,5-cyclooctadiene is the twist-boat (C_2 symmetry). Ermer²³⁶ has performed the force-field calculations of the interconversion paths of *cis,cis*-1,5-cyclooctadiene and has shown that three conformations correspond to potential energy minima: the most stable twist-boat form and two forms of C_{2h} (chair) and C_s symmetry with energies higher than that of twist-boat by about 17 kJ mol⁻¹. This result agrees well with the interpretation of nuclear magnetic resonance spectra by Anet and Kozerski²³⁴ but less well with Allinger and Sprague's force-field calculation.²³⁵ In the latter work five conformations correspond to energy minima, the relative energy of chair form being ~6 kJ mol⁻¹. No significant evidence for the presence of the chair form was found in the electron diffraction study of Hagen *et al.*,²³³ but amounts less than about 10% in an equilibrium mixture with the twist-boat form cannot be ruled out. On the basis of the above experimental and theoretical data on *cis,cis*-1,5-cyclooctadiene, three stable conformations of this molecule (twist-boat, chair, and C_s form) were considered in the present work (Table 8). The relative energy of chair and C_s forms, 1500 cm⁻¹ (18 kJ mol⁻¹), was adopted according to the force-field calculations of Ermer.²³⁶

The product of the principal moments of inertia for the twist-boat conformer, given in Table 8, was calculated from structural data obtained by electron diffraction.²³³

Infrared and Raman spectra of the liquid and solid 1,5-cyclooctadiene were reported^{237,238} but no vibrational assignments were presented except for $\nu(C=C)$. To estimate a complete set of fundamentals, the normal coordinate calculations were carried out in this work. The similarity of the force fields for cyclohexene,¹⁵¹ cycloheptene,¹⁸⁹ and *trans,trans*-cyclododecatriene²³⁹ permits the transfer of the force constants from these molecules to 1,5-cyclooctadiene. The fundamental frequencies, adopted in Table 9, were selected on the basis of normal coordinate calculations. For

this purpose we used the NCA program by Novikov.⁸³ The uncertainties of calculated frequencies are estimated to be 20–50 cm⁻¹.

Ideal gas thermodynamic properties for *cis,cis*-1,5-cyclooctadiene are listed in Table 29. No experimental data are available for comparisons. The value of S° (298.15 K) reported by Ermer²³⁶ is about 2 J K⁻¹ mol⁻¹ less than that calculated in the present work (Table 32). We believe our value to be more reliable than that obtained by Ermer²³⁶ from molecular mechanics calculations.

22. 1,3,5,7-Cyclooctatetraene

Cyclooctatetraene has a nonplanar equilibrium conformation (boat form of D_{2d} symmetry), as shown by electron diffraction^{240,241} and nuclear magnetic resonance²⁴² measurements as well as by *ab initio*²⁴³ and force-field^{244–246} calculations. A planar structure (D_{4h} symmetry) was found to be a transition state on the energy surface for interconversion of boat forms.^{242–244,246} Electron diffraction structural parameters of Traetteberg²⁴¹ were used to calculate the product of the principal moments of inertia (Table 8). Bond lengths and angles obtained by Traetteberg²⁴¹ are consistent with those determined in another electron diffraction study²⁴⁰ and calculated theoretically,^{243–246} but Traetteberg's degree of nonplanarity is 10°–25° less than that obtained from theoretical calculations. It is difficult to offer an explanation for this discrepancy. However, it should be emphasized that the electron diffraction model gives a better description of the electronic spectrum than theoretical models.²⁴⁷

Lippincott *et al.*²⁴⁸ observed the infrared and Raman spectra of liquid and gaseous cyclooctatetraene and have made a complete vibrational assignment. Later, Popov and Kogan²⁴⁹ and Traetteberg *et al.*²⁵⁰ carried out a normal coordinate analysis and partly reassigned the observed spectra. The vibrational fundamentals given in Table 9 are those by Popov and Kogan.²⁴⁹ Their vibrational assignment gives a better agreement with experimental entropy than the others. The adopted values of frequencies were taken from the Raman spectrum of the liquid (A_1 and B_1 modes), from in-

Table 29. Ideal gas thermodynamic properties for *cis,cis*-1,5-cyclooctadiene

T	C_p°	$-(G^\circ-H_0^\circ)/T$	S°	$H^\circ-H_0^\circ$	$\log K_f$
K	J K ⁻¹ mol ⁻¹			kJ mol ⁻¹	
100	53.367	217.343	257.120	3.978	-60.758
200	91.777	249.596	305.409	11.163	-40.830
298.15	138.304	275.377	350.508	22.400	-35.277
300	139.226	275.843	351.367	22.657	-35.214
400	188.010	300.563	398.183	39.048	-32.862
500	230.946	324.771	444.885	60.057	-31.672
600	266.581	348.595	490.267	84.992	-30.988
700	295.898	371.954	533.615	113.163	-30.552
800	320.197	394.757	574.762	144.004	-30.252
900	340.521	416.940	613.684	177.070	-30.030
1000	357.646	438.471	650.473	212.002	-29.855
1100	372.165	459.340	685.260	248.512	-29.711
1200	384.529	479.551	718.187	286.363	-29.587
1300	395.109	499.119	749.395	325.359	-29.478
1400	404.198	518.063	779.016	365.336	-29.379
1500	412.043	536.406	807.177	406.157	-29.287

Table 30. Ideal gas thermodynamic properties for 1,3,5,7-cyclooctatetraene

T	C_p°	$-(G^\circ-H_0^\circ)/T$	S°	$H^\circ-H_0^\circ$	$\log K_f$
K	J K ⁻¹ mol ⁻¹			kJ mol ⁻¹	
100	50.684	203.150	241.425	3.828	-171.052
200	83.798	233.889	286.410	10.505	-90.627
298.15	122.013	257.873	326.986	20.606	-64.737
300	123.301	258.301	327.747	20.834	-64.415
400	161.634	286.785	366.617	35.133	-51.580
500	194.415	302.351	406.349	52.999	-44.014
600	220.958	323.202	446.225	73.813	-39.040
700	242.092	343.356	481.969	97.029	-35.523
800	260.665	362.835	515.580	122.220	-32.906
900	275.677	381.553	547.172	149.057	-30.883
1000	288.444	399.616	576.898	177.282	-29.270
1100	299.286	417.018	604.912	206.683	-27.953
1200	308.552	433.789	631.361	237.087	-26.857
1300	316.500	449.958	656.380	268.369	-25.929
1400	323.345	465.556	680.092	300.350	-25.133
1500	329.261	480.616	702.607	332.988	-24.442

Table 31. The uncertainties of the calculated thermal functions (in $\text{J K}^{-1} \text{mol}^{-1}$) and adopted enthalpies of formation (in kJ mol^{-1})

Molecule	Uncertainties in \underline{S}°		Uncertainties in \underline{C}_p°		Uncertainties in $\Delta_f H^{\circ}(298.15 \text{ K})$
	298.15 K	1000 K	298.15 K	1000 K	
	$\text{J K}^{-1} \text{mol}^{-1}$				kJ mol^{-1}
Cyclopropane	0.5	4.5	1.0	6.0	0.5
Cyclopropene	0.5	3.5	1.0	4.0	2.5
Cyclobutane	1.0	6.5	2.0	8.0	0.5
Cyclobutene	1.0	6.5	2.5	7.0	1.5
1,3-Cyclobutadiene	2.0	9.0	3.0	6.0	20.0
Cyclopentane	1.5	9.0	2.5	11.0	0.8
Cyclopentene	1.5	8.0	2.5	9.0	1.7
1,3-Cyclopentadiene	1.5	8.0	2.5	8.0	3.8
Cyclohexane	2.0	13.0	4.0	14.0	0.3
Cyclohexene	2.0	11.5	3.5	12.0	0.5
1,3-Cyclohexadiene	2.0	10.5	3.5	11.0	0.5
1,4-Cyclohexadiene	2.5	11.0	3.5	11.0	5.0
Cycloheptane	3.5	14.0	6.0	15.0	0.6
Cycloheptene	4.0	15.0	6.5	16.0	0.9
1,3-Cycloheptadiene	6.0	17.0	7.5	18.0	0.9
1,3,5-Cycloheptatriene	3.0	13.5	5.0	12.0	1.2
Cyclooctane	4.5	19.0	8.0	20.0	0.9
Cyclooctene	7.5	21.0	8.0	20.0	1.1
1,3-Cyclooctadiene	6.5	18.5	7.0	17.0	2.0
1,5-Cyclooctadiene	7.0	20.0	10.0	20.0	1.2
1,3,5,7-Cyclooctatetraene	3.0	11.0	4.0	10.0	1.3

Table 32. Comparison of experimental^a entropies with calculated values (in J K⁻¹ mol⁻¹)

S°			Investigator (year)	Reference
298.15 K	500 K	1000 K		
Cyclopropane				
<u>237.4</u>			Ruehrwein and Powell (1946)	21
237.4	275.9	360.5	Kobe and Pennington (1951)	22
238.9	277.3		Chang <i>et al.</i> (1970)	24
237.2	274.7	357.9	Karuppannan <i>et al.</i> (1981)	23
237.4	275.5	359.5	This work	
Cyclopropene				
243.5	277.7	346.5	Yun and Eggers (1979)	30
243.5	277.5	346.2	Adame and Vicharelli (1979)	32
243.5	277.7	346.6	This work	
Cyclobutane				
<u>265.4±0.8</u>			Rathjens and Gwinn (1953)	55 ^b
<u>265.4</u>	315.6	428.9	Rathjens <i>et al.</i> (1953)	50
264.1	313.7		Chang <i>et al.</i> (1970)	24
264.4	313.4	424.7	This work	
Cyclobutene				
263.5	308.9	406.5	Danti (1956)	65
262.0	306.1	403.1	This work	
Cyclopentane				
<u>292.8</u>			Kilpatrick <i>et al.</i> (1947)	88
<u>292.9</u>			McCullough <i>et al.</i> (1959)	101
292.9	352.0	491.4	Kilpatrick <i>et al.</i> (1947)	88
292.9	352.3	492.2	McCullough <i>et al.</i> (1959)	101
293.8	353.3		Chang <i>et al.</i> (1970)	24
292.9	352.0	491.0	This work	
Cyclopentene				
<u>289.7</u>			Beckett <i>et al.</i> (1948)	109
289.7	342.3	464.2	Beckett <i>et al.</i> (1948)	109
289.2			Furuyama <i>et al.</i> (1970)	114
289.1	343.6	466.5	Draeger <i>et al.</i> (1983)	115
291.3	347.5	472.6	This work	
1,3-Cyclopentadiene				
<u>270.3±4.0</u> ^c			Grant and Walsh (1969)	124
274.4	325.0	434.3	Turnbull (1967)	125
274.5			Furuyama <i>et al.</i> (1970)	114
274.0	325.8	436.9	This work	
Cyclohexane				
<u>298.8±0.8</u>			Aston <i>et al.</i> (1943)	262
<u>298.2</u>			Beckett <i>et al.</i> (1947)	145
298.2	373.4	550.6	Beckett <i>et al.</i> (1947)	145
299.6	374.6		Chang <i>et al.</i> (1970)	24
297.3	371.8	547.5	This work	

Table 32. Comparison of experimental^a entropies with calculated values (in J K⁻¹ mol⁻¹)--Continued

Cyclohexene				
<u>310.5</u>			Beckett <i>et al.</i> (1948)	109
310.7	383.4	543.1	Beckett <i>et al.</i> (1948)	109
310.5	380.6	537.1	This work	
Cycloheptane				
<u>342.3±1.3</u>			Finke <i>et al.</i> (1956)	188
343.0	431.9		Chang <i>et al.</i> (1970)	24
342.3	434.2		This work	
1,3,5-Cycloheptatriene				
<u>315.6±1.0</u>			Finke <i>et al.</i> (1956)	188
318.0			Evans and Lord (1960)	213
316.3			This work	
Cyclooctane				
<u>366.8±1.3</u>			Finke <i>et al.</i> (1956)	188
373.0	455.3		Chang <i>et al.</i> (1970)	24
366.7	468.2		This work	
1,5-Cyclooctadiene				
347.9			Ermer (1976)	236
350.3			This work	
1,3,5,7-Cyclooctatetraene				
<u>326.8±1.5</u>			Scott <i>et al.</i> (1949)	251
325.3	406.8	575.0	Lippincott and Lord (1951)	252
327.0	408.3	576.9	This work	

^a Experimental values of \bar{S}^0 (298.15 K) are underlined.^b From calorimetric measurements Rathjens and Gwinn⁵⁵ obtained the value of \bar{S}^0 (285.67 K). On the basis of this value Stull *et al.*²⁵³ calculated \bar{S}^0 (298.15 K).^c The entropy value was calculated from equilibrium data.

Table 33. Comparison of experimental heat capacities with calculated values (in $\text{J K}^{-1} \text{mol}^{-1}$)

Experimental C_p° data		Calculated C_p° data	
Cyclopropane			
T, K	Ref. 254	Ref. 254	This work
272.15	50.6	50.8	50.5
300.48	56.5	56.4	56.0
333.70	63.2	63.2	62.8
368.46	70.2	70.4	69.8
T, K	Ref. 255 ^a	Ref. 255	This work
157.6	35.3	35.0	35.0
223.4	42.6, 43.1	42.1	42.1
258.4	48.7, 49.6	48.1	47.9
295.4	55.5, 57.1	55.2	55.0
313.9	59.3, 61.3	59.1	58.7
325.3	61.3, 63.6	61.5	61.1
338.9	64.2	64.2	63.8
Cyclopentane			
T, K	Ref. 258	Ref. 88	This work
353	102.0	102.0	101.8
372	108.2		108.5
395	117.3		116.4
424	126.3		126.2
463	138.7		138.8
503	150.7	150.9	150.9
539	160.1	161.2	161.2
T, K	Ref. 101	Ref. 101	This work
329.05	94.0	93.9	93.4
395.05	117.1	117.1	116.5
463.10	139.5	139.6	138.8
Cyclohexane			
T, K	Ref. 258		This work
384	143.1		141.9
428	161.8		160.3
460	174.5		173.2
495	189.5		186.8
521	196.6		196.4
544	206.3		204.6
Cyclohexene			
T, K	Ref. 109	Ref. 109	This work
400	145.0	144.9	139.7

^a Only a part of experimental values is included in this table

frared spectrum of the gas (B_2 and E modes), and from calculation (unobserved A_2 modes).

Ideal gas thermodynamic properties for 1,3,5,7-cyclooctatetraene are given in Table 30. The calculated value of S° (298.15 K), as shown in Table 32, agrees well with the calorimetric entropy.²⁵¹ Lippincott and Lord²⁵² have calculated thermal functions for the crown structure of D_4 symmetry. Calculated S° values are somewhat lower than values obtained in this work (Table 32), while the C_p° values are close to those given in Table 30. The discrepancies in the entropy values are due to the different vibrational frequencies adopted for calculations (the discrepancy in symmetry numbers is compensated for the presence of optical isomers in the case of D_4 symmetry).

23. References

- ¹L. V. Gurvich, I. V. Veyts, and V. A. Medvedev, *Termodinamicheskie Svoistva Individualnykh Veshchestv*, Tom I (Nauka, Moskva, 1978). [English translation: *Thermodynamic Properties of Individual Substances* (to be published by Hemisphere)].
- ²J. B. Pedley and J. Rylance, *Computer Analysed Thermochemical Data: Organic and Organometallic Compounds* (University of Sussex, England, 1977).
- ³E. P. Vedenev, *Programme resheniya prikladnykh matematicheskikh zadach*, Vip. 11 (MGU, Moskva, 1970). [Programs for Applied Mathematical Problems].
- ⁴N. B. Sherbak and V. S. Jorish, in *Razrabotka i primeneniye dialogovykh sistem dlya resheniya prikladnykh zadach na mini EVM*, Vip. 1 (IVTAN Moskva, 1981). [Development and Application of Dialogue Systems for Solution of Applied Problems by Using Mini-Computers].
- ⁵H. H. Günthard, R. C. Lord, and T. K. McCubbin, Jr., *J. Chem. Phys.* **25**, 768 (1956).
- ⁶J. L. Duncan, *J. Mol. Spectrosc.* **25**, 451 (1968).
- ⁷T. K. McCubbin, Jr., V. Withstandly, and S. R. Polo, *J. Mol. Spectrosc.* **31**, 95 (1969).
- ⁸P. M. Mathai, G. G. Shepherd, and H. L. Welsh, *Can. J. Phys.* **34**, 1448 (1956).
- ⁹W. J. Jones and B. P. Stoicheff, *Can. J. Phys.* **42**, 2259 (1964).
- ¹⁰R. J. Butcher and W. J. Jones, *J. Mol. Spectrosc.* **47**, 64 (1973).
- ¹¹O. Bastiansen, F. N. Fritsch, and K. Hedberg, *Acta Crystallogr.* **17**, 538 (1964).
- ¹²A. W. Baker and R. C. Lord, *J. Chem. Phys.* **23**, 1636 (1955).
- ¹³C. Brecher, E. Krikorian, J. Blanc, and R. C. Halford, *J. Chem. Phys.* **35**, 1097 (1961).
- ¹⁴J. L. Duncan and D. C. McKean, *J. Mol. Spectrosc.* **27**, 117 (1968).
- ¹⁵G. J. Cartwright and I. M. Mills, *J. Mol. Spectrosc.* **34**, 415 (1970).
- ¹⁶J. L. Duncan and G. R. Burns, *J. Mol. Spectrosc.* **30**, 253 (1969).
- ¹⁷T. Hirokawa, M. Hayashi, and H. Murata, *J. Sci. Hiroshima Univ. Ser. A: Math. Phys. Chem.* **37**, 271 (1973); **39**, 161 (1975).
- ¹⁸J. R. Robins, S. J. Daunt, and H. F. Shurvell, *J. Raman Spectrosc.* **5**, 411 (1976).
- ¹⁹M. Spiekermann, D. Bougeard, and B. Schrader, *J. Mol. Struct.* **60**, 55 (1980).
- ²⁰S. J. Daunt and H. F. Shurvell, *J. Raman Spectrosc.* **2**, 463 (1974).
- ²¹R. A. Ruehrwein and T. M. Powell, *J. Am. Chem. Soc.* **68**, 1063 (1946).
- ²²K. A. Kobe and R. E. Pennington, *Pet. Refiner* **29**, 93 (1950).
- ²³T. A. Karuppahnan, R. Ramasamy, and K. Venkateswarlu, *J. Inorg. Nucl. Chem.* **43**, 1555 (1981).
- ²⁴S. Chang, D. McNally, S. Shary-Tehrany, M. J. Hickey, and R. H. Boyd, *J. Am. Chem. Soc.* **92**, 3109 (1970).
- ²⁵J. F. Chiang, *J. Chin. Chem. Soc. (Taipei)* **17**, 65 (1970).
- ²⁶P. H. Kasai, R. J. Myers, D. F. Eggers, and K. B. Wieberg, *J. Chem. Phys.* **30**, 512 (1959).
- ²⁷W. M. Stigliani, V. W. Laurie, and C. L. James, *J. Chem. Phys.* **62**, 1890 (1975).
- ²⁸D. F. Eggers, J. W. Schultz, K. B. Wieberg, E. K. Wagner, L. M. Jackman, and R. L. Erskine, *J. Chem. Phys.* **47**, 946 (1967).
- ²⁹R. W. Mitchell, E. A. Dorko, and J. A. Merritt, *J. Mol. Spectrosc.* **26**, 197 (1968).
- ³⁰T. Y. Yum and D. F. Eggers, Jr., *J. Phys. Chem.* **83**, 501 (1979).
- ³¹K. B. Wieberg and J. J. Wendoloski, *J. Phys. Chem.* **83**, 497 (1979).
- ³²I. E. Adame and P. A. Vicharelli, *Thermochim. Acta* **31**, 203 (1979).
- ³³D. A. Dows and N. Rich, *J. Chem. Phys.* **47**, 333 (1967).
- ³⁴T. Ueda and T. Schimanouchi, *J. Chem. Phys.* **49**, 470 (1968).
- ³⁵J. M. R. Stone and I. M. Mills, *Mol. Phys.* **18**, 631 (1970).
- ³⁶F. A. Miller and R. J. Capwell, *Spectrochim. Acta A* **27**, 947 (1971).
- ³⁷T. B. Malloy, Jr. and W. J. Lafferty, *J. Mol. Spectrosc.* **54**, 20 (1975).
- ³⁸S. Meiboom and L. C. Snyder, *J. Chem. Phys.* **52**, 3857 (1970).
- ³⁹J. D. Dunitz and V. Schomaker, *J. Chem. Phys.* **20**, 1703 (1952).
- ⁴⁰A. Almenningen, O. Bastiansen, and P. N. Skancke, *Acta Chem. Scand.* **15**, 711 (1961).
- ⁴¹P. N. Skancke, Ph.D. thesis (University of Oslo, Norway, 1960).
- ⁴²F. Takabayashi, H. Kambara, and K. Kuchitsu, in *Seventh Austin Symposium on Gas Phase Molecular Structure*, Austin, Texas, 1978, p. 63.
- ⁴³J. S. Wright and L. Salem, *J. Am. Chem. Soc.* **94**, 322 (1972).
- ⁴⁴D. Cremer, *J. Am. Chem. Soc.* **99**, 1307 (1977).
- ⁴⁵J. S. Wright and L. Salem, *J. Chem. Soc. D* (23), 1370 (1969).
- ⁴⁶J. L. Nelson and A. A. Frost, *J. Am. Chem. Soc.* **94**, 3727 (1972).
- ⁴⁷R. Pasternak and A. Y. Meyer, *J. Mol. Struct.* **13**, 201 (1972).
- ⁴⁸L. S. Bartell and B. Andersen, *J. Chem. Soc. D* (20), 786 (1973).
- ⁴⁹R. Pasternak and A. Y. Meyer, *J. Mol. Struct.* **20**, 351 (1974).
- ⁵⁰G. W. Rathjens, Jr., N. K. Freeman, W. D. Gwinn, and K. S. Pitzer, *J. Am. Chem. Soc.* **75**, 5634 (1953).
- ⁵¹R. C. Lord and I. Nakagawa, *J. Chem. Phys.* **39**, 2951 (1963).
- ⁵²F. A. Miller, R. J. Capwell, R. C. Lord, and D. G. Rea, *Spectrochim. Acta Part A* **28**, 603 (1972).
- ⁵³V. T. Aleksanyan and B. G. Antipov, *J. Mol. Struct.* **89**, 15 (1982).
- ⁵⁴G. Bánhegyi, G. Fogarasi, and P. Pulay, *J. Mol. Struct.* **89**, 1 (1982).
- ⁵⁵G. W. Rathjens, Jr. and W. D. Gwinn, *J. Am. Chem. Soc.* **75**, 5629 (1953).
- ⁵⁶E. Goldish, K. Hedberg, and V. Schomaker, *J. Am. Chem. Soc.* **78**, 2714 (1956).
- ⁵⁷H. Kim and D. W. Gwinn, *J. Chem. Phys.* **42**, 3728 (1965).
- ⁵⁸B. Bak, J. J. Led, L. Nygaard, J. Rastrup-Andersen, and G. O. Sørensen, *J. Mol. Struct.* **3**, 369 (1969).
- ⁵⁹K. B. Wieberg and J. J. Wendoloski, *J. Am. Chem. Soc.* **104**, 5679 (1982).
- ⁶⁰R. C. Lord and D. G. Rea, *J. Am. Chem. Soc.* **79**, 2 (1957).
- ⁶¹L. M. Sverdlov and E. P. Krainov, *Opt. Spektrosk.* **13**, 169 (1962).
- ⁶²E. M. Suzuki and J. M. Nibler, *Spectrochim. Acta Part A* **30**, 15 (1974).
- ⁶³V. T. Aleksanyan, and O. G. Garkusha, *Izv. Akad. Nauk SSSR Ser. Khim.* (10), 2227 (1976).
- ⁶⁴J. R. Villarreal, J. Laane, S. F. Bush, and W. C. Harris, *Spectrochim. Acta Part A* **35**, 331 (1979).
- ⁶⁵A. Danti, *J. Chem. Phys.* **27**, 1227 (1956).
- ⁶⁶O. L. Chapman, C. L. McIntosh, and J. Pacansky, *J. Am. Chem. Soc.* **95**, 614 (1973).
- ⁶⁷O. L. Chapman, D. De La Cruz, R. Roth, and J. Pacansky, *J. Am. Chem. Soc.* **95**, 1337 (1973).
- ⁶⁸C. Y. Lin, and A. Krantz, *J. Chem. Soc. D* (19), 1111 (1972).
- ⁶⁹A. Krantz, C. Y. Lin, and M. D. Newton, *J. Am. Chem. Soc.* **95**, 2744 (1973).
- ⁷⁰M. J. S. Dewar and A. Komornicki, *J. Am. Chem. Soc.* **99**, 6174 (1977).
- ⁷¹R. C. Haddon and G. R. J. Williams, *J. Am. Chem. Soc.* **97**, 6582 (1975).
- ⁷²K. Jug and D. N. Nanda, *Theor. Chim. Acta* **57**, 107 (1980).
- ⁷³H. Kollmar, F. Carrion, M. J. S. Dewar, and R. C. Bingham, *J. Am. Chem. Soc.* **103**, 5293 (1981).
- ⁷⁴H. Kollmar and V. Staemmler, *J. Am. Chem. Soc.* **99**, 3583 (1977).
- ⁷⁵W. T. Borden, E. R. Davidson, and P. Hart, *J. Am. Chem. Soc.* **100**, 388 (1978).
- ⁷⁶J. A. Jafri and M. D. Newton, *J. Am. Chem. Soc.* **100**, 5012 (1978).
- ⁷⁷L. J. Schaad, B. A. Hess, Jr., and C. S. Ewig, *J. Am. Chem. Soc.* **101**, 2281 (1979).
- ⁷⁸H. Kollmar, *J. Am. Chem. Soc.* **102**, 2617 (1980).
- ⁷⁹L. J. Schaad, B. A. Hess, Jr., and C. S. Ewig, *J. Org. Chem.* **47**, 2904 (1982).
- ⁸⁰B. A. Hess, Jr., P. Čársky, and L. J. Schaad, *J. Am. Chem. Soc.* **105**, 695 (1983).
- ⁸¹S. Masamune, F. A. Souto-Bachiller, T. Machiguchi, and J. E. Bertie, *J. Am. Chem. Soc.* **100**, 4889 (1978).
- ⁸²H. Kollmar and V. Staemmler, *J. Am. Chem. Soc.* **100**, 4304 (1978).
- ⁸³V. P. Novikov and A. I. Malyshev, *Zh. Prikl. Spektrosk.* **33**, 545 (1980).
- ⁸⁴E. Castellucci, P. Manzelli, B. Fortunato, E. Gallinella, and P. Mirone, *Spectrochim. Acta Part A* **31**, 451 (1975).

- ⁸⁵C. Di Lauro, N. Neto, and S. Califano, *J. Mol. Struct.* **3**, 219 (1969).
- ⁸⁶A. Ansmann and B. Schrader, *J. Raman Spectrosc.* **5**, 281 (1976).
- ⁸⁷A. Schweig and W. Thiel, *J. Am. Chem. Soc.* **101**, 4742 (1979).
- ⁸⁸J. E. Kilpatrick, K. S. Pitzer, and R. Spitzer, *J. Am. Chem. Soc.* **69**, 2483 (1947).
- ⁸⁹K. S. Pitzer and W. E. Donath, *J. Am. Chem. Soc.* **81**, 3213 (1959).
- ⁹⁰J. B. Hendrickson, *J. Am. Chem. Soc.* **83**, 4537 (1961); **85**, 4059 (1963).
- ⁹¹N. L. Allinger, J. A. Hirsch, M. A. Miller, I. J. Tyminski, and F. A. Van Catledge, *J. Am. Chem. Soc.* **90**, 1199 (1968).
- ⁹²S. Lifson and A. Warshel, *J. Chem. Phys.* **49**, 5116 (1968).
- ⁹³J. R. Hoyland, *J. Chem. Phys.* **50**, 2775 (1969).
- ⁹⁴J. R. Durig and D. W. Wertz, *J. Chem. Phys.* **49**, 2118 (1968).
- ⁹⁵L. A. Carreira, G. J. Jiang, and W. B. Person, *J. Chem. Phys.* **56**, 1440 (1972).
- ⁹⁶T. H. Chao and J. Laane, *J. Mol. Spectrosc.* **70**, 357 (1978).
- ⁹⁷W. J. Adams, H. J. Geise, and L. S. Bartell, *J. Am. Chem. Soc.* **92**, 5013 (1970).
- ⁹⁸F. A. Miller and R. G. Inskeep, *J. Chem. Phys.* **18**, 1519 (1950).
- ⁹⁹L. M. Sverdlov and N. I. Prokofjeva, *Opt. Spektrosk.* **7**, 363 (1959).
- ¹⁰⁰F. H. Kurse and D. W. Scott, *J. Mol. Spectrosc.* **20**, 276 (1966).
- ¹⁰¹J. P. McCullough, R. E. Pennington, J. C. Smith, I. A. Hossenlopp, and G. Waddington, *J. Am. Chem. Soc.* **81**, 5880 (1959).
- ¹⁰²G. W. Rathjens, Jr., *J. Chem. Phys.* **36**, 2401 (1962).
- ¹⁰³S. S. Butcher and C. C. Costain, *J. Mol. Spectrosc.* **15**, 40 (1965).
- ¹⁰⁴L. H. Scharpen, *J. Chem. Phys.* **48**, 3552 (1968).
- ¹⁰⁵M. I. Davis and T. W. Muecke, *J. Phys. Chem.* **74**, 1104 (1970).
- ¹⁰⁶J. Laane and R. C. Lord, *J. Chem. Phys.* **47**, 4941 (1967).
- ¹⁰⁷T. B. Malloy, Jr. and L. A. Carreira, *J. Chem. Phys.* **71**, 2488 (1979).
- ¹⁰⁸L. E. Bauman, P. M. Killough, J. M. Cooke, J. R. Villarreal, and J. Laane, *J. Phys. Chem.* **86**, 2000 (1982).
- ¹⁰⁹G. W. Beckett, N. K. Freeman, and K. S. Pitzer, *J. Am. Chem. Soc.* **70**, 4227 (1948).
- ¹¹⁰L. M. Sverdlov and E. P. Krainov, *Opt. Spektrosk.* **6**, 214 (1959).
- ¹¹¹A. LeRoy and J. C. Thonvenout, *C. R. Acad. Sci. Ser. B* **37**, 545 (1967).
- ¹¹²W. C. Harris and C. T. Longshore, *J. Mol. Struct.* **16**, 187 (1973).
- ¹¹³O. Ermer and S. Lifson, *J. Mol. Spectrosc.* **51**, 261 (1974).
- ¹¹⁴S. Furuyama, D. M. Golden, and S. W. Benson, *J. Chem. Thermodyn.* **2**, 161 (1970).
- ¹¹⁵J. A. Draeger, R. H. Harrison, and W. D. Good, *J. Chem. Thermodyn.* **15**, 367 (1983).
- ¹¹⁶L. H. Scharpen and V. W. Laurie, *J. Chem. Phys.* **43**, 2765 (1965).
- ¹¹⁷D. Damiani, L. Ferretti, and E. Gallinella, *Chem. Phys. Lett.* **37**, 265 (1976).
- ¹¹⁸J. Kao and L. Radom, *J. Am. Chem. Soc.* **100**, 379 (1978).
- ¹¹⁹E. V. Sobolev, V. T. Aleksanyan, and V. A. Mironov, *Dokl. Akad. Nauk SSSR* **152**, 923 (1963).
- ¹²⁰H. P. Fritz and L. Schäfer, *Spectrochim. Acta* **21**, 211 (1965).
- ¹²¹L. A. Evseva and L. M. Sverdlov, *Izv. Vyssh. Uchebn. Zaved. Ser. Fiz.* **11**, 91 (1968).
- ¹²²E. Gallinella, B. Fortunato, and P. Mirone, *J. Mol. Spectrosc.* **24**, 345 (1967).
- ¹²³K. E. Blick, J. W. de Haan, and K. Niedenzu, *Spectrochim. Acta Part A* **26**, 2319 (1970).
- ¹²⁴C. J. Grant and R. Walsh, *J. Chem. Soc. Chem. Commun.* (12), 667 (1969).
- ¹²⁵A. G. Turnbull, *Aust. J. Chem.* **20**, 2059 (1967).
- ¹²⁶M. I. Davis and O. Hassel, *Acta Chem. Scand.* **17**, 1181 (1963); **18**, 813 (1964).
- ¹²⁷H. R. Buys and H. J. Geise, *Tetrahedron Lett.* **34**, 2991 (1970).
- ¹²⁸H. J. Geise, H. R. Buys, and F. C. Mijlhoff, *J. Mol. Struct.* **9**, 447 (1971).
- ¹²⁹O. Bastiansen, L. Fernholt, H. M. Seip, H. Kambara, and K. Kuchitsu, *J. Mol. Struct.* **18**, 163 (1973).
- ¹³⁰J. D. Ewbank, G. Kirsch, and L. Schäfer, *J. Mol. Struct.* **31**, 39 (1976).
- ¹³¹H. Takahashi, T. Shimanouchi, K. Fukushima, and T. Miyazawa, *J. Mol. Spectrosc.* **13**, 43 (1964).
- ¹³²R. G. Snyder and J. H. Schachtschneider, *Spectrochim. Acta* **21**, 169 (1965).
- ¹³³K. B. Wieberg and A. Shrake, *Spectrochim. Acta Part A* **27**, 1139 (1971); **29**, 583 (1973).
- ¹³⁴R. L. Hilderbrandt, *Comput. Chem.* **1**, 179 (1977).
- ¹³⁵A. Hiraya, Y. Udagawa, and M. Ito, *Chem. Lett.* 433 (1979).
- ¹³⁶J. L. Offenbach, L. Fredin, and H. L. Strauss, *J. Am. Chem. Soc.* **103**, 1001 (1981).
- ¹³⁷J. P. Huvenne, G. Vergoten, and G. Fleury, *Can. J. Chem.* **60**, 1347 (1982).
- ¹³⁸F. A. L. Anet and A. J. R. Bourn, *J. Am. Chem. Soc.* **89**, 760 (1967).
- ¹³⁹E. M. Engler, J. D. Andose, and P. von R. Schleyer, *J. Am. Chem. Soc.* **95**, 8005 (1973).
- ¹⁴⁰N. L. Allinger, M. T. Tribble, M. A. Miller, and D. H. Wertz, *J. Am. Chem. Soc.* **93**, 1637 (1971).
- ¹⁴¹N. L. Allinger, *Adv. Phys. Org. Chem.* **13**, 1 (1976).
- ¹⁴²N. L. Allinger, *J. Am. Chem. Soc.* **99**, 8127 (1977).
- ¹⁴³D. N. J. White and M. J. Bovill, *J. Chem. Soc. Perkin Trans. 2* (12), 1610 (1977).
- ¹⁴⁴M. Squillacote, R. S. Sheridan, O. L. Chapman, and F. A. L. Anet, *J. Am. Chem. Soc.* **97**, 3244 (1975).
- ¹⁴⁵C. W. Beckett, K. S. Pitzer, and R. Spitzer, *J. Am. Chem. Soc.* **69**, 2488 (1947).
- ¹⁴⁶V. A. Naumov, V. G. Dashevsky, and N. M. Zaripov, *Zh. Strukt. Khim.* **11**, 793 (1970).
- ¹⁴⁷F. Chiang and S. H. Bauer, *J. Am. Chem. Soc.* **91**, 1898 (1969).
- ¹⁴⁸H. J. Geise and H. R. Buys, *Recl. Trav. Chim. Pays-Bas* **89**, 1147 (1970).
- ¹⁴⁹L. H. Schrapen, J. E. Wollrab, and D. P. Ames, *J. Chem. Phys.* **49**, 2368 (1968).
- ¹⁵⁰T. Ogata and K. Kozima, *Bull. Chem. Soc. Jpn.* **42**, 1263 (1969).
- ¹⁵¹N. Neto, C. Di Lauro, E. Castellucci, and S. Califano, *Spectrochim. Acta Part A* **23**, 1763 (1967).
- ¹⁵²J. R. Durig, R. O. Carter, and L. A. Carreira, *J. Chem. Phys.* **60**, 3098 (1974).
- ¹⁵³T. L. Smithson and H. Wieser, *J. Chem. Phys.* **72**, 2340 (1980).
- ¹⁵⁴F. A. L. Anet and M. Z. Haq, *J. Am. Chem. Soc.* **87**, 3147 (1965).
- ¹⁵⁵S. Saebo and J. E. Boggs, *J. Mol. Struct.* **73**, 137 (1981).
- ¹⁵⁶N. L. Allinger, J. A. Hirsch, M. A. Miller, and I. J. Tyminski, *J. Am. Chem. Soc.* **90**, 5773 (1968).
- ¹⁵⁷N. L. Allinger and J. T. Sprague, *J. Am. Chem. Soc.* **94**, 5734 (1972).
- ¹⁵⁸R. Bucourt and D. Hainaut, *Bull. Soc. Chim. Fr.* (5), 1366 (1965).
- ¹⁵⁹V. G. Dashevsky and A. A. Lugovskoy, *J. Mol. Struct.* **12**, 39 (1972).
- ¹⁶⁰F. A. L. Anet and I. Yavari, *Tetrahedron* **34**, 2879 (1978).
- ¹⁶¹G. Dallinga and L. H. Toneman, *J. Mol. Struct.* **1**, 11 (1967).
- ¹⁶²M. Traetteberg, *Acta Chem. Scand.* **22**, 2305 (1968).
- ¹⁶³H. Oberhammer and S. H. Bauer, *J. Am. Chem. Soc.* **91**, 10 (1969).
- ¹⁶⁴A. Warshel and M. Karplus, *J. Am. Chem. Soc.* **94**, 5612 (1972).
- ¹⁶⁵N. L. Allinger and J. T. Sprague, *J. Am. Chem. Soc.* **95**, 3893 (1973).
- ¹⁶⁶A. J. Birch, A. L. Hinde, and L. Radom, *J. Am. Chem. Soc.* **103**, 284 (1981).
- ¹⁶⁷S. S. Butcher, *J. Chem. Phys.* **42**, 1830 (1965).
- ¹⁶⁸G. Luss and M. D. Harmony, *J. Chem. Phys.* **43**, 3768 (1965).
- ¹⁶⁹L. A. Carreira, R. O. Carter, and J. R. Durig, *J. Chem. Phys.* **59**, 812 (1973).
- ¹⁷⁰H. Gerding and F. A. Haak, *Recl. Trav. Chim. Pays-Bas* **68**, 293 (1949).
- ¹⁷¹H. D. Stidham, *Spectrochim. Acta* **21**, 23 (1965).
- ¹⁷²G. Dallinga and L. H. Toneman, *J. Mol. Struct.* **1**, 117 (1967).
- ¹⁷³E. W. Barbsich, Jr. and M. G. Griffith, *J. Am. Chem. Soc.* **90**, 3590 (1968).
- ¹⁷⁴G. Ahlgren, G. Akermark, and J.-E. Bäckvall, *Tetrahedron Lett.* (40), 3501 (1975).
- ¹⁷⁵D. J. Atkinson and M. J. Perkins, *Tetrahedron Lett.* (28), 2335 (1969).
- ¹⁷⁶F. Herbstein, *J. Chem. Soc.* 2292 (1959).
- ¹⁷⁷J. Laane and R. C. Lord, *J. Mol. Spectrosc.* **39**, 340 (1971).
- ¹⁷⁸K. B. Lipkowitz, P. W. Rabideau, D. J. Raber, L. E. Hardee, P. von R. Schleyer, A. J. Kos, and R. A. Kahn, *J. Org. Chem.* **47**, 1002 (1982).
- ¹⁷⁹J. L. Marshall and L. Hall, *Tetrahedron* **37**, 1271 (1981).
- ¹⁸⁰O. Gebhardt and S. J. Cyvin, *J. Mol. Struct.* **27**, 345 (1975).
- ¹⁸¹R. Shaw, D. M. Golden, and S. W. Benson, *J. Phys. Chem.* **81**, 1716 (1977).
- ¹⁸²J. B. Hendrickson, *J. Am. Chem. Soc.* **89**, 7036, 7047 (1967).
- ¹⁸³M. Bixon and S. Lifson, *Tetrahedron* **23**, 769 (1967).
- ¹⁸⁴D. F. Bocian and H. L. Strauss, *J. Am. Chem. Soc.* **99**, 2876 (1977).
- ¹⁸⁵D. F. Bocian and H. L. Strauss, *J. Am. Chem. Soc.* **99**, 2866 (1977).
- ¹⁸⁶J. Dillen and H. J. Geise, *J. Chem. Phys.* **70**, 425 (1979).
- ¹⁸⁷O. V. Dorofeeva, L. V. Gurvich, and V. S. Mastryukov, in *X Vsesoyuznaja Konferentsiya po Kalorimetrii i Khimicheskoi Termodinamike*, Moskva, 1984, Vol. 1, Part 2, p. 241 [Tenth Soviet Conference on Calorimetry and Chemical Thermodynamics]; *J. Mol. Struct.* **129**, 165 (1985).
- ¹⁸⁸H. L. Finke, D. W. Scott, M. E. Gross, J. F. Messerly, and G. Waddington, *J. Am. Chem. Soc.* **78**, 5469 (1956).
- ¹⁸⁹N. Neto, C. Di Lauro, and S. Califano, *Spectrochim. Acta Part A* **26**, 1489 (1970).
- ¹⁹⁰M. St.-Jacques and C. Vaziri, *Can. J. Chem.* **49**, 1256 (1971).
- ¹⁹¹G. Favini, G. Buemi, and M. Raimondi, *J. Mol. Struct.* **2**, 137 (1968).

- ¹⁹²G. Favini and A. Nava, *Theor. Chim. Acta* **31**, 261 (1973).
- ¹⁹³O. Ermer and S. Lifson, *J. Am. Chem. Soc.* **95**, 4121 (1973).
- ¹⁹⁴J. F. Chiang and S. H. Bauer, *J. Am. Chem. Soc.* **88**, 420 (1966).
- ¹⁹⁵K. Hagen and M. Traetteberg, *Acta Chem. Scand.* **26**, 3643 (1972).
- ¹⁹⁶P. Crews, *J. Chem. Soc. D* (11), 583 (1971).
- ¹⁹⁷B. Schrader and A. Ansmann, *Angew. Chem. Int. Ed. Engl.* **14**, 364 (1975).
- ¹⁹⁸T. K. Avirah, T. B. Malloy, and R. L. Cook, *J. Chem. Phys.* **71**, 2194 (1979).
- ¹⁹⁹G. Favini, F. Zuccarello, and G. Buemi, *J. Mol. Struct.* **3**, 385 (1969).
- ²⁰⁰F. Zuccarello, G. Buemi, and G. Favini, *J. Mol. Struct.* **18**, 295 (1973).
- ²⁰¹G. Favini, C. Rubino, and R. Todeschini, *J. Mol. Struct.* **41**, 305 (1977).
- ²⁰²N. L. Allinger and J. T. Sprague, *Tetrahedron* **29**, 3811 (1973).
- ²⁰³S. Saebø and J. E. Boggs, *J. Mol. Struct.* **87**, 365 (1982).
- ²⁰⁴M. Traetteberg, *J. Am. Chem. Soc.* **86**, 4265 (1964).
- ²⁰⁵S. S. Butcher, *J. Chem. Phys.* **42**, 1833 (1965).
- ²⁰⁶C. La Lau and H. De Ruyter, *Spectrochim. Acta* **19**, 1559 (1963).
- ²⁰⁷F. A. L. Anet, *J. Am. Chem. Soc.* **86**, 458 (1964).
- ²⁰⁸F. R. Jensen and L. A. Smith, *J. Am. Chem. Soc.* **86**, 956 (1964).
- ²⁰⁹G. Buemi, F. Zuccarello, and D. Grasso, *J. Mol. Struct.* **42**, 195 (1977).
- ²¹⁰W. Paulick, Ch. Jung, U. Kempka, J. Sühnel, and K. Gustav, *J. Mol. Struct.* **85**, 235 (1981).
- ²¹¹H. J. Linder, *Tetrahedron* **37**, 535 (1981).
- ²¹²J. M. Schulman, R. L. Disch, and M. L. Sabio, *J. Am. Chem. Soc.* **104**, 3785 (1982).
- ²¹³M. V. Evans and R. C. Lord, *J. Am. Chem. Soc.* **82**, 1876 (1960).
- ²¹⁴K. B. Wieberg, *J. Am. Chem. Soc.* **87**, 1070 (1965).
- ²¹⁵F. A. L. Anet and J. Krane, *Tetrahedron Lett.* **50**, 5029 (1973).
- ²¹⁶P. W. Pakes, T. C. Rounds, and H. L. Strauss, *J. Phys. Chem.* **85**, 2469 (1981).
- ²¹⁷F. A. L. Anet and V. T. Basus, *J. Am. Chem. Soc.* **95**, 4424 (1973).
- ²¹⁸S. Meiboom, R. C. Hewitt, and Z. Luz, *J. Chem. Phys.* **66**, 4041 (1977).
- ²¹⁹A. Almenningen, O. Bastiansen, and N. Jensen, *Acta Chem. Scand.* **20**, 2689 (1966).
- ²²⁰O. V. Dorofeeva, V. S. Mastryukov, N. L. Allinger, and A. Almenningen, *J. Phys. Chem.* **89**, 252 (1985).
- ²²¹H. E. Bellis and E. J. Slowinski, Jr., *Spectrochim. Acta* **12**, 1103 (1959).
- ²²²P. W. Pakes, T. C. Rounds, and H. L. Strauss, *J. Phys. Chem.* **85**, 2476 (1981).
- ²²³F. A. L. Anet and I. Yavari, *J. Am. Chem. Soc.* **99**, 6986 (1977).
- ²²⁴R. M. Gavin, Jr. and Z. F. Wang, *J. Am. Chem. Soc.* **95**, 1425 (1973).
- ²²⁵G. Buemi, G. Favini, and F. Zuccarello, *J. Mol. Struct.* **5**, 101 (1970).
- ²²⁶M. Traetteberg, *Acta Chem. Scand. Ser. B* **29**, 29 (1975).
- ²²⁷M. Traetteberg, P. Bakken, and A. Almenningen, *J. Mol. Struct.* **74**, 321 (1981).
- ²²⁸E. A. Bruade, *Chem. Ind. (London)* (51), 1577 (1954).
- ²²⁹M. Traetteberg, *Acta Chem. Scand.* **24**, 2285 (1970).
- ²³⁰N. L. Allinger, J. F. Viskocil, Jr., U. Burkert, and Y. Yuh, *Tetrahedron* **32**, 33 (1976).
- ²³¹F. A. L. Anet and I. Yavari, *J. Am. Chem. Soc.* **100**, 7814 (1978).
- ²³²M. P. Kozina, V. S. Mastryukov, and E. M. Milvitskaya, *Usp. Khim.* **51**, 1337 (1982) [*Russ. Chem. Rev.* **51**, 765 (1982)].
- ²³³K. Hagen, L. Hedberg, and K. Hedberg, *J. Phys. Chem.* **86**, 117 (1982).
- ²³⁴F. A. L. Anet and L. Kozerski, *J. Am. Chem. Soc.* **95**, 3407 (1973).
- ²³⁵N. L. Allinger and J. T. Sprague, *Tetrahedron* **31**, 21 (1975).
- ²³⁶O. Ermer, *J. Am. Chem. Soc.* **98**, 3964 (1976).
- ²³⁷G. G. Barna and I. S. Butler, *J. Raman Spectrosc.* **7**, 168 (1978).
- ²³⁸D. W. Wertz and M. A. Moseley, *Inorg. Chem.* **19**, 705 (1980).
- ²³⁹N. Neto, C. Di Lauro, and S. Califano, *Spectrochim. Acta Part A* **24**, 385 (1968).
- ²⁴⁰O. Bastiansen, L. Hedberg, and K. Hedberg, *J. Chem. Phys.* **27**, 1311 (1957).
- ²⁴¹M. Traetteberg, *Acta Chem. Scand.* **20**, 1724 (1966).
- ²⁴²F. A. L. Anet, *J. Am. Chem. Soc.* **84**, 671 (1962).
- ²⁴³G. Wipff, U. Wahlgren, E. Kochanski, and J. M. Lehn, *Chem. Phys. Lett.* **11**, 350 (1971).
- ²⁴⁴N. L. Allinger, J. T. Sprague, and C. J. Finder, *Tetrahedron* **29**, 2519 (1973).
- ²⁴⁵Z. B. Maksic and A. Rubčić, *J. Am. Chem. Soc.* **99**, 4233 (1977).
- ²⁴⁶G. Buemi, A. Nicolosi, F. Zuccarello, and D. Grasso, *J. Mol. Struct.* **55**, 39 (1979).
- ²⁴⁷F. Zuccarello, G. Buemi, and A. Raudino, *J. Comput. Chem.* **1**, 341 (1980).
- ²⁴⁸E. R. Lippincott, R. C. Lord, and R. S. McDonald, *J. Am. Chem. Soc.* **73**, 3370 (1951).
- ²⁴⁹E. M. Popov and G. A. Kogan, *Zh. Strukt. Khim.* **81**, 476 (1967).
- ²⁵⁰M. Traetteberg, G. Hagen, and S. J. Cyvin, *Z. Naturforsch. Teil B* **25**, 134 (1970).
- ²⁵¹D. W. Scott, M. E. Gross, G. D. Oliver, and H. M. Huffman, *J. Am. Chem. Soc.* **71**, 1634 (1949).
- ²⁵²E. R. Lippincott and R. C. Lord, *J. Am. Chem. Soc.* **73**, 3889 (1951).
- ²⁵³D. R. Stull, E. F. Westrum, Jr., and G. C. Sinke, *The Chemical Thermodynamics of Organic Compounds* (Wiley, New York, 1969).
- ²⁵⁴G. B. Kistiakowsky and W. W. Rice, *J. Chem. Phys.* **8**, 610 (1940).
- ²⁵⁵G. B. Kistiakowsky, J. R. Lacher, and W. W. Ransom, *J. Chem. Phys.* **8**, 970 (1940).
- ²⁵⁶K. B. Wieberg, R. C. Dempsey, and J. J. Wendoloski, *J. Phys. Chem.* **88**, 5596 (1984).
- ²⁵⁷S. Saebø, F. R. Cordell, and J. E. Boggs, *J. Mol. Struct.* **104**, 221 (1983).
- ²⁵⁸R. Spitzer and K. S. Pitzer, *J. Am. Chem. Soc.* **68**, 2537 (1946).
- ²⁵⁹J. G. Aston, S. C. Schumann, H. L. Fink, and P. M. Doty, *J. Am. Chem. Soc.* **63**, 2029 (1941).
- ²⁶⁰T. Ueda and T. Shimanouchi, *J. Chem. Phys.* **47**, 5018 (1967).
- ²⁶¹J. R. Villarreal, L. E. Bauman, J. Laane, W. C. Harris, and S. F. Bush, *J. Chem. Phys.* **63**, 3727 (1975).
- ²⁶²J. G. Aston, G. J. Szasz, and H. L. Fink, *J. Am. Chem. Soc.* **65**, 1135 (1943).
- ²⁶³P. M. Ivanov, and E. Ōsawa, *J. Comput. Chem.* **5**, 307 (1984).
- ²⁶⁴G. Favini, A. Maggi, and R. Todeschini, *J. Mol. Struct.* **105**, 17 (1983).