

# Cross Sections for K-shell X-ray Production by Hydrogen and Helium Ions in Elements from Beryllium to Uranium

Cite as: Journal of Physical and Chemical Reference Data **18**, 111 (1989); <https://doi.org/10.1063/1.555838>  
Submitted: 19 November 1986 . Published Online: 15 October 2009

G. Lapicki



[View Online](#)



[Export Citation](#)

## ARTICLES YOU MAY BE INTERESTED IN


[Present status of L-shell X-ray production cross sections by protons with energies below 1 MeV](#)  
AIP Conference Proceedings **576**, 100 (2001); <https://doi.org/10.1063/1.1395259>

[A Review, Bibliography, and Tabulation of K, L, and Higher Atomic Shell X-Ray Fluorescence Yields](#)

Journal of Physical and Chemical Reference Data **23**, 339 (1994); <https://doi.org/10.1063/1.555955>

[Calculation of Intermolecular Force Constants from Polarizabilities](#)

The Journal of Chemical Physics **24**, 501 (1956); <https://doi.org/10.1063/1.1742536>



# Cross Sections for K-shell X-ray Production by Hydrogen and Helium Ions in Elements from Beryllium to Uranium

G. Lapicki

Department of Physics, East Carolina University, Greenville, North Carolina 27858

Received November 19, 1986; revised manuscript received August 8, 1988

Experimental cross sections for *K*-shell x-ray production by hydrogen and helium ions ( $Z_1 = 1,2$ ) in target atoms from beryllium to uranium ( $Z_2 = 4-92$ ) are tabulated as compiled (7418 cross sections) from the literature (161 references were found) with the search for the data terminated in January 1988. These cross sections are compared with predictions of the first Born approximation and ECPSSR theory for inner-shell ionization. The ECPSSR accounts for the energy loss (E) and Coulomb deflection (C) of the projectile ion as well as for the perturbed stationary state (PSS) and relativistic (R) nature of the target's inner-shell electron. While the first Born approximation generally overestimates the data by orders of magnitude, the ECPSSR theory is confirmed to be, on the average, in agreement with the experiment to within 10%–20%. For light and heavy target atoms, however, systematic and opposite deviations are found in the low projectile-velocity regime. These deviations are associated with the influence of multiple outer-shell ionizations on the fluorescence yields of light elements, particularly in ionization by helium ions, and with the inaccuracy of the ECPSSR theory in the reproduction of relativistic calculations for ionization of heavy elements. The remaining discrepancies at moderate projectile velocities are *prima facie* attributed to inadequacies of a screened hydrogenic description for the *K*-shell electron.

Key words: *K*-shell x-ray production cross sections; *K*-shell ionization; Born approximation; ECPSSR theory; H ions; He ions.

## CONTENTS

1. Introduction .....	112	3.3. Comparison of the experimental and ECPSSR cross sections .....	118
2. Experimental data base .....	113	4. Conclusions .....	121
2.1. Search procedures .....	113	5. Acknowledgments .....	121
2.2. Summary of data base .....	113	6. References .....	211
2.2. a. Compiled <i>K</i> -shell x-ray production cross sections .....	113	6.1. Text references .....	211
2.2. b. Units .....	113	6.2. References to cross-section data compiled in Tables 2–5 .....	212
2.3. Growth and decline in annual publication of data .....	114	6.3. Author index for the data base references in Sec. 6.2 .....	215
3. Data analysis .....	114		
3.1. Ionization cross sections .....	114		
3.1. a. Conversion of ionization cross section to x-ray production cross section .....	114	<b>List of Tables</b>	
3.1. b. Ionization, as the sum of direct ionization and electron capture, in the first Born and ECPSSR theories .....	115	1. Distribution by target element of <i>K</i> -shell x-ray production cross sections compiled from the literature .....	122
3.2. Choice of the ECPSSR for theoretical analysis of the data .....	115	2. <i>K</i> -shell x-ray production by protons in target elements from beryllium to uranium .....	124
3.2. a. Review and general scaling of the ECPSSR .....	115	3. <i>K</i> -shell x-ray production by deuterons in target elements from beryllium to gold .....	170
3.2. b. Current status, alternatives, advantages and shortcomings of the ECPSSR .....	116	4. <i>K</i> -shell x-ray production by ${}^3\text{He}$ in target elements from aluminum to silver .....	177
		5. <i>K</i> -shell x-ray production by ${}^4\text{He}$ in target elements from beryllium to uranium .....	181
		6. Number of <i>K</i> -shell x-ray production cross sections compiled for each projectile and tabulated for each target element .....	202

©1989 by the U. S. Secretary of Commerce on behalf of the United States. This copyright is assigned to the American Institute of Physics and the American Chemical Society. Reprints available from ACS; see Reprints List at back of issue.

7. Contribution of electron capture to ionization according to the ECPSSR theory .....	210
--	-----

### List of Figures

1. Histogram of data for $K$ -shell x-ray production by H and He ions .....	114
2. $K$ -shell x-ray production in nickel by protons as a function of the projectile's velocity scaled by the electron velocity in $K$ -shell orbit of the target .....	114
3. Ratios of experimental cross sections to the first Born approximation for protons incident on nickel .....	116
4. Ratios of experimental cross sections to the ECPSSR for protons incident on nickel.....	116

5. Averaged ratios of experimental cross sections to the first Born calculations for protons .....	118
6. Averaged ratios of experimental cross sections to the ECPSSR predictions for protons .....	118
7. Averaged ratios of experimental cross sections to the ECPSSR predictions for deuterons .....	119
8. Averaged ratios of experimental cross sections to the ECPSSR predictions for $^3\text{He}$ ions .....	119
9. Averaged ratios of experimental cross sections to the ECPSSR predictions for $^4\text{He}$ ions .....	119
10. Same as Fig. 6 but versus the variable $\xi_K^R/\xi_K$ according to which the ECPSSR ionization cross sections scale in the slow collision regime.....	120
11. Same as Fig. 9 but versus the variable $\xi_K^R/\xi_K$ according to which the ECPSSR ionization cross sections scale in the slow collision regime.....	120

## 1. Introduction

Fifteen years ago, Rutledge and Watson<sup>1</sup> originated extensive tabulations of inner-shell cross sections by ionic projectiles in target atoms which cover most of the periodic table. Their compilation was restricted to  $K$ -shell ionization by H and He ions and reported some 600 x-ray production cross sections in 1973. In a 1978 sequel to it, Gardner and Gray<sup>2</sup> extended this compilation to  $\sim 1200$  x-ray production cross sections by H and He ions. This extension covered  $K$ -shell ionization cross sections by heavier ions than helium as well; compilations of  $L$ -shell ionization data also exist.<sup>3</sup>

One could hence speculate that the number of  $K$ -shell x-ray production cross sections by H and He as reported in the literature doubled in a five-year period. To an extent that a constant fraction of all publications on inner-shell ionization phenomena contains such data, we could confirm this speculation. A histogram of publications cited in a 1973 thesis<sup>4</sup> on inner-shell ionization showed an exponential increase in these articles per annum since 1960; the growth rate was constant and indeed such that the number of publications per year has doubled every half of a decade.

Continued updates of these data, as carried almost single-handedly by Paul and co-workers<sup>5-14</sup> since 1978, appear also to be characterized by a rapid increase in their amount. In his 1984 analysis,<sup>8</sup> Paul uses some 3200 cross sections from the literature for protons alone. In an attempt to unravel systematic trends in such a mass of experimental data, Paul *et al.*<sup>5-14</sup> normalize the data to theoretical predictions of the ECPSSR theory for direct  $K$ -shell ionization.<sup>15</sup> The ECPSSR theory for both direct ionization<sup>15</sup> and electron capture<sup>16</sup> accounts for the energy loss (E) and Coulomb deflection (C) of the projectile, and for the perturbed stationary state (PSS) and relativistic (R) changes in the description of the inner-shell electron that undergoes ionization. In our original analysis,<sup>15</sup> we scaled  $\sim 2600$   $K$ -shell x-ray production cross sections to the results of this theory. The devia-

tions of experiment from the ECPSSR theory<sup>15</sup> were found to be within 10% once all the data were considered equally and averaged in the preselected equal intervals of the effective projectile energy-loss variable. Such discrepancies, being comparable to experimental uncertainties, appeared to be acceptable. Analyses<sup>5-14</sup> analogous to the analysis of Ref. 15 subsequently revealed, however, that the ECPSSR theory systematically overestimates the data in the slow collision regime after the proton measurements from 21 out of 77 references were rejected. Similar deviations were observed for deuteron and helium data<sup>8</sup> after the data from 22 out of 55 references were discarded according to the adopted rejection criterion.<sup>6,7</sup> This finding was confirmed with an updated 1986 compilation<sup>12</sup> that contains almost 4000 proton cross sections from 101 references and nearly 1800 alpha particle cross sections from 47 references.

Previous authors either reported compiled cross sections in a tabular form without theoretical scrutiny<sup>1-3</sup> or analyzed them, without listing of the data, through graphical comparisons with the predictions of theories.<sup>5-14</sup> In this work, both a compilation (Sec. 2 and Tables) and an analysis (Sec. 3 and Figs.) are given. Two motivating goals for the present article are: (i) the need for an update of the last tabular report of the data<sup>2</sup> because the number of available cross sections has multiplied sixfold since then, (ii) the desire for an evaluation of the ECPSSR theory vis-à-vis an expanded data base; this evaluation, being independent and methodologically slightly different than critical analyses by Paul *et al.*,<sup>5-15</sup> might be of interest to those who choose to compare the ECPSSR theory with experiment. Also, brief comments that go beyond raw data presentation and their conventional evaluation in the framework of chosen theories are made; the data growth is a good indicator of the dynamic evolution in the field of inner-shell ionization. Such a discussion could offer a useful glimpse at the changing status of this field to those readers who may not be directly involved in it and might be even of vital interest to the researchers who are immersed in this field.

## 2. Experimental data base

### 2.1. Search procedures

All compiled cross sections were taken from the tables from referenced articles or privately communicated by authors of the article. When the tables and authors were not available, the data were read off graphs with the accuracy of two significant figures. All cross sections are reported in this work in a three-digit format even though occasionally original sources published them in larger formats. Uncertainties of the order of 10% in the modern day measurements of these cross sections restrict their significance to, at most, a three-digit accuracy. Errors, found in the original literature by Paul and Muhr,<sup>12</sup> were corrected prior to the accumulation of present data base. All compiled data were stored on disk files in the chronological order for easy updates. These files were spot checked against source papers for possible misprints in transfer to computer files; a few coauthors of source references have kindly provided this author with a check of his printout of their data. The last update of the data was made during the summer of 1988 with January 1988 terminating the data search.

### 2.2. Summary of data base

#### 2.2.a. Compiled K-shell x-ray production cross sections

Table 1 gives a summary of the distribution of 7418 compiled cross sections with respect to the target atomic number  $Z_2$  for each of the projectiles (protons, deuterons,  $^3\text{He}$ ,  $^4\text{He}$ ) separately as well as, cumulatively, for all projectiles ( $Z_1 = 1, 2$ ). It allows a global assessment of the availability of the data for a specific projectile-target combination as well as for a given target and all projectiles. In particular, this table identifies (by contrast with the bold print used for the  $Z_2$  targets that appear in the compilation) the 15 elements for which no data were found in the  $4 \leq Z_2 \leq 92$  range and it singles out copper as the most often (9% of all data) used target for  $K$ -shell x-ray measurements by H and He ions.  $K$ -shell x-ray production cross sections induced by protons, deuterons,  $^3\text{He}$ , and  $^4\text{He}$  ions are compiled in Tables 2–5, respectively. They are listed with the increasing atomic number  $Z_2$  of the target atom which is also identified by name. For each element, the data appear according to the chronological order of the reference of their origin and, for each reference, they are listed with the increasing energy<sup>17</sup> of the projectile.

Tables 2–5 contain 7418 cross sections of which 63% are by protons, 26% by  $^4\text{He}$ , 7% by deuterons, and 4% by  $^3\text{He}$ . The data are from 161 references that are listed chrono-

logically in a separate reference section which lists these source references (see Sec. 6.2). A contact between the references and Table 1, which provides only a summary of the data base content, is made in Table 6. Table 1 shows a distribution of all compiled cross sections with respect to the projectile and target atoms. Table 6 presents this information by identifying the reference from which the data were obtained; the correlation of the number of reported cross sections for a given projectile-target system with the reference number serves a twofold purpose: (1) to exhibit the rate of growth in accumulation of the data with time since references are arranged chronologically and (2) to find all references pertaining to the given projectile-target combination. This overview of data distribution gives a quick perspective on the dynamics with which the data appear in the literature for a selected projectile-target combination. It offers a detailed look at the regions of the periodic table that remain almost uncharted to experimental studies of  $K$  x-ray production by light ions; references identify the researchers who pioneered investigations in these nearly *tabula rasa* regions.

This article ends with an author index (see Sec. 6.3), which is keyed to the reference numbers appearing in Tables 2–6 so that an easy reference exists to the names of all of those who reported the compiled data. The reference numbers which follow given names of particular authors place their research activity in a historical context since the references are ordered chronologically. Anyone interested in contributions of a particular author to the compiled data can trace them easily with the aid of Table 6.

#### 2.2.b. Units

In Tables 2–5, each data set from a given reference consists of pairs: the energy of the projectile in MeV ( $1.6 \times 10^{-13} \text{ J}$ ) and the experimental x-ray production cross section in barn ( $10^{-28} \text{ m}^2$ ). The conventional units of the accelerator-based physics are used to report the data in these Tables because such units are employed in the source literature (SI equivalents of these units are stated in the parentheses). Velocities of the projectile and of the target  $K$ -shell electron are calculated in terms of  $v_0 = e^2/\hbar$ , the Bohr velocity ( $2.2 \times 10^6 \text{ m/s}$ ) of the electron in the ground state of the H atom. In this atomic unit of velocity, the target  $K$ -shell electrons orbit at  $v_{2K} = Z_{2K}$  where  $Z_{2K} = Z_2 - 0.3$  is the electric charge, in units of 1 because one is the magnitude of the electron charge in atomic units, of the target nucleus diminished by Slater's screening constant. In Figs. 3–11, the choice of units is immaterial because dimensionless ratios are plotted along each axis. The parameters that define  $\xi_K^R/\zeta_K$ , the scaling variable of the ECPSSR theory, are dimensionless (See Sec. 3.2.a. and Figs. 10 and 11).

### 2.3. Growth and decline in annual publication of data

Figure 1 shows a histogram of the data compiled in this article: the annual number of cross sections published in a given year is shown. It appears that the rapid rate of growth of the 1960s and early 1970s rose to a maximum in the late 1970's. The annual rate at which the cross sections were reported in the current decade is on the decline. If this trend continues, the total cumulative number of cross sections is destined to reach a saturated value of some 10 000.

This forecast does not mean that the research on inner-shell ionization processes slides down toward its nadir; the annual number of publications in this field continues to double every five years. Rather it is the specialized area of inner-shell ionization research, as measured by the amount of new  $K$ -shell x-ray production cross sections by light ions, that shrinks. Experimental and theoretical interests shift now toward problems of inner-shell ionization in which  $Z_1/Z_2$ , the ratio of projectile-to-target atomic numbers, approaches 1. Also, as investigations of the  $K$  shell in very asymmetric ( $Z_1/Z_2 \ll 1$ ) collisions become less fashionable, the current research on such collisions gives more prominence to studies of  $L$ - and  $M$ -shell ionizations.

### 3. Data analysis

No attempt is made here to report the experimental errors as stated in the original papers. Often estimates of such errors are not consistent, ranging from 5% to 35% amongst various experimental groups even though the experiments were performed apparently under similar conditions. Less often, but most shockingly, the data for the iden-




FIG. 1. Histogram of data for  $K$ -shell x-ray production by  $H$  and  $He$  ions (see Sec. 2.3). The vertical lines indicate the annual number of published cross sections as compiled in this work; the solid circles correspond to the cumulative number of these cross sections as they appeared up to a given year.




FIG. 2.  $K$ -shell x-ray production in nickel by protons as a function of the projectile's velocity scaled by the electron velocity in  $K$ -shell orbit of the target. Data are from Refs. 5, 20, 36, 47, 52, 55, 57, 69, 73, 76, 77, 84, 89, 94, 97, 108, 113, 114, 115, 120, 122, 132, 137, and 151 from the list of source references (see Sec. 6.2). The curves are based on the first Born (Refs. 20 and 21: dashed curve) and the ECPSSR (Refs. 15 and 16: solid curve) theories.

tical collision systems are found to differ by a significantly larger margin of error than the claimed experimental uncertainties would imply<sup>18</sup>; in rare instances such data disagree by even more than a factor of 2. Hence, although, justifiably due to constant improvements in data gathering techniques, 25% uncertainties are quoted in older references and 10% uncertainties are claimed in recent articles, we assign equal weights to all data at the outset of our analysis.

Figure 2 shows the cross sections for protons on nickel, one of the most often used materials in the  $K$ -shell x-ray production measurements. These cross sections increase by as much as nine orders of magnitude with the projectile energy, labeled at the top of the figure in MeV. They exhibit a general trend in all data in that the cross sections peak where the velocity of the projectile  $v_1$ , matches approximately the orbital velocity of the  $K$ -shell electron in the target atom  $v_{2K} = Z_2 - 0.3$ .

#### 3.1. Ionization cross sections

##### 3.1.a. Conversion of ionization cross section to x-ray production cross section

Experimental x-ray production cross sections  $\sigma_{KK}^{Exper.}$  can be compared with theoretical x-ray production cross sections  $\sigma_{KK}$ , after the ionization cross section  $\sigma_K$  is multiplied by the fluorescence yield  $\omega_K$ , i.e.,  $\sigma_{KK} = \sigma_K \omega_K$ . Throughout this work we use the single-vacancy fluorescence yields and employ for them the values as recommend-

ed by Krause<sup>19</sup> and listed in Tables 2–5. Multiple ionizations increase  $\omega_K$  with the increasing  $Z_1/v_1$ . They do this, however, insignificantly (less than a percent) in K-shell ionization of heavier elements by light (hydrogen, helium) ions, in which  $Z_1/v_1 \approx (Z_1/Z_2)/(v_1/v_{2K})$  is small even at low projectile velocities. Only small fluorescence yields ( $\omega_K < 0.02$  for  $Z_2 < 10$ ) are appreciably altered due to multiple ionization, more so in ionization by helium ( $Z_1 = 2$ ) ions for which the condition of, say,  $Z_1/Z_2 \geq 0.15$  covers twice as large a range of light elements. For such collision systems, theoretical x-ray production cross sections will be somewhat underestimated because the use of single-hole  $\omega_K$  values. It should be noted that even single-hole fluorescence yields are in 10%–40% error for these relatively light target atoms.<sup>19</sup> The deviations become dramatic with increasing  $Z_1/Z_2$  so that comparison of the theoretical predictions with the  $^2\text{He}$  on  $^4\text{Be}$  data ( $Z_1/Z_2 = 1/2$ ) is the most problematic.

### 3.1.b. Ionization, as the sum of direct ionization and electron capture, in the first Born and ECPSSR theories

Ionization cross sections are obtained according to the first Born approximation<sup>20,21</sup> [ $\sigma_K^{\text{FBORN}}$ , as shown by dashed curve in Fig. 2, consists of direct ionization and electron capture calculated in the plane wave Born approximation (PWBA)<sup>20</sup> and the Oppenheimer-Brinkman-Kramers treatment,<sup>21</sup> respectively], and the ECPSSR theory<sup>15,16</sup> ( $\sigma_K^{\text{ECPSSR}}$ , solid curve in Fig. 2). In both calculations, ionization cross sections  $\sigma_K$  are taken as a sum of the cross sections for direct ionization to the target atom continuum plus electron capture to all bound states on the projectile. Although electron capture gives an additional contribution to ionization, the confusion in the literature exists because many authors still refer to ionization cross sections when only direct ionization cross sections are calculated. This unfortunate error of terminology can be found in particular in the most recent references to ECPSSR calculations.<sup>12,13,22,23</sup> We define and, as a matter of principle, calculate the ECPSSR ionization cross sections always as a sum of the direct ionization<sup>15</sup> and electron capture<sup>16</sup> cross sections, i.e.,

$$\begin{aligned} \sigma_K^{\text{ECPSSR}} = & \sigma_K^{\text{ECPSSR}}(\text{DIRECT IONIZATION}) \\ & + \sigma_K^{\text{ECPSSR}}(\text{ELECTRON CAPTURE}). \quad (1) \end{aligned}$$

Although electron capture has negligible contribution to ionization when  $Z_1/Z_2$  is small, we evaluate the ECPSSR ionization cross sections using Eq. (1) for all  $Z_1/Z_2$  projectile-target combinations. Table 7 states the percentage contributions of electron capture to ionization as calculated in the ECPSSR theory.<sup>15,16</sup> Electron capture can contribute more than 1% when  $Z_1/Z_2 \geq 1/15$  and the projectile energy per its mass is below 10 MeV/u. Table 7 lists these percentages only for protons and alpha particles because the electron capture contributions are essentially independent of the isotope nature of the projectile at a given velocity. The projectile is assumed to be fully stripped in these calculations (all its states are unoccupied and there are no electrons to screen it); this represents the condition under which most of the data were gathered. Some data were specifically reported for  $\text{He}^+$ ; in many articles, however, the charge was unspeci-

fied. Contribution of electron capture to total ionization cross sections is calculated in the ECPSSR theory to be at most 5% when  $Z_1/Z_2 \leq 0.15$  and for fully stripped projectiles, and it would be approximately one-half of that if the projectile were assumed to carry an electron into the collision. Hence, calculations which always presume a fully stripped projectile overestimate the ionization process by no more than a few percents if  $Z_1/Z_2 \leq 0.15$ . For protons on nickel ( $Z_1/Z_2 = 0.036$ ) data of Fig. 2 electron capture contributes less than 0.1% to ionization. For  $Z_1/Z_2 > 0.15$  collision systems, theoretical x-ray production cross sections used in this work are underestimated because single-hole  $\omega_K$  values were employed and, sometimes, these cross sections are overestimated because a fully stripped projectile was always assumed. These deviations become dramatic with the increasing  $Z_1/Z_2$  so that the comparison of the theoretical predictions with the  $^2\text{He}$  on  $^4\text{Be}$  data ( $Z_1/Z_2 = 1/2$ ) is the most problematic. We assume, however, that the ignored effect of multiple ionization and an overestimated<sup>16,24</sup> contribution of electron capture at  $Z_1/Z_2 \rightarrow 1/2$  tend to cancel each other to a great extent.

### 3.2. Choice of the ECPSSR for theoretical analysis of the data

Figure 2 demonstrates that, while the first Born approximation  $\sigma_K^{\text{FBORN}}$  overestimates the proton on nickel data by as much as three orders of magnitude at lowest proton velocities,  $\sigma_K^{\text{ECPSSR}}$  appears to be in good agreement with the measured cross sections. To exhibit these findings in a more refined way, unobscured by the artifact of a log–log graphical comparison, we plot the same data as the ratios of experimental cross sections  $\sigma_{KX}^{\text{Exper.}}$  to theoretical predictions in Fig. 3 for the first Born approximation and in Fig. 4 for the ECPSSR theory. To make a complete and compact comparison with all compiled data, the data are grouped in equal (0.1 in length) intervals on the  $\log(v_1/v_{2K})$  scale. An arithmetic average of all cross sections in each group so defined is calculated, all data within the group that differ from this average by more than a factor of 2 are rejected, a new average for the group is found, and the rejection is made again from all the data in the group (including previously eliminated data) on the basis of the same criterion. Typically in two but no more than three iterations of this procedure the averages converge to constant values which are plotted in Figs. 3 and 4 for our example of K-shell x-ray production by protons in just one target element.

The success and relative ease in the implementation<sup>23</sup> of the ECPSSR theory, lead to its adoption as a theoretical benchmark for further analysis of the compiled data. A self-contained and critical<sup>25</sup> review of this theory is in order; development, scaling properties, and current status of the ECPSSR theory with alternative treatments is presented to justify a selection of this particular approach to inner-shell ionizations. The ECPSSR theory is reviewed *vis à vis* the first Born approximation and more *ab initio* theoretical approaches to inner-shell ionization.




FIG. 3. Ratios of experimental cross sections to the first Born approximation for protons incident on nickel. Each step in the staircase curve represents the arithmetic average of all ratios found in the corresponding interval of  $v_1/v_{2K}$ .

### 3.2.a. Review and general scaling of the ECPSSR

A reduction of the discrepancies between the first Born approximation and the experiment occurs because the ECPSSR theory accounts for the binding effect that, being important at low projectile velocities and for large  $Z_1/Z_2$ , inhibits ionization and results in lower cross sections than the first Born approximation. Also, the ECPSSR approach corrects for the Coulomb-deflection of the projectile from a straight-line trajectory and considers the projectile energy loss exactly in the minimum momentum transfer; both corrections lead to smaller cross sections. The underestimation of the data in the first Born approximation for ionization of heavy target elements (large  $Z_2$ 's mean small  $Z_1/Z_2$ ) stems from its nonrelativistic treatment of the  $K$ -shell electron. The ECPSSR theory attempts to remedy this shortcoming by accounting for the relativistic effect and indeed by bringing the calculations in closer agreement with the data.

The ECPSSR theory originates with the work of Brandt, Laubert, and Sellin<sup>26</sup> who accounted for the increased binding and Coulomb-deflection effects in  $K$ -shell ionization. An extension of this work to the  $L$  shell was made<sup>27</sup> and subsequently, after a theoretical basis for the perturbed stationary-state (PSS) approach was established,<sup>28</sup> polarization<sup>29,30</sup> and relativistic<sup>30</sup> effects were included in the CPSSR theory<sup>30</sup> as a precursor of the ECPSSR approach<sup>15</sup> which also accounts for the projectile-energy loss. This theory was developed for electron capture in Ref. 16 in a similar manner as for direct ionization in Ref. 15. The ECPSSR theory for  $K$ - and  $L$ -shell ionization has been also extended to the  $M$  shell.<sup>31</sup>




FIG. 4. Ratios of experimental cross sections to the ECPSSR for protons incident on nickel. Each step in the solid curve represents the arithmetic average of all ratios found in the corresponding interval of  $v_1/v_{2K}$ ; the mean value for all proton on nickel ratios is 0.96.

In the slow collision limit, the calculations of the first Born approximation—for direct ionization which generally dominates over electron capture—scales with  $\xi_K = 2v_1/v_{2K}\theta_K$  where  $\theta_K$  is defined as the observed binding energy in terms of screened hydrogenic value  $\frac{1}{2}Z_{2K}^2$ . In the ECPSSR theory,<sup>15</sup>  $\xi_K$  is replaced by  $\xi_K^R/\xi_K$  to correct<sup>30</sup> the first Born approximation for the relativistic and perturbed stationary-state effects;  $\xi_K$  is replaced with  $\xi_K^R = [m_K^R(\xi_K/\xi_K)]^{1/2}\xi_K$  to simulate the relativistic effect<sup>32</sup> and  $\xi_K$  accounts for the PSS effect according to Eq.(20) of Ref. 30. After the analytically known<sup>15</sup> functions that correct for the projectile's energy loss and Coulomb deflection are factored out, all cross sections are reduced<sup>27,33</sup> in the slow collision limit to  $F_K$ , a universal function of  $\xi_K^R/\xi_K$ . For  $\xi_K^R/\xi_K > 1$ ,  $F_K$  diverges from this form depending on  $\xi_K\theta_K$ . However, to the extent that  $\xi_K\theta_K$  does not (except for very light targets) vary significantly, the ionization cross section remains to a good approximation a universal function of  $\xi_K^R/\xi_K$  in all collisional regimes. This enables us to group  $K$ -shell x-ray production cross sections according to the  $\xi_K^R/\xi_K$  parameter for a comprehensive analysis of the compiled data against the predictions of the ECPSSR theory.

### 3.2.b. Current status, alternatives, advantages and shortcomings of the ECPSSR

The strength of the ECPSSR calculations lies in the relative ease with which this approach allows to incorporate analytically relevant physical effects into formulas of the first Born approximation for the ionization cross section; the

role that these effects play can be recognized without being entangled in intricacies of the second or distorted Born approximation which requires a considerable numerical effort. Nevertheless, as an approximate description of an inelastic collision process, the ECPSSR theory is yet to be fully tested by more involved numerical procedures. It is hoped that with the phenomenal progress in computerized techniques such procedures will emerge as an penultimate check of the ECPSSR theory as well as its sophisticated replacements. The ultimate test for any theory will be in comparison of its predictions with experimental results.

Coulomb-deflection and PSS factors derived in the ECPSSR treatment have been utilized to modify first-order perturbation theories such as the binary encounter approximation (BEA).<sup>34</sup> We have stated previously<sup>4,35</sup> that the incorporation of the essentially quantum-mechanically derived correction factors into the BEA cross section, which equals the PWBA cross section under very restrictive conditions,<sup>36</sup> is not proper. Even in semiclassical and quantum approximations a selective use of just one of the ECPSSR factors might be questioned, especially when corrections for other effects are made on the basis of older<sup>26,27,29,30</sup> or different<sup>14,37-39</sup> accountings for the C, PSS, and R effects. An obvious example of misapplication<sup>40</sup> of the ECPSSR theory has been discussed elsewhere.<sup>41</sup> The Coulomb-deflection factor of the ECPSSR approach has been extensively used by Chen and Crasemann<sup>22,43-45</sup> in calculations that employ the united atom binding energy to simulate the PSS effect but take the energy loss and relativistic effects into account *ab initio*. These numerical calculations allow for exact limits for the momentum transfers and use relativistic wavefunctions based on the screened hydrogenic<sup>42</sup> or Hartree-Slater<sup>22,43-45</sup> potential. The *K*- and *L*-shell direct ionization calculations<sup>22,43</sup> were extended to the *M* shell<sup>44</sup> and even to the *N* shell.<sup>45</sup> The ECPSSR theory has been utilized in numerous comparisons with experimental inner-shell ionization cross sections. Predictions of the ECPSSR approach and its predecessors<sup>26,27,29,30</sup> were also used in (i) generation of proton-induced x-ray emission (PIXE) spectra,<sup>46</sup> (ii) calculation of relative *L*-shell x-ray intensities,<sup>47</sup> (iii) absolute calibration of the efficiency for semiconductor detectors,<sup>48</sup> (iv) alignment studies,<sup>49</sup> (v) semiempirical extraction of *L*-shell fluorescence yields,<sup>50</sup> and (vi) discussion of the feasibility of an antiproton detector.<sup>51</sup> The ECPSSR theory was employed in the determination of semiempirical formulas for *K*-shell ionization.<sup>14,52</sup>

In this work we calculate the ECPSSR ionization cross sections as stated in Refs. 15 and 16, although some improvements have been suggested since these references were published. Rigorous, numerical *ab initio* calculations and comprehensive comparisons with all inner-shell ionization data will decide whether nonadiabatic extinctions<sup>53</sup> of the PSS approach are warranted. Coupled-state calculations are still in development. Their reliance always hinges on a clever choice of a set of basis states. Optimal selections have to be large enough to account for the physics of a collision and yet sufficiently small to be computationally manageable. A coupled-state calculation by Reading *et al.*<sup>54</sup> that utilizes the so-called

forced impulse approximation and claims to conquer the slow collision regime has been carried out only at the first Born approximation level.

Unfortunately, the suggestion<sup>10</sup> that one should "investigate various effects theoretically since it is much easier to turn an effect on or off in a computer experiment than in nature" cannot be as yet carried out in practice. A "highly sophisticated computer program"<sup>10</sup> that could control all ECPSSR effects *ab initio* in any collision regime does not exist. While some calculations from the outset incorporate the E and R effects<sup>22,42-45</sup> and also account semiclassically for the Coulomb deflection,<sup>55,56</sup> they treat the PSS effect using sometimes<sup>55</sup> the old prescription of Ref. 26 or making<sup>56</sup> the united atom approximation which applies only in the strict limit of low projectile velocities. While other schemes<sup>57</sup> perform admirably to test the E, C and PSS effects, they were implemented only with nonrelativistic wave functions. Perhaps the closest to rigorous numerical test of all E, C, PSS, and R factors are the codes of Trautmann and co-workers<sup>58</sup>; they still, however, make ad hoc modifications to simulate the PSS effect. This effect is clearly seen in the *ab initio* coupled-state calculation Mehler *et al.*<sup>59</sup> that uses relativistic wave functions and offers promise; however, it is difficult to judge the outcome of this scheme because only one graph for *K*-shell ionization of silver by 0.9-MeV protons was presented<sup>59</sup> and in the subsequent paper only the probability for *K*-shell ionization is reported.<sup>60</sup> In accounting for PSS effects, this calculation gives a 20% reduction of the direct-ionization cross section as opposed to the ECPSSR approach that predicts only a few percent decrease of  $\sigma_K$  for the analyzed collision. This would be in agreement with Kocbach, who has concluded<sup>61</sup> that the ECPSSR treatment underestimates the role of the binding effect.<sup>26</sup> Mukoyama and Lin,<sup>62</sup> with an expansion of the relativistic wave function into Slater-type orbitals, have evaluated cross sections for *K*-shell ionization of copper by 0.5–2 MeV protons. These calculations, just as those of Refs. 59 and 60, lie ~15% below the ECPSSR results. Anholt *et al.*<sup>63</sup> have recommended that the cutoff impact parameter below which binding occurs be doubled; this would lower the ECPSSR cross sections, especially around their maxima, and thus would bring them in agreement with Refs. 59–62.

Sarkadi,<sup>64</sup> accounting for the nonadiabaticity of PSS states, finds contrary to Anholt's recipe<sup>63</sup> that the binding effect should have been deemphasized outside the slow collision regime; when  $v_1$  approaches  $v_{2K}$ , the *K*-shell does not adjust adiabatically and hence the binding effect should not be as large as the ECPSSR has it. This would increase the ECPSSR cross section around its peak, and thus widen the existing disagreement with Refs. 59–62. The coupled-state calculations of Mehler *et al.*<sup>59,60</sup> explain an enhancement of ionization, which counters the effect of the increased binding, as an effect of interaction among the continuum states, while the approach of Brandt *et al.*<sup>29,30</sup> traces the increase in ionization cross sections to the polarization of the bound state. A variational PSS description<sup>65</sup> of the polarization effect<sup>29,30</sup> determines that the ECPSSR underestimates as well this antibinding effect. Modifications suggested in Refs. 61

and 65 appear to cancel each other and thus they mask possible overall inadequacies in the ECPSSR treatment of the PSS (combined account for binding and polarization) effect. We now turn to the ultimate test of any theory, i.e., a broad comparison of its predictions with experimental observations.

### 3.3. Comparison of experimental and ECPSSR cross sections

In the pursuit of systematic discrepancies between the data and the predictions of the ECPSSR theory as  $Z_2$ -dependent deviations, we classify somewhat arbitrarily all elements as: light ( $4 \leq Z_2 \leq 13$ ), medium ( $14 \leq Z_2 \leq 66$ ), and heavy ( $67 \leq Z_2 \leq 92$ ). Note that this classification assigns  $Z_1/Z_2 > 0.15$  for the light atoms and  $Z_1/Z_2 < 0.03$  for the heavy atoms bombarded by helium ions. The ratios of  $\sigma_{KX}^{\text{exper.}}$  to  $\sigma_K^{\text{BORN}}$  or  $\sigma_{KX}^{\text{ECPSSR}}$  exhibit a substantial and erratic dependence on  $Z_2$  for the lightest target atoms ( $4 \leq Z_2 \leq 9$ ) which lack a fully filled  $L$  shell. Fluorescence yields for these elements could be uncertain by more than 40%.<sup>19</sup> These small  $K$ -shell x-ray fluorescence yields are indeed greatly affected by multiple ionizations. They are also changed by chemical and morphological changes in the incomplete  $L$  shell depending on the molecular composition and physical phase of the target. Finally, even in monatomic gas targets, the ionization cross section in itself is affected by relatively strong correlation effects in the very structure of the lightest atoms; the screened hydrogenic wave functions, on which our cal-

culations are based, or even Hartree-Slater schemes become inappropriate because the independent electron model of an atom breaks down. For the lightest atoms, the experiment-to-theory ratios are not shown at all in Fig. 5 since their erratic behavior detracts from the main impression that this figure conveys, e.g., predictions of first Born approximation can be as much as three orders of magnitude above the data. The erratic behavior among the lightest atoms can be easily observed in Figs. 6, 7, and 9, where ratios for the  $4 \leq Z_2 \leq 9$  elements are displayed separately with every element identified by its atomic number. We exclude these lightest target atoms from further statistical analysis: the rejection criterion will be applied to some 7000 data only in the  $10 \leq Z_2 \leq 92$  range of elements.

Figures 5–9 show the experimental-to-theoretical cross section ratios as horizontal bars for all data with  $10 \leq Z_2 \leq 92$  and as circles for three groups of data in the preselected  $Z_2$  ranges. For moderately light ( $10 \leq Z_2 \leq 13$ ) elements, which are predominantly (81%) based on aluminum cross sections, these ratios are drawn as the open circles. The half-open circles represent similar ratios for medium elements of which titanium, chromium, iron, cobalt, nickel, copper, silver, and tin amount to nearly a one-half of all data in the  $14 \leq Z_2 \leq 66$  range. The solid circles are drawn for heavy elements ( $67 \leq Z_2 \leq 92$ ) of which tantalum, gold, and lead are most typical, accounting for almost a one-half of all data in the  $67 \leq Z_2 \leq 92$  range. Aluminum and gold were chosen, in fact, as representative of light and heavy elements by Chadwick,<sup>66</sup> after the 1912 discovery of x rays from iron bombard-




FIG. 5. Averaged [within the 0.1 intervals of  $\log(v_1/v_{2K})$ ] ratios of experimental cross sections to the first Born calculations for the relatively light ( $10 \leq Z_2 \leq 13$ : open circles), medium ( $14 \leq Z_2 \leq 66$ : half-open circles), and heavy ( $67 \leq Z_2 \leq 92$ : closed circles) target elements bombarded by protons. The solid curve is based on the averaged ratios for the  $10 \leq Z_2 \leq 92$  targets.




FIG. 6. Averaged [within the 0.1 intervals of  $\log(v_1/v_{2K})$ ] ratios of experimental cross sections to the ECPSSR predictions for relatively light (open circles), medium (half-open circles), and heavy (closed circles) target elements bombarded by protons. The solid curve is based on the averaged ratios for the  $10 \leq Z_2 \leq 92$  targets; ratios for the  $4 \leq Z_2 \leq 9$  elements are identified by the atomic numbers of these targets. The mean value of the solid curve is 0.96.

ed by alpha particles. The trends of Fig. 2, the failure of the first Born approximation (illustrated in Fig. 5 for protons only since these trends are similar for other projectiles) and the relative success of the ECPSSR theory, are confirmed and well documented by Figs. 6–9.

The rejected data, i.e., the measurements which differ by more than a factor of 2 from other experimental cross sections in comparable collision regimes, are listed in Tables 2–5 in the bold print for easy recognition. Their identification may serve as a guide for experimentalists into trouble areas in which more measurements would be needed and worthwhile. Our criterion rejects 227 cross sections out of 7007 data. Such a large rejection would be anticipated if the standard deviation  $\sigma$  in the normal distribution of these data was such that  $2.14\sigma$  were comparable to the measured cross sections. Experimental uncertainties, however, rarely exceed 25%. The ratios, which are more than a factor of 2 different from the mean values, typically lie no less than  $4\sigma$  from these averages: at most five such ratios would be statistically expected in a sample of 7000 data, while 98% of all rejected ratios is most probably due to truly bad experiments.

In addition, new information emerges from this comprehensive and detailed experiment-to-theory comparison. The first Born approximation overestimates the data by orders of magnitude for the elements in the middle of the periodic table when projectiles are slow. It does it even more dramatically for light elements where  $Z_1/Z_2$  is relatively large. On the other hand, when  $Z_1/Z_2$  is small the first Born




FIG. 7. Averaged [within the 0.1 intervals of  $\log(v_1/v_{2K})$ ] ratios of experimental cross sections to the ECPSSR predictions for relatively light (open circles), medium (half-open circles), and heavy (closed circles) target elements bombarded by deuterons. The solid curve is based on the averaged ratios for the  $11 < Z_2 < 79$  targets; ratios for beryllium are identified by its atomic number. The mean value of the solid curve is 0.92.




FIG. 8. Averaged [within the 0.1 intervals of  $\log(v_1/v_{2K})$ ] ratios of experimental cross sections to the ECPSSR predictions for aluminum (open circles) and medium (half-open circles) target elements bombarded by  ${}^3\text{He}$  ions. The solid curve is based on the averaged ratios for the  $13 < Z_2 < 47$  targets. The mean value of the solid curve is 1.01.




FIG. 9. Averaged [within the 0.1 intervals of  $\log(v_1/v_{2K})$ ] ratios of experimental cross sections to the ECPSSR predictions for light (open circles), medium (half-open circles), and heavy (closed circles) target elements bombarded by  ${}^4\text{He}$  ions. The solid curve is based on the averaged ratios for the  $10 < Z_2 < 29$  targets; ratios for the  $4 < Z_2 < 9$  elements identified by the atomic numbers of these targets. The mean value of the solid curve is 1.00.

approximation on the average underestimates the data by nearly a factor of 3 when  $v_1/v_{2K} \approx 0.1$ . The ECPSSR removes these discrepancies so that the average ratios of experiment to theory are within 20% of the ideal ratio of 1 for protons. A similar conclusion was made in Ref. 15 for identical (protons and targets with  $10 < Z_2 < 92$ ) collision systems but merely a one-half of the current data base for proton-induced x-ray production cross sections. For deuterons the agreement is within 25%, except at the lowest projectile velocities where ECPSSR overestimates the measured cross sections by a factor of 2.

For  ${}^3\text{He}$  ions, the discrepancies are much more pronounced. They are, however, less significant due to the relatively small (4% of all compiled cross sections) and limited (to  $13 < Z_2 < 47$  targets) amount of data that is available. As opposed to general trends at small  $v_1/v_{2K}$  for any other target-projectile combination, the experimental  $K$  x-ray production cross section from aluminum bombarded by  ${}^3\text{He}$  is up to 70% larger than the ECPSSR predictions; these data, however, are from only a few references. The agreement of ECPSSR with the compiled  ${}^4\text{He}$  data is comparable to its concord with the proton data on the average. Yet the divergence in agreement with the light versus heavy target data is more evident in helium-induced cross sections because  $Z_1/Z_2$  is twice as large.

Experiment-to-theory comparisons, such as presented in this work and most recently by Paul and his collaborators,<sup>5-14</sup> are interpreted as tests of theories to be gauged by massive empirical collections of data. It is amusing to recall Cork<sup>67</sup> who, in a reversal of this procedure, argued that his experiment was acceptable because its deviation from the theory was comparable to theoretical uncertainties. Cork concluded that the measured cross section for  $K$ -shell x-ray production in iron by deuterons was "10 to 100 times greater than the theoretical value, but the difference could not be regarded as outside the limit of error in the calculation." Ironically, this particular calculation agrees (well within a factor of 2) with the predictions of current theories for  $\sigma_{KX}$  in iron by 10-MeV deuterons.

We continue to use our latest formulation of the ECPSSR theory.<sup>15,16</sup> Residual deviations of this theory from the data are present and are indeed statistically significant. While the data for moderately heavy and light target elements are in basic agreement with the averages for all data, the cross sections for the lightest and heaviest target atoms oscillate in opposite directions around these averages. In the slow collision limit, the measured cross sections are overpredicted when  $Z_1/Z_2$  is small but they appear to be underpredicted when  $Z_1/Z_2$  is large. Similar trends are noticed in recent work of Paul *et al.*<sup>8-14</sup> The overprediction of the experimental cross sections in heaviest targets is connected with a crude way in which the ECPSSR theory accounts for the relativistic effect; this theory indeed overestimates the importance of the relativistic treatment of the  $K$ -shell electron as proven<sup>42</sup> by numerical calculations that use the Dirac wave functions. The underprediction of the data for  $Z_1/Z_2 > 0.15$  has been discussed above in terms of the influence of multiple ionizations on  $\sigma_K$ . This underprediction seems to contradict the pronouncements<sup>59-61</sup> that the




FIG. 10. Same as Fig. 6 but vs the variable  $\xi_K^R/\xi_K$  according to which the ECPSSR ionization cross sections scale in the slow collision regime. Correspondingly, the averages are within the 0.1 intervals of  $\log(\xi_K^R/\xi_K)$ .




FIG. 11. Same as Fig. 9 but vs the variable  $\xi_K^R/\xi_K$  according to which the ECPSSR ionization cross sections scale in the slow collision regime. Correspondingly, the averages are within the 0.1 intervals of  $\log(\xi_K^R/\xi_K)$ .

ECPSSR theory underestimates the PSS effect, smaller ionization cross sections that Refs. 59 and 61 suggest would accentuate the discrepancy with experiments. On the other hand, revised accounts for the binding<sup>64</sup> or for the polarization<sup>65</sup> effects could perhaps remove some of this discrepancy. As discussed in Sec. 3.2.a, the ECPSSR theory exhibits a nearly universal scaling with respect to  $\xi_K^R/\xi_K$ . Hence the ratios of Figs. 6 (for protons) and 9 (for  ${}^4\text{He}$ ) are, respectively, replotted as Figs. 10 and 11 in terms of this variable; the deuteron and  ${}^3\text{He}$  ratios remain essentially the same because their relative scarcity prevents a statistically meaningful differentiation. Since  $\xi_K^R/\xi_K$  is more natural than  $v_1/v_{2K}$  in the scaling of the ECPSSR calculations, the replotted ratios are somewhat smoother and, especially at low velocities, the dichotomy between the light and heavy targets is more evident. Also, for large  $Z_1/Z_2$ , the discrepancy between the theory and the data is larger in Figs. 10 and 11. The deviations detected in Figs. 6 and 9 are now seen in the sharpest focus; they still persist and a fortiori reflect real discrepancies between experiment and the ECPSSR theory.

#### 4. Conclusions

This analysis supports the main conclusions of Ref. 15: for the  $10 < Z_2 < 92$  targets, theory and experiment agree, on the average, to within  $\pm 10\%$  to  $20\%$ . With one standard deviation of  $\pm 0.20$ , the mean ratio of  $\sigma_{KK}^{\text{Exper.}}/\sigma_{KK}^{\text{ECPSSR}}$  for these targets and all projectiles equals 0.97. For  ${}^1\text{H}$ ,  ${}^2\text{H}$ ,  ${}^3\text{He}$ , and  ${}^4\text{He}$ , this ratio is, respectively,  $0.96 \pm 0.19$ ,  $0.92 \pm 0.19$ ,  $1.01 \pm 0.24$ , and  $1.00 \pm 0.23$ . The residual deviations are nevertheless genuine and systematic. Only a comprehensive survey of all the data allows to isolate these deviations as a fine structure superimposed on the billionfold change of cross sections with the projectile velocity.

Perhaps the Coulomb deflection factor of the ECPSSR theory should be reconsidered.<sup>66</sup> A quantum mechanical derivation<sup>69</sup> of this factor might be fundamentally more correct.<sup>70</sup> The ECPSSR could be faulty in its treatment of PSS effects. The discrepancy between it and multistate calculations<sup>59,60,62</sup>, however, might reflect differences in the description of the  $K$  shell rather than inadequacies in the PSS formulation. The calculations of Refs. 59, 60, and 62 span a short interval of  $v_1/v_{2K}$  from 0.12 to 0.31; an extension of this interval with calculations that employ and do not employ a screened Coulomb potential for the  $K$ -shell electron would be of interest.

For now, the deviations found in our analysis are attributed primarily to inadequacies of a screened hydrogenic description of the target electron on which the ionization calculations<sup>15,16,20,21</sup> rest; this explanation of the observed deviations seems to be particularly valid when  $K$  shells of relatively light targets are considered. The ratios of the cross section based on Hartree-Slater wave functions<sup>22,29,71</sup> to the cross section evaluated with the screened hydrogenic wave functions show (see Fig. 3 of Ref. 15) remarkable resemblance to the ratios displayed in Figs. 10 and 11. Hence, we speculate that, provided the relativistic effect will be better accounted for in the theory and the multiple-ionization effect considered, almost perfect agreement with the data

would result if the ECPSSR cross sections were calculated with better wavefunctions for atomic  $K$  shells.

Known disagreements between the ECPSSR predictions and  $L$ -subshell data appear to make this conclusion very speculative indeed. Attempts have been made to explain some of these discrepancies in terms of a two-step mechanism in which a vacancy decay in an ionized subshell is followed by intrashell transitions during the same collision.<sup>72</sup> These corrections have been made, however, in terms of the second order transition probabilities (instead of amplitudes) that were evaluated using the straight-line approximation and without account for PSS effects. An inclusion of PSS effects in the second Born approximation has been advocated by Sarkadi.<sup>73</sup> Strong inter-subshell couplings influence  $L$ -subshell ionization probabilities<sup>74</sup> and affect ionization cross sections.<sup>75</sup> It is hoped that rigorous numerical calculations—which extend beyond the first Born approximation, treat the E, C, PSS, and R effects concomitantly, and are *ab initio* in all collisional regimes—will become available in a near future. Ultimately, comprehensive compilations and analyses of the  $L$ - and  $M$ -shell data are needed to convert our tentative deductions, on the shortcomings of the ECPSSR treatment of  $K$ -shell ionization in particular, to more firm conclusions on inadequacies of this theory in general.

Aside from open questions of theoretical interpretation of the compiled data, the present compilation appears to have its own merits as an assessment of worthwhile experiments and, perhaps, as a stimulant for further measurements. It identifies the target elements for which  $K$ -shell x-ray production cross sections have never or seldom been measured with light ion bombardment. It points to the projectiles for which more measurements would be desirable. The compiled data exhibit particularly large scatter among the deuteron and  ${}^3\text{He}$  induced cross sections; possible bad measurements cannot be reliably recognized because of the relatively small (11% of all compiled cross sections) amount of these data. All helium-induced x-ray production cross sections should be reported with the He charge state; especially, for light target elements and at low-projectile velocities where electron capture contributes significantly to  $K$ -shell ionization (see Table 7). An extension of proton measurements at relatively high velocities,  $v_1 > v_{2K}$ , to other fast projectiles would be beneficial in understanding of relatively large discrepancies between lighter and heavier elements that appear (see Figs. 6 and 10) in the proton data at high velocities. It remains to be seen whether experimentalists will be prompted to a revival of  $K$ -shell x-ray measurements in asymmetric collisions. Such a resurgence could slow down the current rapid decline in the rate with which new data are reported (see Fig. 1) and it might force a quantitative revision of our present forecast about the total number of compiled cross sections saturating at 10 000.

#### 5. Acknowledgments

This work was supported by the National Institute of Standards and Technology Grant No. NB82NADA3033, as a part of an interagency program supported by the National Science Foundation and the Office of Basic Energy Sciences, Department of Energy.

TABLE 1. Distribution of compiled K-shell x-ray production cross sections, for each target of atomic number  $Z_2=4-92$ , with respect to the type of projectile ( $Z_1=1,2$ : ions of H-1, H-2, He-3, He-4).  $Z_2$  of the elements, for which data are listed in Tables 2-5, is highlighted in the bold print. A summary of the compiled data for all target elements appears at the bottom of this table

$Z_2$	Protons	+	Deuterons	+	He-3	+	He-4	=	All Ions
<b>4</b>	43		7		0		22		72
5	0		0		0		0		0
<b>6</b>	164		0		0		52		216
<b>7</b>	21		0		0		18		39
<b>8</b>	54		0		0		0		54
<b>9</b>	18		0		0		12		30
<b>10</b>	22		0		0		12		34
<b>11</b>	8		3		0		0		11
<b>12</b>	41		0		0		12		53
<b>13</b>	200		45		70		104		419
<b>14</b>	16		2		7		6		31
<b>15</b>	29		13		10		0		52
<b>16</b>	34		13		0		4		51
<b>17</b>	31		13		0		19		63
<b>18</b>	37		0		10		5		52
<b>19</b>	32		13		0		4		49
<b>20</b>	87		6		0		37		130
<b>21</b>	86		14		0		9		109
<b>22</b>	286		42		33		169		530
<b>23</b>	110		10		0		63		183
<b>24</b>	162		6		32		61		261
<b>25</b>	98		6		0		35		139
<b>26</b>	267		38		0		110		415
<b>27</b>	126		10		23		56		215
<b>28</b>	222		45		14		89		370
<b>29</b>	420		38		19		178		655
<b>30</b>	162		0		0		32		194
<b>31</b>	34		13		0		11		58
<b>32</b>	73		4		23		51		151
<b>33</b>	16		0		0		9		25
<b>34</b>	56		0		13		54		123
<b>35</b>	14		0		0		4		18
<b>36</b>	23		0		0		0		23
<b>37</b>	31		10		0		26		67
<b>38</b>	21		11		0		0		32
<b>39</b>	53		0		0		30		83
<b>40</b>	38		10		9		12		69
<b>41</b>	32		8		8		27		75
<b>42</b>	118		0		0		51		169
43	0		0		0		0		0
<b>44</b>	1		0		0		0		1
<b>45</b>	10		0		0		16		26
<b>46</b>	44		0		11		13		68
<b>47</b>	268		18		8		82		376
<b>48</b>	54		10		0		24		88

TABLE 1. Distribution of compiled *K*-shell x-ray production cross sections, for each target of atomic number  $Z_2=4\text{--}92$ , with respect to the type of projectile ( $Z_1=1,2$ : ions of H-1, H-2, He-3, He-4).  $Z_2$  of the elements, for which data are listed in Tables 2-5, is highlighted in the bold print. A summary of the compiled data for all target elements appears at the bottom of this table — Continued

$Z_2$	Protons	+	Deuterons	+	He-3	+	He-4	=	All Ions
<b>49</b>	70		8		0		7		85
<b>50</b>	120		8		0		83		211
<b>51</b>	31		16		0		16		63
<b>52</b>	17		0		0		10		27
<b>53</b>	39		0		0		10		49
<b>54</b>	2		0		0		0		2
<b>55</b>	15		0		0		4		19
<b>56</b>	40		0		0		8		48
<b>57</b>	12		0		0		0		12
<b>58</b>	44		5		0		7		56
<b>59</b>	17		0		0		7		24
<b>60</b>	59		0		0		9		68
<b>61</b>	1		0		0		0		1
<b>62</b>	58		0		0		12		70
<b>63</b>	12		0		0		0		12
<b>64</b>	28		11		0		26		65
<b>65</b>	25		0		0		0		25
<b>66</b>	0		0		0		0		0
<b>67</b>	57		0		0		21		78
<b>68</b>	0		0		0		0		0
<b>69</b>	26		0		0		22		48
<b>70</b>	10		0		0		0		10
<b>71</b>	0		0		0		5		5
<b>72</b>	6		0		0		7		13
<b>73</b>	66		11		0		18		95
<b>74</b>	32		15		0		23		70
<b>75</b>	6		0		0		6		12
<b>76</b>	0		0		0		0		0
<b>77</b>	0		0		0		0		0
<b>78</b>	8		0		0		12		20
<b>79</b>	90		14		0		39		143
<b>80</b>	0		0		0		0		0
<b>81</b>	0		0		0		0		0
<b>82</b>	56		0		0		31		87
<b>83</b>	7		0		0		16		23
<b>84</b>	0		0		0		0		0
<b>85</b>	0		0		0		0		0
<b>86</b>	0		0		0		0		0
<b>87</b>	0		0		0		0		0
<b>88</b>	0		0		0		0		0
<b>89</b>	0		0		0		0		0
<b>90</b>	26		0		0		21		47
<b>91</b>	0		0		0		0		0
<b>92</b>	45		0		0		6		51

Z2=4-92    Protons    Deuterons    He-3    He-4    All Data  
Targets    4687(63%)    496(7%)    290(4%)    1945(26%)    7418

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
<b>4 Beryllium      Fluorescence yield = 0.00033</b>									
1.50-2	8.70+0	4.76-1	2.00-2	3.00+1	4.71-1	2.50-2	7.50+1	5.02-1	35
3.00-2	1.60+2	5.83-1	4.00-2	4.00+2	6.60-1	5.00-2	6.50+2	6.64-1	
6.00-2	1.00+3	7.50-1	7.00-2	1.30+3	7.91-1	8.00-2	1.60+3	8.42-1	
9.00-2	1.80+3	8.54-1	1.00-1	2.00+3	8.82-1	1.20-1	2.30+3	9.24-1	
1.40-1	2.70+3	1.03+0	1.60-1	3.00+3	1.13+0	1.80-1	3.20+3	1.20+0	
2.00-1	3.40+3	1.28+0							
5.00-1	1.51+3	7.72-1	7.50-1	1.43+3	9.08-1	1.00+0	1.32+3	9.96-1	71
1.20+0	1.23+3	1.12+0	1.40+0	1.14+3	1.15+0	1.60+0	1.07+3	1.17+0	
1.80+0	1.01+3	1.20+0	2.00+0	9.78+2	1.24+0				
3.00-1	2.53+3	1.04+0	5.00-1	2.26+3	1.16+0	7.00-1	2.00+3	1.22+0	92
1.00+0	1.68+3	1.27+0	1.20+0	1.61+3	1.47+0	1.50+0	1.52+3	1.60+0	
1.80+0	1.42+3	1.68+0							
1.00-2	3.17-1	1.25-1	1.20-2	8.69-1	1.37-1	1.50-2	2.78+0	1.52-1	119
2.00-2	1.16+1	1.82-1	2.50-2	3.33+1	2.23-1	3.00-2	7.49+1	2.73-1	
4.00-2	2.23+2	3.68-1	5.00-2	4.47+2	4.56-1	6.00-2	7.33+2	5.50-1	
8.00-2	1.28+3	6.74-1	1.00-1	1.84+3	8.12-1	1.20-1	2.17+3	8.72-1	
<b>6 Carbon      Fluorescence yield = 0.0028</b>									
1.50+0	3.00+3	1.37+0							10
1.50-2	5.76-2	1.03-1	2.00-2	2.00-1	9.24-2	2.50-2	5.58-1	9.55-2	11
3.00-2	1.83+0	1.45-1	4.00-2	6.33+0	1.60-1	5.00-2	1.53+1	1.74-1	
6.00-2	3.40+1	2.13-1	7.00-2	5.94+1	2.36-1	8.00-2	1.17+2	3.25-1	
9.00-2	1.66+2	3.46-1	1.00-1	2.07+2	3.41-1	1.10-1	2.74+2	3.71-1	
4.99-1	2.17+3	7.65-1	5.95-1	2.24+3	7.87-1	6.98-1	2.28+3	8.11-1	
7.75-1	2.28+3	8.26-1	9.10-1	2.29+3	8.61-1	1.02+0	2.24+3	8.73-1	
1.10+0	2.18+3	8.70-1	1.20+0	2.17+3	8.95-1	1.27+0	2.10+3	8.88-1	
1.36+0	2.09+3	9.10-1	1.51+0	2.02+3	9.21-1	1.66+0	1.94+3	9.28-1	
1.91+0	1.89+3	9.73-1							
2.00-2	9.50-1	4.39-1	3.00-2	4.30+0	3.40-1	4.00-2	2.00+1	5.07-1	16
5.00-2	4.40+1	5.00-1	6.00-2	8.20+1	5.15-1	7.00-2	1.20+2	4.77-1	
8.00-2	2.20+2	6.11-1							
1.50-2	4.80-1	8.58-1	2.00-2	1.80+0	8.31-1	2.50-2	5.40+0	9.25-1	23
3.00-2	1.10+1	8.70-1	3.50-2	2.10+1	8.90-1	4.00-2	3.60+1	9.12-1	
5.00-2	8.10+1	9.20-1							

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
2.00-2	1.70+0	7.85-1	2.50-2	5.00+0	8.56-1	3.00-2	1.20+1	9.49-1	26
4.00-2	3.60+1	9.12-1	5.00-2	7.60+1	8.64-1	6.00-2	1.30+2	8.16-1	
7.00-2	2.00+2	7.95-1	8.00-2	2.80+2	7.78-1	9.00-2	3.70+2	7.72-1	
1.00-1	4.70+2	7.75-1	1.20-1	6.60+2	7.59-1	1.40-1	9.10+2	8.07-1	
1.60-1	1.20+3	8.76-1	1.80-1	1.30+3	8.18-1	2.00-1	1.50+3	8.40-1	
2.00+0	7.60+2	4.01-1	3.00+0	8.60+2	5.79-1	4.00+0	8.40+2	6.85-1	37
6.00+0	7.20+2	7.84-1	1.00+1	4.60+2	7.35-1	1.20+1	2.80+2	5.15-1	
1.40+1	3.30+2	6.85-1							
2.80-2	7.56+0	7.97-1	3.70-2	2.02+1	6.89-1	4.80-2	5.54+1	7.24-1	46
5.80-2	9.94+1	6.94-1	6.80-2	1.55+2	6.69-1	7.70-2	2.44+2	7.49-1	
8.70-2	3.32+2	7.51-1	9.70-2	4.40+2	7.75-1	1.06-1	5.42+2	7.90-1	
1.16-1	6.67+2	8.15-1	1.26-1	7.82+2	8.24-1	1.35-1	8.99+2	8.44-1	
1.45-1	1.03+3	8.66-1							
1.00-1	3.50+2	5.77-1	1.10-1	5.00+2	6.78-1	1.20-1	6.00+2	6.90-1	50
1.30-1	7.00+2	7.00-1	1.40-1	7.50+2	6.65-1	1.50-1	8.00+2	6.39-1	
1.60-1	1.00+3	7.30-1	1.70-1	1.10+3	7.43-1	2.00-1	1.20+3	6.72-1	
2.50-1	1.50+3	6.89-1	3.00-1	1.60+3	6.55-1	4.00-1	2.00+3	7.31-1	
5.00-1	2.10+3	7.40-1	6.00-1	2.10+3	7.38-1	7.00-1	2.00+3	7.11-1	
8.00-1	2.00+3	7.28-1	9.00-1	1.90+3	7.12-1	1.00+0	1.80+3	6.97-1	
1.00-2	7.00-2	1.01+0	1.25-2	1.80-1	7.97-1	1.50-2	4.00-1	7.15-1	51
1.75-2	7.80-1	6.67-1	2.00-2	1.30+0	6.00-1	2.50-2	3.30+0	5.65-1	
3.00-2	6.90+0	5.46-1	3.50-2	1.30+1	5.51-1	4.00-2	2.20+1	5.57-1	
4.50-2	3.20+1	5.26-1	5.00-2	5.10+1	5.80-1				
1.00+0	2.97+3	1.15+0	2.00+0	2.04+3	1.08+0	3.00+0	1.65+3	1.11+0	70
4.00+0	1.34+3	1.09+0	6.00+0	1.04+3	1.13+0	8.00+0	8.40+2	1.13+0	
1.00+1	7.00+2	1.12+0	1.20+1	6.20+2	1.14+0	1.40+1	5.60+2	1.16+0	
1.60+1	4.50+2	1.04+0	1.80+1	4.20+2	1.07+0				
2.90-1	2.21+3	9.21-1	5.20-1	2.80+3	9.83-1	7.20-1	2.77+3	9.90-1	81
1.02+0	2.45+3	9.54-1	2.00+0	2.01+3	1.06+0	3.00+0	1.46+3	9.83-1	
4.00+0	1.23+3	1.00+0	5.00+0	1.08+3	1.03+0	6.00+0	9.30+2	1.01+0	
7.00+0	8.30+2	1.01+0	8.00+0	8.10+2	1.09+0	9.00+0	7.20+2	1.06+0	
1.00+1	6.70+2	1.07+0	1.10+1	6.50+2	1.12+0	1.20+1	6.30+2	1.16+0	
1.30+1	5.50+2	1.08+0	1.40+1	5.30+2	1.10+0	1.50+1	5.30+2	1.16+0	
1.60+1	5.80+2	1.34+0							
1.00-1	4.15+2	6.84-1	1.25-1	6.50+2	6.94-1	1.50-1	9.10+2	7.27-1	82
1.75-1	1.15+3	7.48-1	2.00-1	1.36+3	7.62-1	2.50-1	1.77+3	8.14-1	
3.00-1	2.04+3	8.35-1	4.00-1	2.22+3	8.11-1	5.00-1	2.33+3	8.21-1	
6.00-1	2.35+3	8.26-1	7.00-1	2.28+3	8.11-1	8.00-1	2.09+3	7.61-1	
1.00+0	2.10+3	8.13-1							
1.80+1	4.28+2	1.09+0	1.90+1	3.79+2	1.00+0	2.00+1	3.69+2	1.02+0	85
2.10+1	3.59+2	1.03+0	2.20+1	3.39+2	1.01+0	2.30+1	3.30+2	1.02+0	
2.40+1	3.10+2	9.92-1	2.50+1	3.10+2	1.03+0	2.60+1	3.10+2	1.06+0	

TABLE 2. *K*-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
6.00-1	2.45+3	8.61-1	8.00-1	2.34+3	8.52-1	1.00+0	2.27+3	8.78-1	157
1.20+0	2.19+3	9.05-1	1.40+0	2.12+3	9.35-1	1.60+0	2.02+3	9.49-1	
1.80+0	1.97+3	9.83-1	2.00+0	1.91+3	1.01+0				
<b>7 Nitrogen</b>						<b>Fluorescence yield = 0.0052</b>			
2.80-2	2.69+0	1.06+0	3.70-2	7.18+0	8.78-1	4.80-2	1.81+1	7.97-1	46
5.80-2	3.38+1	7.47-1	6.80-2	5.91+1	7.59-1	7.70-2	9.19+1	7.95-1	
8.70-2	1.26+2	7.56-1	9.70-2	1.68+2	7.43-1	1.06-1	2.20+2	7.71-1	
1.16-1	2.73+2	7.66-1	1.26-1	3.26+2	7.58-1	1.35-1	3.84+2	7.67-1	
1.45-1	4.28+2	7.38-1							
1.25-1	3.17+2	7.51-1	1.50-1	4.70+2	7.56-1	1.75-1	6.42+2	7.82-1	82
2.00-1	8.21+2	8.08-1	2.50-1	1.09+3	7.97-1	3.00-1	1.36+3	8.20-1	
4.00-1	1.81+3	8.74-1	5.00-1	2.05+3	8.86-1				
<b>8 Oxygen</b>						<b>Fluorescence yield = 0.0083</b>			
2.00-2	1.86-1	1.08+0	2.50-2	4.60-1	9.14-1	3.00-2	9.40-1	8.14-1	18
3.50-2	1.72+0	7.62-1	4.00-2	2.90+0	7.34-1	4.50-2	4.70+0	7.36-1	
5.00-2	7.00+0	7.23-1	5.50-2	1.01+1	7.24-1	6.00-2	1.40+1	7.25-1	
6.50-2	1.88+1	7.28-1	7.00-2	2.44+1	7.26-1	7.50-2	3.10+1	7.27-1	
8.00-2	3.86+1	7.29-1	8.50-2	4.75+1	7.36-1	9.00-2	5.70+1	7.37-1	
9.50-2	6.70+1	7.34-1	1.00-1	7.90+1	7.39-1				
1.50-1	2.40+2	7.64-1							29
2.00-2	5.70-1	3.32+0	2.50-2	1.10+0	2.19+0	3.00-2	2.10+0	1.82+0	33
3.50-2	2.90+0	1.28+0	4.00-2	4.50+0	1.14+0	4.50-2	6.70+0	1.05+0	
5.00-2	1.00+1	1.03+0	5.50-2	1.50+1	1.07+0	6.00-2	2.00+1	1.04+0	
6.50-2	2.40+1	9.29-1	7.00-2	3.00+1	8.93-1	7.50-2	4.20+1	9.85-1	
8.00-2	5.50+1	1.04+0							
5.00-2	7.00+0	7.23-1	6.00-2	1.40+1	7.25-1	7.00-2	2.40+1	7.14-1	34
8.00-2	4.00+1	7.55-1	9.00-2	6.00+1	7.76-1	1.00-1	8.00+1	7.48-1	
1.50-1	2.30+2	7.32-1	2.00-1	4.50+2	7.86-1	2.50-1	7.00+2	8.39-1	
3.00-1	9.30+2	8.64-1	3.50-1	1.20+3	9.30-1	4.00-1	1.30+3	8.82-1	
2.00-2	1.80-1	1.05+0	2.50-2	4.60-1	9.14-1	3.00-2	9.50-1	8.23-1	51
3.50-2	1.65+0	7.31-1	4.00-2	3.00+0	7.59-1	4.50-2	4.70+0	7.36-1	
5.00-2	7.00+0	7.23-1	5.50-2	1.00+1	7.17-1	6.00-2	1.40+1	7.25-1	
6.50-2	1.90+1	7.36-1	7.00-2	2.50+1	7.44-1				
<b>9 Fluorine</b>						<b>Fluorescence yield = 0.013</b>			
5.00-1	1.10+3	8.29-1	8.45-1	1.45+3	7.72-1	1.00+0	1.50+3	7.53-1	110
3.00-1	3.14+2	4.48-1	3.50-1	4.15+2	4.72-1	4.00-1	5.14+2	4.92-1	116
5.00-1	7.54+2	5.68-1	6.00-1	1.02+3	6.60-1	7.00-1	1.27+3	7.41-1	
8.00-1	1.49+3	8.12-1	9.00-1	1.68+3	8.72-1	1.00+0	1.87+3	9.39-1	
1.10+0	1.94+3	9.51-1	1.20+0	2.02+3	9.76-1	1.40+0	2.06+3	9.82-1	
1.60+0	2.02+3	9.66-1	1.80+0	1.92+3	9.27-1	2.00+0	1.83+3	8.98-1	

TABLE 2. *K*-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
<b>10 Neon</b>					<b>Fluorescence yield = 0.018</b>				
4.80-2	1.41+0	1.13+0	5.80-2	2.67+0	9.73-1	6.80-2	4.68+0	9.11-1	46
7.70-2	7.83+0	9.50-1	9.70-2	1.79+1	9.45-1	1.16-1	3.29+1	9.51-1	
1.35-1	5.40+1	9.66-1							
5.00+0	1.40+3	1.00+0							58
1.25-1	3.20+1	7.27-1	1.50-1	5.70+1	7.47-1	2.00-1	1.25+2	7.50-1	82
2.50-1	2.09+2	7.47-1	3.00-1	3.18+2	7.86-1	4.00-1	5.63+2	8.56-1	
5.00-1	7.04+2	7.94-1	6.00-1	8.37+2	7.74-1	7.00-1	9.60+2	7.72-1	
8.00-1	1.05+3	7.64-1	9.00-1	1.20+3	8.10-1	1.00+0	1.30+3	8.31-1	
1.10+0	1.42+3	8.73-1	1.20+0	1.50+3	8.92-1				
<b>11 Sodium</b>					<b>Fluorescence yield = 0.023</b>				
2.00-2	8.00-3	1.72+0	2.50-2	2.50-2	1.38+0	3.00-2	6.00-2	1.21+0	101
4.00-2	1.20-1	5.79-1	5.00-2	2.50-1	4.39-1	5.50-2	6.00-1	6.97-1	
6.00-2	8.50-1	6.86-1	6.50-2	1.30+0	7.54-1				
<b>12 Magnesium</b>					<b>Fluorescence yield = 0.03</b>				
6.00-2	3.50-1	6.64-1	1.00-1	2.80+0	6.78-1	1.50-1	1.20+1	6.95-1	9
2.00-1	2.80+1	6.58-1	3.00-1	9.40+1	7.44-1	4.00-1	1.80+2	7.57-1	
5.00-1	2.30+2	6.38-1							
2.50-2	4.31-3	7.40-1	3.00-2	1.53-2	8.80-1	4.00-2	7.34-2	9.12-1	11
5.00-2	2.27-1	9.71-1	6.00-2	4.45-1	8.44-1	7.00-2	8.52-1	8.41-1	
8.00-2	1.85+0	1.06+0	9.00-2	2.41+0	8.71-1	1.00-1	3.46+0	8.38-1	
6.02-1	4.25+2	8.81-1	6.86-1	4.49+2	7.77-1	8.00-1	5.06+2	7.29-1	
9.00-1	5.77+2	7.35-1	1.00+0	6.42+2	7.42-1	1.10+0	7.27+2	7.77-1	
1.20+0	7.73+2	7.75-1	1.31+0	8.79+2	8.33-1	1.40+0	9.46+2	8.64-1	
1.50+0	1.03+3	9.08-1	1.60+0	1.07+3	9.16-1	1.70+0	1.14+3	9.55-1	
1.25-1	8.30+0	8.95-1	1.50-1	1.50+1	8.69-1	1.75-1	2.60+1	9.17-1	12
2.00-1	4.00+1	9.40-1							
3.00+0	1.10+3	8.39-1							94
2.00-2	6.00-4	<b>4.70-1</b>	2.50-2	4.00-3	6.87-1	3.00-2	1.20-2	6.90-1	101
3.50-2	3.00-2	7.40-1	4.00-2	6.00-2	7.46-1	4.50-2	1.00-1	7.00-1	
5.00-2	2.00-1	8.56-1	6.00-2	5.00-1	9.48-1				
<b>13 Aluminum</b>					<b>Fluorescence yield = 0.039</b>				
6.00-2	1.70-1	7.05-1	1.00-1	1.50+0	7.31-1	1.32-1	3.90+0	6.74-1	9
1.50-1	5.60+0	6.17-1	2.00-1	1.50+1	6.42-1	3.00-1	5.50+1	7.34-1	
4.00-1	1.20+2	7.97-1	5.00-1	2.30+2	9.64-1				
1.50+0	1.00+3	1.07+0							10
2.50-2	1.43-3	7.43-1	3.00-2	5.01-3	7.87-1	4.00-2	2.98-2	8.98-1	11
5.00-2	9.19-2	8.93-1	6.00-2	2.11-1	8.75-1	7.00-2	4.09-1	8.58-1	

TABLE 2. *K*-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
8.00-2	6.97-1	8.31-1	9.00-2	1.06+0	7.83-1	1.00-1	1.54+0	7.50-1	
4.84-1	1.92+2	8.58-1	5.98-1	2.72+2	8.24-1	7.00-1	3.44+2	8.14-1	
8.00-1	4.17+2	8.19-1	9.00-1	4.86+2	8.24-1	1.00+0	5.46+2	8.23-1	
1.10+0	6.17+2	8.45-1	1.20+0	6.66+2	8.43-1	1.30+0	7.19+2	8.53-1	
1.40+0	7.66+2	8.60-1	1.50+0	8.28+2	8.87-1	1.60+0	8.80+2	9.07-1	
1.70+0	9.24+2	9.21-1							
3.00-2	1.30-2	<b>2.04+0</b>	5.00-2	1.20-1	1.17+0	8.00-2	7.40-1	8.83-1	12
1.00-1	1.50+0	7.31-1	1.25-1	4.80+0	1.01+0	1.50-1	7.80+0	8.60-1	
1.75-1	1.60+1	1.05+0	2.00-1	2.90+1	1.24+0				
4.80-2	8.00-2	9.49-1	5.00-2	8.60-2	8.36-1	6.00-2	2.00-1	8.29-1	13
6.00-2	2.20-1	9.12-1	7.00-2	4.00-1	8.39-1	7.20-2	4.20-1	7.81-1	
8.00-2	6.50-1	7.75-1	8.40-2	7.70-1	7.52-1	9.00-2	1.00+0	7.38-1	
9.60-2	1.11+0	6.35-1	1.00-1	1.54+0	7.50-1				
2.50-2	2.00-3	1.04+0	3.00-2	6.10-3	9.58-1	4.00-2	3.10-2	9.34-1	17
5.00-2	1.20-1	1.17+0	6.00-2	2.20-1	9.12-1	7.00-2	4.40-1	9.23-1	
8.00-2	7.60-1	9.06-1	9.00-2	1.20+0	8.86-1	1.20-1	3.60+0	8.80-1	
1.40-1	6.20+0	8.69-1	1.80-1	1.00+1	5.98-1	1.90-1	1.40+1	7.04-1	
2.00-1	1.90+1	8.13-1							
8.00-2	5.60-1	6.68-1	9.00-2	8.40-1	6.20-1	1.00-1	1.40+0	6.82-1	22
1.10-1	2.00+0	6.76-1	1.20-1	2.80+0	6.84-1	1.30-1	4.00+0	7.31-1	
1.40-1	4.60+0	6.45-1	1.50-1	6.00+0	6.61-1	1.60-1	7.20+0	6.37-1	
1.70-1	8.00+0	5.77-1	1.80-1	1.00+1	5.98-1	1.90-1	1.20+1	6.03-1	
2.00-1	1.40+1	5.99-1							
9.00-2	1.05+0	7.75-1	1.00-1	1.49+0	7.26-1	1.40-1	4.67+0	6.55-1	25
1.80-1	1.06+1	6.34-1							
1.75-2	1.55-4	<b>1.45+0</b>	2.00-2	4.60-4	1.32+0	2.50-2	2.02-3	1.05+0	45
3.00-2	6.54-3	1.03+0	3.50-2	1.61-2	1.01+0	4.00-2	3.26-2	9.82-1	
5.00-2	1.07-1	1.04+0	6.00-2	2.27-1	9.41-1	7.00-2	4.33-1	9.08-1	
8.00-2	7.33-1	8.74-1	9.00-2	1.14+0	8.42-1	1.00-1	1.68+0	8.19-1	
1.10-1	2.35+0	7.95-1	1.20-1	3.46+0	8.46-1	1.40-1	6.44+0	9.03-1	
1.53-1	7.42+0	7.64-1	1.60-1	9.14+0	8.08-1	1.79-1	1.22+1	7.43-1	
2.00-1	1.79+1	7.66-1	2.04-1	1.84+1	7.40-1	2.50-1	3.11+1	6.82-1	
2.55-1	3.28+1	6.81-1	3.06-1	5.34+1	6.77-1	3.57-1	7.81+1	6.74-1	
4.08-1	9.81+1	6.23-1	4.59-1	1.38+2	6.84-1	5.10-1	1.70+2	6.86-1	
6.12-1	2.34+2	6.82-1	7.14-1	2.95+2	6.78-1	8.16-1	3.50+2	6.69-1	
9.18-1	4.01+2	6.64-1	1.02+0	4.46+2	6.59-1	1.12+0	4.84+2	6.52-1	
1.22+0	5.21+2	6.51-1	1.33+0	5.48+2	6.39-1	1.43+0	5.76+2	6.37-1	
1.53+0	6.04+2	6.39-1	1.63+0	6.27+2	6.39-1	1.73+0	6.50+2	6.42-1	
1.84+0	6.70+2	6.42-1	1.94+0	6.89+2	6.45-1	2.04+0	7.05+2	6.49-1	
2.14+0	7.23+2	6.54-1	2.24+0	7.39+2	6.59-1	2.35+0	7.54+2	6.62-1	
2.45+0	7.68+2	6.69-1	2.55+0	7.81+2	6.74-1	2.65+0	7.96+2	6.82-1	
2.75+0	8.09+2	6.89-1	2.85+0	8.23+2	6.98-1	2.96+0	8.37+2	7.07-1	
2.00-2	4.60-4	1.32+0	2.25-2	1.20-3	1.35+0	2.50-2	2.30-3	1.20+0	51
2.75-2	4.60-3	1.25+0	3.00-2	7.50-3	1.18+0	3.25-2	1.20-2	1.16+0	
3.50-2	1.50-2	9.42-1	4.00-2	3.80-2	1.14+0	4.50-2	7.00-2	1.14+0	
5.00-2	1.20-1	1.17+0	5.50-2	1.70-1	1.05+0				

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
7.50-1	2.96+2	6.34-1	1.00+0	4.78+2	7.21-1	1.25+0	6.16+2	7.54-1	75
1.50+0	6.98+2	7.48-1	2.00+0	8.10+2	7.50-1	2.50+0	9.47+2	8.21-1	
3.00+0	9.93+2	8.38-1	3.50+0	9.98+2	8.37-1	4.00+0	9.73+2	8.24-1	
2.00-1	2.28+1	9.75-1							88
3.00+0	1.09+3	9.18-1							94
2.50+0	9.55+2	8.28-1	3.00+0	9.88+2	8.33-1	5.00+0	9.16+2	8.05-1	97
7.50+0	7.80+2	7.81-1	9.00+0	7.19+2	7.81-1	9.75+0	6.83+2	7.72-1	
1.50-2	8.00-6	3.58-1	1.60-2	2.00-5	4.51-1	1.70-2	5.00-5	6.16-1	101
1.80-2	9.00-5	6.47-1	1.90-2	1.60-4	7.10-1	2.00-2	3.50-4	1.00+0	
2.20-2	9.00-4	1.20+0	2.60-2	1.80-3	7.13-1	2.80-2	3.70-3	8.94-1	
3.00-2	6.00-3	9.42-1	3.50-2	1.20-2	7.54-1	4.00-2	2.40-2	7.23-1	
4.50-2	3.70-2	6.05-1	5.00-2	6.50-2	6.32-1	5.50-2	1.20-1	7.41-1	
6.00-2	2.00-1	8.29-1							
3.26-1	9.43+1	1.02+0	3.60-1	1.20+2	1.02+0	3.85-1	1.40+2	1.01+0	110
5.00-1	2.44+2	1.02+0	1.00+0	6.67+2	1.01+0				
7.50-1	2.96+2	6.34-1	1.00+0	4.78+2	7.21-1	1.25+0	6.16+2	7.54-1	117
1.50+0	6.98+2	7.48-1	2.00+0	8.10+2	7.50-1	2.50+0	9.47+2	8.21-1	
2.99+0	9.69+2	8.18-1	3.00+0	9.93+2	8.38-1	3.94+0	9.92+2	8.39-1	
6.07+0	8.79+2	8.14-1	9.13+0	7.33+2	8.02-1	1.22+1	6.13+2	7.83-1	
1.83+1	4.69+2	7.69-1	2.40+1	3.94+2	7.81-1	3.01+1	3.42+2	8.00-1	
3.56+1	2.97+2	7.86-1	3.96+1	2.77+2	7.94-1				
3.00-1	6.85+1	9.14-1	6.00-1	3.00+2	9.04-1	1.00+0	6.05+2	9.12-1	149
<b>14</b>	Silcon								
		Fluorescence yield = 0.05							
7.50-1	2.47+2	7.31-1	1.00+0	3.95+2	7.79-1	1.25+0	4.85+2	7.47-1	75
1.50+0	6.35+2	8.31-1	2.00+0	7.68+2	8.31-1	2.50+0	8.61+2	8.48-1	
3.00+0	9.34+2	8.75-1	3.50+0	9.55+2	8.75-1	4.00+0	9.60+2	8.74-1	
3.00+0	1.00+3	9.37-1							94
2.50-2	3.00-4	4.91-1	3.00-2	1.30-3	5.60-1	3.50-2	5.00-3	7.97-1	101
4.00-2	1.40-2	1.01+0	5.00-2	6.00-2	1.29+0	6.00-2	1.40-1	1.23+0	
<b>15</b>	Phosphorus								
		Fluorescence yield = 0.063							
1.40+0	5.27+2	9.12-1	1.50+0	5.51+2	8.91-1	1.60+0	6.12+2	9.32-1	122
1.70+0	6.33+2	9.15-1	1.80+0	6.70+2	9.25-1	1.90+0	6.95+2	9.22-1	
2.00+0	7.25+2	9.28-1							
6.00-1	1.59+2	1.01+0	8.00-1	3.49+2	1.29+0	1.00+0	4.56+2	1.19+0	123
1.20+0	5.59+2	1.15+0	1.40+0	6.72+2	1.16+0	1.60+0	7.92+2	1.21+0	
1.80+0	9.12+2	1.26+0	2.00+0	9.24+2	1.18+0	2.20+0	9.72+2	1.17+0	
2.40+0	9.66+2	1.11+0	2.60+0	9.28+2	1.03+0				

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_i$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_i$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_i$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
1.00+0	4.81+2	1.25+0	1.30+0	6.50+2	1.22+0	1.90+0	8.46+2	1.12+0	156
2.25+0	8.54+2	1.02+0	2.55+0	9.73+2	1.09+0	2.80+0	1.01+3	1.09+0	
3.00+0	1.00+3	1.05+0	3.50+0	9.69+2	9.80-1	4.00+0	1.09+3	1.08+0	
4.50+0	1.13+3	1.11+0	5.00+0	1.13+3	1.11+0				
<b>16 Sulfur</b> Fluorescence yield = 0.078									
8.30-1	1.80+2	8.50-1	1.58+0	5.40+2	1.02+0	2.56+0	7.50+2	9.64-1	66
3.28+0	1.08+3	1.24+0							
3.00+0	8.46+2	1.00+0							94
1.40+0	4.62+2	1.00+0	1.50+0	4.56+2	9.12-1	1.60+0	4.91+2	9.17-1	122
1.70+0	5.26+2	9.24-1	1.80+0	5.58+2	9.29-1	1.90+0	5.65+2	8.97-1	
2.00+0	5.97+2	9.08-1							
6.00-1	1.55+2	1.41+0	8.00-1	2.95+2	1.49+0	1.00+0	4.04+2	1.39+0	123
1.20+0	5.30+2	1.39+0	1.40+0	6.43+2	1.39+0	1.60+0	7.45+2	1.39+0	
1.80+0	8.09+2	1.35+0	2.00+0	8.77+2	1.33+0	2.20+0	9.68+2	1.37+0	
2.40+0	9.84+2	1.31+0	2.60+0	1.03+3	1.31+0				
1.00+0	3.14+2	1.08+0	1.30+0	4.57+2	1.08+0	1.90+0	6.76+2	1.07+0	156
2.25+0	7.28+2	1.01+0	2.55+0	8.23+2	1.06+0	2.80+0	8.17+2	1.00+0	
3.00+0	8.98+2	1.07+0	3.50+0	9.17+2	1.03+0	4.00+0	1.02+3	1.11+0	
4.50+0	1.12+3	1.19+0	5.00+0	1.13+3	1.19+0				
<b>17 Chlorine</b> Fluorescence yield = 0.097									
9.50-1	2.60+2	1.27+0							76
3.00+0	6.95+2	9.21-1							94
1.40+0	3.63+2	9.69-1	1.50+0	3.74+2	9.14-1	1.60+0	4.02+2	9.09-1	122
1.70+0	4.23+2	8.92-1	1.80+0	5.25+2	1.04+0	1.90+0	5.15+2	9.66-1	
2.00+0	5.34+2	9.54-1							
8.00-1	1.63+2	1.11+0	1.00+0	2.64+2	1.18+0	1.20+0	2.80+2	9.31-1	123
1.40+0	3.31+2	8.84-1	1.60+0	4.18+2	9.45-1	1.80+0	4.80+2	9.52-1	
2.00+0	5.32+2	9.51-1	2.20+0	5.46+2	8.97-1	2.40+0	5.33+2	8.17-1	
2.60+0	5.71+2	8.26-1	2.80+0	5.53+2	7.63-1				
1.00+0	2.69+2	1.20+0	1.30+0	4.00+2	1.18+0	1.90+0	6.09+2	1.14+0	156
2.25+0	6.63+2	1.07+0	2.55+0	7.22+2	1.06+0	2.80+0	7.62+2	1.05+0	
3.00+0	8.08+2	1.07+0	3.50+0	8.42+2	1.04+0	4.00+0	9.35+2	1.10+0	
4.50+0	9.86+2	1.12+0	5.00+0	1.02+3	1.14+0				
<b>18 Argon</b> Fluorescence yield = 0.118									
1.50+0	2.80+2	8.53-1	2.00+0	4.20+2	8.98-1	2.50+0	5.20+2	9.00-1	40
3.00+0	6.10+2	9.22-1	3.50+0	6.80+2	9.38-1	4.00+0	6.90+2	8.95-1	
4.50+0	7.60+2	9.47-1							

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
6.80-2	1.85-2	1.37+0	7.70-2	3.01-2	1.14+0	8.70-2	5.54-2	1.12+0	46
9.70-2	1.04-1	1.25+0	1.06-1	1.48-1	1.17+0	1.16-1	2.27-1	1.20+0	
1.26-1	3.05-1	1.13+0	1.35-1	4.01-1	1.10+0				
1.50+0	3.19+2	9.72-1	2.00+0	4.57+2	9.77-1	2.50+0	5.53+2	9.58-1	48
3.00+0	6.38+2	9.64-1	3.50+0	7.25+2	1.00+0	4.00+0	7.61+2	9.87-1	
4.50+0	7.99+2	9.96-1	5.00+0	8.68+2	1.05+0				
3.00+0	7.03+2	1.06+0							65
2.00+0	5.23+2	1.12+0							74
1.25-1	3.12-1	1.19+0	1.50-1	5.92-1	1.05+0	2.00-1	1.59+0	9.20-1	82
2.50-1	3.57+0	9.21-1	3.00-1	6.51+0	9.03-1	4.00-1	1.58+1	8.83-1	
5.00-1	3.00+1	8.83-1	6.00-1	4.79+1	8.72-1	7.00-1	6.85+1	8.58-1	
8.00-1	9.35+1	8.68-1	9.00-1	1.18+2	8.56-1	1.00+0	1.40+2	8.29-1	
<b>19</b>	<b>Potassium</b>		<b>Fluorescence yield = 0.14</b>						
9.50-1	1.21+2	1.06+0							76
3.00+0	5.35+2	9.38-1							94
1.40+0	2.20+2	9.46-1	1.50+0	2.27+2	8.75-1	1.60+0	2.48+2	8.69-1	122
1.70+0	2.70+2	8.68-1	1.80+0	3.43+2	1.02+0	1.90+0	3.55+2	9.86-1	
2.00+0	3.72+2	9.70-1							
6.00-1	4.04+1	1.06+0	8.00-1	8.65+1	1.11+0	1.00+0	1.23+2	9.73-1	123
1.20+0	1.79+2	1.00+0	1.40+0	2.46+2	1.06+0	1.60+0	2.98+2	1.04+0	
1.80+0	3.41+2	1.02+0	2.00+0	4.31+2	1.12+0	2.20+0	5.22+2	1.22+0	
2.40+0	5.56+2	1.19+0	2.60+0	5.73+2	1.13+0	2.80+0	6.04+2	1.12+0	
1.00+0	1.57+2	1.24+0	1.30+0	2.45+2	1.19+0	1.90+0	3.93+2	1.09+0	156
2.25+0	4.62+2	1.05+0	2.55+0	5.26+2	1.06+0	2.80+0	5.66+2	1.05+0	
3.00+0	6.30+2	1.11+0	3.50+0	6.75+2	1.06+0	4.00+0	7.24+2	1.06+0	
4.50+0	7.90+2	1.09+0	5.00+0	8.34+2	1.11+0				
<b>20</b>	<b>Calcium</b>		<b>Fluorescence yield = 0.163</b>						
2.00+0	2.34+2	7.54-1	3.00+0	3.95+2	8.17-1	4.00+0	4.94+2	8.25-1	20
5.00+0	5.76+2	8.58-1	6.00+0	5.93+2	8.34-1	7.00+0	6.32+2	8.60-1	
8.00+0	6.29+2	8.44-1	9.00+0	6.35+2	8.51-1	1.00+1	6.12+2	8.25-1	
1.10+1	6.17+2	8.42-1	1.20+1	6.18+2	8.58-1	1.30+1	6.17+2	8.71-1	
1.65+1	5.39+2	8.19-1	2.00+1	5.46+2	8.96-1	2.50+1	4.49+2	8.21-1	
3.00+0	4.85+2	1.00+0	5.00+0	7.47+2	1.11+0	7.00+0	8.12+2	1.10+0	94
9.00+0	7.95+2	1.07+0	1.10+1	7.76+2	1.06+0				
5.00-1	1.09+1	6.91-1	5.50-1	1.58+1	7.59-1	6.00-1	1.95+1	7.33-1	98
6.50-1	2.55+1	7.72-1	7.00-1	3.16+1	7.89-1	7.50-1	3.77+1	7.87-1	
8.00-1	4.36+1	7.78-1	8.50-1	5.07+1	7.82-1	9.00-1	5.87+1	7.93-1	
9.50-1	6.61+1	7.91-1	1.00+0	7.50+1	8.02-1	1.05+0	8.41+1	8.11-1	
1.10+0	9.47+1	8.29-1	1.15+0	1.01+2	8.09-1	1.20+0	1.10+2	8.10-1	
1.25+0	1.17+2	7.99-1	1.30+0	1.29+2	8.20-1	1.35+0	1.38+2	8.19-1	

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
1.40+0	1.44+2	8.00-1	1.45+0	1.55+2	8.11-1	1.50+0	1.67+2	8.26-1	
1.55+0	1.75+2	8.21-1	1.60+0	1.89+2	8.42-1	1.65+0	1.97+2	8.35-1	
1.70+0	2.00+2	8.11-1	1.75+0	2.16+2	8.39-1	1.80+0	2.23+2	8.31-1	
1.85+0	2.25+2	8.06-1	1.90+0	2.46+2	8.50-1	1.95+0	2.51+2	8.37-1	
2.00+0	2.66+2	8.58-1							
2.00-1	6.30-1	9.51-1	2.25-1	8.10-1	7.73-1	2.50-1	1.32+0	8.47-1	106
2.75-1	1.71+0	7.75-1	3.00-1	2.38+0	7.91-1	3.25-1	3.16+0	7.96-1	
3.50-1	3.72+0	7.29-1	3.75-1	5.00+0	7.80-1	4.00-1	5.88+0	7.43-1	
4.25-1	7.62+0	7.95-1	4.50-1	8.86+0	7.73-1	4.75-1	1.10+1	8.14-1	
5.00-1	1.28+1	8.12-1							
6.00-1	2.64+1	9.92-1	8.00-1	5.66+1	1.01+0	1.00+0	9.16+1	9.79-1	123
1.20+0	1.40+2	1.03+0	1.40+0	1.81+2	1.01+0	1.60+0	2.23+2	9.93-1	
1.80+0	2.61+2	9.73-1	2.00+0	2.93+2	9.45-1	2.20+0	3.73+2	1.07+0	
2.40+0	4.24+2	1.10+0	2.60+0	4.24+2	1.00+0	2.80+0	4.56+2	1.00+0	
1.00+0	1.16+2	1.24+0	1.30+0	2.18+2	1.39+0	1.90+0	3.19+2	1.10+0	156
2.25+0	3.85+2	1.07+0	2.55+0	4.43+2	1.07+0	2.80+0	4.81+2	1.06+0	
3.00+0	5.09+2	1.05+0	3.50+0	5.69+2	1.04+0	4.00+0	6.47+2	1.08+0	
4.50+0	7.01+2	1.10+0	5.00+0	7.47+2	1.11+0				
<b>21 Scandium      Fluorescence yield = 0.188</b>									
6.67-1	2.37+1	9.44-1	1.00+0	6.83+1	9.84-1	1.33+0	1.29+2	1.01+0	64
1.67+0	2.14+2	1.13+0	2.00+0	2.68+2	1.07+0	2.33+0	3.17+2	1.03+0	
2.67+0	3.79+2	1.05+0	3.00+0	4.44+2	1.09+0	3.33+0	4.85+2	1.08+0	
3.67+0	5.97+2	1.22+0	4.00+0	5.76+2	1.10+0	4.33+0	6.05+2	1.10+0	
4.67+0	6.37+2	1.11+0	5.00+0	6.69+2	1.12+0	5.33+0	6.81+2	1.11+0	
5.67+0	6.99+2	1.11+0							
8.30-1	4.20+1	9.43-1	1.58+0	1.79+2	1.04+0	2.56+0	3.21+2	9.30-1	66
3.28+0	4.71+2	1.06+0							
5.00-1	9.88+0	9.11-1	5.50-1	1.32+1	9.13-1	6.00-1	1.72+1	9.23-1	98
6.50-1	2.14+1	9.16-1	7.00-1	2.62+1	9.15-1	7.50-1	3.13+1	9.13-1	
8.00-1	3.72+1	9.19-1	8.50-1	4.33+1	9.17-1	9.00-1	4.97+1	9.17-1	
9.50-1	5.70+1	9.24-1	1.00+0	6.56+1	9.46-1	1.05+0	7.33+1	9.47-1	
1.10+0	8.27+1	9.64-1	1.15+0	8.98+1	9.53-1	1.20+0	9.71+1	9.43-1	
1.25+0	1.06+2	9.48-1	1.30+0	1.15+2	9.52-1	1.35+0	1.22+2	9.38-1	
1.40+0	1.32+2	9.50-1	1.45+0	1.40+2	9.45-1	1.50+0	1.48+2	9.40-1	
1.55+0	1.57+2	9.40-1	1.60+0	1.65+2	9.34-1	1.65+0	1.69+2	9.10-1	
1.70+0	1.80+2	9.23-1	1.75+0	1.89+2	9.25-1	1.80+0	1.99+2	9.30-1	
1.85+0	2.06+2	9.22-1	1.90+0	2.15+2	9.26-1	1.95+0	2.26+2	9.37-1	
2.00+0	2.30+2	9.19-1							
2.00-1	3.60-1	8.62-1	2.25-1	5.70-1	8.53-1	2.50-1	9.70-1	9.66-1	106
2.75-1	1.32+0	9.19-1	3.00-1	1.87+0	9.48-1	3.25-1	2.15+0	8.20-1	
3.50-1	3.20+0	9.43-1	3.75-1	4.17+0	9.71-1	4.00-1	4.07+0	7.64-1	
4.25-1	5.13+0	7.90-1	4.50-1	6.24+0	8.00-1	4.75-1	7.33+0	7.92-1	
5.00-1	9.82+0	9.06-1							

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
2.00-1	3.76-1	9.00-1	3.00-1	1.82+0	9.22-1	4.00-1	4.98+0	9.35-1	121
5.00-1	9.96+0	9.19-1	6.00-1	1.74+1	9.34-1	7.00-1	2.62+1	9.15-1	
8.00-1	3.53+1	8.72-1	9.00-1	4.81+1	8.88-1				
5.00-1	9.65+0	8.90-1	6.00-1	1.71+1	9.18-1	7.00-1	2.70+1	9.43-1	137
8.00-1	3.91+1	9.66-1	1.00+0	6.73+1	9.70-1	1.20+0	1.01+2	9.81-1	
1.40+0	1.39+2	1.00+0	1.60+0	1.72+2	9.73-1	1.80+0	2.07+2	9.68-1	
2.00+0	2.78+2	1.11+0	2.20+0	2.85+2	9.98-1	2.30+0	3.06+2	1.01+0	
2.40+0	3.14+2	9.83-1	2.50+0	3.27+2	9.74-1				
<b>22</b>	Titanium				Fluorescence yield = 0.214				
1.60+0	1.70+2	1.23+0	1.80+0	2.20+2	1.29+0	2.10+0	2.60+2	1.20+0	5
2.30+0	3.30+2	1.33+0	2.70+0	3.90+2	1.28+0	2.80+0	4.40+2	1.38+0	
3.20+0	5.10+2	1.38+0	3.50+0	5.80+2	1.44+0	3.70+0	6.40+2	1.51+0	
3.90+0	6.80+2	1.54+0							
2.30-1	1.10-1	<b>2.32-1</b>	3.25-1	4.70-1	<b>2.67-1</b>	4.20-1	1.20+0	<b>2.80-1</b>	7
4.54-1	1.63+0	<b>2.94-1</b>	5.10-1	2.10+0	<b>2.62-1</b>				
1.50+0	1.10+2	8.93-1							10
2.00+0	2.21+2	1.10+0	3.00+0	3.69+2	1.07+0	4.00+0	4.90+2	1.08+0	20
5.00+0	5.53+2	1.05+0	6.00+0	6.26+2	1.08+0	7.00+0	6.63+2	1.08+0	
8.00+0	6.95+2	1.10+0	9.00+0	6.97+2	1.08+0	1.00+1	7.17+2	1.10+0	
1.10+1	6.92+2	1.06+0	1.20+1	6.75+2	1.04+0	1.30+1	6.44+2	1.00+0	
1.65+1	6.94+2	1.14+0	2.00+1	6.12+2	1.06+0	2.50+1	5.76+2	1.10+0	
9.00-2	5.65-3	1.07+0	1.10-1	1.88-2	1.15+0	1.30-1	4.54-2	1.19+0	25
1.50-1	9.01-2	1.19+0	1.70-1	1.65-1	1.24+0				
1.00-1	9.00-3	9.27-1	1.25-1	2.60-2	8.25-1	1.50-1	6.40-2	8.46-1	36
1.50+0	1.50+2	1.22+0	2.00+0	2.30+2	1.14+0	2.50+0	3.25+2	1.17+0	38
3.00+0	4.40+2	1.28+0	3.50+0	5.00+2	1.24+0	4.00+0	5.80+2	1.28+0	
4.50+0	6.40+2	1.30+0	5.00+0	6.10+2	1.16+0	5.50+0	6.70+2	1.21+0	
7.00-1	6.31+0	<b>3.07-1</b>	9.00-1	1.13+1	<b>2.83-1</b>	1.10+0	2.03+1	<b>3.15-1</b>	44
1.30+0	2.48+1	<b>2.68-1</b>	1.50+0	3.11+1	<b>2.53-1</b>	1.70+0	3.55+1	<b>2.30-1</b>	
1.90+0	4.28+1	<b>2.30-1</b>	2.10+0	4.68+1	<b>2.15-1</b>				
1.00+0	3.70+1	7.16-1	2.25+0	1.85+2	7.70-1	3.00+0	2.92+2	8.49-1	47
1.50+0	1.30+2	1.06+0	2.00+0	2.15+2	1.07+0	2.50+0	3.12+2	1.13+0	55
3.00+0	3.83+2	1.11+0	4.96+0	5.82+2	1.11+0	5.96+0	6.55+2	1.13+0	
6.96+0	7.08+2	1.16+0	8.94+0	7.48+2	1.16+0	1.09+1	7.62+2	1.17+0	
1.30-1	2.68-2	7.00-1	1.50-1	5.31-2	7.02-1	1.95-1	1.75-1	7.28-1	57
2.50-1	4.74-1	7.21-1	2.95-1	8.15-1	6.60-1	3.30-1	1.26+0	6.78-1	
3.60-1	1.70+0	6.72-1	3.80-1	1.90+0	6.23-1	4.15-1	2.26+0	5.49-1	
4.00-1	4.60+0	1.27+0	5.00-1	9.40+0	1.25+0	6.00-1	1.60+1	1.21+0	69
7.00-1	2.50+1	1.22+0	8.00-1	3.70+1	1.25+0	9.00-1	4.80+1	1.20+0	
1.00+0	6.40+1	1.24+0	1.10+0	8.00+1	1.24+0	1.20+0	9.70+1	1.24+0	

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_i$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_i$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_i$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
1.30+0	1.13+2	1.22+0	1.40+0	1.33+2	1.24+0	1.50+0	1.57+2	1.27+0	
1.60+0	1.82+2	1.32+0	1.70+0	1.94+2	1.26+0	1.80+0	2.22+2	1.30+0	
1.90+0	2.38+2	1.28+0	2.00+0	2.59+2	1.28+0	2.10+0	2.74+2	1.26+0	
2.20+0	3.03+2	1.30+0	2.30+0	3.20+2	1.29+0				
6.00-1	1.30+1	9.86-1	7.00-1	2.10+1	1.02+0	8.00-1	3.20+1	1.08+0	72
9.00-1	4.50+1	1.13+0	1.00+0	6.10+1	1.18+0	1.50+0	1.40+2	1.14+0	
2.00+0	2.10+2	1.04+0	2.50+0	3.10+2	1.12+0	3.00+0	4.00+2	1.16+0	
1.50-1	8.30-2	1.10+0	2.00-1	2.98-1	1.11+0	2.50-1	6.20-1	9.43-1	73
3.00-1	1.24+0	9.44-1	3.50-1	2.00+0	8.74-1	4.00-1	2.80+0	7.70-1	
4.50-1	3.80+0	7.06-1	5.50-1	6.60+0	6.51-1	6.50-1	1.09+1	6.54-1	
7.50-1	1.83+1	7.36-1	8.50-1	2.63+1	7.62-1	9.50-1	3.68+1	8.05-1	
1.05+0	4.14+1	7.14-1							
1.00+0	5.45+1	1.06+0	1.10+0	6.83+1	1.06+0	1.20+0	8.37+1	1.07+0	77
1.30+0	1.02+2	1.10+0	1.40+0	1.18+2	1.10+0	1.50+0	1.36+2	1.10+0	
1.60+0	1.55+2	1.12+0	1.70+0	1.74+2	1.13+0	1.80+0	1.93+2	1.13+0	
1.90+0	2.08+2	1.12+0	2.00+0	2.30+2	1.14+0	2.10+0	2.50+2	1.15+0	
2.20+0	2.67+2	1.15+0	2.30+0	2.85+2	1.15+0	2.40+0	3.00+2	1.14+0	
2.50+0	3.18+2	1.15+0	2.60+0	3.31+2	1.14+0	2.70+0	3.46+2	1.13+0	
2.80+0	3.59+2	1.13+0	2.90+0	3.72+2	1.12+0	3.00+0	3.85+2	1.12+0	
1.00-1	7.45-3	7.67-1	1.10-1	1.45-2	8.89-1	1.20-1	2.64-2	1.03+0	79
1.30-1	4.58-2	1.20+0	1.40-1	7.63-2	1.39+0	1.50-1	1.23-1	1.63+0	
1.00+0	3.70+1	7.16-1	2.00+0	1.54+2	7.63-1	5.00+0	5.20+2	9.86-1	84
3.00+0	3.70+2	1.08+0							94
5.00-1	6.65+0	8.82-1	5.50-1	9.80+0	9.66-1	6.00-1	1.32+1	1.00+0	98
6.50-1	1.63+1	9.79-1	7.00-1	1.99+1	9.68-1	7.50-1	2.32+1	9.33-1	
8.00-1	2.82+1	9.55-1	8.50-1	3.30+1	9.57-1	9.00-1	3.78+1	9.46-1	
9.50-1	4.41+1	9.65-1	1.00+0	4.89+1	9.47-1	1.05+0	5.43+1	9.36-1	
1.10+0	6.11+1	9.47-1	1.15+0	6.77+1	9.50-1	1.20+0	7.34+1	9.37-1	
1.25+0	8.06+1	9.44-1	1.30+0	8.97+1	9.68-1	1.35+0	9.61+1	9.59-1	
1.40+0	1.05+2	9.75-1	1.45+0	1.12+2	9.71-1	1.50+0	1.18+2	9.58-1	
1.55+0	1.27+2	9.72-1	1.60+0	1.34+2	9.69-1	1.65+0	1.41+2	9.65-1	
1.70+0	1.49+2	9.66-1	1.75+0	1.55+2	9.54-1	1.80+0	1.61+2	9.45-1	
1.85+0	1.72+2	9.66-1	1.90+0	1.78+2	9.58-1	1.95+0	1.90+2	9.81-1	
2.00+0	1.96+2	9.71-1							
3.80-2	4.00-6	1.11+0	4.00-2	1.00-5	1.49+0	4.50-2	2.50-5	1.02+0	101
5.00-2	6.00-5	8.64-1	5.50-2	1.40-4	8.60-1	6.00-2	3.00-4	9.00-1	
7.00-2	1.00-3	9.50-1							
4.00+0	4.80+2	1.06+0	6.00+0	6.81+2	1.18+0	8.00+0	7.16+2	1.13+0	105
1.00+1	7.56+2	1.16+0	1.20+1	7.40+2	1.14+0	1.40+1	7.16+2	1.13+0	
1.60+1	6.70+2	1.09+0	1.80+1	6.13+2	1.03+0	2.00+1	5.93+2	1.03+0	
2.20+1	5.65+2	1.02+0							
2.00-1	2.70-1	1.01+0	2.25-1	3.90-1	9.00-1	2.50-1	6.20-1	9.43-1	106
2.75-1	8.20-1	8.65-1	3.00-1	1.32+0	1.01+0	3.25-1	1.43+0	8.13-1	

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
3.50-1	2.01+0	8.78-1	3.75-1	2.48+0	8.51-1	4.00-1	3.17+0	8.72-1	
4.25-1	3.63+0	8.15-1	4.50-1	4.56+0	8.48-1	4.75-1	5.57+0	8.69-1	
5.00-1	6.64+0	8.80-1							
6.00-2	2.09-4	6.27-1	7.00-2	6.16-4	5.85-1	8.00-2	1.57-3	6.08-1	108
9.00-2	3.57-3	6.75-1	1.00-1	7.45-3	7.67-1	1.10-1	1.45-2	8.89-1	
1.20-1	2.64-2	1.03+0	1.30-1	4.58-2	1.20+0	1.40-1	7.63-2	1.39+0	
1.50-1	1.23-1	1.63+0							
1.00+0	5.28+1	1.02+0	1.25+0	8.53+1	9.99-1	1.50+0	1.30+2	1.06+0	112
1.75+0	1.68+2	1.03+0	2.00+0	2.11+2	1.05+0	2.25+0	2.48+2	1.03+0	
2.50+0	2.88+2	1.04+0	2.75+0	3.33+2	1.07+0	3.00+0	3.71+2	1.08+0	
2.00-1	2.74-1	1.02+0	3.00-1	1.48+0	1.13+0	4.00-1	3.99+0	1.10+0	113
5.00-1	8.00+0	1.06+0	6.00-1	1.32+1	1.00+0	8.00-1	3.00+1	1.02+0	
1.00+0	5.15+1	9.97-1	1.20+0	8.12+1	1.04+0	1.60+0	1.43+2	1.03+0	
2.00+0	2.15+2	1.07+0	2.40+0	2.72+2	1.04+0				
1.00-1	3.77-3	<b>3.88-1</b>	1.25-1	1.56-2	4.95-1	1.50-1	4.34-2	5.73-1	118
1.75-1	9.50-2	6.29-1	2.00-1	1.77-1	6.61-1	3.00-1	9.57-1	7.29-1	
4.00-1	2.89+0	7.95-1	5.00-1	6.08+0	8.06-1	6.00-1	1.08+1	8.19-1	
7.00-1	1.70+1	8.27-1	8.00-1	2.50+1	8.47-1				
3.00-1	1.20+0	9.14-1	4.00-1	3.37+0	9.27-1	5.00-1	7.10+0	9.41-1	132
6.00-1	1.26+1	9.55-1	7.00-1	1.90+1	9.24-1	8.00-1	2.87+1	9.72-1	
1.00+0	4.88+1	9.45-1	1.20+0	7.67+1	9.79-1	1.40+0	1.08+2	1.00+0	
1.60+0	1.38+2	9.98-1	1.80+0	1.73+2	1.02+0	2.00+0	2.08+2	1.03+0	
2.20+0	2.36+2	1.02+0	2.40+0	2.69+2	1.02+0				
5.00-1	6.88+0	9.12-1	6.00-1	1.24+1	9.40-1	7.00-1	1.94+1	9.44-1	137
8.00-1	2.80+1	9.49-1	1.00+0	4.88+1	9.45-1	1.20+0	7.29+1	9.31-1	
1.40+0	1.02+2	9.47-1	1.60+0	1.32+2	9.54-1	1.80+0	1.65+2	9.69-1	
2.00+0	1.97+2	9.76-1	2.20+0	2.26+2	9.72-1	2.30+0	2.50+2	1.01+0	
2.40+0	2.61+2	9.95-1	2.50+0	2.76+2	9.97-1				
1.50+0	8.95+1	7.27-1	2.10+0	2.13+2	9.80-1	2.60+0	2.85+2	9.78-1	148
3.10+0	3.64+2	1.02+0	3.60+0	4.40+2	1.07+0				
1.00+0	4.94+1	9.57-1							149
<b>23</b>	<b>Vanadium</b>				<b>Fluorescence yield = 0.243</b>				
1.00-1	5.50-3	9.90-1	1.25-1	1.70-2	8.93-1	1.50-1	4.30-2	9.10-1	36
2.25+0	1.43+2	7.27-1							47
4.00-1	3.00+0	1.19+0	5.00-1	6.30+0	1.19+0	6.00-1	1.60+1	1.70+0	69
7.00-1	1.73+1	1.16+0	8.00-1	2.60+1	1.20+0	9.00-1	3.50+1	1.18+0	
1.00+0	4.50+1	1.16+0	1.10+0	5.80+1	1.19+0	1.20+0	7.00+1	1.17+0	
1.30+0	8.70+1	1.22+0	1.40+0	1.00+2	1.19+0	1.50+0	1.18+2	1.22+0	
1.60+0	1.34+2	1.22+0	1.70+0	1.47+2	1.20+0	1.80+0	1.69+2	1.25+0	
1.90+0	1.81+2	1.21+0	2.00+0	1.97+2	1.21+0	2.10+0	2.13+2	1.21+0	
2.20+0	2.19+2	1.15+0	2.30+0	2.44+2	1.20+0				

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
1.50-1	5.80-2	1.23+0	2.00-1	1.65-1	9.51-1	2.50-1	4.30-1	9.84-1	73
3.00-1	8.80-1	9.93-1	3.50-1	1.20+0	7.66-1	4.00-1	2.30+0	9.14-1	
4.50-1	3.00+0	7.98-1	5.50-1	5.00+0	6.94-1	6.50-1	8.80+0	7.33-1	
7.50-1	1.60+1	8.81-1	8.50-1	1.78+1	6.96-1	9.50-1	2.55+1	7.49-1	
9.50-1	2.27+1	6.66-1							76
1.00+0	3.52+1	9.07-1	1.10+0	4.38+1	8.96-1	1.30+0	6.63+1	9.27-1	86
1.50+0	8.60+1	8.92-1	1.70+0	1.11+2	9.06-1	1.90+0	1.37+2	9.17-1	
2.00+0	1.51+2	9.26-1	2.10+0	1.64+2	9.30-1	2.30+0	1.90+2	9.35-1	
2.50+0	2.14+2	9.34-1	2.70+0	2.35+2	9.24-1	2.80+0	2.45+2	9.19-1	
2.90+0	2.53+2	9.08-1	3.00+0	2.63+2	9.06-1				
5.00-1	4.80+0	9.03-1	5.50-1	6.92+0	9.60-1	6.00-1	8.75+0	9.28-1	98
6.50-1	1.13+1	9.41-1	7.00-1	1.40+1	9.39-1	7.50-1	1.66+1	9.15-1	
8.00-1	2.06+1	9.49-1	8.50-1	2.52+1	9.85-1	9.00-1	2.92+1	9.85-1	
9.50-1	3.39+1	9.95-1	1.00+0	3.84+1	9.89-1	1.05+0	4.29+1	9.80-1	
1.10+0	4.72+1	9.66-1	1.15+0	5.43+1	1.00+0	1.20+0	5.91+1	9.87-1	
1.25+0	6.57+1	1.00+0	1.30+0	7.17+1	1.00+0	1.35+0	7.79+1	1.00+0	
1.40+0	8.35+1	9.98-1	1.45+0	9.05+1	1.01+0	1.50+0	9.48+1	9.83-1	
1.55+0	1.05+2	1.02+0	1.60+0	1.10+2	1.01+0	1.65+0	1.16+2	9.99-1	
1.70+0	1.23+2	1.00+0	1.75+0	1.28+2	9.92-1	1.80+0	1.34+2	9.88-1	
1.85+0	1.40+2	9.82-1	1.90+0	1.45+2	9.70-1	1.95+0	1.56+2	9.97-1	
2.00+0	1.56+2	9.57-1							
2.00-1	1.45-1	8.36-1	2.25-1	2.42-1	8.49-1	2.50-1	3.61-1	8.26-1	106
2.75-1	4.71-1	7.41-1	3.00-1	7.80-1	8.80-1	3.25-1	1.09+0	9.11-1	
3.50-1	1.42+0	9.06-1	3.75-1	1.90+0	9.48-1	4.00-1	2.23+0	8.86-1	
4.25-1	2.82+0	9.11-1	4.50-1	3.31+0	8.81-1	4.75-1	4.30+0	9.57-1	
5.00-1	5.17+0	9.73-1							
5.00-1	5.63+0	1.06+0	6.00-1	1.10+1	1.17+0	8.00-1	2.04+1	9.40-1	113
1.00+0	4.00+1	1.03+0	1.20+0	5.75+1	9.60-1	1.60+0	1.10+2	1.01+0	
2.00+0	1.54+2	9.45-1	2.40+0	2.21+2	1.02+0				
1.00+0	4.13+1	1.06+0	1.20+0	6.20+1	1.04+0	1.40+0	8.40+1	1.00+0	137
1.60+0	1.09+2	9.96-1	1.80+0	1.35+2	9.95-1	2.00+0	1.71+2	1.05+0	
2.20+0	1.89+2	9.95-1							
<b>24 Chromium</b> <b>Fluorescence yield = 0.275</b>									
1.00-1	2.80-3	8.78-1	1.25-1	1.00-2	8.65-1	1.50-1	2.70-2	9.08-1	36
5.00-1	4.90+0	1.29+0	6.00-1	8.70+0	1.28+0	7.00-1	1.37+1	1.26+0	69
8.00-1	2.09+1	1.30+0	9.00-1	2.70+1	1.21+0	1.00+0	3.60+1	1.23+0	
1.10+0	4.70+1	1.26+0	1.20+0	5.50+1	1.20+0	1.30+0	6.80+1	1.23+0	
1.40+0	8.20+1	1.26+0	1.50+0	9.40+1	1.24+0	1.60+0	1.08+2	1.25+0	
1.70+0	1.24+2	1.27+0	1.80+0	1.35+2	1.24+0	1.90+0	1.49+2	1.24+0	
2.00+0	1.67+2	1.27+0	2.10+0	1.77+2	1.23+0	2.20+0	1.98+2	1.28+0	
2.00-1	6.80-2	5.97-1	2.50-1	2.10-1	7.15-1	3.00-1	4.10-1	6.77-1	73
3.50-1	7.80-1	7.19-1	4.00-1	1.13+0	6.42-1	4.50-1	1.62+0	6.10-1	
5.50-1	2.80+0	5.41-1	6.50-1	4.90+0	5.61-1	7.50-1	8.20+0	6.14-1	
8.50-1	1.16+1	6.09-1	9.50-1	1.80+1	7.00-1	1.05+0	2.15+1	6.48-1	

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
9.50-1	1.73+1	6.73-1							76
1.00-1	3.60-3	1.13+0	1.10-1	6.20-3	1.10+0	1.20-1	1.07-2	1.16+0	79
1.30-1	1.80-2	1.26+0	1.40-1	2.80-2	1.33+0	1.50-1	4.00-2	1.35+0	
9.00-1	1.55+1	6.96-1	9.50-1	1.73+1	6.73-1	1.00+0	2.50+1	8.54-1	80
1.10+0	2.73+1	7.31-1	1.20+0	3.60+1	7.83-1	1.30+0	4.17+1	7.52-1	
1.40+0	5.13+1	7.85-1	1.60+0	6.06+1	7.00-1	1.70+0	7.19+1	7.36-1	
1.80+0	8.46+1	7.75-1	1.90+0	8.18+1	6.80-1	2.00+0	8.60+1	6.53-1	
2.10+0	1.20+2	8.35-1	2.20+0	1.13+2	7.28-1	2.30+0	1.28+2	7.68-1	
2.40+0	1.37+2	7.68-1	2.50+0	1.71+2	9.00-1	2.60+0	1.54+2	7.66-1	
2.70+0	1.68+2	7.91-1	2.80+0	1.69+2	7.56-1	3.00+0	2.07+2	8.46-1	
3.40+0	2.50+2	8.76-1	3.60+0	2.65+2	8.71-1	3.80+0	2.79+2	8.65-1	
4.00+0	3.53+2	1.04+0							
3.00+0	2.81+2	1.15+0	5.00+0	4.86+2	1.18+0	7.00+0	6.19+2	1.22+0	94
9.00+0	6.63+2	1.20+0	1.10+1	6.78+2	1.19+0				
5.00-1	3.52+0	9.30-1	5.50-1	4.84+0	9.36-1	6.00-1	6.57+0	9.64-1	98
6.50-1	8.19+0	9.38-1	7.00-1	1.02+1	9.35-1	7.50-1	1.24+1	9.28-1	
8.00-1	1.49+1	9.27-1	8.50-1	1.79+1	9.40-1	9.00-1	2.07+1	9.30-1	
9.50-1	2.38+1	9.26-1	1.00+0	2.80+1	9.56-1	1.05+0	3.13+1	9.44-1	
1.10+0	3.60+1	9.65-1	1.15+0	3.87+1	9.31-1	1.20+0	4.40+1	9.57-1	
1.25+0	4.92+1	9.72-1	1.30+0	5.29+1	9.53-1	1.35+0	5.77+1	9.57-1	
1.40+0	6.31+1	9.66-1	1.45+0	6.74+1	9.55-1	1.50+0	7.26+1	9.58-1	
1.55+0	7.82+1	9.64-1	1.60+0	8.49+1	9.81-1	1.65+0	8.93+1	9.69-1	
1.70+0	9.46+1	9.69-1	1.75+0	1.00+2	9.68-1	1.80+0	1.06+2	9.71-1	
1.85+0	1.13+2	9.86-1	1.90+0	1.17+2	9.73-1	1.95+0	1.22+2	9.69-1	
2.00+0	1.28+2	9.72-1							
2.00-1	1.07-1	9.39-1	2.25-1	1.69-1	8.93-1	2.50-1	2.77-1	9.44-1	106
2.75-1	4.05-1	9.39-1	3.00-1	5.80-1	9.57-1	3.25-1	7.80-1	9.48-1	
3.50-1	1.02+0	9.40-1	3.75-1	1.31+0	9.38-1	4.00-1	1.68+0	9.55-1	
4.25-1	2.02+0	9.28-1	4.50-1	2.47+0	9.31-1	4.75-1	2.98+0	9.34-1	
5.00-1	3.66+0	9.66-1							
7.00+0	3.80+2	7.51-1							114
7.00+0	3.95+2	7.81-1							125
3.00-1	6.37-1	1.05+0	4.00-1	1.71+0	9.72-1	5.00-1	3.86+0	1.02+0	132
6.00-1	6.97+0	1.02+0	7.00-1	1.05+1	9.62-1	8.00-1	1.59+1	9.89-1	
1.00+0	2.85+1	9.73-1	1.20+0	4.48+1	9.74-1	1.40+0	6.37+1	9.75-1	
1.60+0	8.46+1	9.77-1	1.80+0	1.09+2	9.99-1	2.00+0	1.27+2	9.64-1	
2.20+0	1.52+2	9.79-1	2.40+0	1.78+2	9.98-1				
8.00-1	1.50+1	9.33-1	1.00+0	2.76+1	9.42-1	1.20+0	4.31+1	9.37-1	137
1.40+0	6.18+1	9.46-1	1.60+0	8.29+1	9.57-1	1.80+0	1.08+2	9.90-1	
2.00+0	1.29+2	9.79-1	2.20+0	1.48+2	9.53-1	2.30+0	1.61+2	9.66-1	
2.40+0	1.72+2	9.64-1	2.50+0	1.79+2	9.42-1				

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
1.00+0	2.83+1	9.66-1	1.10+0	3.79+1	1.02+0	1.20+0	4.90+1	1.07+0	152
1.30+0	5.74+1	1.03+0	1.40+0	7.27+1	1.11+0	1.50+0	8.83+1	1.16+0	
1.60+0	9.93+1	1.15+0	1.70+0	1.18+2	1.21+0	1.80+0	1.40+2	1.28+0	
1.90+0	1.64+2	1.36+0	2.00+0	1.82+2	1.38+0	2.10+0	2.09+2	1.46+0	
2.20+0	2.17+2	1.40+0	2.30+0	2.41+2	1.45+0	2.40+0	2.70+2	1.51+0	
2.50+0	3.03+2	1.59+0	2.60+0	3.34+2	1.66+0	2.70+0	3.57+2	1.68+0	
2.80+0	3.82+2	1.71+0	2.90+0	4.30+2	1.84+0	3.00+0	4.53+2	1.85+0	
<b>25</b>	<b>Manganese</b>				<b>Fluorescence yield = 0.308</b>				
1.00-1	1.10-3	6.02-1	1.25-1	4.30-3	6.13-1	1.50-1	1.30-2	6.94-1	36
2.50+0	1.56+2	9.99-1	3.00+0	2.26+2	1.10+0	4.50+0	3.56+2	1.08+0	52
6.00+0	4.46+2	1.07+0	6.50+0	4.37+2	9.98-1	7.00+0	5.19+2	1.14+0	
8.00+0	5.28+2	1.09+0	8.50+0	5.97+2	1.20+0	9.00+0	5.68+2	1.12+0	
1.00+1	5.47+2	1.05+0	1.05+1	5.64+2	1.07+0	1.10+1	6.19+2	1.16+0	
1.20+1	6.56+2	1.22+0							
1.30-1	8.20-3	9.39-1	1.50-1	1.23-2	6.57-1	1.95-1	5.39-2	8.07-1	57
2.50-1	1.65-1	8.33-1	3.00-1	2.70-1	6.49-1	3.30-1	3.64-1	6.04-1	
3.60-1	5.13-1	6.12-1	3.85-1	7.02-1	6.53-1	4.15-1	8.15-1	5.77-1	
4.00-1	1.55+0	1.25+0	5.00-1	3.20+0	1.18+0	6.00-1	5.70+0	1.15+0	69
7.00-1	9.30+0	1.16+0	8.00-1	1.45+1	1.22+0	9.00-1	1.98+1	1.19+0	
1.00+0	2.60+1	1.17+0	1.10+0	3.40+1	1.20+0	1.20+0	4.10+1	1.16+0	
1.30+0	5.10+1	1.19+0	1.40+0	6.00+1	1.18+0	1.50+0	6.90+1	1.16+0	
1.60+0	8.20+1	1.20+0	1.70+0	9.30+1	1.20+0	1.80+0	1.06+2	1.22+0	
1.90+0	1.19+2	1.23+0	2.00+0	1.33+2	1.25+0	2.10+0	1.40+2	1.21+0	
2.20+0	1.59+2	1.26+0	2.30+0	1.71+2	1.25+0				
3.00+0	2.18+2	1.06+0							94
5.00-1	2.64+0	9.76-1	5.50-1	3.64+0	9.78-1	6.00-1	4.72+0	9.56-1	98
6.50-1	6.12+0	9.62-1	7.00-1	7.44+0	9.30-1	7.50-1	9.14+0	9.28-1	
8.00-1	1.11+1	9.32-1	8.50-1	1.32+1	9.31-1	9.00-1	1.55+1	9.31-1	
9.50-1	1.81+1	9.37-1	1.00+0	2.10+1	9.47-1	1.05+0	2.38+1	9.45-1	
1.10+0	2.68+1	9.45-1	1.15+0	3.00+1	9.45-1	1.20+0	3.35+1	9.48-1	
1.25+0	3.75+1	9.61-1	1.30+0	4.03+1	9.42-1	1.35+0	4.43+1	9.47-1	
1.40+0	4.83+1	9.48-1	1.45+0	5.37+1	9.75-1	1.50+0	5.62+1	9.47-1	
1.55+0	6.12+1	9.60-1	1.60+0	6.58+1	9.62-1	1.65+0	7.04+1	9.67-1	
1.70+0	7.29+1	9.41-1	1.75+0	7.93+1	9.65-1	1.80+0	8.29+1	9.53-1	
1.85+0	8.69+1	9.47-1	1.90+0	9.13+1	9.45-1	1.95+0	9.92+1	9.77-1	
2.00+0	1.03+2	9.68-1							
2.00-1	4.80-2	6.40-1	2.25-1	8.90-2	7.05-1	2.50-1	1.55-1	7.82-1	106
2.75-1	2.21-1	7.53-1	3.00-1	3.45-1	8.29-1	3.25-1	4.60-1	8.09-1	
3.50-1	6.60-1	8.75-1	3.75-1	8.50-1	8.71-1	4.00-1	1.11+0	8.98-1	
4.25-1	1.43+0	9.30-1	4.50-1	1.73+0	9.20-1	4.75-1	2.15+0	9.47-1	
5.00-1	2.60+0	9.61-1							
5.00-1	2.37+0	8.76-1	6.00-1	4.24+0	8.59-1	8.00-1	1.08+1	9.07-1	113
1.00+0	2.18+1	9.83-1	1.20+0	3.52+1	9.96-1	1.60+0	6.63+1	9.70-1	
2.00+0	1.03+2	9.68-1	2.40+0	1.59+2	1.09+0				

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
<b>26 Iron</b> <b>Fluorescence yield = 0.34</b>									
1.55+0	4.40+1	8.82-1	3.90+0	2.70+2	1.12+0				2
1.40-1	7.40-3	9.16-1	1.50-1	1.10-2	9.37-1	1.60-1	1.40-2	8.49-1	6
1.80-1	2.50-2	8.39-1	2.00-1	4.00-2	8.12-1	2.20-1	5.80-2	7.60-1	
2.40-1	8.30-2	7.41-1	2.60-1	1.10-1	6.97-1	2.80-1	1.50-1	6.96-1	
3.00-1	2.00-1	7.01-1	3.20-1	2.50-1	6.76-1	3.40-1	3.00-1	6.40-1	
3.60-1	3.80-1	6.50-1	4.00-1	5.70-1	6.56-1	4.40-1	8.00-1	6.50-1	
7.40-1	3.60+0	5.17-1	9.35-1	6.80+0	4.91-1	1.04+0	1.10+1	5.92-1	
1.20+0	1.40+1	5.20-1	1.30+0	1.70+1	5.16-1				
4.54-1	5.53-1	<b>4.02-1</b>							7
1.50+0	3.80+1	8.21-1							10
7.00-1	2.96+0	5.05-1	9.00-1	6.21+0	5.00-1	1.10+0	1.13+1	5.24-1	27
1.30+0	1.59+1	<b>4.82-1</b>	1.50+0	2.22+1	<b>4.80-1</b>	1.70+0	2.61+1	<b>4.28-1</b>	
1.90+0	3.31+1	<b>4.30-1</b>	2.10+0	4.19+1	<b>4.48-1</b>	2.30+0	4.92+1	<b>4.47-1</b>	
2.50+0	6.42+1	5.03-1							
1.60+2	2.29+2	1.51+0							30
1.00-1	1.20-3	1.17+0	1.25-1	4.90-3	1.16+0	1.50-1	1.20-2	1.02+0	36
1.00+0	1.40+1	8.39-1	1.50+0	4.40+1	9.51-1	2.00+0	8.10+1	9.52-1	38
2.50+0	1.20+2	9.40-1	3.00+0	1.85+2	1.09+0	3.50+0	2.20+2	1.04+0	
4.00+0	2.50+2	1.00+0	4.50+0	2.80+2	9.88-1	5.00+0	3.10+2	9.86-1	
5.50+0	3.30+2	9.65-1							
1.00+0	1.10+1	6.59-1	2.25+0	7.40+1	6.98-1	3.00+0	1.25+2	7.33-1	47
2.50+0	1.19+2	9.32-1	3.00+0	1.77+2	1.04+0	4.00+0	2.68+2	1.08+0	52
4.50+0	2.75+2	9.70-1	5.00+0	3.12+2	9.93-1	6.00+0	4.05+2	1.11+0	
6.50+0	3.86+2	9.95-1	7.00+0	4.29+2	1.05+0	8.00+0	4.49+2	1.03+0	
8.50+0	4.91+2	1.09+0	9.00+0	5.00+2	1.09+0	1.00+1	4.90+2	1.03+0	
1.05+1	4.86+2	1.01+0	1.10+1	4.99+2	1.02+0	1.20+1	4.94+2	9.92-1	
5.00-1	1.40+0	7.25-1	6.00-1	4.40+0	1.23+0	7.00-1	7.20+0	1.23+0	53
8.00-1	9.50+0	1.08+0	9.00-1	1.42+1	1.14+0	1.00+0	1.69+1	1.01+0	
1.10+0	2.40+1	1.11+0	1.20+0	2.96+1	1.10+0	1.30+0	3.44+1	1.04+0	
1.40+0	3.92+1	9.96-1	1.50+0	5.02+1	1.08+0	1.60+0	5.69+1	1.06+0	
1.70+0	6.57+1	1.08+0	1.80+0	6.92+1	1.01+0	1.90+0	7.27+1	9.45-1	
2.00+0	7.65+1	8.99-1							
1.50+0	4.66+1	1.01+0	2.00+0	8.68+1	1.02+0	2.50+0	1.32+2	1.03+0	55
3.00+0	1.75+2	1.03+0	4.97+0	3.24+2	1.04+0	5.96+0	3.88+2	1.06+0	
6.94+0	4.39+2	1.09+0	6.96+0	4.32+2	1.07+0	8.94+0	4.95+2	1.08+0	
1.09+1	5.29+2	1.08+0							
1.30-1	8.00-3	1.51+0	1.50-1	1.56-2	1.33+0	1.95-1	4.45-2	1.02+0	57
2.55-1	1.12-1	7.70-1	3.00-1	1.86-1	6.51-1	3.30-1	2.76-1	6.61-1	
3.60-1	3.46-1	5.92-1	3.90-1	4.22-1	5.34-1	4.15-1	5.26-1	5.28-1	

TABLE 2. *K*-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
8.30-1	8.10+0	8.25-1	1.58+0	5.90+1	1.13+0	2.56+0	1.34+2	1.01+0	66
3.28+0	2.27+2	1.17+0							
4.00-1	1.24+0	1.43+0	5.00-1	2.30+0	1.19+0	6.00-1	4.20+0	1.17+0	69
7.00-1	6.70+0	1.14+0	8.00-1	1.05+1	1.19+0	9.00-1	1.39+1	1.12+0	
1.00+0	1.87+1	1.12+0	1.10+0	2.50+1	1.16+0	1.20+0	3.10+1	1.15+0	
1.30+0	3.90+1	1.18+0	1.40+0	4.70+1	1.19+0	1.50+0	5.40+1	1.17+0	
1.60+0	6.10+1	1.14+0	1.70+0	7.00+1	1.15+0	1.80+0	8.20+1	1.19+0	
1.90+0	9.30+1	1.21+0	2.00+0	9.90+1	1.16+0	2.10+0	1.11+2	1.19+0	
2.20+0	1.20+2	1.18+0	2.30+0	1.28+2	1.16+0				
2.00-1	2.80-2	5.68-1	2.50-1	9.40-2	7.04-1	3.00-1	1.90-1	6.65-1	73
3.50-1	3.50-1	6.67-1	4.00-1	5.40-1	6.22-1	4.50-1	8.90-1	6.67-1	
5.50-1	1.88+0	7.02-1	6.50-1	3.26+0	7.04-1	7.50-1	5.50+0	7.59-1	
8.50-1	8.80+0	8.35-1	9.50-1	1.35+1	9.32-1	1.05+0	1.40+1	7.34-1	
9.50-1	9.30+0	6.42-1							76
1.00-1	4.73-4	4.61-1	1.10-1	9.95-4	5.14-1	1.20-1	1.96-3	5.93-1	79
1.30-1	3.66-3	6.89-1	1.40-1	6.50-3	8.05-1	1.50-1	1.11-2	9.45-1	
1.00+0	1.26+1	7.55-1	2.00+0	6.37+1	7.48-1	5.00+0	2.37+2	7.54-1	84
1.00+0	1.49+1	8.93-1	1.20+0	2.32+1	8.61-1	1.50+0	3.90+1	8.43-1	86
1.80+0	5.92+1	8.60-1	2.00+0	7.39+1	8.68-1	2.30+0	9.71+1	8.81-1	
2.50+0	1.12+2	8.78-1	2.80+0	1.29+2	8.42-1	2.90+0	1.34+2	8.27-1	
3.00+0	1.76+2	1.03+0							94
5.00-1	2.04+0	1.06+0	5.50-1	2.82+0	1.05+0	6.00-1	3.48+0	9.74-1	98
6.50-1	4.48+0	9.67-1	7.00-1	6.62+0	1.13+0	7.50-1	6.83+0	9.43-1	
8.00-1	8.47+0	9.62-1	8.50-1	1.13+1	1.07+0	9.00-1	1.21+1	9.74-1	
9.50-1	1.40+1	9.67-1	1.00+0	1.57+1	9.40-1	1.05+0	1.81+1	9.50-1	
1.10+0	2.19+1	1.01+0	1.15+0	2.49+1	1.03+0	1.20+0	2.71+1	1.01+0	
1.25+0	3.02+1	1.01+0	1.30+0	3.23+1	9.80-1	1.35+0	3.67+1	1.02+0	
1.40+0	4.10+1	1.04+0	1.45+0	4.36+1	1.02+0	1.50+0	4.51+1	9.75-1	
1.55+0	4.77+1	9.56-1	1.60+0	5.22+1	9.76-1	1.65+0	5.53+1	9.67-1	
1.70+0	6.02+1	9.86-1	1.75+0	6.26+1	9.63-1	1.80+0	6.60+1	9.59-1	
1.85+0	6.94+1	9.53-1	1.90+0	7.53+1	9.79-1	1.95+0	7.79+1	9.61-1	
2.00+0	8.31+1	9.76-1							
4.00+0	2.67+2	1.07+0	6.00+0	4.00+2	1.09+0	8.00+0	5.00+2	1.14+0	105
1.00+1	5.48+2	1.15+0	1.20+1	6.04+2	1.21+0	1.40+1	5.69+2	1.12+0	
1.60+1	5.86+2	1.16+0	1.80+1	6.18+2	1.23+0	2.00+1	5.83+2	1.18+0	
2.20+1	5.34+2	1.11+0							
2.50-1	1.02-1	7.63-1	2.75-1	1.93-1	9.65-1	3.00-1	2.44-1	8.55-1	106
3.25-1	3.47-1	8.82-1	3.50-1	4.70-1	8.96-1	3.75-1	6.60-1	9.67-1	
4.00-1	7.90-1	9.10-1	4.25-1	1.03+0	9.49-1	4.50-1	1.26+0	9.45-1	
4.75-1	1.46+0	9.04-1	5.00-1	1.83+0	9.47-1				
7.00-2	2.85-5	4.76-1	8.00-2	8.16-5	4.24-1	9.00-2	2.06-4	4.27-1	108
1.00-1	4.73-4	4.61-1	1.10-1	9.95-4	5.14-1	1.20-1	1.96-3	5.93-1	

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_i$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_i$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_i$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
1.30-1	3.66-3	6.89-1	1.40-1	6.50-3	8.05-1	1.50-1	1.11-2	9.45-1	
5.00-1	1.94+0	1.00+0	6.00-1	3.82+0	1.07+0	8.00-1	9.23+0	1.05+0	113
1.00+0	1.68+1	1.01+0	1.20+0	2.74+1	1.02+0	1.60+0	5.00+1	9.35-1	
2.00+0	8.57+1	1.01+0	2.40+0	1.13+2	9.51-1				
1.00-1	3.09-4	<b>3.01-1</b>	1.25-1	1.67-3	<b>3.96-1</b>	1.50-1	5.71-3	4.86-1	118
1.75-1	1.44-2	5.55-1	2.00-1	2.98-2	6.05-1	2.50-1	9.15-2	6.85-1	
3.00-1	2.13-1	7.46-1	4.00-1	7.21-1	8.30-1	5.00-1	1.56+0	8.07-1	
7.00-1	4.86+0	8.30-1	8.00-1	7.31+0	8.30-1				
2.00-1	5.78-2	1.17+0	2.50-1	1.46-1	1.09+0	3.00-1	3.09-1	1.08+0	121
4.00-1	8.96-1	1.03+0	5.00-1	2.05+0	1.06+0	6.00-1	3.73+0	1.04+0	
7.00-1	5.99+0	1.02+0	8.00-1	9.10+0	1.03+0	9.00-1	1.27+1	1.02+0	
1.00+0	1.76+1	1.05+0	1.10+0	2.05+1	9.50-1	1.20+0	2.76+1	1.02+0	152
1.30+0	3.11+1	9.43-1	1.40+0	3.81+1	9.68-1	1.50+0	4.25+1	9.18-1	
1.60+0	5.19+1	9.71-1	1.70+0	5.86+1	9.60-1	1.80+0	6.50+1	9.44-1	
1.90+0	6.77+1	8.80-1	2.00+0	6.86+1	8.06-1	2.10+0	7.41+1	7.93-1	
2.20+0	7.76+1	7.62-1	2.30+0	9.02+1	8.19-1	2.40+0	9.76+1	8.22-1	
2.50+0	1.03+2	8.07-1	2.60+0	1.05+2	7.70-1	2.70+0	1.10+2	7.60-1	
2.80+0	1.18+2	7.70-1	2.90+0	1.23+2	7.59-1	3.00+0	1.29+2	7.57-1	
<b>27 Cobalt                   Fluorescence yield = 0.373</b>									
1.00-1	5.00-4	8.70-1	1.25-1	2.30-3	9.03-1	1.50-1	6.80-3	9.19-1	36
1.00+0	7.60+0	6.03-1	1.50+0	4.20+1	1.16+0	2.00+0	7.80+1	1.15+0	38
2.50+0	1.10+2	1.06+0	3.00+0	1.50+2	1.06+0	3.50+0	1.90+2	1.07+0	
4.00+0	2.10+2	9.92-1	4.50+0	2.40+2	9.86-1	5.00+0	2.70+2	9.91-1	
5.50+0	3.10+2	1.04+0	6.00+0	3.20+2	9.94-1				
5.00-1	1.60+0	1.15+0	6.00-1	2.70+0	1.04+0	7.00-1	4.20+0	9.74-1	53
8.00-1	6.40+0	9.78-1	9.00-1	9.50+0	1.02+0	1.00+0	1.27+1	1.01+0	
1.10+0	1.65+1	1.00+0	1.20+0	1.99+1	9.60-1	1.30+0	2.47+1	9.73-1	
1.40+0	2.79+1	9.12-1	1.50+0	3.34+1	9.25-1	1.60+0	3.91+1	9.32-1	
1.70+0	4.42+1	9.18-1	1.80+0	4.60+1	8.43-1	1.90+0	5.35+1	8.73-1	
2.00+0	5.93+1	8.71-1							
1.50+0	3.72+1	1.03+0	2.00+0	7.02+1	1.03+0	2.50+0	1.09+2	1.05+0	55
3.00+0	1.46+2	1.04+0	4.96+0	2.67+2	9.89-1	5.96+0	3.20+2	1.00+0	
6.94+0	3.66+2	1.02+0	6.96+0	3.64+2	1.01+0	8.94+0	4.25+2	1.02+0	
1.09+1	4.61+2	1.03+0							
5.00-1	1.90+0	1.37+0	6.00-1	3.50+0	1.34+0	7.00-1	5.70+0	1.32+0	69
8.00-1	9.20+0	1.41+0	9.00-1	1.21+1	1.30+0	1.00+0	1.66+1	1.32+0	
1.10+0	2.10+1	1.28+0	1.20+0	2.60+1	1.25+0	1.30+0	3.20+1	1.26+0	
1.40+0	3.90+1	1.28+0	1.50+0	4.50+1	1.25+0	1.60+0	5.20+1	1.24+0	
1.70+0	5.90+1	1.22+0	1.80+0	6.70+1	1.23+0	1.90+0	8.00+1	1.30+0	
2.00+0	8.60+1	1.26+0	2.10+0	9.60+1	1.28+0	2.20+0	1.03+2	1.25+0	
2.30+0	1.10+2	1.23+0							
2.00-1	3.10-2	9.50-1	2.50-1	8.70-2	9.56-1	3.00-1	1.60-1	8.08-1	73
3.50-1	3.10-1	8.42-1	4.00-1	5.70-1	9.25-1	4.50-1	7.20-1	7.55-1	

TABLE 2. *K*-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
2.00+0	5.13+1	7.54-1	5.00+0	1.99+2	7.31-1				84
3.00+0	1.35+2	9.57-1							94
2.00-1	3.57-2	1.09+0	2.50-1	9.94-2	1.09+0	3.00-1	2.06-1	1.04+0	115
3.50-1	3.75-1	1.02+0	4.00-1	6.29-1	1.02+0	4.50-1	9.94-1	1.04+0	
5.00-1	1.35+0	9.70-1	6.00-1	2.57+0	9.87-1	7.00-1	4.04+0	9.37-1	
8.00-1	6.02+0	9.20-1	9.00-1	8.57+0	9.20-1	1.00+0	1.13+1	8.96-1	
1.10+0	1.48+1	9.01-1	1.20+0	1.88+1	9.07-1	1.30+0	2.34+1	9.22-1	
1.40+0	2.78+1	9.09-1	1.50+0	3.22+1	8.92-1	1.60+0	3.68+1	8.77-1	
1.70+0	4.27+1	8.86-1	1.80+0	4.84+1	8.87-1	1.90+0	5.52+1	9.00-1	
2.00+0	6.25+1	9.18-1							
3.00-1	2.03-1	1.03+0	4.00-1	6.21-1	1.01+0	5.00-1	1.41+0	1.01+0	132
6.00-1	2.64+0	1.01+0	7.00-1	4.34+0	1.01+0	8.00-1	6.48+0	9.90-1	
1.00+0	1.25+1	9.91-1	1.20+0	2.08+1	1.00+0	1.40+0	3.05+1	9.97-1	
1.60+0	4.31+1	1.03+0	1.80+0	5.67+1	1.04+0	2.00+0	6.82+1	1.00+0	
2.20+0	8.38+1	1.02+0	2.30+0	8.99+1	1.00+0	2.40+0	9.79+1	1.01+0	
6.00-1	2.48+0	9.52-1	7.00-1	4.11+0	9.53-1	8.00-1	6.25+0	9.55-1	137
1.00+0	1.23+1	9.75-1	1.20+0	2.00+1	9.65-1	1.40+0	2.93+1	9.58-1	
1.60+0	4.08+1	9.72-1	1.80+0	5.30+1	9.72-1	2.00+0	6.59+1	9.68-1	
2.20+0	8.00+1	9.73-1							
1.00+0	1.21+1	9.60-1	1.20+0	1.63+1	7.87-1	1.40+0	2.69+1	8.80-1	152
1.60+0	3.32+1	7.91-1	1.80+0	4.28+1	7.85-1	2.00+0	5.86+1	8.61-1	
2.20+0	6.51+1	7.92-1	2.40+0	8.06+1	8.33-1	2.60+0	8.49+1	7.63-1	
2.80+0	1.04+2	8.22-1	3.00+0	1.16+2	8.23-1				
<b>28 Nickel</b> <b>Fluorescence yield = 0.406</b>									
1.50+0	2.00+1	7.11-1	1.70+0	2.40+1	6.34-1	1.80+0	2.70+1	6.25-1	5
1.90+0	3.20+1	6.58-1	2.30+0	4.40+1	6.09-1	2.40+0	5.00+1	6.38-1	
2.50+0	5.90+1	6.96-1	2.80+0	7.60+1	7.34-1	3.10+0	9.20+1	7.49-1	
3.50+0	1.10+2	7.41-1	3.70+0	1.20+2	7.47-1				
5.00+0	2.94+2	1.26+0	8.00+0	4.09+2	1.17+0	1.00+1	4.77+2	1.21+0	20
1.40+1	5.24+2	1.21+0	1.70+1	5.17+2	1.17+0	2.00+1	5.46+2	1.24+0	
2.40+1	5.10+2	1.19+0	2.80+1	5.06+2	1.23+0				
1.00-1	2.50-4	7.90-1	1.25-1	1.40-3	9.15-1	1.50-1	4.50-3	9.71-1	36
1.00+0	7.20+0	7.56-1	2.25+0	5.40+1	7.81-1	3.00+0	9.20+1	7.88-1	47
2.50+0	7.89+1	9.31-1	3.00+0	1.23+2	1.05+0	4.00+0	1.82+2	1.02+0	52
4.50+0	2.12+2	1.02+0	5.00+0	2.49+2	1.06+0	6.00+0	2.67+2	9.50-1	
6.50+0	2.56+2	8.50-1	7.00+0	3.53+2	1.11+0	8.00+0	3.18+2	9.09-1	
8.50+0	4.18+2	1.15+0	9.00+0	3.62+2	9.67-1	1.00+1	4.06+2	1.03+0	
1.05+1	4.55+2	1.13+0	1.10+1	4.54+2	1.11+0	1.20+1	4.93+2	1.17+0	
1.50+0	2.75+1	9.78-1	2.00+0	5.52+1	1.02+0	2.50+0	8.64+1	1.02+0	55
2.98+0	1.14+2	9.88-1	3.00+0	1.20+2	1.03+0	4.97+0	2.33+2	1.00+0	
5.96+0	2.82+2	1.01+0	6.94+0	3.21+2	1.01+0	6.95+0	3.23+2	1.02+0	
8.94+0	3.84+2	1.03+0	1.09+1	4.17+2	1.02+0				

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
9.00-2	1.22-4	9.00-1	1.00-1	2.27-4	7.17-1	1.10-1	4.19-4	6.54-1	57
1.20-1	7.75-4	6.63-1	1.30-1	1.47-3	7.47-1	1.40-1	2.73-3	8.85-1	
1.50-1	5.07-3	1.09+0	2.00-1	2.01-2	9.33-1	2.50-1	5.12-2	8.27-1	
2.95-1	9.36-2	7.34-1	3.30-1	1.43-1	7.02-1	3.60-1	1.92-1	6.64-1	
3.90-1	2.55-1	6.44-1	4.15-1	2.68-1	5.33-1				
5.00-1	1.50+0	1.50+0	6.00-1	2.60+0	1.37+0	7.00-1	4.30+0	1.35+0	69
8.00-1	6.20+0	1.27+0	9.00-1	8.30+0	1.19+0	1.00+0	1.16+1	1.22+0	
1.10+0	1.53+1	1.22+0	1.20+0	1.82+1	1.15+0	1.30+0	2.40+1	1.22+0	
1.40+0	2.70+1	1.14+0	1.50+0	3.50+1	1.24+0	1.60+0	4.10+1	1.25+0	
1.70+0	4.70+1	1.24+0	1.80+0	5.30+1	1.23+0	1.90+0	5.80+1	1.19+0	
2.00+0	6.80+1	1.25+0	2.10+0	7.70+1	1.28+0				
2.00-1	2.10-2	9.75-1	2.50-1	5.40-2	8.73-1	3.00-1	1.08-1	7.89-1	73
3.50-1	2.20-1	8.52-1	4.00-1	3.90-1	8.93-1	4.50-1	5.20-1	7.63-1	
5.50-1	1.20+0	8.53-1	6.50-1	2.14+0	8.59-1	7.50-1	3.16+0	7.96-1	
8.50-1	4.20+0	7.15-1							
9.50-1	5.30+0	6.46-1							76
1.00+0	1.00+1	1.05+0	1.10+0	1.28+1	1.02+0	1.20+0	1.60+1	1.01+0	77
1.30+0	1.97+1	1.00+0	1.40+0	2.38+1	1.01+0	1.50+0	2.84+1	1.01+0	
1.60+0	3.35+1	1.02+0	1.70+0	3.90+1	1.03+0	1.80+0	4.47+1	1.03+0	
1.90+0	5.09+1	1.05+0	2.00+0	5.71+1	1.05+0	2.10+0	6.33+1	1.05+0	
2.20+0	7.02+1	1.06+0	2.30+0	7.61+1	1.05+0	2.40+0	8.23+1	1.05+0	
2.50+0	8.82+1	1.04+0	2.60+0	9.36+1	1.03+0	2.70+0	9.81+1	1.01+0	
2.80+0	1.03+2	9.95-1	2.90+0	1.06+2	9.63-1	3.00+0	1.09+2	9.34-1	
1.00+0	7.38+0	7.74-1	2.00+0	4.06+1	7.47-1	5.00+0	1.98+2	8.46-1	84
1.25+0	9.50+0	5.37-1							89
3.00+0	1.19+2	1.02+0							94
2.50+0	7.55+1	8.91-1	3.00+0	1.02+2	8.74-1	5.00+0	2.12+2	9.05-1	97
7.50+0	3.07+2	9.15-1	9.00+0	3.43+2	9.16-1	9.75+0	3.48+2	8.94-1	
8.00-2	2.67-5	5.61-1	9.00-2	7.59-5	5.60-1	1.00-1	1.93-4	6.10-1	108
1.10-1	4.48-4	6.99-1	1.20-1	9.63-4	8.24-1	1.30-1	1.95-3	9.90-1	
1.40-1	3.72-3	1.21+0	1.50-1	6.81-3	1.47+0				
5.00-1	9.85-1	9.82-1	6.00-1	1.81+0	9.51-1	8.00-1	4.35+0	8.93-1	113
1.00+0	9.03+0	9.48-1	1.20+0	1.51+1	9.52-1	1.40+0	2.18+1	9.22-1	
1.60+0	3.05+1	9.28-1	1.80+0	4.05+1	9.37-1	2.00+0	5.01+1	9.22-1	
2.20+0	6.17+1	9.34-1	2.40+0	7.33+1	9.35-1				
7.00+0	3.10+2	9.71-1							114
2.00-1	2.13-2	9.89-1	2.50-1	6.09-2	9.84-1	3.00-1	1.30-1	9.49-1	115
3.50-1	2.50-1	9.68-1	4.00-1	3.82-1	8.75-1	4.50-1	6.29-1	9.22-1	
5.00-1	8.90-1	8.87-1	6.00-1	1.69+0	8.88-1	7.00-1	2.85+0	8.96-1	
8.00-1	4.30+0	8.83-1	9.00-1	6.17+0	8.84-1	1.00+0	8.40+0	8.81-1	
1.10+0	1.11+1	8.89-1	1.20+0	1.42+1	8.95-1	1.30+0	1.78+1	9.07-1	

TABLE 2. *K*-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	Ref.
1.40+0	2.10+1	8.88-1	1.50+0	2.48+1	8.82-1	1.60+0	2.98+1	9.07-1	
1.70+0	3.37+1	8.90-1	1.80+0	3.81+1	8.82-1	1.90+0	4.43+1	9.11-1	
2.00+0	4.97+1	9.15-1							
5.50-2	7.10-7	1.13+0	6.00-2	2.30-6	1.13+0	6.50-2	6.70-6	1.24+0	120
7.00-2	1.40-5	1.13+0	7.50-2	2.40-5	9.43-1	8.00-2	4.40-5	9.25-1	
9.00-2	1.40-4	1.03+0	1.00-1	3.40-4	1.07+0	1.10-1	6.80-4	1.06+0	
1.20-1	1.20-3	1.03+0	1.30-1	1.90-3	9.65-1	1.40-1	3.20-3	1.04+0	
1.50-1	4.30-3	9.28-1	1.60-1	5.70-3	8.53-1	1.80-1	1.10-2	8.73-1	
2.00-1	1.70-2	7.89-1	2.20-1	2.90-2	8.47-1	2.40-1	4.30-2	8.36-1	
2.60-1	6.10-2	8.27-1	2.80-1	8.80-2	8.63-1	3.00-1	1.30-1	9.49-1	
1.40+0	2.54+1	1.07+0	1.50+0	3.00+1	1.07+0	1.60+0	3.30+1	1.00+0	122
1.70+0	3.99+1	1.05+0	1.80+0	4.33+1	1.00+0	1.90+0	4.88+1	1.00+0	
2.00+0	5.40+1	9.94-1							
3.00-1	1.44-1	1.05+0	4.00-1	4.43-1	1.01+0	5.00-1	1.01+0	1.01+0	132
6.00-1	1.97+0	1.04+0	7.00-1	3.28+0	1.03+0	8.00-1	5.13+0	1.05+0	
1.00+0	9.69+0	1.02+0	1.20+0	1.64+1	1.03+0	1.40+0	2.41+1	1.02+0	
1.60+0	3.35+1	1.02+0	1.80+0	4.39+1	1.02+0	2.00+0	5.55+1	1.02+0	
2.20+0	6.75+1	1.02+0	2.30+0	7.37+1	1.02+0	2.40+0	7.95+1	1.01+0	
1.00+0	9.00+0	9.44-1	1.20+0	1.49+1	9.39-1	1.40+0	2.27+1	9.60-1	137
1.60+0	3.13+1	9.52-1	1.80+0	4.22+1	9.76-1	2.00+0	5.22+1	9.61-1	
2.20+0	6.29+1	9.52-1							
1.60+0	3.53+1	1.07+0	1.80+0	4.21+1	9.74-1	2.00+0	5.48+1	1.01+0	151
2.20+0	7.23+1	1.09+0	2.40+0	8.09+1	1.03+0				
<b>29 Copper                  Fluorescence yield = 0.44</b>									
2.00-1	4.50-3	<b>3.13-1</b>	2.50-1	1.00-2	<b>2.35-1</b>	3.00-1	2.10-2	<b>2.19-1</b>	3
3.50-1	4.00-2	<b>2.18-1</b>	4.00-1	8.50-2	<b>2.71-1</b>	4.60-1	2.00-1	<b>3.73-1</b>	
7.00-1	2.00+0	8.46-1	1.00+0	8.60+0	1.19+0	1.22+0	1.60+1	1.27+0	
4.00-1	1.59-1	5.07-1	5.00-1	3.28-1	<b>4.49-1</b>	6.00-1	6.84-1	4.89-1	4
7.00-1	1.07+0	<b>4.53-1</b>	8.00-1	1.71+0	<b>4.68-1</b>	9.00-1	3.01+0	5.71-1	
1.00+0	4.36+0	6.02-1							
1.40-1	1.90-3	9.87-1	1.50-1	2.80-3	9.59-1	1.60-1	4.50-3	1.05+0	6
1.80-1	8.20-3	9.94-1	2.00-1	1.40-2	9.75-1	2.20-1	2.20-2	9.51-1	
2.40-1	3.20-2	9.11-1	2.60-1	4.60-2	9.04-1	2.80-1	6.60-2	9.30-1	
3.00-1	9.00-2	9.39-1	3.20-1	1.20-1	9.51-1	3.40-1	1.60-1	9.83-1	
3.60-1	2.00-1	9.73-1	3.80-1	3.00-1	1.17+0	4.00-1	3.10-1	9.89-1	
4.40-1	3.80-1	8.40-1	6.00-1	1.20+0	8.57-1	7.40-1	2.20+0	7.75-1	
9.35-1	4.50+0	7.59-1	1.04+0	6.30+0	7.75-1	1.14+0	7.80+0	7.46-1	
1.20+0	9.40+0	7.69-1							
2.30-1	2.10-2	7.32-1	3.25-1	7.10-2	5.27-1	4.20-1	1.90-1	5.02-1	7
4.54-1	2.88-1	5.65-1	5.10-1	4.70-1	5.99-1				
1.50-1	3.00-3	1.03+0	2.00-1	1.30-2	9.05-1	3.00-1	8.80-2	9.18-1	9
4.00-1	3.10-1	9.89-1	5.00-1	6.30-1	8.63-1				

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
1.50+0	2.00+1	9.10-1							10
6.40+0	2.75+2	1.06+0	6.70+0	2.85+2	1.05+0	7.00+0	3.00+2	1.07+0	21
7.50+0	3.10+2	1.04+0	7.80+0	3.25+2	1.06+0	8.20+0	3.35+2	1.06+0	
8.40+0	3.40+2	1.06+0	8.50+0	3.50+2	1.08+0	8.80+0	3.60+2	1.09+0	
9.00+0	3.65+2	1.09+0	9.20+0	3.70+2	1.09+0	9.40+0	3.75+2	1.09+0	
9.60+0	3.75+2	1.08+0	9.90+0	3.70+2	1.05+0				
7.00-1	1.77+0	7.49-1	9.00-1	4.09+0	7.76-1	1.10+0	7.11+0	7.44-1	27
1.30+0	1.09+1	7.18-1	1.50+0	1.47+1	6.69-1	1.70+0	1.80+1	6.01-1	
1.90+0	2.29+1	5.91-1	2.10+0	3.09+1	6.40-1	2.30+0	3.64+1	6.23-1	
2.50+0	4.79+1	6.94-1							
1.60+2	2.20+2	1.50+0							30
1.25-1	9.50-4	1.04+0	1.50-1	3.10-3	1.06+0				36
1.00+0	8.80+0	1.21+0	2.00+0	5.10+1	1.17+0				39
1.00+0	5.40+0	7.45-1	2.25+0	4.20+1	7.52-1	3.00+0	7.50+1	7.81-1	47
2.50+0	6.64+1	9.63-1	3.00+0	9.81+1	1.02+0	4.00+0	1.56+2	1.03+0	52
4.50+0	1.80+2	1.02+0	5.00+0	1.86+2	9.24-1	6.00+0	2.53+2	1.03+0	
6.50+0	2.46+2	9.31-1	7.00+0	2.99+2	1.06+0	8.00+0	2.90+2	9.31-1	
8.50+0	3.46+2	1.07+0	9.00+0	3.25+2	9.67-1	1.00+1	3.79+2	1.07+0	
1.05+1	3.62+2	9.95-1	1.10+1	4.05+2	1.09+0	1.20+1	4.25+2	1.11+0	
5.00-1	5.00-1	6.85-1	6.00-1	1.00+0	7.14-1	7.00-1	1.80+0	7.61-1	53
8.00-1	2.80+0	7.67-1	9.00-1	4.10+0	7.77-1	1.00+0	6.00+0	8.28-1	
1.10+0	8.00+0	8.37-1	1.20+0	1.07+1	8.76-1	1.30+0	1.34+1	8.82-1	
1.40+0	1.63+1	8.82-1	1.50+0	1.98+1	9.01-1	1.60+0	2.34+1	9.06-1	
1.70+0	2.68+1	8.95-1	1.80+0	3.22+1	9.42-1	1.90+0	3.59+1	9.26-1	
2.00+0	3.92+1	9.03-1							
1.50+0	2.14+1	9.74-1	2.00+0	4.27+1	9.83-1	2.50+0	6.87+1	9.96-1	55
3.00+0	9.51+1	9.90-1	4.96+0	2.06+2	1.03+0	5.96+0	2.52+2	1.04+0	
6.96+0	2.97+2	1.06+0	8.94+0	3.55+2	1.06+0	1.09+1	3.98+2	1.07+0	
1.30-1	1.14-3	9.56-1	1.50-1	2.35-3	8.05-1	2.00-1	1.28-2	8.91-1	57
2.50-1	4.54-2	1.07+0	2.95-1	7.21-2	8.08-1	3.30-1	9.81-2	6.83-1	
3.60-1	1.26-1	6.13-1	3.85-1	1.46-1	5.42-1	4.15-1	1.72-1	4.76-1	
3.00+0	8.70+1	9.06-1	5.00+0	1.80+2	8.94-1				59
6.67-1	2.00+0	9.94-1	1.00+0	7.30+0	1.01+0	1.33+0	1.54+1	9.48-1	64
1.67+0	2.79+1	9.75-1	2.00+0	4.26+1	9.81-1	2.33+0	5.45+1	9.07-1	
2.67+0	7.46+1	9.56-1	3.00+0	9.05+1	9.42-1	3.33+0	1.16+2	1.01+0	
3.67+0	1.34+2	1.00+0	4.00+0	1.44+2	9.55-1	4.33+0	1.85+2	1.10+0	
4.67+0	1.99+2	1.07+0	5.00+0	2.13+2	1.06+0	5.33+0	2.31+2	1.06+0	
5.67+0	2.47+2	1.07+0							

TABLE 2. *K*-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
8.30-1	4.30+0	1.05+0	1.58+0	2.90+1	1.16+0	2.56+0	6.80+1	9.42-1	66
	3.28+0	1.23+2	1.10+0						
5.00-1	9.00-1	1.23+0	6.00-1	1.80+0	1.29+0	7.00-1	2.90+0	1.23+0	69
8.00-1	4.70+0	1.29+0	9.00-1	6.40+0	1.21+0	1.00+0	8.70+0	1.20+0	
1.10+0	1.16+1	1.21+0	1.20+0	1.44+1	1.18+0	1.30+0	1.82+1	1.20+0	
1.40+0	2.20+1	1.19+0	1.50+0	2.60+1	1.18+0	1.60+0	3.20+1	1.24+0	
1.70+0	3.60+1	1.20+0	1.80+0	4.10+1	1.20+0	1.90+0	4.80+1	1.24+0	
2.00+0	5.30+1	1.22+0	2.10+0	5.70+1	1.18+0	2.20+0	6.00+1	1.12+0	
2.30+0	6.60+1	1.13+0							
2.00+0	2.30+1	5.30-1							74
9.50-1	4.20+0	6.76-1							76
1.00+0	7.17+0	9.89-1	1.20+0	1.21+1	9.90-1	1.40+0	1.88+1	1.02+0	77
1.50+0	2.27+1	1.03+0	1.60+0	2.69+1	1.04+0	1.80+0	3.62+1	1.06+0	
2.00+0	4.63+1	1.07+0	2.20+0	5.61+1	1.05+0	2.40+0	6.50+1	1.02+0	
2.50+0	6.94+1	1.01+0	2.60+0	7.30+1	9.82-1	2.80+0	7.97+1	9.36-1	
3.00+0	8.41+1	8.75-1							
9.00-1	5.05+0	9.58-1	1.00+0	7.89+0	1.09+0	1.10+0	1.27+1	1.33+0	80
1.20+0	1.75+1	1.43+0	1.30+0	1.85+1	1.22+0	1.40+0	2.08+1	1.13+0	
1.50+0	2.24+1	1.02+0	1.60+0	2.80+1	1.08+0	1.70+0	3.30+1	1.10+0	
1.80+0	3.54+1	1.04+0	1.90+0	4.44+1	1.15+0	2.00+0	4.63+1	1.07+0	
2.10+0	4.62+1	9.57-1	2.20+0	5.38+1	1.01+0	2.30+0	6.10+1	1.04+0	
2.40+0	7.42+1	1.17+0	2.50+0	6.95+1	1.01+0	2.60+0	6.04+1	8.13-1	
2.70+0	8.95+1	1.12+0	2.80+0	8.90+1	1.04+0	2.90+0	7.49+1	8.27-1	
3.00+0	8.15+1	8.48-1	3.10+0	1.17+2	1.15+0	3.20+0	1.24+2	1.15+0	
3.30+0	1.25+2	1.11+0	3.40+0	1.15+2	9.71-1	3.50+0	1.16+2	9.36-1	
3.60+0	1.29+2	9.97-1	3.70+0	1.22+2	9.03-1	3.80+0	2.22+2	1.58+0	
3.90+0	2.14+2	1.47+0	4.00+0	2.20+2	1.46+0				
3.00+0	9.72+1	1.01+0	5.00+0	2.12+2	1.05+0	7.00+0	2.98+2	1.06+0	94
9.00+0	3.57+2	1.06+0	1.10+1	3.92+2	1.06+0				
2.49-1	4.68-2	1.12+0	3.00-1	1.08-1	1.13+0	3.53-1	2.11-1	1.11+0	99
4.03-1	3.51-1	1.09+0	4.84-1	6.77-1	1.05+0	5.52-1	1.06+0	1.02+0	
6.35-1	1.66+0	9.75-1	7.20-1	2.44+0	9.40-1	8.10-1	3.47+0	9.14-1	
9.00-1	4.70+0	8.91-1	1.10+0	8.18+0	8.56-1	1.31+0	1.30+1	8.38-1	
1.52+0	1.91+1	8.41-1	1.72+0	2.61+1	8.48-1	1.91+0	3.40+1	8.67-1	
1.50-1	1.46-3	5.00-1	2.00-1	9.46-3	6.59-1	3.00-1	7.58-2	7.90-1	103
4.00-1	2.77-1	8.84-1	5.00-1	6.60-1	9.04-1	6.00-1	1.26+0	9.00-1	
7.00-1	2.20+0	9.31-1	8.00-1	3.48+0	9.53-1				
2.20+1	4.47+2	1.09+0	3.10+1	3.91+2	1.02+0	4.40+1	3.28+2	9.84-1	104
4.00+0	1.56+2	1.03+0	6.00+0	2.80+2	1.14+0	8.00+0	3.69+2	1.18+0	105
1.00+1	4.09+2	1.15+0	1.20+1	4.49+2	1.17+0	1.40+1	4.81+2	1.20+0	
1.60+1	4.72+2	1.15+0	1.80+1	4.94+2	1.20+0	2.00+1	4.85+2	1.18+0	
2.20+1	4.76+2	1.16+0							

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
1.00+0	7.25+0	1.00+0	1.25+0	1.38+1	1.01+0	1.50+0	2.40+1	1.09+0	112
1.75+0	3.41+1	1.06+0	2.00+0	4.49+1	1.03+0	2.25+0	5.83+1	1.04+0	
2.50+0	7.03+1	1.02+0	2.75+0	8.68+1	1.05+0	3.00+0	1.03+2	1.07+0	
5.00-1	7.65-1	1.05+0	6.00-1	1.46+0	1.04+0	8.00-1	3.80+0	1.04+0	113
1.00+0	7.12+0	9.83-1	1.20+0	1.16+1	9.49-1	1.40+0	1.79+1	9.68-1	
1.60+0	2.51+1	9.72-1	1.80+0	3.31+1	9.68-1	2.00+0	4.28+1	9.85-1	
2.20+0	5.03+1	9.43-1	2.40+0	6.19+1	9.73-1				
7.00+0	3.10+2	1.10+0							114
2.00-1	1.65-2	1.15+0	2.50-1	4.45-2	1.05+0	3.00-1	9.52-2	9.93-1	115
3.50-1	1.78-1	9.70-1	4.00-1	3.03-1	9.67-1	4.50-1	4.81-1	9.76-1	
5.00-1	6.81-1	9.33-1	6.00-1	1.38+0	9.86-1	7.00-1	2.30+0	9.73-1	
8.00-1	3.59+0	9.84-1	9.00-1	5.12+0	9.71-1	1.00+0	7.48+0	1.03+0	
1.10+0	9.79+0	1.02+0	1.20+0	1.21+1	9.90-1	1.30+0	1.48+1	9.74-1	
1.40+0	1.78+1	9.63-1	1.50+0	2.15+1	9.78-1	1.60+0	2.51+1	9.72-1	
1.70+0	2.92+1	9.75-1	1.80+0	3.35+1	9.80-1	1.90+0	3.78+1	9.75-1	
2.00+0	4.24+1	9.76-1							
5.00-1	6.99-1	9.58-1	6.25-1	1.35+0	8.38-1	7.50-1	2.47+0	8.33-1	117
8.75-1	3.95+0	8.17-1	1.00+0	6.01+0	8.29-1	1.25+0	1.19+1	8.71-1	
1.50+0	1.93+1	8.78-1	1.75+0	2.92+1	9.12-1	2.00+0	3.95+1	9.09-1	
2.25+0	5.47+1	9.79-1	2.50+0	6.85+1	9.93-1	2.75+0	8.41+1	1.02+0	
3.00+0	9.70+1	1.01+0	2.99+0	9.22+1	9.68-1	3.94+0	1.45+2	9.81-1	
6.07+0	2.46+2	9.93-1	9.13+0	3.50+2	1.03+0	1.22+1	4.06+2	1.05+0	
1.83+1	4.23+2	1.02+0	2.40+1	4.12+2	1.02+0	3.01+1	3.96+2	1.03+0	
3.56+1	3.74+2	1.03+0	3.96+1	3.56+2	1.02+0				
1.25-1	3.34-4	3.65-1	1.50-1	1.33-3	<b>4.56-1</b>	1.75-1	4.02-3	5.68-1	118
2.00-1	8.89-3	6.19-1	2.50-1	2.98-2	7.01-1	3.00-1	7.30-2	7.61-1	
4.00-1	2.68-1	8.55-1	5.00-1	6.42-1	8.80-1	6.00-1	1.24+0	8.86-1	
7.00-1	2.09+0	8.84-1	8.00-1	3.24+0	8.88-1				
1.40+0	1.98+1	1.07+0	1.50+0	2.16+1	9.83-1	1.60+0	2.51+1	9.72-1	122
1.70+0	2.94+1	9.82-1	1.80+0	3.40+1	9.94-1	1.90+0	3.58+1	9.24-1	
2.00+0	4.15+1	9.55-1							
7.00+0	3.29+2	1.17+0							125
1.00-1	1.37-4	7.87-1	1.20-1	5.92-4	8.57-1	1.40-1	1.69-3	8.78-1	126
1.60-1	3.82-3	8.95-1	1.80-1	7.60-3	9.22-1	2.00-1	1.33-2	9.26-1	
2.50-1	3.91-2	9.20-1	3.00-1	8.48-2	8.84-1				
5.00-1	7.65-1	1.05+0	7.07-1	2.54+0	1.04+0	1.00+0	7.16+0	9.88-1	130
1.41+0	1.84+1	9.71-1	2.00+0	4.23+1	9.74-1	2.50+0	6.55+1	9.50-1	
3.00-1	9.40-2	9.80-1	4.00-1	3.16-1	1.01+0	5.00-1	7.57-1	1.04+0	132
6.00-1	1.44+0	1.03+0	7.00-1	2.30+0	9.73-1	8.00-1	3.67+0	1.01+0	
1.00+0	7.25+0	1.00+0	1.20+0	1.22+1	9.99-1	1.40+0	1.86+1	1.01+0	
1.60+0	2.64+1	1.02+0	1.80+0	3.50+1	1.02+0	2.00+0	4.26+1	9.81-1	
2.20+0	5.34+1	1.00+0	2.30+0	5.74+1	9.83-1	2.40+0	6.32+1	9.93-1	

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_i$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_i$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_i$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
2.80-1	9.20-2	1.30+0	3.30-1	1.70-1	1.18+0	3.70-1	2.50-1	1.09+0	135
4.20-1	3.70-1	9.77-1	5.00-1	6.60-1	9.04-1	5.60-1	8.80-1	8.01-1	
6.10-1	1.10+0	7.42-1	6.50-1	1.30+0	7.05-1	6.90-1	1.60+0	7.10-1	
7.40-1	1.90+0	6.69-1	7.90-1	2.40+0	6.84-1	8.30-1	2.80+0	6.83-1	
8.70-1	3.30+0	6.94-1	9.10-1	4.40+0	8.06-1	1.01+0	5.70+0	7.64-1	
1.11+0	7.90+0	8.05-1	1.21+0	9.70+0	7.76-1	1.31+0	1.30+1	8.38-1	
1.41+0	1.60+1	8.50-1	1.51+0	1.90+1	8.50-1	1.61+0	2.20+1	8.39-1	
1.71+0	2.40+1	7.90-1	1.81+0	3.00+1	8.66-1				
5.00-1	6.76-1	9.26-1	6.00-1	1.32+0	9.43-1	7.00-1	2.31+0	9.77-1	137
8.00-1	3.54+0	9.70-1	1.00+0	6.90+0	9.52-1	1.20+0	1.15+1	9.41-1	
1.40+0	1.74+1	9.41-1	1.60+0	2.44+1	9.45-1	1.80+0	3.25+1	9.50-1	
2.00+0	4.15+1	9.55-1	2.20+0	5.07+1	9.50-1				
1.50+0	2.06+1	9.37-1	2.00+0	4.61+1	1.06+0	2.25+0	5.57+1	9.97-1	143
2.50+0	6.88+1	9.98-1	2.75+0	7.99+1	9.68-1	3.00+0	9.68+1	1.01+0	
1.00+0	6.96+0	9.60-1	2.00+0	4.26+1	9.81-1				149
1.60+0	2.61+1	1.01+0	1.80+0	3.46+1	1.01+0	2.00+0	4.55+1	1.05+0	151
2.20+0	5.58+1	1.05+0	2.40+0	6.57+1	1.03+0				
<b>30 Zinc</b> Fluorescence yield = 0.47									
7.00-1	1.49+0	8.49-1	9.00-1	3.54+0	8.91-1	1.10+0	5.33+0	7.31-1	27
1.30+0	8.31+0	7.09-1	1.50+0	1.41+1	8.20-1	1.70+0	1.73+1	7.36-1	
1.90+0	2.54+1	8.27-1	2.10+0	2.75+1	7.12-1	2.30+0	3.01+1	6.40-1	
2.50+0	3.45+1	6.18-1							
2.50+0	5.34+1	9.56-1	3.00+0	7.80+1	9.86-1	4.00+0	1.34+2	1.05+0	52
4.50+0	1.52+2	1.01+0	5.00+0	1.71+2	9.93-1	6.00+0	2.12+2	1.00+0	
6.50+0	2.17+2	9.43-1	7.00+0	2.51+2	1.02+0	8.00+0	2.81+2	1.02+0	
8.50+0	3.23+2	1.12+0	9.00+0	3.03+2	1.01+0	1.00+1	3.37+2	1.06+0	
1.05+1	3.39+2	1.03+0	1.10+1	3.58+2	1.07+0	1.20+1	3.82+2	1.10+0	
4.50-1	3.00-1	8.44-1	5.60-1	5.00-1	6.22-1	6.60-1	1.60+0	1.12+0	53
7.70-1	2.60+0	1.08+0	8.70-1	3.60+0	1.01+0	9.70-1	5.30+0	1.06+0	
1.08+0	7.00+0	1.01+0	1.20+0	9.10+0	9.71-1	1.30+0	1.14+1	9.72-1	
1.40+0	1.40+1	9.77-1	1.50+0	1.66+1	9.66-1	1.60+0	1.94+1	9.60-1	
1.70+0	2.22+1	9.44-1	1.80+0	2.52+1	9.30-1	1.90+0	2.87+1	9.35-1	
2.00+0	3.19+1	9.23-1							
7.00-1	2.63+0	1.50+0	1.00+0	6.42+0	1.17+0	1.40+0	1.57+1	1.10+0	56
1.60+0	2.00+1	9.89-1	1.80+0	2.67+1	9.85-1	2.00+0	3.31+1	9.58-1	
2.20+0	4.25+1	9.95-1	2.40+0	5.32+1	1.03+0	2.60+0	6.47+1	1.07+0	
2.80+0	7.62+1	1.09+0	3.00+0	8.72+1	1.10+0	3.20+0	1.01+2	1.14+0	
3.40+0	1.11+2	1.13+0	3.60+0	1.23+2	1.14+0	3.80+0	1.36+2	1.16+0	
4.00+0	1.54+2	1.21+0	4.20+0	1.66+2	1.22+0	4.40+0	1.78+2	1.22+0	
1.50-1	1.72-3	9.39-1	1.95-1	7.35-3	8.86-1	2.55-1	2.90-2	9.09-1	57
3.00-1	5.39-2	8.05-1	3.35-1	7.63-2	7.08-1	3.60-1	9.87-2	6.77-1	
3.90-1	1.15-1	5.69-1	4.15-1	1.53-1	5.90-1				

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
2.00-1	1.40-2	1.47+0	2.50-1	3.80-2	1.31+0	3.00-1	7.80-2	1.17+0	73
3.50-1	1.50-1	1.16+0	4.00-1	2.80-1	1.25+0	4.50-1	3.90-1	1.10+0	
1.00+0	5.60+0	1.02+0	1.10+0	7.50+0	1.03+0	1.20+0	9.70+0	1.03+0	86
1.40+0	1.50+1	1.05+0	1.60+0	2.16+1	1.07+0	1.80+0	2.92+1	1.08+0	
2.00+0	3.78+1	1.09+0	2.20+0	4.68+1	1.10+0	2.40+0	5.56+1	1.08+0	
2.60+0	6.42+1	1.06+0	2.80+0	7.14+1	1.02+0	3.00+0	7.76+1	9.80-1	
1.00-1	6.08-5	6.51-1	1.10-1	1.34-4	6.51-1	1.20-1	2.74-4	6.84-1	108
1.30-1	5.30-4	7.45-1	1.40-1	9.73-4	8.28-1	1.50-1	1.72-3	9.39-1	
7.00+0	2.30+2	9.33-1							114
2.00-1	1.30-2	1.37+0	2.50-1	3.41-2	1.17+0	3.00-1	7.71-2	1.15+0	115
3.50-1	1.41-1	1.09+0	4.00-1	2.34-1	1.05+0	4.50-1	3.83-1	1.08+0	
5.00-1	5.70-1	1.07+0	6.00-1	9.63-1	9.36-1	7.00-1	1.61+0	9.18-1	
8.00-1	2.49+0	9.12-1	9.00-1	3.61+0	9.08-1	1.00+0	5.13+0	9.33-1	
1.10+0	7.38+0	1.01+0	1.20+0	9.25+0	9.87-1	1.30+0	1.20+1	1.02+0	
1.40+0	1.38+1	9.63-1	1.50+0	1.60+1	9.31-1	1.60+0	1.84+1	9.10-1	
1.70+0	2.21+1	9.40-1	1.80+0	2.61+1	9.63-1	1.90+0	2.95+1	9.61-1	
2.00+0	3.33+1	9.63-1							
7.00+0	2.06+2	8.35-1							125
3.00-1	6.37-2	9.52-1	4.00-1	2.23-1	9.97-1	5.00-1	5.22-1	9.84-1	132
6.00-1	1.01+0	9.82-1	7.00-1	1.69+0	9.63-1	8.00-1	2.68+0	9.82-1	
1.00+0	5.46+0	9.93-1	1.20+0	9.24+0	9.86-1	1.40+0	1.43+1	9.98-1	
1.60+0	2.03+1	1.00+0	1.80+0	2.78+1	1.03+0	2.00+0	3.47+1	1.00+0	
2.20+0	4.28+1	1.00+0	2.30+0	4.93+1	1.05+0	2.40+0	5.17+1	1.01+0	
5.00-1	4.84-1	9.13-1	6.00-1	9.48-1	9.21-1	7.00-1	1.62+0	9.23-1	137
8.00-1	2.52+0	9.23-1	1.00+0	5.27+0	9.59-1	1.20+0	8.77+0	9.35-1	
1.40+0	1.35+1	9.42-1	1.60+0	1.96+1	9.70-1	1.80+0	2.55+1	9.41-1	
2.00+0	3.27+1	9.46-1	2.20+0	4.06+1	9.50-1				
1.00+0	5.31+0	9.66-1	1.10+0	7.31+0	1.00+0	1.20+0	8.34+0	8.90-1	152
1.30+0	1.02+1	8.70-1	1.40+0	1.25+1	8.72-1	1.50+0	1.53+1	8.90-1	
1.60+0	1.78+1	8.81-1	1.70+0	1.97+1	8.38-1	1.80+0	2.35+1	8.67-1	
1.90+0	2.66+1	8.66-1	2.00+0	2.77+1	8.01-1	2.10+0	3.24+1	8.38-1	
2.20+0	3.64+1	8.52-1	2.30+0	4.01+1	8.53-1	2.40+0	4.13+1	8.03-1	
2.50+0	4.31+1	7.72-1	2.60+0	4.68+1	7.75-1	2.70+0	4.95+1	7.61-1	
2.80+0	5.44+1	7.80-1	2.90+0	5.98+1	8.03-1	3.00+0	6.20+1	7.83-1	
<b>31</b>	<b>Gallium</b>				<b>Fluorescence yield = 0.507</b>				
5.00-1	3.00-1	7.79-1	6.00-1	6.00-1	7.93-1	7.00-1	1.10+0	8.46-1	53
8.00-1	1.80+0	8.81-1	9.00-1	2.70+0	9.01-1	1.00+0	4.10+0	9.84-1	
1.10+0	5.30+0	9.53-1	1.20+0	7.60+0	1.06+0	1.30+0	8.90+0	9.85-1	
1.40+0	1.11+1	1.00+0	1.50+0	1.36+1	1.02+0	1.60+0	1.61+1	1.02+0	
1.70+0	1.88+1	1.02+0	1.80+0	2.19+1	1.03+0	1.90+0	2.48+1	1.02+0	
2.00+0	2.83+1	1.03+0							

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
1.00+0	4.00+0	9.60-1	1.20+0	6.40+0	8.90-1	1.40+0	9.20+0	8.29-1	86
1.60+0	1.28+1	8.08-1	1.80+0	1.74+1	8.18-1	2.00+0	2.29+1	8.34-1	
2.20+0	2.89+1	8.45-1	2.40+0	3.47+1	8.40-1	2.60+0	3.94+1	8.05-1	
2.80+0	4.21+1	7.41-1	3.00+0	4.21+1	6.50-1				
1.40+0	1.03+1	9.28-1	1.50+0	1.22+1	9.12-1	1.60+0	1.50+1	9.46-1	122
1.70+0	1.74+1	9.43-1	1.80+0	2.01+1	9.45-1	1.90+0	2.25+1	9.25-1	
2.00+0	2.48+1	9.03-1							
<b>32 Germanium</b> <b>Fluorescence yield = 0.535</b>									
1.00+0	2.80+0	8.92-1	1.50+0	1.00+1	9.68-1	2.00+0	2.10+1	9.72-1	38
2.50+0	3.50+1	9.71-1	3.00+0	5.40+1	1.03+0	3.50+0	7.00+1	1.00+0	
4.00+0	9.70+1	1.10+0	4.50+0	1.05+2	9.92-1	5.00+0	1.20+2	9.74-1	
5.50+0	1.50+2	1.07+0	6.00+0	1.60+2	1.03+0				
1.00+0	2.30+0	7.33-1	2.25+0	2.20+1	7.72-1	3.00+0	4.00+1	7.61-1	47
5.00-1	3.00-1	1.08+0	6.00-1	5.00-1	9.04-1	7.00-1	9.00-1	9.36-1	53
8.00-1	1.40+0	9.21-1	9.00-1	2.00+0	8.92-1	1.00+0	2.80+0	8.92-1	
1.10+0	3.70+0	8.78-1	1.20+0	4.80+0	8.77-1	1.30+0	6.00+0	8.68-1	
1.40+0	7.20+0	8.43-1	1.50+0	8.70+0	8.42-1	1.60+0	1.10+1	8.95-1	
1.70+0	1.22+1	8.46-1	1.80+0	1.41+1	8.47-1	1.90+0	1.60+1	8.41-1	
2.00+0	1.85+1	8.56-1							
3.00+0	5.44+1	1.04+0							94
2.00-1	3.66-3	8.93-1	3.00-1	3.07-2	9.48-1	3.50-1	5.56-2	8.59-1	115
4.00-1	1.08-1	9.47-1	4.50-1	1.71-1	9.28-1	5.00-1	2.67-1	9.57-1	
6.00-1	5.45-1	9.85-1	7.00-1	9.07-1	9.43-1	8.00-1	1.43+0	9.41-1	
9.00-1	2.10+0	9.36-1	1.00+0	2.94+0	9.37-1	1.10+0	3.97+0	9.42-1	
1.20+0	5.17+0	9.45-1	1.30+0	6.37+0	9.21-1	1.40+0	8.10+0	9.49-1	
1.50+0	1.02+1	9.88-1	1.60+0	1.18+1	9.60-1	1.70+0	1.38+1	9.58-1	
1.80+0	1.59+1	9.55-1	1.90+0	1.83+1	9.61-1	2.00+0	2.06+1	9.53-1	
3.30-1	8.30-2	1.66+0	3.70-1	1.20-1	1.46+0	4.20-1	1.70-1	1.22+0	135
4.60-1	2.10-1	1.04+0	5.20-1	3.40-1	1.05+0	5.60-1	4.30-1	1.00+0	
6.00-1	5.30-1	9.58-1	6.50-1	6.40-1	8.66-1	7.10-1	9.10-1	9.01-1	
7.50-1	1.10+0	9.02-1	7.90-1	1.30+0	8.92-1	9.10-1	1.80+0	7.75-1	
1.01+0	2.30+0	7.10-1	1.11+0	3.40+0	7.85-1	1.21+0	4.60+0	8.20-1	
1.31+0	5.40+0	7.64-1	1.41+0	7.50+0	8.61-1	1.51+0	8.60+0	8.18-1	
1.61+0	1.10+1	8.80-1	1.71+0	1.20+1	8.20-1	1.81+0	1.40+1	8.29-1	
<b>33 Arsenic</b> <b>Fluorescence yield = 0.562</b>									
4.70-1	1.00-1	6.32-1	5.70-1	3.00-1	8.95-1	6.80-1	5.00-1	7.79-1	53
7.80-1	9.00-1	8.66-1	8.90-1	1.40+0	8.63-1	9.90-1	2.10+0	9.15-1	
1.10+0	2.80+0	8.74-1	1.20+0	3.80+0	9.09-1	1.30+0	4.80+0	9.05-1	
1.40+0	6.00+0	9.12-1	1.50+0	7.40+0	9.26-1	1.60+0	8.80+0	9.22-1	
1.70+0	1.04+1	9.26-1	1.80+0	1.22+1	9.35-1	1.90+0	1.41+1	9.42-1	
2.00+0	1.59+1	9.36-1							
<b>34 Selenium</b> <b>Fluorescence yield = 0.589</b>									

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
1.00+0	2.00+0	1.11+0	2.00+0	1.50+1	1.11+0				39
4.00-1	7.16-2	1.22+0	6.00-1	2.91-1	9.70-1	8.00-1	8.49-1	9.97-1	61
1.00+0	1.86+0	1.03+0	1.20+0	2.81+0	8.77-1	1.40+0	4.41+0	8.68-1	
1.60+0	6.54+0	8.80-1	1.80+0	8.57+0	8.37-1	2.00+0	1.15+1	8.54-1	
3.00+0	3.42+1	9.92-1							94
1.00+1	2.26+2	1.17+0	1.20+1	2.74+2	1.24+0	1.40+1	2.80+2	1.16+0	
1.60+1	2.98+2	1.17+0	1.80+1	3.22+2	1.22+0	2.00+1	3.28+2	1.21+0	
2.20+1	3.10+2	1.12+0							
6.00-1	3.19-1	1.06+0	8.00-1	8.28-1	9.72-1	1.00+0	1.79+0	9.93-1	113
1.20+0	3.18+0	9.92-1	1.40+0	4.89+0	9.62-1	1.60+0	7.21+0	9.70-1	
1.80+0	1.04+1	1.02+0	2.00+0	1.34+1	9.95-1	2.20+0	1.62+1	9.52-1	
2.40+0	1.90+1	9.04-1							
7.00+0	1.50+2	1.10+0							114
2.00-1	2.08-3	1.19+0	2.50-1	6.20-3	9.97-1	3.00-1	1.44-2	9.16-1	115
3.50-1	3.59-2	1.11+0	4.00-1	5.86-2	1.00+0	4.50-1	9.12-2	9.48-1	
5.00-1	1.52-1	1.03+0	6.00-1	3.06-1	1.02+0	7.00-1	5.46-1	1.03+0	
8.00-1	8.70-1	1.02+0	9.00-1	1.28+0	1.01+0	1.00+0	1.79+0	9.93-1	
1.10+0	2.39+0	9.78-1	1.20+0	3.15+0	9.83-1	1.30+0	4.02+0	9.84-1	
1.40+0	5.05+0	9.94-1	1.50+0	5.88+0	9.48-1	1.60+0	7.39+0	9.94-1	
1.70+0	8.46+0	9.63-1	1.80+0	9.83+0	9.60-1	1.90+0	1.13+1	9.57-1	
2.00+0	1.28+1	9.51-1							
7.00+0	1.54+2	1.13+0							125
<b>35</b>	Bromine	Fluorescence yield = 0.615							
6.00-1	3.34-1	1.50+0	8.00-1	7.93-1	1.24+0	1.00+0	1.56+0	1.14+0	61
1.20+0	2.71+0	1.10+0	1.40+0	4.47+0	1.13+0	1.60+0	6.03+0	1.04+0	
1.80+0	8.31+0	1.03+0	2.00+0	1.04+1	9.75-1				
1.50+0	4.00+0	8.29-1	2.00+0	9.95+0	9.33-1	2.25+0	1.29+1	9.00-1	143
2.50+0	1.74+1	9.37-1	2.75+0	2.18+1	9.45-1	3.00+0	2.95+1	1.05+0	
<b>36</b>	Krypton	Fluorescence yield = 0.643							
1.50+0	5.90+0	1.56+0	2.00+0	1.30+1	1.54+0	2.50+0	2.10+1	1.41+0	40
3.00+0	4.30+1	1.89+0	3.50+0	4.70+1	1.48+0	4.00+0	6.20+1	1.50+0	
4.50+0	7.70+1	1.50+0							
1.50+0	6.23+0	1.65+0	2.00+0	1.32+1	1.56+0	2.50+0	2.25+1	1.51+0	48
3.00+0	3.66+1	1.61+0	3.50+0	4.74+1	1.49+0	4.00+0	6.38+1	1.55+0	
4.50+0	7.92+1	1.55+0	5.00+0	1.02+2	1.67+0				
3.00+0	3.15+1	1.38+0							65
5.00-1	7.10-2	8.88-1	8.16-1	5.20-1	9.97-1	9.15-1	7.00-1	8.99-1	68
1.00+0	1.00+0	9.50-1	1.29+0	2.20+0	9.20-1	1.63+0	4.30+0	8.95-1	
2.00+0	6.80+0	8.03-1							

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
<b>37 Rubidium</b> <b>Fluorescence yield = 0.667</b>									
1.00+0	5.80-1	7.19-1	2.25+0	7.50+0	8.17-1	3.00+0	1.50+1	8.12-1	47
4.00-1	2.77-2	1.25+0	6.00-1	1.24-1	9.99-1	8.00-1	4.08-1	1.11+0	61
1.00+0	9.27-1	1.15+0	1.20+0	1.32+0	8.95-1	1.40+0	2.37+0	9.90-1	
1.60+0	3.62+0	1.01+0	1.80+0	4.47+0	8.91-1	2.00+0	5.43+0	8.08-1	
1.00+0	8.00-1	9.92-1	1.50+0	2.90+0	9.83-1	2.00+0	6.90+0	1.03+0	72
2.50+0	1.10+1	9.20-1	3.00+0	2.00+1	1.08+0				
2.20+1	2.21+2	1.05+0	3.10+1	2.08+2	9.49-1	4.40+1	2.09+2	9.90-1	104
3.50-1	5.75-3	4.86-1	4.00-1	1.15-2	5.18-1	4.50-1	2.32-2	6.16-1	111
5.00-1	3.51-2	5.94-1	6.00-1	8.00-2	6.44-1	7.00-1	1.70-1	7.55-1	
1.50+0	2.67+0	9.05-1	2.10+0	7.41+0	9.68-1	2.60+0	1.23+1	9.35-1	148
3.10+0	1.89+1	9.50-1	3.60+0	2.45+1	8.90-1				
<b>38 Strontium</b> <b>Fluorescence yield = 0.69</b>									
6.00-1	7.97-2	8.56-1	8.00-1	3.64-1	1.30+0	1.00+0	7.02-1	1.13+0	61
1.20+0	1.18+0	1.03+0	1.40+0	1.81+0	9.68-1	1.60+0	2.69+0	9.58-1	
1.80+0	3.94+0	9.94-1	2.00+0	5.28+0	9.90-1				
3.00+0	1.23+1	8.18-1							94
4.00-1	9.88-3	6.11-1	4.50-1	2.21-2	7.99-1	5.00-1	3.42-2	7.81-1	111
6.00-1	7.39-2	7.93-1	7.00-1	1.44-1	8.45-1	8.00-1	2.06-1	7.33-1	
9.00-1	3.32-1	7.73-1	1.00+0	5.19-1	8.36-1	1.10+0	7.26-1	8.47-1	
1.20+0	1.03+0	9.01-1	1.40+0	1.52+0	8.13-1	1.50+0	1.81+0	7.83-1	
<b>39 Yttrium</b> <b>Fluorescence yield = 0.71</b>									
2.50+0	7.44+0	9.62-1	3.00+0	1.25+1	1.03+0	4.00+0	2.54+1	1.10+0	52
4.50+0	3.20+1	1.09+0	5.00+0	3.56+1	9.92-1	6.00+0	5.22+1	1.06+0	
6.50+0	6.09+1	1.09+0	7.00+0	5.84+1	9.36-1	8.00+0	7.27+1	9.67-1	
8.50+0	9.60+1	1.18+0	9.00+0	9.50+1	1.09+0	1.00+1	1.10+2	1.12+0	
1.05+1	1.02+2	9.84-1	1.10+1	1.11+2	1.02+0	1.20+1	1.03+2	8.73-1	
3.00+0	9.30+0	7.66-1	5.00+0	2.80+1	7.81-1				59
6.00-1	7.20-2	1.03+0	8.00-1	2.20-1	1.03+0	1.00+0	4.80-1	1.01+0	61
1.20+0	8.81-1	9.93-1	1.40+0	1.43+0	9.80-1	1.60+0	2.11+0	9.56-1	
1.80+0	3.09+0	9.87-1	2.00+0	3.94+0	9.30-1				
1.00+0	4.10-1	8.59-1	1.20+0	7.70-1	8.68-1	1.40+0	1.30+0	8.91-1	86
1.60+0	1.90+0	8.61-1	1.80+0	2.80+0	8.95-1	2.00+0	3.70+0	8.74-1	
2.20+0	4.70+0	8.53-1	2.40+0	5.80+0	8.34-1	2.60+0	7.00+0	8.18-1	
2.80+0	8.10+0	7.89-1	3.00+0	9.20+0	7.58-1				

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
5.00-1	3.60-2	1.11+0	7.50-1	1.80-1	1.07+0	1.00+0	5.50-1	1.15+0	95
1.25+0	1.20+0	1.18+0	1.50+0	2.20+0	1.22+0	2.00+0	5.20+0	1.23+0	
2.50+0	9.20+0	1.19+0							
4.00+0	2.13+1	9.20-1	6.00+0	5.69+1	1.16+0	8.00+0	8.53+1	1.13+0	105
1.00+1	1.14+2	1.16+0	1.20+1	1.49+2	1.26+0	1.40+1	1.56+2	1.16+0	
1.60+1	1.85+2	1.26+0	1.80+1	1.92+2	1.22+0	2.00+1	2.13+2	1.29+0	
2.20+1	2.06+2	1.20+0							
<b>40</b>	Zirconium		Fluorescence yield = 0.73						
2.30-1	1.20-4	<b>3.33-1</b>	3.25-1	1.00-3	<b>3.39-1</b>	4.20-1	4.00-3	<b>3.68-1</b>	7
4.54-1	8.75-3	5.59-1	5.10-1	1.20-2	<b>4.55-1</b>				
1.60+2	1.35+2	1.39+0							30
1.00+0	3.10-1	8.38-1	2.25+0	3.60+0	7.66-1	3.00+0	7.50+0	7.59-1	47
4.00-1	3.51-3	<b>4.09-1</b>	4.50-1	5.69-3	<b>3.79-1</b>	5.00-1	1.75-2	7.23-1	111
7.00-1	6.28-2	6.40-1	9.00-1	2.10-1	8.28-1	1.00+0	3.24-1	8.76-1	
1.10+0	4.56-1	8.85-1	1.20+0	5.88-1	8.50-1	1.30+0	7.74-1	8.59-1	
1.40+0	9.49-1	8.28-1	1.50+0	1.28+0	8.98-1				
1.00+0	3.77-1	1.02+0	2.00+0	3.34+0	9.89-1				113
7.00+0	6.00+1	1.13+0							114
2.75-1	5.69-4	5.03-1	3.00-1	1.15-3	6.09-1	3.50-1	2.91-3	6.65-1	118
3.80-1	4.38-3	6.57-1	4.00-1	6.34-3	7.38-1	5.00-1	1.88-2	7.77-1	
6.00-1	4.26-2	8.09-1	7.00-1	8.03-2	8.19-1	8.00-1	1.35-1	8.24-1	
7.00+0	6.35+1	1.19+0							125
1.50+0	1.15+0	8.07-1	2.10+0	3.20+0	8.25-1	2.60+0	5.39+0	7.80-1	148
3.10+0	8.67+0	8.11-1	3.60+0	1.14+1	7.52-1				
<b>41</b>	Niobium		Fluorescence yield = 0.74						
9.00-1	8.60-1	<b>4.40+0</b>	1.10+0	1.60+0	<b>3.99+0</b>	1.30+0	3.10+0	<b>4.39+0</b>	44
1.50+0	5.20+0	<b>4.63+0</b>	1.70+0	1.10+1	<b>6.63+0</b>	1.90+0	1.30+1	<b>5.61+0</b>	
2.10+0	1.50+1	<b>4.84+0</b>	2.30+0	1.70+1	<b>4.25+0</b>	2.50+0	2.50+1	<b>4.98+0</b>	
1.13+0	3.60-1	8.18-1	1.34+0	6.50-1	8.32-1	1.55+0	1.00+0	8.03-1	87
1.76+0	1.60+0	8.67-1	1.97+0	2.10+0	8.15-1	2.18+0	3.00+0	8.71-1	
2.39+0	3.60+0	8.10-1	2.60+0	4.50+0	8.07-1	2.70+0	4.80+0	7.80-1	
3.00+0	6.80+0	8.45-1							94
2.00-1	7.07-5	8.49-1	2.40-1	3.15-4	9.75-1	2.80-1	9.61-4	1.10+0	126
3.20-1	2.14-3	1.11+0	3.60-1	4.28-3	1.17+0	4.00-1	7.65-3	1.22+0	
4.50-1	1.39-2	1.25+0	5.00-1	2.32-2	1.28+0				

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
1.50+0	9.38-1	8.35-1	2.10+0	2.59+0	8.36-1	2.60+0	4.50+0	8.07-1	148
3.10+0	7.18+0	8.25-1	3.60+0	1.04+1	8.35-1				
<b>42 Molybdenum      Fluorescence yield = 0.765</b>									
2.40+0	8.00+0	<b>2.20+0</b>							1
2.50-1	1.90-4	6.55-1	3.00-1	5.50-4	5.86-1	3.50-1	1.60-3	7.01-1	3
4.00-1	3.30-3	7.14-1	7.00-1	6.50-2	1.13+0	1.00+0	2.30-1	1.03+0	
1.22+0	4.40-1	9.92-1	1.61+0	1.20+0	1.08+0				
2.40-1	5.50-4	<b>2.53+0</b>	2.60-1	8.20-4	<b>2.16+0</b>	2.80-1	1.20-3	<b>1.95+0</b>	6
3.00-1	1.80-3	<b>1.92+0</b>	3.20-1	2.50-3	1.82+0	3.40-1	3.60-3	1.85+0	
3.60-1	4.70-3	1.77+0	3.80-1	6.30-3	1.78+0	4.00-1	8.10-3	1.75+0	
4.40-1	1.30-2	1.75+0	6.00-1	3.40-2	1.12+0	7.40-1	8.50-2	1.18+0	
9.35-1	2.10-1	1.20+0	1.03+0	2.90-1	1.17+0	1.20+0	5.30-1	1.25+0	
4.54-1	5.62-3	6.50-1							7
1.60+2	1.18+2	1.34+0							30
2.50+0	3.13+0	7.70-1	3.00+0	5.38+0	8.18-1	4.00+0	1.00+1	7.66-1	52
4.50+0	1.31+1	7.76-1	5.00+0	1.71+1	8.15-1	6.00+0	2.36+1	7.96-1	
6.50+0	2.17+1	6.38-1	7.00+0	2.21+1	5.74-1	8.00+0	3.80+1	8.01-1	
8.50+0	4.12+1	7.94-1	9.00+0	3.73+1	6.65-1	1.00+1	5.34+1	8.29-1	
1.05+1	4.67+1	6.84-1	1.10+1	6.35+1	8.79-1	1.20+1	7.11+1	8.95-1	
4.00-1	4.72-3	1.02+0	6.00-1	2.94-2	9.68-1	8.00-1	9.68-2	9.95-1	61
1.00+0	2.18-1	9.74-1	1.20+0	4.02-1	9.46-1	1.40+0	6.79-1	9.52-1	
1.60+0	1.13+0	1.03+0	1.80+0	1.48+0	9.39-1	2.00+0	2.05+0	9.49-1	
1.00+0	2.35-1	1.05+0	1.20+0	4.21-1	9.90-1	1.40+0	6.95-1	9.75-1	77
1.50+0	8.71-1	9.77-1	1.60+0	1.10+0	1.00+0	1.80+0	1.58+0	1.00+0	
2.00+0	2.20+0	1.02+0	2.20+0	2.92+0	1.03+0	2.40+0	3.68+0	1.01+0	
2.50+0	4.06+0	9.98-1	2.60+0	4.42+0	9.78-1	2.80+0	5.07+0	9.21-1	
3.00+0	5.52+0	8.40-1							
5.00-1	1.40-2	1.03+0	7.50-1	7.50-2	9.91-1	1.00+0	2.30-1	1.03+0	95
1.25+0	5.20-1	1.06+0	1.50+0	9.90-1	1.11+0	2.00+0	2.40+0	1.11+0	
2.50+0	4.50+0	1.11+0							
7.00+0	3.64+1	9.45-1							125
3.90-1	4.10-3	1.01+0	4.30-1	6.80-3	1.02+0	4.80-1	1.10-2	9.78-1	135
5.50-1	1.90-2	9.12-1	6.00-1	2.90-2	9.55-1	6.60-1	3.70-2	8.19-1	
7.10-1	5.00-2	8.21-1	7.70-1	6.90-2	8.23-1	8.00-1	9.90-2	1.02+0	
8.50-1	1.10-1	8.95-1	9.00-1	1.40-1	9.20-1	1.01+0	2.20-1	9.48-1	
1.11+0	3.10-1	9.55-1	1.21+0	4.30-1	9.83-1	1.31+0	5.40-1	9.45-1	
1.41+0	6.80-1	9.32-1	1.51+0	7.70-1	8.46-1	1.61+0	1.10+0	9.85-1	
1.71+0	1.20+0	8.90-1	1.81+0	1.50+0	9.36-1				

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_i$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_i$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_i$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
5.00-1	1.31-2	9.65-1	6.00-1	3.17-2	1.04+0	7.00-1	6.01-2	1.05+0	144
8.00-1	1.04-1	1.07+0	9.00-1	1.56-1	1.03+0	1.00+0	2.39-1	1.07+0	
1.20+0	4.54-1	1.07+0	1.40+0	7.52-1	1.05+0	1.60+0	1.16+0	1.06+0	
1.80+0	1.64+0	1.04+0	2.00+0	2.24+0	1.04+0	2.20+0	2.98+0	1.05+0	
2.40+0	3.73+0	1.03+0	2.60+0	4.63+0	1.02+0	2.80+0	5.62+0	1.02+0	
3.00+0	6.66+0	1.01+0	3.20+0	7.80+0	1.01+0	3.50+0	9.64+0	1.01+0	
4.00+0	1.32+1	1.01+0	4.50+0	1.68+1	9.95-1	5.00+0	2.16+1	1.03+0	
5.50+0	2.52+1	9.98-1	6.00+0	3.01+1	1.02+0				
1.50+0	7.09-1	7.95-1	2.10+0	1.97+0	7.91-1	2.60+0	3.40+0	7.52-1	148
3.10+0	5.55+0	7.77-1							
<b>44</b>	Ruthenium	Fluorescence yield = 0.794							
7.00+0	2.63+1	9.43-1							125
<b>45</b>	Rhodium	Fluorescence yield = 0.808							
1.60+2	1.16+2	1.54+0							30
1.03+0	1.00-1	8.29-1	1.24+0	2.10-1	8.92-1	1.45+0	3.40-1	8.46-1	87
1.66+0	5.20-1	8.29-1	1.87+0	7.70-1	8.42-1	2.07+0	1.10+0	8.81-1	
2.29+0	1.50+0	8.89-1	2.50+0	1.90+0	8.75-1	2.70+0	2.20+0	8.16-1	
<b>46</b>	Palladium	Fluorescence yield = 0.82							
6.00-1	8.12-3	7.76-1	8.00-1	3.28-2	9.17-1	1.00+0	7.14-2	8.35-1	61
1.20+0	1.52-1	9.11-1	1.40+0	2.49-1	8.71-1	1.60+0	3.86-1	8.64-1	
1.80+0	5.41-1	8.27-1	2.00+0	7.55-1	8.29-1				
2.15-1	8.50-6	5.19-1	2.30-1	1.90-5	6.33-1	2.45-1	4.20-5	8.22-1	120
2.60-1	6.50-5	7.93-1	2.75-1	9.70-5	7.74-1	3.00-1	1.70-4	7.30-1	
5.00-1	4.12-3	9.32-1	6.00-1	1.07-2	1.02+0	7.00-1	2.12-2	1.03+0	144
8.00-1	3.85-2	1.08+0	9.00-1	6.10-2	1.07+0	1.00+0	9.27-2	1.08+0	
1.20+0	1.84-1	1.10+0	1.40+0	3.14-1	1.10+0	1.60+0	4.86-1	1.09+0	
1.80+0	7.06-1	1.08+0	2.00+0	9.76-1	1.07+0	2.20+0	1.31+0	1.08+0	
2.40+0	1.69+0	1.07+0	2.60+0	2.10+0	1.06+0	2.80+0	2.53+0	1.04+0	
3.00+0	3.09+0	1.05+0	3.20+0	3.55+0	1.01+0	3.50+0	4.53+0	1.02+0	
4.00+0	6.37+0	1.04+0	4.50+0	8.36+0	1.03+0	5.00+0	1.06+1	1.03+0	
5.50+0	1.30+1	1.03+0	6.00+0	1.53+1	1.01+0				
1.50+0	2.99-1	8.28-1	2.10+0	8.58-1	8.12-1	2.60+0	1.70+0	8.58-1	148
3.10+0	2.43+0	7.55-1	3.60+0	4.11+0	8.65-1	3.70+0	4.46+0	8.77-1	
3.80+0	4.51+0	8.30-1							
<b>47</b>	Silver	Fluorescence yield = 0.831							
1.70+0	5.30-1	1.21+0	1.92+0	1.00+0	1.54+0	2.17+0	1.60+0	1.69+0	1
2.40+0	2.30+0	1.79+0	2.64+0	3.30+0	1.95+0	2.88+0	6.30+0	2.91+0	
6.00-1	3.20-3	<b>3.97-1</b>	7.00-1	7.83-3	4.89-1	8.00-1	1.35-2	4.81-1	4
9.00-1	2.58-2	5.72-1	1.00+0	3.58-2	5.29-1				

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
1.80+0	4.40-1	8.32-1	2.00+0	4.90-1	6.64-1	2.10+0	5.70-1	6.64-1	5
2.10+0	7.50-1	8.74-1	2.30+0	8.50-1	7.52-1	2.40+0	9.20-1	7.17-1	
2.40+0	1.00+0	7.79-1	2.60+0	1.20+0	7.41-1	2.70+0	1.40+0	7.76-1	
2.80+0	1.60+0	8.00-1	2.90+0	2.10+0	9.52-1	3.20+0	2.70+0	9.36-1	
3.40+0	3.00+0	8.86-1	3.60+0	3.40+0	8.65-1				
2.60-1	1.30-4	<b>2.34+0</b>	2.80-1	2.00-4	<b>2.03+0</b>	3.00-1	3.30-4	<b>2.02+0</b>	6
3.20-1	4.90-4	<b>1.92+0</b>	3.40-1	7.40-4	<b>1.96+0</b>	3.60-1	1.10-3	<b>2.03+0</b>	
3.80-1	1.50-3	<b>2.00+0</b>	4.00-1	2.10-3	<b>2.07+0</b>	6.00-1	1.10-2	1.36+0	
7.40-1	2.40-2	1.18+0	9.35-1	6.60-2	1.26+0	1.04+0	9.90-2	1.26+0	
1.20+0	1.70-1	1.28+0							
1.60+2	1.09+2	1.61+0							30
2.00+0	8.10-1	1.10+0	3.00+0	2.60+0	1.07+0	4.00+0	6.10+0	1.19+0	32
5.00+0	1.00+1	1.16+0	6.00+0	1.40+1	1.10+0	7.00+0	1.90+1	1.10+0	
8.00+0	2.40+1	1.10+0	9.00+0	2.90+1	1.09+0	1.00+1	3.60+1	1.15+0	
1.10+1	3.90+1	1.08+0	1.20+1	4.40+1	1.09+0	1.30+1	5.00+1	1.12+0	
1.40+1	5.70+1	1.17+0	1.50+1	5.90+1	1.12+0	1.70+1	7.40+1	1.24+0	
1.80+1	8.20+1	1.30+0	1.90+1	7.90+1	1.20+0	2.00+1	7.90+1	1.15+0	
2.10+1	8.60+1	1.21+0	2.20+1	9.80+1	1.33+0	2.30+1	9.80+1	1.29+0	
2.40+1	8.40+1	1.08+0	2.50+1	8.50+1	1.06+0	2.60+1	9.50+1	1.16+0	
2.70+1	9.60+1	1.15+0	2.80+1	9.50+1	1.12+0	3.00+1	9.20+1	1.05+0	
1.50+0	3.50-1	1.21+0	2.00+0	1.10+0	1.49+0	2.50+0	1.70+0	1.18+0	38
3.00+0	3.10+0	1.28+0	3.50+0	4.30+0	1.18+0	4.00+0	5.50+0	1.08+0	
4.50+0	7.80+0	1.15+0	5.00+0	9.50+0	1.10+0	5.50+0	1.10+1	1.03+0	
1.00+0	8.00-2	1.18+0	2.00+0	8.10-1	1.10+0				39
1.00+0	6.50-2	9.60-1	2.25+0	1.00+0	9.45-1	3.00+0	2.30+0	9.50-1	47
2.50+0	1.07+0	7.40-1	3.00+0	1.88+0	7.76-1	4.00+0	4.18+0	8.17-1	52
4.50+0	5.49+0	8.10-1	5.00+0	6.81+0	7.90-1	6.00+0	8.88+0	6.97-1	
6.50+0	1.25+1	8.36-1	7.00+0	1.41+1	8.18-1	8.00+0	1.88+1	8.58-1	
8.50+0	2.05+1	8.46-1	9.00+0	2.14+1	8.04-1	1.00+1	2.60+1	8.30-1	
1.05+1	2.45+1	7.28-1	1.10+1	3.09+1	8.57-1	1.20+1	3.69+1	9.13-1	
1.50+0	2.85-1	9.83-1	2.00+0	7.52-1	1.02+0	2.50+0	1.44+0	9.96-1	55
2.98+0	2.19+0	9.21-1	3.00+0	2.42+0	9.99-1	4.97+0	7.68+0	9.03-1	
5.96+0	1.12+1	8.92-1	6.94+0	1.54+1	9.08-1	6.96+0	1.48+1	8.66-1	
8.94+0	2.36+1	8.97-1	1.09+1	3.22+1	9.02-1				
6.00-1	6.70-3	8.31-1	8.00-1	2.70-2	9.63-1	1.00+0	6.40-2	9.45-1	63
1.20+0	1.30-1	9.77-1	1.40+0	2.20-1	9.60-1	1.60+0	3.40-1	9.45-1	
1.80+0	4.70-1	8.89-1	2.00+0	6.70-1	9.08-1				
1.00+0	7.06-2	1.04+0	1.10+0	9.88-2	1.02+0	1.20+0	1.34-1	1.01+0	77
1.30+0	1.78-1	1.01+0	1.40+0	2.32-1	1.01+0	1.50+0	2.96-1	1.02+0	
1.60+0	3.70-1	1.03+0	1.70+0	4.57-1	1.04+0	1.80+0	5.54-1	1.05+0	
1.90+0	6.63-1	1.06+0	2.00+0	7.82-1	1.06+0	2.10+0	9.13-1	1.06+0	
2.20+0	1.05+0	1.06+0	2.30+0	1.20+0	1.06+0	2.40+0	1.34+0	1.04+0	
2.50+0	1.49+0	1.03+0	2.60+0	1.63+0	1.01+0	2.70+0	1.77+0	9.81-1	
2.80+0	1.89+0	9.45-1	2.90+0	2.00+0	9.07-1	3.00+0	2.10+0	8.67-1	

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
4.00-1	9.00-4	8.88-1	6.00-1	9.40-3	1.17+0	8.00-1	3.30-2	1.18+0	87
1.00+0	7.90-2	1.17+0	1.20+0	1.40-1	1.05+0	1.40+0	2.30-1	1.00+0	
1.60+0	3.60-1	1.00+0	1.80+0	5.20-1	9.84-1	2.00+0	7.50-1	1.02+0	
2.20+0	9.90-1	1.00+0	2.40+0	1.30+0	1.01+0				
3.00+0	1.93+0	7.97-1							94
5.00-1	2.80-3	8.34-1	7.50-1	1.80-2	8.38-1	1.00+0	6.30-2	9.30-1	95
1.25+0	1.50-1	9.74-1	1.50+0	3.00-1	1.03+0	2.00+0	7.90-1	1.07+0	
2.50+0	1.70+0	1.18+0							
2.49-1	1.76-5	4.53-1	3.00-1	9.84-5	6.02-1	3.53-1	4.09-4	8.53-1	99
4.03-1	9.34-4	8.83-1	4.84-1	2.37-3	8.33-1	5.52-1	4.51-3	8.27-1	
6.35-1	8.93-3	8.56-1	7.20-1	1.54-2	8.52-1	8.10-1	2.58-2	8.74-1	
9.00-1	4.02-2	8.92-1	1.10+0	8.60-2	8.88-1	1.52+0	2.64-1	8.70-1	
1.72+0	4.15-1	9.09-1	1.91+0	5.80-1	9.08-1				
4.00+0	5.00+0	9.77-1	7.50+0	1.69+1	8.64-1	8.00+0	1.83+1	8.35-1	100
9.00+0	2.24+1	8.42-1	1.00+1	2.62+1	8.36-1	1.10+1	3.24+1	8.99-1	
1.20+1	3.58+1	8.86-1	1.30+1	3.93+1	8.79-1	1.40+1	4.29+1	8.79-1	
1.50+1	4.87+1	9.26-1							
4.00-1	3.22-4	<b>3.18-1</b>	5.00-1	1.62-3	4.82-1	6.00-1	4.64-3	5.75-1	103
7.00-1	9.58-3	5.98-1	8.00-1	1.93-2	6.88-1				
2.20+1	7.22+1	9.80-1	3.10+1	8.20+1	9.25-1	4.40+1	8.34+1	8.64-1	104
3.50-1	3.30-4	7.26-1	4.00-1	8.17-4	8.06-1	5.00-1	3.00-3	8.93-1	113
6.00-1	7.50-3	9.30-1	8.00-1	2.77-2	9.88-1	1.00+0	7.08-2	1.05+0	
1.20+0	1.38-1	1.04+0	1.40+0	2.40-1	1.05+0	1.60+0	3.70-1	1.03+0	
1.80+0	5.23-1	9.89-1	2.00+0	7.52-1	1.02+0	2.20+0	1.00+0	1.01+0	
3.50-1	1.45-4	<b>3.19-1</b>	3.75-1	2.53-4	<b>3.65-1</b>	4.00-1	4.42-4	<b>4.36-1</b>	118
4.50-1	9.89-4	5.09-1	5.00-1	1.87-3	5.57-1	6.00-1	5.10-3	6.32-1	
7.00-1	1.06-2	6.62-1	8.00-1	1.93-2	6.88-1				
2.00-1	2.80-6	5.60-1	2.15-1	5.50-6	5.31-1	2.30-1	1.00-5	5.13-1	120
2.45-1	1.90-5	5.60-1	2.60-1	3.50-5	6.31-1	2.75-1	5.90-5	6.86-1	
3.00-1	1.40-4	8.57-1							
2.50-1	1.89-5	4.70-1	3.00-1	1.05-4	6.43-1	3.50-1	3.26-4	7.17-1	121
4.00-1	7.01-4	6.92-1	5.00-1	2.46-3	7.33-1	6.00-1	6.62-3	8.21-1	
7.00-1	1.36-2	8.49-1	8.00-1	2.40-2	8.56-1	9.00-1	4.05-2	8.98-1	
1.00+0	6.18-2	9.13-1							
7.00+0	1.70+1	9.86-1							125
2.50-1	1.96-5	4.87-1	3.00-1	1.16-4	7.10-1	3.50-1	3.75-4	8.25-1	126
4.00-1	1.00-3	9.87-1	4.50-1	2.10-3	1.08+0	5.00-1	3.64-3	1.08+0	
5.50-1	6.08-3	1.13+0	6.00-1	9.57-3	1.19+0				
7.07-1	1.70-2	1.02+0	1.00+0	7.30-2	1.08+0	1.41+0	2.40-1	1.01+0	130
2.00+0	7.50-1	1.02+0	2.50+0	1.50+0	1.04+0				

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_i$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_i$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_i$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
4.20-1	1.60-3	1.20+0	4.60-1	2.80-3	1.28+0	5.10-1	4.30-3	1.16+0	135
5.50-1	6.50-3	1.21+0	6.00-1	9.10-3	1.13+0	6.40-1	1.20-2	1.11+0	
6.90-1	1.70-2	1.13+0	7.40-1	2.20-2	1.08+0	8.20-1	3.20-2	1.03+0	
8.80-1	4.40-2	1.07+0	1.01+0	7.60-2	1.08+0	1.11+0	1.20-1	1.20+0	
1.21+0	1.50-1	1.09+0	1.31+0	1.90-1	1.05+0	1.41+0	2.60-1	1.11+0	
1.51+0	3.20-1	1.08+0	1.61+0	4.00-1	1.09+0	1.71+0	4.60-1	1.03+0	
1.81+0	5.90-1	1.10+0							
1.50+0	2.62-1	9.04-1	2.10+0	6.68-1	7.78-1	2.60+0	1.21+0	7.47-1	148
3.10+0	1.95+0	7.36-1	3.30+0	2.44+0	7.79-1	3.60+0	3.06+0	7.79-1	
3.80+0	3.57+0	7.92-1							
1.60+0	3.66-1	1.02+0	1.80+0	5.72-1	1.08+0	2.00+0	7.15-1	9.69-1	151
2.20+0	1.07+0	1.08+0	2.40+0	1.32+0	1.03+0				
<b>48 Cadmium</b> <b>Fluorescence yield = 0.843</b>									
2.50+0	9.07-1	7.65-1	3.00+0	1.65+0	8.26-1	4.00+0	3.86+0	9.05-1	52
4.50+0	4.31+0	7.61-1	5.00+0	5.44+0	7.49-1	6.00+0	8.70+0	8.06-1	
6.50+0	8.53+0	6.71-1	7.00+0	1.17+1	7.96-1	8.00+0	1.49+1	7.92-1	
8.50+0	1.82+1	8.71-1	9.00+0	1.77+1	7.71-1	1.00+1	2.20+1	8.08-1	
1.05+1	2.12+1	7.25-1	1.10+1	2.70+1	8.62-1	1.20+1	2.99+1	8.46-1	
6.00-1	4.80-3	7.68-1	8.00-1	1.80-2	8.13-1	1.00+0	4.90-2	9.09-1	63
1.20+0	1.00-1	9.38-1	1.40+0	1.70-1	9.21-1	1.60+0	2.70-1	9.27-1	
1.80+0	3.90-1	9.09-1	2.00+0	5.40-1	8.99-1				
1.00+0	4.60-2	8.53-1	1.20+0	9.00-2	8.44-1	1.40+0	1.60-1	8.67-1	86
1.60+0	2.50-1	8.58-1	1.80+0	3.80-1	8.85-1	2.00+0	5.30-1	8.82-1	
2.20+0	7.10-1	8.79-1	2.40+0	9.00-1	8.57-1	2.60+0	1.10+0	8.27-1	
2.80+0	1.30+0	7.90-1	3.00+0	1.60+0	8.01-1				
3.00+0	1.47+0	7.36-1							94
6.00-1	2.27-3	<b>3.63-1</b>	7.00-1	6.14-3	4.90-1	8.00-1	1.57-2	7.10-1	111
9.00-1	2.52-2	7.06-1	1.00+0	3.46-2	6.42-1	1.10+0	5.46-2	7.05-1	
1.20+0	7.90-2	7.41-1	1.30+0	1.08-1	7.60-1	1.40+0	1.40-1	7.59-1	
1.50+0	1.70-1	7.27-1	1.60+0	2.25-1	7.73-1	1.70+0	2.73-1	7.67-1	
1.80+0	3.18-1	7.41-1	1.90+0	3.97-1	7.78-1	2.00+0	4.74-1	7.89-1	
2.10+0	5.25-1	7.50-1	2.20+0	6.33-1	7.84-1	2.30+0	6.62-1	7.16-1	
7.00+0	1.32+1	8.98-1							125
<b>49 Indium</b> <b>Fluorescence yield = 0.853</b>									
3.00+0	1.51+0	9.16-1	5.00+0	5.20+0	8.53-1				59
9.00-1	3.74-2	1.32+0	1.00+0	6.12-2	1.42+0	1.10+0	9.35-2	1.51+0	80
1.20+0	1.28-1	1.50+0	1.30+0	1.70-1	1.49+0	1.40+0	2.30-1	1.55+0	
1.50+0	2.55-1	1.35+0	1.60+0	3.40-1	1.44+0	1.70+0	3.91-1	1.35+0	
1.80+0	4.08-1	1.17+0	1.90+0	5.10-1	1.23+0	2.00+0	4.34-1	8.87-1	
2.10+0	6.89-1	1.21+0	2.20+0	7.06-1	1.07+0	2.30+0	9.86-1	1.30+0	
2.40+0	1.31+0	1.52+0	2.50+0	1.49+0	1.53+0				

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
1.00+0	3.60-2	8.37-1	1.20+0	7.30-2	8.53-1	1.40+0	1.30-1	8.74-1	86
1.60+0	2.00-1	8.49-1	1.80+0	3.10-1	8.89-1	2.00+0	4.30-1	8.78-1	
2.20+0	5.90-1	8.94-1	2.40+0	7.60-1	8.83-1	2.60+0	9.00-1	8.24-1	
2.80+0	1.10+0	8.12-1	3.00+0	1.30+0	7.89-1				
6.00-1	5.50-3	1.13+0	8.00-1	1.90-2	1.09+0	1.00+0	5.00-2	1.16+0	87
1.20+0	8.50-2	9.93-1	1.40+0	1.50-1	1.01+0	1.60+0	2.30-1	9.76-1	
1.80+0	3.50-1	1.00+0	2.00+0	5.00-1	1.02+0	2.20+0	6.60-1	1.00+0	
2.40+0	8.50-1	9.87-1							
3.00+0	1.12+0	6.80-1	5.00+0	3.62+0	5.94-1	7.00+0	9.20+0	7.34-1	94
9.00+0	1.62+1	8.18-1	1.10+1	2.30+1	8.45-1				
6.00-1	1.73-3	<b>3.57-1</b>	7.00-1	4.25-3	<b>4.32-1</b>	8.00-1	8.50-3	4.86-1	111
9.00-1	1.88-2	6.63-1	1.00+0	2.64-2	6.14-1	1.10+0	3.90-2	6.30-1	
1.20+0	5.70-2	6.66-1	1.30+0	7.23-2	6.32-1	1.40+0	9.27-2	6.23-1	
1.50+0	1.22-1	6.45-1	1.60+0	1.49-1	6.32-1	1.70+0	1.98-1	6.85-1	
1.80+0	2.39-1	6.86-1	1.90+0	2.53-1	6.08-1	2.00+0	3.03-1	6.19-1	
2.10+0	3.71-1	6.49-1							
3.00-1	3.36-5	4.19-1	3.50-1	1.32-4	5.49-1	4.00-1	3.47-4	6.23-1	129
5.00-1	1.46-3	7.48-1	6.00-1	3.77-3	7.78-1	7.00-1	8.26-3	8.40-1	
8.00-1	1.50-2	8.58-1	9.00-1	2.48-2	8.75-1	1.00+0	3.77-2	8.76-1	
<b>50</b>	Tin		Fluorescence yield = 0.862						
2.60-1	5.70-5	<b>3.39+0</b>	2.80-1	9.70-5	<b>3.03+0</b>	3.20-1	2.10-4	<b>2.29+0</b>	6
3.60-1	5.20-4	<b>2.47+0</b>	3.80-1	7.70-4	<b>2.58+0</b>	4.00-1	1.10-3	<b>2.66+0</b>	
4.40-1	2.00-3	<b>2.73+0</b>	4.50-1	2.30-3	<b>2.76+0</b>	6.00-1	5.60-3	1.49+0	
7.40-1	1.40-2	1.42+0	9.35-1	3.80-2	1.44+0	1.04+0	5.90-2	1.47+0	
1.60+2	9.41+1	1.63+0							30
1.00+0	3.20-2	9.29-1	2.25+0	5.50-1	9.48-1	3.00+0	1.30+0	9.54-1	47
1.80+0	3.13-1	1.10+0	2.20+0	5.82-1	1.08+0	2.60+0	9.19-1	1.02+0	60
3.00+0	1.32+0	9.69-1	3.20+0	1.71+0	1.05+0	3.60+0	2.25+0	9.99-1	
4.00+0	3.31+0	1.11+0	4.40+0	4.03+0	1.07+0				
6.00-1	3.20-3	8.50-1	8.00-1	8.20-3	5.92-1	1.00+0	3.20-2	9.29-1	63
1.20+0	5.80-2	8.41-1	1.40+0	1.00-1	8.30-1	1.60+0	1.60-1	8.36-1	
1.80+0	2.40-1	8.45-1	2.00+0	3.20-1	7.99-1				
2.33+0	6.10-1	9.41-1	2.67+0	9.41-1	9.70-1	3.00+0	1.38+0	1.01+0	64
3.33+0	1.76+0	9.64-1	3.67+0	2.39+0	1.01+0	4.00+0	2.96+0	9.96-1	
4.33+0	3.69+0	1.01+0	4.67+0	4.40+0	1.01+0	5.00+0	5.25+0	1.02+0	
5.33+0	6.05+0	1.01+0	5.67+0	7.03+0	1.03+0				
1.00+0	2.60-2	7.55-1	1.20+0	5.60-2	8.12-1	1.40+0	1.00-1	8.30-1	86
1.60+0	1.60-1	8.36-1	1.80+0	2.40-1	8.45-1	2.00+0	3.40-1	8.49-1	
2.20+0	4.50-1	8.31-1	2.40+0	5.80-1	8.20-1	2.60+0	7.30-1	8.12-1	
2.80+0	9.00-1	8.05-1	2.90+0	1.00+0	8.08-1	3.00+0	1.20+0	8.81-1	
3.00+0	9.40-1	6.90-1							94

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
5.00-1	1.00-3	6.71-1	7.50-1	8.00-3	7.63-1	1.00+0	2.90-2	8.42-1	95
1.25+0	7.30-2	9.12-1	1.50+0	1.50-1	9.78-1	2.00+0	4.00-1	9.99-1	
2.50+0	7.60-1	9.50-1							
8.00-1	4.50-3	<b>3.25-1</b>	9.00-1	1.00-2	<b>4.43-1</b>	1.00+0	1.67-2	4.85-1	111
1.10+0	2.31-2	4.65-1	1.20+0	3.55-2	5.15-1	1.30+0	5.11-2	5.53-1	
1.40+0	6.45-2	5.36-1	1.50+0	8.00-2	5.22-1	1.60+0	9.80-2	5.12-1	
1.70+0	1.21-1	5.15-1	1.80+0	1.60-1	5.63-1	1.90+0	1.91-1	5.63-1	
2.00+0	2.18-1	5.45-1	2.10+0	2.68-1	5.74-1				
7.00+0	1.07+1	1.00+0							125
4.20-1	8.60-4	1.55+0	5.10-1	1.80-3	1.09+0	6.00-1	3.00-3	7.97-1	135
7.00-1	5.70-3	7.38-1	7.50-1	7.80-3	7.44-1	7.90-1	1.00-2	7.62-1	
8.30-1	1.50-2	9.27-1	8.80-1	2.10-2	1.02+0	1.01+0	3.40-2	9.49-1	
1.11+0	4.80-2	9.33-1	1.21+0	6.80-2	9.56-1	1.31+0	9.50-2	1.00+0	
1.41+0	1.30-1	1.05+0	1.51+0	1.60-1	1.02+0	1.61+0	1.90-1	9.72-1	
1.71+0	2.10-1	8.76-1	1.81+0	2.70-1	9.33-1				
5.00-1	1.36-3	9.12-1	6.00-1	3.59-3	9.53-1	7.00-1	7.63-3	9.87-1	144
8.00-1	1.46-2	1.05+0	9.00-1	2.43-2	1.08+0	1.00+0	3.70-2	1.07+0	
1.10+0	5.44-2	1.09+0	1.20+0	7.50-2	1.09+0	1.30+0	1.00-1	1.08+0	
1.40+0	1.32-1	1.10+0	1.60+0	2.12-1	1.11+0	1.80+0	3.14-1	1.11+0	
2.00+0	4.37-1	1.09+0	2.20+0	5.92-1	1.09+0	2.40+0	7.62-1	1.08+0	
2.60+0	9.74-1	1.08+0	2.80+0	1.19+0	1.06+0	3.00+0	1.46+0	1.07+0	
3.20+0	1.72+0	1.05+0	3.50+0	2.22+0	1.06+0	4.00+0	3.11+0	1.05+0	
4.50+0	4.17+0	1.04+0	5.00+0	5.26+0	1.03+0	5.50+0	6.64+0	1.04+0	
6.00+0	7.93+0	1.02+0							
<b>51</b>	<b>Antimony</b>				<b>Fluorescence yield = 0.87</b>				
1.00+0	2.60-2	9.41-1	2.25+0	5.40-1	1.13+0	3.00+0	1.20+0	1.06+0	47
6.00-1	1.40-3	4.77-1	8.00-1	1.00-2	9.09-1	1.00+0	2.60-2	9.41-1	63
1.20+0	5.50-2	9.88-1	1.40+0	1.00-1	1.02+0	1.60+0	1.60-1	1.03+0	
1.80+0	2.30-1	9.90-1	2.00+0	3.40-1	1.04+0				
3.00-1	1.86-5	4.77-1	3.50-1	7.25-5	5.77-1	4.00-1	2.03-4	6.62-1	126
4.50-1	5.49-4	8.73-1	5.00-1	1.12-3	9.82-1	5.50-1	1.82-3	9.62-1	
6.00-1	3.00-3	1.02+0	6.50-1	4.77-3	1.10+0				
3.50-1	7.50-5	5.96-1	4.00-1	1.92-4	6.26-1	5.00-1	8.93-4	7.83-1	129
6.00-1	2.28-3	7.77-1	7.00-1	5.12-3	8.41-1	8.00-1	8.93-3	8.12-1	
9.00-1	1.48-2	8.20-1	1.00+0	2.57-2	9.31-1				
2.10+0	3.15-1	8.21-1	2.60+0	6.06-1	8.17-1	3.10+0	9.55-1	7.70-1	148
3.30+0	1.32+0	8.93-1							
<b>52</b>	<b>Tellurium</b>				<b>Fluorescence yield = 0.877</b>				
6.00-1	2.20-3	9.60-1	8.00-1	9.00-3	1.03+0	1.00+0	1.90-2	8.55-1	63
1.20+0	3.30-2	7.32-1	1.40+0	5.90-2	7.42-1	1.60+0	9.20-2	7.22-1	
1.80+0	1.20-1	6.31-1	2.00+0	1.80-1	6.68-1				

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
7.00+0	7.06+0	9.05-1							125
1.60+0	6.14-2	4.82-1	1.80+0	8.26-2	<b>4.34-1</b>	2.00+0	1.25-1	<b>4.64-1</b>	138
2.20+0	1.79-1	4.89-1	2.40+0	2.42-1	5.03-1	2.60+0	2.82-1	<b>4.59-1</b>	
2.80+0	4.17-1	5.44-1	3.00+0	5.19-1	5.54-1				
<b>53 Iodine</b> Fluorescence yield = 0.884									
6.00-1	1.50-3	8.36-1	8.00-1	6.20-3	8.85-1	1.00+0	1.40-2	7.80-1	87
1.20+0	3.20-2	8.73-1	1.40+0	5.90-2	9.08-1	1.60+0	9.20-2	8.81-1	
1.80+0	1.50-1	9.59-1	2.00+0	2.10-1	9.45-1	2.20+0	3.00-1	9.92-1	
2.40+0	3.60-1	9.05-1							
3.00+0	5.50-1	7.04-1	5.00+0	2.36+0	7.64-1	7.00+0	5.14+0	7.71-1	94
9.00+0	8.70+0	7.91-1	1.10+1	1.18+1	7.57-1				
6.11-1	2.01-3	1.02+0	8.11-1	6.01-3	8.07-1	1.02+0	1.62-2	8.37-1	127
1.20+0	3.32-2	9.05-1	1.40+0	5.79-2	8.91-1	1.60+0	9.60-2	9.19-1	
1.80+0	1.34-1	8.56-1	2.00+0	1.96-1	8.82-1	2.20+0	2.70-1	8.92-1	
2.40+0	3.58-1	9.00-1	2.60+0	4.78-1	9.39-1	2.80+0	5.48-1	8.60-1	
3.00+0	6.23-1	7.97-1	3.20+0	7.52-1	7.99-1	3.40+0	9.45-1	8.46-1	
3.60+0	1.11+0	8.47-1	3.80+0	1.32+0	8.68-1	3.85+0	1.36+0	8.64-1	
1.50+0	8.07-2	9.69-1	2.00+0	2.02-1	9.09-1	2.25+0	2.95-1	9.08-1	143
2.50+0	4.56-1	1.01+0	2.75+0	4.82-1	7.98-1	3.00+0	7.88-1	1.01+0	
<b>54 Xenon</b> Fluorescence yield = 0.891									
4.50+0	4.73+0	<b>2.36+0</b>	5.00+0	6.64+0	<b>2.54+0</b>				48
<b>55 Cesium</b> Fluorescence yield = 0.897									
1.13+0	1.60-2	8.26-1	1.34+0	2.80-2	7.52-1	1.55+0	5.00-2	7.90-1	87
1.76+0	7.80-2	7.89-1	1.97+0	1.10-1	7.60-1	2.18+0	1.70-1	8.41-1	
2.39+0	2.10-1	7.73-1	2.60+0	2.90-1	8.21-1	2.70+0	3.00-1	7.56-1	
1.50+0	5.96-2	1.06+0	2.00+0	1.66-1	1.09+0	2.25+0	2.26-1	1.01+0	143
2.50+0	3.56-1	1.14+0	2.75+0	4.34-1	1.03+0	3.00+0	6.06-1	1.11+0	
<b>56 Barium</b> Fluorescence yield = 0.902									
1.60+2	4.22+1	1.02+0							30
1.00+0	8.40-3	8.70-1	1.20+0	1.40-2	6.96-1	1.40+0	2.00-2	5.54-1	63
1.60+0	3.00-2	5.12-1	1.80+0	4.40-2	4.97-1	2.00+0	4.90-2	<b>3.87-1</b>	
7.00+0	3.75+0	8.92-1							125
1.50+0	4.65-2	1.00+0	2.00+0	1.07-1	8.45-1	2.25+0	2.05-1	1.10+0	143
2.50+0	3.26-1	1.25+0	2.75+0	4.04-1	1.15+0	3.00+0	5.71-1	1.25+0	

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
6.00-1	7.64-4	8.78-1	7.00-1	1.72-3	8.93-1	8.00-1	3.50-3	9.62-1	144
9.00-1	6.29-3	1.02+0	1.00+0	1.03-2	1.07+0	1.10+0	1.52-2	1.07+0	
1.20+0	2.18-2	1.08+0	1.30+0	2.94-2	1.08+0	1.40+0	3.93-2	1.09+0	
1.50+0	5.06-2	1.09+0	1.60+0	6.67-2	1.14+0	1.70+0	8.22-2	1.13+0	
1.80+0	9.65-2	1.09+0	2.00+0	1.42-1	1.12+0	2.20+0	1.90-1	1.10+0	
2.40+0	2.57-1	1.12+0	2.60+0	3.27-1	1.11+0	2.80+0	4.10-1	1.10+0	
3.00+0	4.98-1	1.09+0	3.20+0	6.09-1	1.10+0	3.50+0	7.96-1	1.11+0	
4.00+0	1.16+0	1.11+0	4.50+0	1.53+0	1.07+0	5.00+0	2.00+0	1.06+0	
5.50+0	2.56+0	1.07+0	6.00+0	3.07+0	1.04+0				
<b>57 Lanthanum</b>									
Fluorescence yield = 0.907									
8.00-1	2.70-3	9.17-1	1.00+0	7.40-3	9.37-1	1.20+0	1.40-2	8.45-1	63
1.40+0	2.50-2	8.37-1	1.60+0	3.10-2	6.37-1	1.80+0	3.90-2	5.29-1	
2.00+0	5.20-2	4.92-1							
2.00+0	1.21-1	1.14+0	2.25+0	1.96-1	1.25+0	2.50+0	2.47-1	1.13+0	143
2.75+0	3.57-1	1.21+0	3.00+0	4.93-1	1.28+0				
<b>58 Cerium</b>									
Fluorescence yield = 0.912									
6.00-1	3.20-4	5.89-1	8.00-1	1.70-3	7.12-1	1.00+0	5.10-3	7.86-1	87
1.20+0	1.10-2	8.03-1	1.40+0	2.10-2	8.46-1	1.60+0	3.10-2	7.63-1	
1.80+0	5.60-2	9.08-1	2.00+0	7.40-2	8.36-1	2.20+0	1.10-1	9.04-1	
2.40+0	1.40-1	8.66-1							
7.00+0	2.64+0	8.50-1							125
5.00-1	9.57-5	5.26-1	6.00-1	3.43-4	6.31-1	7.00-1	8.82-4	7.11-1	129
8.00-1	1.97-3	8.25-1	9.00-1	3.87-3	9.44-1	1.00+0	5.69-3	8.77-1	
6.00-1	4.38-4	8.06-1	6.50-1	6.89-4	8.18-1	7.00-1	1.06-3	8.55-1	144
8.00-1	2.23-3	9.34-1	9.00-1	4.01-3	9.78-1	1.00+0	6.51-3	1.00+0	
1.10+0	1.00-2	1.04+0	1.20+0	1.46-2	1.07+0	1.30+0	2.02-2	1.08+0	
1.40+0	2.70-2	1.09+0	1.50+0	3.48-2	1.08+0	1.60+0	4.48-2	1.10+0	
1.70+0	5.62-2	1.11+0	1.80+0	6.87-2	1.11+0	2.00+0	9.94-2	1.12+0	
2.20+0	1.34-1	1.10+0	2.40+0	1.82-1	1.13+0	2.60+0	2.34-1	1.12+0	
2.80+0	2.93-1	1.11+0	3.00+0	3.62-1	1.11+0	3.20+0	4.40-1	1.12+0	
3.50+0	5.73-1	1.12+0	4.00+0	8.30-1	1.11+0	4.50+0	1.14+0	1.10+0	
5.00+0	1.50+0	1.10+0	5.50+0	1.90+0	1.09+0	6.00+0	2.34+0	1.08+0	
<b>59 Praseodymium</b>									
Fluorescence yield = 0.917									
6.00-1	2.90-4	6.72-1	8.00-1	1.31-3	6.74-1	1.00+0	3.70-3	6.92-1	87
1.20+0	8.30-3	7.30-1	1.40+0	1.60-2	7.72-1	1.60+0	2.70-2	7.94-1	
1.80+0	3.90-2	7.54-1	2.00+0	5.70-2	7.66-1	2.20+0	8.10-2	7.90-1	
2.40+0	1.00-1	7.33-1							
5.00-1	3.80-5	<b>2.70-1</b>	7.50-1	6.20-4	<b>4.37-1</b>	1.00+0	3.30-3	6.17-1	95
1.25+0	9.20-3	6.88-1	1.50+0	1.90-2	7.08-1	2.00+0	6.40-2	8.60-1	
2.50+0	1.50-1	9.65-1							
<b>60 Neodymium</b>									
Fluorescence yield = 0.921									
3.00+0	2.10-1	8.99-1	5.00+0	8.40-1	8.35-1				59

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	Ref.
6.00-1	2.10-4	6.13-1	8.00-1	1.10-3	6.94-1	1.00+0	3.40-3	7.68-1	87
1.20+0	8.10-3	8.55-1	1.40+0	1.60-2	9.22-1	1.60+0	2.60-2	9.11-1	
1.80+0	3.90-2	8.96-1	2.00+0	5.30-2	8.44-1	2.20+0	7.50-2	8.65-1	
2.40+0	9.70-2	8.41-1							
7.00-1	6.83-4	8.47-1	8.00-1	1.50-3	9.46-1	9.00-1	2.66-3	9.62-1	144
1.00+0	4.43-3	1.00+0	1.10+0	6.86-3	1.03+0	1.20+0	1.02-2	1.08+0	
1.30+0	1.41-2	1.08+0	1.40+0	1.91-2	1.10+0	1.50+0	2.49-2	1.11+0	
1.60+0	3.19-2	1.12+0	1.80+0	4.92-2	1.13+0	2.00+0	7.06-2	1.12+0	
2.20+0	9.95-2	1.15+0	2.40+0	1.30-1	1.13+0	2.60+0	1.69-1	1.13+0	
2.80+0	2.11-1	1.12+0	3.00+0	2.63-1	1.13+0	3.20+0	3.23-1	1.14+0	
3.50+0	4.21-1	1.13+0	4.00+0	6.06-1	1.11+0	4.50+0	8.32-1	1.10+0	
5.00+0	1.09+0	1.08+0	5.50+0	1.41+0	1.09+0	6.00+0	1.73+0	1.08+0	
1.00+0	1.86-3	<b>4.20-1</b>	1.20+0	5.61-3	5.92-1	1.40+0	1.27-2	7.32-1	154
1.60+0	2.34-2	8.20-1	1.80+0	3.96-2	9.10-1	2.00+0	5.61-2	8.93-1	
2.20+0	8.19-2	9.45-1	2.40+0	1.10-1	9.53-1	2.60+0	1.42-1	9.51-1	
2.60+0	1.55-1	1.04+0	2.80+0	1.79-1	9.48-1	3.00+0	2.43-1	1.04+0	155
3.20+0	3.04-1	1.07+0	3.40+0	3.73-1	1.09+0	3.60+0	4.23-1	1.05+0	
2.70+0	1.35-1	8.02-1	3.00+0	2.13-1	9.12-1	4.00+0	5.02-1	9.20-1	159
4.50+0	7.95-1	1.05+0	4.90+0	1.05+0	1.10+0	5.40+0	1.26+0	1.03+0	
6.00+0	1.84+0	1.15+0	6.50+0	2.14+0	1.10+0				
<b>61</b>	Promethium	Fluorescence yield = 0.925							
7.00+0	1.74+0	8.65-1							125
<b>62</b>	Samarium	Fluorescence yield = 0.929							
1.60+2	4.55+1	1.54+0							30
7.50-1	4.30-4	5.63-1	1.00+0	2.10-3	6.89-1	1.25+0	6.30-3	8.04-1	95
1.50+0	1.40-2	8.77-1	2.00+0	4.30-2	9.54-1	2.50+0	8.90-2	9.35-1	
7.00-1	3.97-4	7.52-1	8.00-1	9.25-4	8.70-1	9.00-1	1.74-3	9.23-1	144
1.00+0	2.96-3	9.71-1	1.10+0	4.69-3	1.02+0	1.20+0	7.06-3	1.06+0	
1.30+0	9.85-3	1.08+0	1.40+0	1.36-2	1.11+0	1.50+0	1.77-2	1.11+0	
1.60+0	2.30-2	1.13+0	1.70+0	2.99-2	1.18+0	1.80+0	3.57-2	1.15+0	
2.00+0	5.08-2	1.13+0	2.20+0	7.12-2	1.14+0	2.40+0	9.57-2	1.15+0	
2.60+0	1.24-1	1.15+0	2.80+0	1.57-1	1.15+0	3.00+0	1.95-1	1.15+0	
3.20+0	2.34-1	1.13+0	3.50+0	3.11-1	1.15+0	4.00+0	4.51-1	1.13+0	
4.50+0	6.27-1	1.12+0	5.00+0	8.22-1	1.10+0	5.50+0	1.05+0	1.10+0	
6.00+0	1.32+0	1.10+0							
8.00-1	5.66-4	5.33-1	1.00+0	2.22-3	7.28-1	1.20+0	5.24-3	7.89-1	154
1.40+0	1.05-2	8.57-1	1.60+0	1.75-2	8.62-1	1.80+0	2.88-2	9.25-1	
2.00+0	4.36-2	9.67-1	2.20+0	6.12-2	9.82-1	2.40+0	8.37-2	1.00+0	
2.60+0	1.10-1	1.02+0							
2.60+0	1.24-1	1.15+0	2.80+0	1.54-1	1.13+0	3.00+0	1.82-1	1.07+0	155
3.20+0	2.39-1	1.15+0	3.40+0	2.81-1	1.13+0	3.60+0	3.45-1	1.17+0	
3.80+0	3.87-1	1.12+0							

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
2.70+0	1.14-1	9.35-1	3.00+0	1.71-1	1.01+0	4.00+0	4.25-1	1.06+0	159
4.50+0	5.68-1	1.02+0	5.00+0	7.48-1	1.01+0	5.50+0	1.01+0	1.06+0	
6.00+0	1.27+0	1.06+0	6.50+0	1.52+0	1.04+0	7.00+0	1.90+0	1.09+0	
<b>63 Europium</b> Fluorescence yield = 0.932									
8.00-1	8.50-4	9.74-1	1.00+0	2.00-3	7.88-1	1.20+0	4.50-3	8.07-1	87
1.40+0	9.10-3	8.80-1	1.60+0	1.40-2	8.15-1	1.80+0	2.30-2	8.70-1	
2.00+0	3.30-2	8.61-1	2.20+0	4.40-2	8.29-1	2.40+0	5.90-2	8.30-1	
2.20+1	1.26+1	9.87-1	3.10+1	1.65+1	8.83-1	4.40+1	2.00+1	8.15-1	104
<b>64 Gadolinium</b> Fluorescence yield = 0.935									
8.00-1	7.10-4	9.89-1	1.00+0	2.20-3	1.04+0	1.20+0	4.90-3	1.04+0	87
1.40+0	8.60-3	9.83-1	1.60+0	1.50-2	1.03+0	1.80+0	2.10-2	9.33-1	
2.00+0	3.30-2	1.01+0	2.20+0	4.40-2	9.70-1	2.40+0	5.60-2	9.22-1	
7.00+0	9.90-1	7.50-1	8.00+0	1.48+0	8.19-1	9.00+0	1.85+0	7.82-1	100
1.00+1	2.48+0	8.37-1	1.10+1	3.06+0	8.47-1	1.20+1	3.75+0	8.75-1	
1.30+1	4.10+0	8.22-1	1.40+1	4.94+0	8.65-1	1.50+1	5.68+0	8.83-1	
4.75+0	2.88-1	5.98-1							107
9.00-1	5.60-4	<b>4.33-1</b>	1.20+0	3.18-3	6.77-1	1.40+0	7.01-3	8.01-1	139
1.60+0	1.31-2	8.97-1	1.80+0	2.05-2	9.11-1	2.00+0	3.15-2	9.64-1	
2.20+0	4.56-2	1.00+0	2.40+0	6.24-2	1.03+0	2.60+0	8.00-2	1.01+0	
<b>65 Terbium</b> Fluorescence yield = 0.938									
1.60+2	2.94+1	1.18+0							30
8.00-1	1.85-4	<b>3.13-1</b>	1.00+0	9.94-4	5.60-1	1.20+0	3.01-3	7.60-1	155
1.40+0	6.47-3	8.71-1	1.60+0	1.17-2	9.40-1	1.80+0	1.74-2	9.06-1	
2.00+0	2.77-2	9.91-1	2.20+0	3.90-2	1.00+0	2.40+0	5.26-2	1.01+0	
2.60+0	7.30-2	1.08+0	2.80+0	9.57-2	1.11+0	3.00+0	1.13-1	1.05+0	
3.20+0	1.43-1	1.09+0	3.40+0	1.60-1	1.01+0	3.60+0	1.96-1	1.05+0	
2.70+0	9.10-2	1.19+0	3.00+0	1.00-1	9.33-1	4.00+0	2.76-1	1.08+0	159
4.50+0	3.51-1	9.79-1	5.00+0	4.77-1	9.93-1	5.50+0	6.67-1	1.07+0	
6.00+0	1.04+0	1.33+0	6.50+0	1.06+0	1.11+0	7.00+0	1.56+0	1.36+0	
<b>67 Holmium</b> Fluorescence yield = 0.944									
8.00-1	3.62-4	8.99-1	9.00-1	7.27-4	9.72-1	1.05+0	1.70-3	1.08+0	91
1.20+0	3.14-3	1.10+0	1.35+0	5.43-3	1.16+0	1.50+0	8.36-3	1.18+0	
1.60+0	1.08-2	1.19+0	1.80+0	1.67-2	1.18+0	2.00+0	2.42-2	1.17+0	
2.10+0	2.79-2	1.14+0	2.20+0	3.35-2	1.16+0	2.40+0	4.32-2	1.12+0	
2.60+0	5.56-2	1.10+0	2.80+0	7.21-2	1.12+0	3.00+0	8.99-2	1.12+0	
3.20+0	1.18-1	1.21+0	3.40+0	1.38-1	1.17+0	3.60+0	1.58-1	1.13+0	
3.80+0	1.85-1	1.12+0	4.00+0	2.06-1	1.07+0				
7.50-1	1.10-4	3.92-1	1.00+0	7.10-4	5.67-1	1.25+0	2.50-3	7.38-1	95
1.50+0	6.00-3	8.46-1	2.00+0	1.90-2	9.19-1	2.50+0	4.20-2	9.48-1	

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	Ref.
7.00+0	1.00+0	1.14+0							125
9.00-1	2.08-4	2.78-1	1.00+0	6.08-4	4.85-1	1.20+0	1.74-3	6.11-1	145
1.20+0	1.89-3	6.63-1	1.40+0	3.90-3	7.22-1	1.60+0	7.45-3	8.18-1	
1.60+0	7.92-3	8.70-1	1.80+0	1.30-2	9.20-1	2.00+0	1.97-2	9.53-1	
2.00+0	2.06-2	9.97-1	2.20+0	2.84-2	9.87-1	2.40+0	3.92-2	1.01+0	
2.40+0	3.95-2	1.02+0	2.60+0	5.32-2	1.05+0	2.60+0	4.95-2	9.81-1	
2.70+0	6.11-2	1.07+0	3.00+0	8.53-2	1.07+0	4.00+0	2.04-1	1.06+0	159
4.50+0	2.84-1	1.05+0	5.00+0	3.87-1	1.07+0	5.50+0	5.14-1	1.09+0	
6.00+0	6.47-1	1.09+0	6.50+0	8.96-1	1.23+0	7.00+0	9.87-1	1.12+0	
7.50+0	1.22+0	1.17+0	8.00+0	1.31+0	1.08+0	8.50+0	1.65+0	1.18+0	
9.00+0	1.96+0	1.23+0	9.50+0	2.01+0	1.12+0	1.00+1	2.47+0	1.22+0	
<b>69 Thulium</b>			<b>Fluorescence yield = 0.949</b>						
3.00+0	5.30-2	8.80-1	5.00+0	2.40-1	8.68-1				59
7.00+0	7.32-1	1.08+0							125
1.00+0	4.63-4	5.20-1	1.20+0	1.46-3	7.07-1	1.40+0	3.11-3	7.84-1	154
1.60+0	5.64-3	8.38-1	1.80+0	9.57-3	9.11-1	2.00+0	1.40-2	9.09-1	
2.20+0	2.27-2	1.05+0	2.40+0	2.85-2	9.82-1	2.60+0	4.58-2	1.21+0	
1.00+0	5.97-4	6.71-1	1.20+0	1.90-3	9.20-1	1.40+0	4.02-3	1.01+0	155
1.60+0	7.21-3	1.07+0	1.80+0	1.21-2	1.15+0	2.00+0	1.74-2	1.13+0	
2.20+0	2.82-2	1.31+0	2.40+0	3.56-2	1.23+0	2.60+0	4.63-2	1.22+0	
2.80+0	5.89-2	1.22+0	3.00+0	8.19-2	1.36+0	3.20+0	1.04-1	1.41+0	
3.40+0	1.31-1	1.47+0	3.60+0	1.43-1	1.35+0				
<b>70 Ytterbium</b>			<b>Fluorescence yield = 0.951</b>						
4.00+0	1.19-1	9.37-1	7.50+0	5.64-1	7.99-1	8.00+0	7.10-1	8.58-1	100
9.00+0	9.06-1	8.27-1	1.00+1	1.10+0	7.92-1	1.10+1	1.39+0	8.08-1	
1.20+1	1.74+0	8.44-1	1.30+1	2.04+0	8.38-1	1.40+1	2.35+0	8.36-1	
1.50+1	2.74+0	8.53-1							
<b>72 Hafnium</b>			<b>Fluorescence yield = 0.955</b>						
7.50-1	5.10-5	4.86-1	1.00+0	3.70-4	6.88-1	1.25+0	1.30-3	8.37-1	95
1.50+0	3.00-3	8.92-1	2.00+0	9.30-3	9.21-1	2.50+0	2.10-2	9.54-1	
<b>73 Tantalum</b>			<b>Fluorescence yield = 0.957</b>						
1.92+0	1.30-2	<b>1.71+0</b>	2.17+0	2.30-2	1.95+0	2.40+0	3.30-2	1.97+0	1
2.64+0	6.00-2	<b>2.59+0</b>	2.88+0	7.70-2	<b>2.50+0</b>	3.15+0	1.10-1	<b>2.68+0</b>	
1.80+0	3.10-3	5.19-1	2.00+0	3.80-3	<b>4.31-1</b>	2.20+0	6.00-3	<b>4.84-1</b>	5
2.30+0	8.00-3	5.52-1	2.40+0	9.70-3	5.78-1	2.60+0	1.10-2	<b>5.00-1</b>	
3.00+0	2.00-2	5.69-1	3.40+0	2.50-2	<b>4.79-1</b>	3.50+0	2.80-2	<b>4.90-1</b>	
3.70+0	3.50-2	<b>5.17-1</b>	3.90+0	4.50-2	5.67-1				

TABLE 2. *K*-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
1.00+0	8.70-4	<b>1.91+0</b>	1.12+0	1.40-3	<b>1.74+0</b>	1.25+0	2.50-3	<b>1.87+0</b>	6
1.60+2	2.28+1	1.41+0							30
7.50-1	3.30-5	3.82-1	1.00+0	3.10-4	6.80-1	1.25+0	1.60-3	1.20+0	95
1.50+0	2.80-3	9.60-1	2.00+0	8.70-3	9.87-1	2.50+0	2.00-2	1.04+0	
7.00+0	3.27-1	8.04-1	8.00+0	5.00-1	8.77-1	9.00+0	6.62-1	8.72-1	100
1.00+1	8.17-1	8.42-1	1.10+1	9.94-1	8.29-1	1.20+1	1.41+0	9.69-1	
1.30+1	1.53+0	8.91-1	1.40+1	1.63+0	8.15-1	1.50+1	1.87+0	8.17-1	
7.00+0	4.57-1	1.12+0							125
1.10+0	4.40-4	5.98-1	1.30+0	1.01-3	6.35-1	1.50+0	2.24-3	7.68-1	158
1.70+0	4.28-3	8.93-1	1.90+0	6.84-3	9.35-1	2.10+0	1.08-2	1.03+0	
2.25+0	1.40-2	1.04+0	2.45+0	1.98-2	1.10+0	2.65+0	2.54-2	1.08+0	
2.80+0	3.47-2	1.23+0	3.00+0	4.14-2	1.18+0	3.20+0	5.13-2	1.19+0	
3.40+0	6.52-2	1.25+0	3.60+0	7.50-2	1.20+0	3.80+0	8.50-2	1.16+0	
2.70+0	2.80-2	1.12+0	3.00+0	3.92-2	1.11+0	3.30+0	5.24-2	1.10+0	159
3.60+0	7.11-2	1.14+0	4.00+0	9.76-2	1.14+0	4.50+0	1.38-1	1.14+0	
5.00+0	1.96-1	1.20+0	5.50+0	2.72-1	1.27+0	6.00+0	3.18-1	1.17+0	
6.50+0	3.81-1	1.14+0	7.00+0	4.93-1	1.21+0	7.50+0	5.98-1	1.23+0	
8.00+0	6.43-1	1.13+0	8.50+0	8.13-1	1.23+0				
<b>74 Tungsten                  Fluorescence yield = 0.958</b>									
4.75+0	1.05-1	8.41-1							107
1.20+0	5.84-4	6.13-1	1.39+0	1.64-3	8.92-1	1.60+0	2.68-3	8.15-1	139
2.00+0	6.60-3	8.56-1	2.22+0	1.10-2	9.81-1	2.40+0	1.59-2	1.08+0	
2.60+0	1.91-2	9.90-1	2.80+0	2.44-2	9.88-1	3.20+0	3.89-2	1.02+0	
3.80+0	6.59-2	1.02+0	4.50+0	1.08-1	1.01+0	5.20+0	1.62-1	1.00+0	
6.00+0	2.14-1	8.95-1	6.90+0	3.03-1	8.75-1	8.00+0	4.32-1	8.56-1	
6.50-1	9.01-6	3.54-1	7.20-1	2.51-5	4.68-1	8.00-1	6.10-5	5.68-1	142
9.00-1	1.44-4	6.63-1	1.00+0	2.79-4	7.21-1	1.25+0	8.99-4	7.82-1	
1.50+0	2.01-3	7.94-1	1.75+0	3.80-3	8.13-1	2.00+0	6.57-3	8.52-1	
2.25+0	1.02-2	8.67-1	2.50+0	1.56-2	9.23-1	2.75+0	2.25-2	9.67-1	
3.00+0	3.18-2	1.03+0	3.25+0	4.27-2	1.07+0	3.50+0	5.29-2	1.05+0	
3.75+0	7.35-2	1.18+0							
<b>75 Rhenium                  Fluorescence yield = 0.959</b>									
7.50-1	2.20-5	3.80-1	1.00+0	1.60-4	4.88-1	1.25+0	7.40-4	7.46-1	95
1.50+0	2.10-3	9.54-1	2.00+0	7.10-3	1.05+0	2.50+0	9.60-3	6.46-1	
<b>78 Platinum                  Fluorescence yield = 0.963</b>									
1.60+2	1.82+1	1.48+0							30

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
7.50-1	1.20-5	3.79-1	1.00+0	1.30-4	6.46-1	1.25+0	5.10-4	7.95-1	95
1.50+0	1.30-3	8.88-1	2.00+0	4.00-3	8.67-1	2.50+0	9.20-3	8.97-1	
7.00+0	2.72-1	1.22+0							125
<b>79 Gold</b>									
									Fluorescence yield = 0.964
2.40+0	1.60-2	2.03+0							1
1.60+2	1.68+1	1.44+0							30
3.00+0	2.00-2	1.19+0	4.00+0	4.20-2	1.02+0	6.00+0	1.30-1	9.86-1	41
7.00+0	2.40-1	1.20+0	8.00+0	2.60-1	9.24-1	9.00+0	4.50-1	1.19+0	
1.00+1	5.80-1	1.19+0	1.10+1	6.50-1	1.07+0	1.20+1	8.20-1	1.11+0	
1.30+1	1.06+0	1.20+0	1.40+1	1.20+0	1.16+0	1.50+1	1.40+0	1.17+0	
1.00+0	9.47-5	5.54-1	1.10+0	1.88-4	6.49-1	1.20+0	3.07-4	6.75-1	91
1.30+0	4.96-4	7.40-1	1.40+0	7.61-4	8.06-1	1.50+0	1.03-3	8.04-1	
1.60+0	1.41-3	8.36-1	1.70+0	1.96-3	9.06-1	1.80+0	2.50-3	9.21-1	
1.90+0	3.16-3	9.43-1	2.00+0	4.04-3	9.91-1	2.10+0	4.95-3	1.01+0	
2.20+0	5.75-3	9.93-1	2.40+0	8.33-3	1.06+0	2.60+0	1.12-2	1.08+0	
2.80+0	1.47-2	1.10+0	3.00+0	1.89-2	1.13+0	3.20+0	2.36-2	1.15+0	
3.40+0	2.96-2	1.18+0	3.60+0	3.70-2	1.24+0	3.80+0	4.35-2	1.23+0	
4.00+0	4.83-2	1.17+0							
1.00+0	9.60-5	5.61-1	1.30+0	3.80-4	5.67-1	1.50+0	9.00-4	7.03-1	93
1.75+0	1.80-3	7.41-1	2.00+0	3.00-3	7.36-1	2.25+0	5.50-3	8.77-1	
2.50+0	7.50-3	8.25-1	2.75+0	1.00-2	7.96-1	2.80+0	1.30-2	9.75-1	
3.00+0	1.90-2	1.13+0	3.50+0	2.50-2	9.13-1	4.00+0	4.00-2	9.70-1	
5.00+0	7.00-2	8.83-1	6.00+0	1.10-1	8.34-1	7.00+0	1.80-1	9.03-1	
8.00+0	2.60-1	9.24-1							
7.50-1	7.70-6	2.99-1	1.00+0	9.40-5	5.50-1	1.25+0	3.90-4	7.02-1	95
1.50+0	9.60-4	7.49-1	2.00+0	3.20-3	7.85-1	2.50+0	7.50-3	8.25-1	
7.50+0	2.25-1	9.43-1	8.00+0	2.46-1	8.74-1	9.00+0	3.23-1	8.55-1	100
1.00+1	4.12-1	8.46-1	1.10+1	5.50-1	9.04-1	1.20+1	6.75-1	9.13-1	
1.30+1	8.03-1	9.12-1	1.40+1	9.41-1	9.09-1	1.50+1	1.16+0	9.72-1	
2.20+1	2.66+0	1.08+0	3.10+1	3.88+0	9.31-1	4.40+1	5.93+0	9.31-1	104
7.00+0	2.12-1	1.06+0							125
7.10-1	4.32-6	<b>2.57-1</b>	7.60-1	8.77-6	3.08-1	8.20-1	1.75-5	3.57-1	142
9.00-1	3.89-5	4.29-1	1.00+0	8.94-5	5.23-1	1.25+0	3.86-4	6.95-1	
1.50+0	1.05-3	8.20-1	1.75+0	2.21-3	9.10-1	2.00+0	3.95-3	9.69-1	
2.25+0	6.30-3	1.00+0	2.50+0	9.24-3	1.02+0	3.00+0	1.60-2	9.55-1	
3.50+0	2.70-2	9.86-1							
1.60+0	1.13-3	6.70-1	2.00+0	3.74-3	9.18-1	2.20+0	5.49-3	9.49-1	145
2.40+0	7.13-3	9.04-1	2.60+0	1.12-2	1.08+0	2.60+0	1.36-2	1.31+0	

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
<b>82 Lead</b>					<b>Fluorescence yield = 0.967</b>				
1.92+0	3.60-3	1.49+0	2.17+0		5.90-3	1.53+0	2.40+0		1.05-2 1.89+0
2.88+0	3.05-2	<b>2.94+0</b>							1
1.60+0	1.10-3	9.58-1	1.70+0	1.50-3	1.01+0	1.80+0	2.00-3	1.07+0	5
1.90+0	2.20-3	9.46-1	2.00+0	2.70-3	9.52-1	2.10+0	3.60-3	1.05+0	
2.20+0	4.30-3	1.06+0	2.30+0	4.50-3	9.43-1	2.40+0	5.10-3	9.18-1	
2.50+0	7.90-3	1.23+0	2.60+0	8.40-3	1.14+0	2.70+0	9.00-3	1.08+0	
2.70+0	9.90-3	1.18+0	2.90+0	1.10-2	1.04+0	3.00+0	1.30-2	1.09+0	
3.20+0	1.40-2	9.55-1	3.20+0	1.50-2	1.02+0	3.30+0	1.60-2	9.90-1	
3.40+0	1.70-2	9.57-1	3.60+0	1.80-2	8.47-1				
1.60+2	1.84+1	1.87+0							30
3.00+0	1.28-2	1.08+0	5.00+0	5.30-2	9.37-1				59
7.50-1	2.60-6	<b>1.90-1</b>	1.00+0	6.10-5	5.83-1	1.25+0	2.40-4	6.62-1	95
1.50+0	6.60-4	7.63-1	2.00+0	2.30-3	8.11-1	2.50+0	5.10-3	7.95-1	
4.00+0	4.07-2	1.39+0	7.50+0	1.57-1	9.18-1	8.00+0	1.86-1	9.22-1	100
9.00+0	2.41-1	8.87-1	1.00+1	2.98-1	8.49-1	1.10+1	3.99-1	9.07-1	
1.20+1	4.78-1	8.90-1	1.30+1	5.68-1	8.86-1	1.40+1	6.75-1	8.97-1	
1.50+1	8.00-1	9.16-1							
1.00+0	4.18-5	4.00-1	1.25+0	2.44-4	6.73-1	1.50+0	6.38-4	7.38-1	113
1.75+0	1.22-3	7.29-1	2.00+0	2.45-3	8.63-1	2.25+0	3.69-3	8.37-1	
3.00+0	9.87-3	8.31-1	4.00+0	2.61-2	8.89-1	5.00+0	5.37-2	9.49-1	
6.00+0	9.13-2	9.69-1	7.00+0	1.35-1	9.45-1	8.00+0	1.85-1	9.17-1	
7.00+0	1.78-1	1.25+0							125
<b>83 Bismuth</b>					<b>Fluorescence yield = 0.968</b>				
7.50-1	1.80-6	<b>1.63-1</b>	1.00+0	5.40-5	6.09-1	1.25+0	2.40-4	7.63-1	95
1.50+0	6.00-4	7.89-1	2.00+0	1.90-3	7.53-1	2.50+0	4.90-3	8.55-1	
7.00+0	1.50-1	1.17+0							125
<b>90 Thorium</b>					<b>Fluorescence yield = 0.971</b>				
7.00+0	7.02-2	1.12+0	8.00+0	1.06-1	1.19+0	9.00+0	1.28-1	1.07+0	100
1.00+1	1.61-1	1.03+0	1.10+1	2.02-1	1.03+0	1.20+1	2.38-1	9.91-1	
1.30+1	2.75-1	9.53-1	1.40+1	2.95-1	8.65-1	1.50+1	3.72-1	9.37-1	
4.75+0	3.02-2	1.41+0	6.80+0	5.75-2	9.88-1	7.20+0	7.69-2	1.14+0	107
8.80+0	1.07-1	9.43-1	1.01+1	1.95-1	1.22+0				
1.20+0	2.00-5	<b>2.21-1</b>	1.60+0	2.21-4	5.17-1	1.83+0	4.45-4	5.65-1	124
2.00+0	8.18-4	7.14-1	2.30+0	1.52-3	7.63-1	2.40+0	2.10-3	8.98-1	
2.80+0	3.60-3	8.87-1	2.81+0	3.64-3	8.86-1	3.30+0	7.38-3	1.05+0	
3.79+0	1.17-2	1.07+0	4.80+0	2.72-2	1.23+0	6.00+0	4.91-2	1.18+0	

TABLE 2. K-shell x-ray production by protons in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
<b>92 Uranium</b> <b>Fluorescence yield = 0.972</b>									
2.25+0	1.50-3	1.01+0	3.75+0	9.20-3	1.05+0				2
2.50+0	5.80-4	<b>2.61-1</b>	2.70+0	8.40-4	<b>2.86-1</b>	2.90+0	1.00-3	<b>2.65-1</b>	5
3.10+0	1.20-3	<b>2.54-1</b>	3.30+0	1.50-3	<b>2.58-1</b>	3.60+0	2.00-3	<b>2.60-1</b>	
3.90+0	2.70-3	<b>2.74-1</b>	4.20+0	3.50-3	<b>2.83-1</b>	4.40+0	4.80-3	<b>3.38-1</b>	
4.60+0	5.80-3	<b>3.58-1</b>	5.00+0	7.30-3	<b>3.54-1</b>	5.60+0	9.50-3	<b>3.34-1</b>	
1.60+2	1.00+1	1.75+0							30
1.40+0	1.05-4	6.29-1	1.50+0	1.53-4	6.34-1	1.60+0	2.42-4	7.23-1	91
1.80+0	4.92-4	8.43-1	2.00+0	9.05-4	9.81-1	2.20+0	1.41-3	1.04+0	
2.40+0	2.04-3	1.07+0	2.60+0	2.82-3	1.10+0	2.80+0	4.28-3	1.28+0	
3.00+0	5.29-3	1.25+0	3.20+0	6.70-3	1.28+0	3.40+0	7.93-3	1.24+0	
3.60+0	8.50-3	1.11+0	3.80+0	1.13-2	1.24+0	4.00+0	1.41-2	1.32+0	
1.25+0	3.60-5	4.17-1	1.50+0	1.30-4	5.38-1	2.00+0	5.90-4	6.40-1	95
2.50+0	2.10-3	9.46-1							
4.75+0	1.46-2	8.21-1							107
1.10+0	1.67-5	4.46-1	1.20+0	3.12-5	4.67-1	1.30+0	5.35-5	4.89-1	142
1.40+0	8.59-5	5.15-1	1.60+0	1.93-4	5.76-1	1.80+0	3.80-4	6.51-1	
2.00+0	6.81-4	7.38-1	2.25+0	1.29-3	8.68-1	2.50+0	2.23-3	1.00+0	

<sup>a</sup>

Cross sections and their ratios are printed in a compressed power of 10  
 $-1$   
 notation, e.g. 4.76-1 means  $4.76 \times 10^{-1}$ .

<sup>b</sup>

The ratios shown in bold print differ by more than a factor of 2 from  
 the averaged ratios and were -- as described in the text -- rejected.  
 This rejection criterion was applied only to the  $Z_2 > 9$  data.

TABLE 3. K-shell x-ray production by deuterons in target elements from beryllium to gold<sup>a,b</sup>

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
<b>4 Beryllium      Fluorescence yield = 0.00033</b>									
3.00-2	5.66+0	2.91-1	4.00-2	2.29+1	3.46-1	5.00-2	6.46+1	4.21-1	119
6.00-2	1.35+2	4.82-1	8.00-2	3.39+2	5.52-1	1.00-1	5.90+2	5.97-1	
1.20-1	8.95+2	6.66-1							
<b>11 Sodium      Fluorescence yield = 0.023</b>									
4.00-2	1.80-2	1.61+0	5.00-2	5.00-2	1.47+0	6.50-2	2.00-1	1.75+0	102
<b>13 Aluminum      Fluorescence yield = 0.039</b>									
1.80-1	1.18+0	7.46-1	2.00-1	1.68+0	7.16-1	2.80-1	5.06+0	6.53-1	25
3.60-1	1.12+1	6.33-1							
4.00-2	1.68-3	1.11+0	4.50-2	3.25-3	1.07+0	5.00-2	5.71-3	1.05+0	45
6.00-2	1.44-2	1.03+0	7.00-2	3.00-2	1.01+0	8.00-2	5.46-2	9.85-1	
9.00-2	9.00-2	9.57-1	1.00-1	1.39-1	9.35-1	1.10-1	2.02-1	9.06-1	
1.20-1	2.80-1	8.76-1	1.30-1	3.81-1	8.59-1	1.40-1	4.98-1	8.34-1	
1.60-1	8.05-1	7.98-1	1.80-1	1.22+0	7.71-1	2.00-1	1.79+0	7.63-1	
2.20-1	2.54+0	7.64-1	2.40-1	3.50+0	7.72-1	2.60-1	4.79+0	7.98-1	
2.80-1	6.42+0	8.29-1							
2.00+0	5.04+2	7.56-1	3.00+0	7.59+2	8.11-1	4.00+0	8.89+2	8.22-1	97
5.00+0	9.46+2	8.19-1	6.00+0	9.79+2	8.25-1	8.00+0	9.51+2	8.05-1	
1.00+1	9.07+2	7.97-1	1.20+1	8.46+2	7.81-1	1.40+1	8.08+2	7.87-1	
1.50+1	7.75+2	7.76-1	1.80+1	7.06+2	7.67-1	1.95+1	6.83+2	7.72-1	
2.40-2	7.00-6	1.84-1	2.80-2	2.50-5	1.87-1	3.00-2	7.00-5	3.12-1	102
3.20-2	1.30-4	3.65-1	3.40-2	3.00-4	5.57-1	3.80-2	7.00-4	6.33-1	
4.00-2	1.40-3	9.23-1	5.00-2	3.50-3	6.42-1	5.50-2	7.00-3	7.73-1	
6.00-2	1.20-2	8.55-1							
<b>14 Silicon      Fluorescence yield = 0.05</b>									
5.00-2	1.20-3	5.29-1	6.00-2	4.60-3	7.35-1				102
<b>15 Phosphorus      Fluorescence yield = 0.063</b>									
1.60+0	2.32+2	8.46-1	1.80+0	2.83+2	8.54-1	2.00+0	3.22+2	8.33-1	122
2.20+0	3.68+2	8.37-1	2.40+0	4.29+2	8.76-1	2.60+0	4.77+2	8.87-1	
2.80+0	5.20+2	8.95-1	3.00+0	5.46+2	8.78-1	3.20+0	6.13+2	9.29-1	
3.40+0	6.31+2	9.08-1	3.60+0	6.70+2	9.22-1	3.80+0	6.77+2	8.95-1	
4.00+0	7.04+2	8.98-1							

TABLE 3. K-shell x-ray production by deuterons in target elements from beryllium to gold<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
<b>16 Sulfur</b> <b>Fluorescence yield = 0.078</b>									
1.60+0	1.64+2	8.19-1	1.80+0	2.15+2	8.70-1	2.00+0	2.68+2	9.12-1	122
2.20+0	2.84+2	8.37-1	2.40+0	3.25+2	8.49-1	2.60+0	3.59+2	8.45-1	
2.80+0	4.04+2	8.69-1	3.00+0	4.29+2	8.53-1	3.20+0	4.62+2	8.58-1	
3.40+0	5.10+2	8.92-1	3.60+0	5.34+2	8.85-1	3.80+0	5.55+2	8.78-1	
4.00+0	5.59+2	8.47-1							
<b>17 Chlorine</b> <b>Fluorescence yield = 0.097</b>									
1.60+0	1.19+2	7.95-1	1.80+0	1.51+2	8.06-1	2.00+0	1.81+2	7.99-1	122
2.20+0	2.18+2	8.21-1	2.40+0	2.28+2	7.51-1	2.60+0	2.72+2	7.97-1	
2.80+0	2.99+2	7.92-1	3.00+0	3.31+2	8.03-1	3.20+0	3.41+2	7.66-1	
3.40+0	3.96+2	8.30-1	3.60+0	4.34+2	8.56-1	3.80+0	4.39+2	8.19-1	
4.00+0	4.57+2	8.13-1							
<b>19 Potassium</b> <b>Fluorescence yield = 0.14</b>									
1.60+0	6.13+1	7.69-1	1.80+0	8.05+1	7.79-1	2.00+0	1.01+2	7.86-1	122
2.20+0	1.23+2	7.96-1	2.40+0	1.32+2	7.29-1	2.60+0	1.62+2	7.78-1	
2.80+0	1.80+2	7.66-1	3.00+0	2.04+2	7.79-1	3.20+0	2.13+2	7.40-1	
3.40+0	2.47+2	7.88-1	3.60+0	2.74+2	8.10-1	3.80+0	2.82+2	7.78-1	
4.00+0	3.01+2	7.80-1							
<b>20 Calcium</b> <b>Fluorescence yield = 0.163</b>									
4.00-1	6.93-1	8.76-1	5.00-1	1.52+0	8.58-1	6.00-1	2.80+0	8.43-1	133
7.00-1	4.73+0	8.56-1	8.00-1	8.15+0	9.65-1	9.00-1	1.05+1	8.67-1	
<b>21 Scandium</b> <b>Fluorescence yield = 0.188</b>									
4.00-1	4.73-1	9.24-1	6.00-1	1.98+0	8.95-1	8.00-1	5.19+0	9.03-1	121
1.00+0	1.04+1	9.07-1	1.20+0	1.78+1	9.15-1	1.40+0	2.62+1	8.84-1	
1.60+0	3.74+1	8.97-1	1.80+0	5.07+1	9.13-1				
4.00-1	4.45-1	8.69-1	5.00-1	1.01+0	8.68-1	6.00-1	1.92+0	8.68-1	133
7.00-1	3.25+0	8.74-1	8.00-1	5.19+0	9.03-1	9.00-1	7.33+0	8.81-1	
<b>22 Titanium</b> <b>Fluorescence yield = 0.214</b>									
1.80-1	1.26-2	1.12+0	2.20-1	3.28-2	1.15+0	2.60-1	7.05-2	1.19+0	25
3.00-1	1.29-1	1.19+0	3.40-1	2.23-1	1.25+0				
6.00+0	3.70+2	1.07+0	1.25+1	6.80+2	1.15+0	1.50+1	7.80+2	1.25+0	31
1.00+0	7.68+0	9.55-1	2.00+0	5.19+1	9.81-1	2.50+0	8.85+1	1.02+0	131
4.00-1	2.98-1	8.81-1	5.00-1	6.96-1	8.94-1	6.00-1	1.35+0	9.02-1	133
7.00-1	2.19+0	8.61-1	8.00-1	3.40+0	8.57-1	9.00-1	5.02+0	8.67-1	
2.00-1	2.42-2	1.31+0	2.50-1	6.79-2	1.35+0	3.00-1	1.40-1	1.30+0	153
3.50-1	2.52-1	1.26+0	4.00-1	4.17-1	1.23+0	4.50-1	6.50-1	1.23+0	

TABLE 3. K-shell x-ray production by deuterons in target elements from beryllium to gold<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
5.00-1	9.63-1	1.24+0	5.50-1	1.36+0	1.24+0	6.00-1	1.86+0	1.24+0	
6.50-1	2.46+0	1.24+0	7.00-1	3.16+0	1.24+0	7.50-1	3.98+0	1.24+0	
8.00-1	4.91+0	1.24+0	8.50-1	5.95+0	1.23+0	9.00-1	7.11+0	1.23+0	
1.00+0	9.80+0	1.22+0	1.10+0	1.30+1	1.21+0	1.20+0	1.67+1	1.21+0	
1.30+0	2.09+1	1.20+0	1.40+0	2.56+1	1.20+0	1.50+0	3.08+1	1.19+0	
1.60+0	3.65+1	1.20+0	1.80+0	4.94+1	1.20+0	2.00+0	6.42+1	1.21+0	
<b>23</b>	<b>Vanadium</b>		<b>Fluorescence yield = 0.243</b>						
1.00+0	5.44+0	9.52-1	2.00+0	4.07+1	1.02+0	2.50+0	6.91+1	1.03+0	131
3.00+0	1.01+2	1.03+0							
4.00-1	2.23-1	9.85-1	5.00-1	5.16-1	9.73-1	6.00-1	9.97-1	9.69-1	133
7.00-1	1.74+0	9.85-1	8.00-1	2.73+0	9.83-1	9.00-1	3.77+0	9.22-1	
<b>24</b>	<b>Chromium</b>		<b>Fluorescence yield = 0.275</b>						
4.00-1	1.53-1	9.92-1	5.00-1	3.54-1	9.69-1	6.00-1	6.79-1	9.47-1	133
7.00-1	1.20+0	9.66-1	8.00-1	1.84+0	9.35-1	9.00-1	2.63+0	9.01-1	
<b>25</b>	<b>Manganese</b>		<b>Fluorescence yield = 0.308</b>						
4.00-1	1.05-1	9.96-1	5.00-1	2.49-1	9.82-1	6.00-1	4.91-1	9.76-1	133
7.00-1	8.61-1	9.80-1	8.00-1	1.34+0	9.56-1	9.00-1	2.00+0	9.56-1	
<b>26</b>	<b>Iron</b>		<b>Fluorescence yield = 0.34</b>						
4.00-1	8.86-2	1.22+0	5.00-1	2.03-1	1.15+0	6.00-1	3.97-1	1.12+0	121
8.00-1	1.14+0	1.14+0	1.00+0	2.38+0	1.11+0	1.20+0	4.14+0	1.07+0	
1.40+0	6.67+0	1.07+0	1.60+0	9.85+0	1.06+0	1.80+0	1.33+1	1.02+0	
4.00-1	7.49-2	1.04+0	5.00-1	1.87-1	1.06+0	6.00-1	3.80-1	1.08+0	133
7.00-1	6.53-1	1.05+0	8.00-1	1.01+0	1.01+0	9.00-1	1.50+0	9.98-1	
2.50-1	1.19-2	1.31+0	3.00-1	2.83-2	1.34+0	3.50-1	5.42-2	1.31+0	153
4.00-1	9.27-2	1.28+0	4.50-1	1.47-1	1.26+0	5.00-1	2.21-1	1.25+0	
5.50-1	3.14-1	1.23+0	6.00-1	4.39-1	1.24+0	6.50-1	5.88-1	1.24+0	
7.00-1	7.66-1	1.23+0	7.50-1	9.77-1	1.23+0	8.00-1	1.22+0	1.22+0	
8.50-1	1.50+0	1.21+0	9.00-1	1.81+0	1.20+0	1.00+0	2.55+0	1.19+0	
1.10+0	3.46+0	1.18+0	1.20+0	4.52+0	1.17+0	1.30+0	5.77+0	1.16+0	
1.40+0	7.19+0	1.15+0	1.50+0	8.81+0	1.15+0	1.60+0	1.06+1	1.14+0	
1.80+0	1.49+1	1.15+0	2.00+0	2.00+1	1.15+0				
<b>27</b>	<b>Cobalt</b>		<b>Fluorescence yield = 0.373</b>						
1.00+0	1.52+0	9.74-1	2.00+0	1.34+1	1.02+0	2.50+0	2.34+1	9.85-1	131
3.00+0	3.71+1	1.00+0							
4.00-1	4.83-2	9.63-1	5.00-1	1.22-1	9.83-1	6.00-1	2.48-1	9.88-1	133
7.00-1	4.43-1	9.94-1	8.00-1	7.03-1	9.75-1	9.00-1	1.01+0	9.27-1	

TABLE 3. K-shell x-ray production by deuterons in target elements from beryllium to gold<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	Ref.
<b>28 Nickel</b>					Fluorescence yield = 0.406				
2.00+0	8.37+0	8.37-1	3.00+0	2.55+1	8.82-1	4.00+0	4.91+1	8.87-1	97
5.00+0	7.70+1	8.96-1	6.00+0	1.07+2	9.07-1	8.00+0	1.64+2	9.10-1	
1.00+1	2.16+2	9.17-1	1.20+1	2.58+2	9.14-1	1.40+1	2.96+2	9.24-1	
1.50+1	3.08+2	9.15-1	1.80+1	3.37+2	8.97-1	1.95+1	3.53+2	9.05-1	
1.00-1	4.20-6	4.77-1	1.10-1	8.70-6	4.23-1	1.20-1	1.90-5	4.52-1	120
1.30-1	4.30-5	5.55-1	1.40-1	7.50-5	5.67-1	1.50-1	1.30-4	6.13-1	
1.60-1	2.00-4	6.18-1	1.80-1	4.40-4	6.58-1	2.00-1	9.30-4	7.57-1	
2.20-1	1.60-3	7.73-1	2.40-1	2.60-3	7.96-1	2.60-1	4.00-3	8.18-1	
2.80-1	6.10-3	8.76-1	3.00-1	8.30-3	8.60-1				
1.60+0	5.13+0	9.86-1	1.80+0	6.77+0	9.17-1	2.00+0	9.65+0	9.65-1	122
2.20+0	1.26+1	9.67-1	2.40+0	1.59+1	9.65-1	2.60+0	1.94+1	9.56-1	
2.80+0	2.38+1	9.76-1	3.00+0	2.78+1	9.62-1	3.20+0	3.19+1	9.46-1	
3.40+0	3.64+1	9.39-1	3.60+0	4.19+1	9.48-1	3.80+0	4.80+1	9.68-1	
4.00+0	5.26+1	9.50-1							
4.00-1	3.27-2	9.40-1	5.00-1	8.32-2	9.52-1	6.00-1	1.74-1	9.75-1	133
7.00-1	3.16-1	9.89-1	8.00-1	5.22-1	1.00+0	9.00-1	7.99-1	1.01+0	
<b>29 Copper</b>					Fluorescence yield = 0.44				
6.00-1	8.59-2	6.68-1	7.00-1	1.70-1	7.33-1	8.00-1	3.46-1	9.09-1	4
9.00-1	4.73-1	8.14-1	1.00+0	7.06-1	8.40-1				
1.60+0	3.63+0	9.25-1	1.80+0	5.42+0	9.66-1	2.00+0	7.69+0	1.01+0	122
2.20+0	9.39+0	9.37-1	2.40+0	1.18+1	9.26-1	2.60+0	1.44+1	9.13-1	
2.80+0	1.79+1	9.36-1	3.00+0	2.07+1	9.13-1	3.20+0	2.43+1	9.15-1	
3.40+0	2.86+1	9.30-1	3.60+0	3.30+1	9.42-1	3.80+0	3.69+1	9.31-1	
4.00+0	3.98+1	8.97-1							
1.60-1	6.90-5	3.50-1	1.80-1	2.25-4	5.35-1	2.00-1	6.09-4	7.70-1	126
2.40-1	2.03-3	9.36-1	2.80-1	4.59-3	9.63-1	3.20-1	9.18-3	1.02+0	
3.60-1	1.63-2	1.06+0	4.00-1	2.58-2	1.06+0	5.00-1	6.59-2	1.06+0	
6.00-1	1.45-1	1.13+0							
1.00+0	7.73-1	9.20-1	2.00+0	7.69+0	1.01+0	2.50+0	1.44+1	1.01+0	131
3.00+0	2.33+1	1.03+0							
4.00-1	2.40-2	9.82-1	5.00-1	6.33-2	1.02+0	6.00-1	1.26-1	9.79-1	133
7.00-1	2.33-1	1.00+0	8.00-1	3.83-1	1.01+0	9.00-1	5.80-1	9.99-1	
<b>31 Gallium</b>					Fluorescence yield = 0.507				
1.60+0	2.14+0	9.59-1	1.80+0	3.10+0	9.60-1	2.00+0	4.16+0	9.36-1	122
2.20+0	5.46+0	9.27-1	2.40+0	7.15+0	9.46-1	2.60+0	9.00+0	9.52-1	
2.80+0	1.11+1	9.60-1	3.00+0	1.30+1	9.37-1	3.20+0	1.58+1	9.63-1	
3.40+0	1.78+1	9.35-1	3.60+0	2.08+1	9.50-1	3.80+0	2.27+1	9.08-1	
4.00+0	2.66+1	9.45-1							

TABLE 3. K-shell x-ray production by deuterons in target elements from beryllium to gold<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
<b>32 Germanium      Fluorescence yield = 0.535</b>									
1.00+0	3.16-1	9.38-1	2.00+0	3.45+0	1.02+0	2.50+0	6.78+0	1.04+0	131
3.00+0	1.08+1	1.00+0							
<b>37 Rubidium      Fluorescence yield = 0.667</b>									
8.00-1	1.89-2	5.69-1	1.00+0	5.13-2	6.50-1	1.20+0	1.06-1	6.83-1	111
1.40+0	1.92-1	7.13-1	1.60+0	3.27-1	7.65-1	1.80+0	4.61-1	7.25-1	
2.00+0	7.15-1	7.96-1	2.20+0	8.88-1	7.28-1	2.40+0	1.27+0	7.92-1	
2.60+0	1.62+0	7.90-1							
<b>38 Strontium      Fluorescence yield = 0.69</b>									
7.00-1	1.04-2	7.21-1	8.00-1	1.91-2	7.65-1	1.00+0	4.78-2	7.97-1	111
1.20+0	1.00-1	8.43-1	1.40+0	1.93-1	9.33-1	1.60+0	3.00-1	9.10-1	
1.80+0	4.38-1	8.90-1	2.00+0	6.23-1	8.93-1	2.20+0	7.95-1	8.37-1	
2.40+0	1.11+0	8.87-1	2.60+0	1.55+0	9.66-1				
<b>40 Zirconium      Fluorescence yield = 0.73</b>									
7.00-1	4.96-3	6.09-1	8.00-1	9.64-3	6.74-1	1.00+0	2.63-2	7.53-1	111
1.20+0	5.07-2	7.26-1	1.40+0	1.01-1	8.22-1	1.60+0	1.59-1	8.05-1	
1.80+0	2.34-1	7.88-1	2.00+0	3.60-1	8.49-1	2.40+0	5.91-1	7.69-1	
2.60+0	6.81-1	6.88-1							
<b>41 Niobium      Fluorescence yield = 0.74</b>									
3.20-1	5.09-5	4.92-1	4.00-1	4.63-4	1.17+0	4.80-1	1.35-3	1.28+0	126
5.60-1	3.02-3	1.35+0	6.40-1	5.60-3	1.35+0	7.20-1	1.03-2	1.48+0	
8.00-1	1.52-2	1.40+0	9.00-1	2.35-2	1.33+0				
<b>47 Silver      Fluorescence yield = 0.831</b>									
5.00-1	1.06-4	4.74-1	6.00-1	4.23-4	7.09-1	7.00-1	1.07-3	8.44-1	121
8.00-1	2.02-3	8.62-1	1.00+0	5.83-3	9.52-1	1.20+0	1.22-2	9.55-1	
1.40+0	2.18-2	9.43-1	1.60+0	3.63-2	9.57-1	1.80+0	5.54-2	9.52-1	
4.00-1	3.06-5	5.35-1	5.00-1	1.82-4	8.13-1	6.00-1	5.26-4	8.81-1	126
7.00-1	1.33-3	1.05+0	8.00-1	2.59-3	1.11+0	9.00-1	4.66-3	1.19+0	
1.00+0	6.64-3	1.08+0	1.10+0	1.09-2	1.21+0	1.20+0	1.64-2	1.28+0	
<b>48 Cadmium      Fluorescence yield = 0.843</b>									
9.00-1	1.01-3	<b>3.25-1</b>	1.00+0	2.35-3	<b>4.83-1</b>	1.20+0	6.89-3	6.75-1	111
1.40+0	1.05-2	5.66-1	1.60+0	2.41-2	7.89-1	1.80+0	3.09-2	6.59-1	
2.00+0	4.43-2	6.51-1	2.20+0	5.48-2	5.77-1	2.40+0	7.90-2	6.19-1	
2.60+0	1.00-1	5.99-1							

TABLE 3. K-shell x-ray production by deuterons in target elements from beryllium to gold<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
<b>49 Indium</b> <b>Fluorescence yield = 0.853</b>									
6.00-1	2.02-4	5.72-1	7.00-1	4.98-4	6.43-1	8.00-1	1.10-3	7.59-1	129
1.00+0	3.46-3	8.95-1	1.20+0	7.38-3	9.03-1	1.40+0	1.33-2	8.92-1	
1.60+0	2.30-2	9.33-1	1.80+0	3.61-2	9.54-1				
<b>50 Tin</b> <b>Fluorescence yield = 0.862</b>									
1.20+0	1.94-3	<b>2.95-1</b>	1.40+0	5.03-3	<b>4.18-1</b>	1.60+0	9.54-3	<b>4.78-1</b>	111
1.80+0	1.38-2	<b>4.50-1</b>	2.00+0	1.78-2	<b>3.97-1</b>	2.20+0	2.65-2	<b>4.24-1</b>	
2.40+0	3.39-2	<b>4.01-1</b>	2.60+0	4.00-2	<b>3.61-1</b>				
<b>51 Antimony</b> <b>Fluorescence yield = 0.87</b>									
5.00-1	4.54-5	6.29-1	6.00-1	1.71-4	8.13-1	7.00-1	4.27-4	8.99-1	126
8.00-1	8.40-4	9.26-1	9.00-1	1.63-3	1.05+0	1.00+0	2.82-3	1.14+0	
1.10+0	4.49-3	1.21+0	1.20+0	6.73-3	1.27+0	1.30+0	8.50-3	1.16+0	
7.00-1	3.12-4	6.57-1	8.00-1	6.74-4	7.43-1	1.00+0	1.83-3	7.39-1	129
1.20+0	4.41-3	8.32-1	1.40+0	8.27-3	8.48-1	1.60+0	1.47-2	9.07-1	
1.80+0	2.27-2	9.08-1							
<b>58 Cerium</b> <b>Fluorescence yield = 0.912</b>									
1.00+0	3.94-4	6.75-1	1.20+0	1.02-3	7.74-1	1.40+0	2.30-3	9.18-1	129
1.60+0	4.19-3	9.86-1	1.80+0	6.72-3	1.01+0				
<b>64 Gadolinium</b> <b>Fluorescence yield = 0.935</b>									
9.00-1	6.91-5	6.21-1	1.10+0	2.12-4	6.90-1	1.20+0	2.30-4	4.99-1	150
1.40+0	5.28-4	5.82-1	1.60+0	1.26-3	8.00-1	1.80+0	1.46-3	5.84-1	
2.00+0	3.75-3	1.01+0	2.20+0	4.64-3	8.77-1	2.40+0	6.72-3	9.30-1	
2.60+0	9.43-3	9.86-1	2.80+0	1.08-2	8.75-1				
<b>73 Tantalum</b> <b>Fluorescence yield = 0.957</b>									
1.20+0	3.82-5	<b>3.43-1</b>	1.40+0	8.32-5	<b>3.51-1</b>	1.60+0	2.18-4	5.05-1	158
1.80+0	3.94-4	5.57-1	2.00+0	7.58-4	7.05-1	2.20+0	1.29-3	8.33-1	
2.40+0	2.19-3	1.03+0	2.60+0	2.72-3	9.57-1	2.80+0	3.24-3	8.80-1	
3.00+0	4.08-3	8.74-1	3.20+0	4.60-3	7.94-1				
<b>74 Tungsten</b> <b>Fluorescence yield = 0.958</b>									
1.30+0	1.07-4	7.44-1	1.44+0	1.89-4	8.04-1	1.60+0	3.13-4	8.29-1	147
1.70+0	4.10-4	8.37-1	1.80+0	5.22-4	8.40-1	1.90+0	6.51-4	8.41-1	
2.00+0	7.93-4	8.36-1	2.25+0	1.22-3	8.20-1	2.50+0	1.79-3	8.19-1	
2.75+0	2.48-3	8.09-1	3.00+0	3.33-3	8.05-1	3.25+0	4.34-3	8.02-1	
3.50+0	5.54-3	8.01-1	3.75+0	6.96-3	8.04-1	4.00+0	8.72-3	8.19-1	

TABLE 3. *K*-shell x-ray production by deuterons in target elements from beryllium to gold<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
<b>79 Gold</b> <b>Fluorescence yield = 0.964</b>									
1.42+0	8.15-5	7.35-1	1.52+0	1.20-4	7.74-1	1.64+0	1.77-4	8.00-1	147
1.80+0	2.73-4	8.17-1	1.90+0	3.43-4	8.15-1	2.00+0	4.22-4	8.12-1	
2.25+0	6.70-4	8.09-1	2.50+0	9.74-4	7.92-1	2.75+0	1.39-3	8.02-1	
3.00+0	1.91-3	8.14-1	3.25+0	2.57-3	8.34-1	3.50+0	3.40-3	8.64-1	
3.75+0	4.36-3	8.85-1	4.00+0	5.64-3	9.31-1				

<sup>a</sup>

Cross sections and their ratios are printed in a compressed power of 10  
 $^{-1}$   
 notation, e.g. 9.31-1 means  $9.31 \times 10^{-1}$ .

<sup>b</sup>

The ratios shown in **bold** print differ by more than a factor of 2 from  
 the averaged ratios and were -- as described in the text -- rejected.  
 This rejection criterion was applied only to the  $Z_2 > 9$  data.

TABLE 4. K-shell x-ray production by helium-3 in target elements from aluminum to silver<sup>a,b</sup>

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.	
<b>13 Aluminum</b> <b>Fluorescence yield = 0.039</b>										
1.00-1 3.80-2 1.98+0 1.75-1 4.40-1 1.36+0										
1.25-1 1.30-1 2.08+0 2.00-1 6.80-1 1.13+0										
4.50-2 1.20-4 1.69+0 7.50-2 3.70-3 1.07+0 1.00-1 3.10-2 1.62+0 1.70-1 2.50-1 8.85-1	5.00-2 3.00-4 1.68+0 8.00-2 9.20-3 1.77+0 1.20-1 7.10-2 1.40+0 1.90-1 3.80-1 8.02-1	6.00-2 1.20-3 1.60+0 9.00-2 2.00-2 1.90+0 1.50-1 1.50-1 9.66-1 2.00-1 6.50-1 1.08+0	1.50-1 2.60-1 1.67+0	1.25-1 2.60-1 1.67+0	1.50-1 2.60-1 1.67+0	1.50-1 2.60-1 1.67+0	1.50-1 2.60-1 1.67+0	1.50-1 2.60-1 1.67+0	12	
1.50-1 1.72-1 1.11+0 2.40-1 1.22+0 9.02-1	1.80-1 3.69-1 9.99-1 2.70-1 2.08+0 9.20-1	2.10-1 7.29-1 9.75-1 3.00-1 3.17+0 8.93-1	28							
4.50-2 1.24-4 1.75+0 7.00-2 3.97-3 1.80+0 1.00-1 3.29-2 1.72+0 1.40-1 1.45-1 1.31+0 2.00-1 5.96-1 9.94-1 6.12-1 3.70+1 6.20-1 1.02+0 2.11+2 6.60-1 1.43+0 5.54+2 7.10-1 2.04+0 1.27+3 7.86-1 2.65+0 2.22+3 9.18-1	5.00-2 3.26-4 1.82+0 8.00-2 9.58-3 1.85+0 1.10-1 5.12-2 1.59+0 1.60-1 2.43-1 1.15+0 4.08-1 8.28+0 6.56-1 7.14-1 6.49+1 6.31-1 1.12+0 2.80+2 6.70-1 1.63+0 7.65+2 7.32-1 2.24+0 1.57+3 8.31-1 2.85+0 2.58+3 9.70-1	6.00-2 1.28-3 1.71+0 9.00-2 1.75-2 1.66+0 1.20-1 7.55-2 1.49+0 1.80-1 3.88-1 1.05+0 5.10-1 1.95+1 6.45-1 8.16-1 1.01+2 6.28-1 1.22+0 3.62+2 6.88-1 1.84+0 1.01+3 7.56-1 2.45+0 1.90+3 8.76-1 2.96+0 2.76+3 9.91-1	6.00-2 1.28-3 1.71+0 9.00-2 1.75-2 1.66+0 1.20-1 7.55-2 1.49+0 1.80-1 3.88-1 1.05+0 5.10-1 1.95+1 6.45-1 8.16-1 1.01+2 6.28-1 1.22+0 3.62+2 6.88-1 1.84+0 1.01+3 7.56-1 2.45+0 1.90+3 8.76-1 2.96+0 2.76+3 9.91-1	6.00-2 1.28-3 1.71+0 9.00-2 1.75-2 1.66+0 1.20-1 7.55-2 1.49+0 1.80-1 3.88-1 1.05+0 5.10-1 1.95+1 6.45-1 8.16-1 1.01+2 6.28-1 1.22+0 3.62+2 6.88-1 1.84+0 1.01+3 7.56-1 2.45+0 1.90+3 8.76-1 2.96+0 2.76+3 9.91-1	6.00-2 1.28-3 1.71+0 9.00-2 1.75-2 1.66+0 1.20-1 7.55-2 1.49+0 1.80-1 3.88-1 1.05+0 5.10-1 1.95+1 6.45-1 8.16-1 1.01+2 6.28-1 1.22+0 3.62+2 6.88-1 1.84+0 1.01+3 7.56-1 2.45+0 1.90+3 8.76-1 2.96+0 2.76+3 9.91-1	6.00-2 1.28-3 1.71+0 9.00-2 1.75-2 1.66+0 1.20-1 7.55-2 1.49+0 1.80-1 3.88-1 1.05+0 5.10-1 1.95+1 6.45-1 8.16-1 1.01+2 6.28-1 1.22+0 3.62+2 6.88-1 1.84+0 1.01+3 7.56-1 2.45+0 1.90+3 8.76-1 2.96+0 2.76+3 9.91-1	6.00-2 1.28-3 1.71+0 9.00-2 1.75-2 1.66+0 1.20-1 7.55-2 1.49+0 1.80-1 3.88-1 1.05+0 5.10-1 1.95+1 6.45-1 8.16-1 1.01+2 6.28-1 1.22+0 3.62+2 6.88-1 1.84+0 1.01+3 7.56-1 2.45+0 1.90+3 8.76-1 2.96+0 2.76+3 9.91-1	6.00-2 1.28-3 1.71+0 9.00-2 1.75-2 1.66+0 1.20-1 7.55-2 1.49+0 1.80-1 3.88-1 1.05+0 5.10-1 1.95+1 6.45-1 8.16-1 1.01+2 6.28-1 1.22+0 3.62+2 6.88-1 1.84+0 1.01+3 7.56-1 2.45+0 1.90+3 8.76-1 2.96+0 2.76+3 9.91-1	6.00-2 1.28-3 1.71+0 9.00-2 1.75-2 1.66+0 1.20-1 7.55-2 1.49+0 1.80-1 3.88-1 1.05+0 5.10-1 1.95+1 6.45-1 8.16-1 1.01+2 6.28-1 1.22+0 3.62+2 6.88-1 1.84+0 1.01+3 7.56-1 2.45+0 1.90+3 8.76-1 2.96+0 2.76+3 9.91-1	45
2.25+0 8.08+2 <b>4.25-1</b> 4.50+0 2.66+3 6.49-1 9.00+0 4.65+3 9.06-1	3.00+0 1.45+3 5.12-1 6.00+0 3.68+3 7.77-1	3.75+0 2.08+3 5.85-1 7.50+0 4.35+3 8.63-1	75							
6.00-1 4.46+1 8.03-1 1.05+0 3.43+2 9.87-1 1.50+0 9.00+2 1.03+0 2.40+0 2.31+3 1.10+0	7.50-1 9.58+1 7.87-1 1.20+0 4.36+2 8.67-1 1.80+0 1.35+3 1.06+0	9.00-1 1.94+2 8.85-1 1.35+0 6.77+2 9.96-1 2.10+0 1.81+3 1.06+0	161							
<b>14 Silicon</b> <b>Fluorescence yield = 0.05</b>										
2.25+0 6.28+2 4.77-1 4.50+0 2.27+3 6.92-1 9.00+0 3.91+3 8.51-1	3.00+0 1.15+3 5.50-1 6.00+0 3.20+3 8.00-1	3.75+0 1.67+3 6.08-1 7.50+0 3.68+3 8.38-1	75							

TABLE 4. K-shell x-ray production by helium-3 in target elements from aluminum to silver<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
<b>15 Phosphorus</b> <b>Fluorescence yield = 0.063</b>									
6.00-1	8.97+0	5.32-1	7.50-1	2.82+1	7.33-1	9.00-1	6.16+1	8.47-1	161
1.05+0	1.00+2	8.27-1	1.20+0	1.57+2	8.59-1	1.35+0	2.35+2	9.08-1	
1.50+0	3.64+2	1.05+0	1.80+0	5.94+2	1.08+0	2.10+0	7.32+2	9.38-1	
2.40+0	1.07+3	1.04+0							
<b>18 Argon</b> <b>Fluorescence yield = 0.118</b>									
6.00-1	2.58+0	6.77-1	7.50-1	7.39+0	8.18-1	9.00-1	1.27+1	7.17-1	161
1.05+0	2.73+1	8.90-1	1.20+0	4.12+1	8.51-1	1.35+0	6.90+1	9.68-1	
1.50+0	9.05+1	9.10-1	1.80+0	1.77+2	1.04+0	2.10+0	2.49+2	9.53-1	
2.40+0	3.63+2	9.87-1							
<b>22 Titanium</b> <b>Fluorescence yield = 0.214</b>									
5.25-1	4.27-1	1.16+0	6.00-1	6.75-1	1.04+0	6.75-1	1.13+0	1.07+0	160
7.50-1	1.61+0	1.00+0	9.00-1	3.13+0	9.57-1	1.05+0	5.52+0	9.44-1	
1.20+0	8.96+0	9.42-1	1.50+0	1.90+1	9.13-1	1.80+0	3.57+1	9.37-1	
2.10+0	5.89+1	9.50-1	2.40+0	8.78+1	9.50-1	2.70+0	1.25+2	9.70-1	
3.00+0	1.68+2	9.77-1	3.30+0	2.13+2	9.71-1	3.60+0	2.80+2	1.03+0	
3.90+0	3.29+2	1.00+0	4.20+0	4.07+2	1.05+0	4.50+0	4.62+2	1.03+0	
4.80+0	5.28+2	1.03+0							
6.00-1	6.83-1	1.05+0	7.50-1	1.80+0	1.12+0	9.00-1	3.06+0	9.36-1	161
1.05+0	5.71+0	9.77-1	1.20+0	9.67+0	1.02+0	1.35+0	1.42+1	9.83-1	
1.50+0	2.04+1	9.80-1	1.80+0	3.79+1	9.94-1	2.10+0	6.06+1	9.78-1	
2.40+0	8.56+1	9.26-1	2.70+0	1.36+2	1.06+0	3.00+0	1.91+2	1.11+0	
3.30+0	2.48+2	1.13+0	3.60+0	3.10+2	1.14+0				
<b>24 Chromium</b> <b>Fluorescence yield = 0.275</b>									
5.25-1	1.85-1	1.12+0	6.00-1	3.30-1	1.11+0	6.75-1	5.50-1	1.13+0	160
7.50-1	7.84-1	1.04+0	9.00-1	1.60+0	1.03+0	1.05+0	2.79+0	9.87-1	
1.20+0	4.51+0	9.67-1	1.50+0	9.93+0	9.53-1	1.80+0	1.84+1	9.45-1	
2.10+0	2.99+1	9.26-1	2.40+0	4.54+1	9.25-1	2.70+0	6.68+1	9.55-1	
3.00+0	9.39+1	9.91-1	3.30+0	1.15+2	9.36-1	3.60+0	1.57+2	1.01+0	
3.90+0	1.89+2	9.95-1	4.20+0	2.26+2	9.90-1	4.50+0	2.68+2	9.99-1	
6.00-1	3.27-1	1.10+0	7.50-1	7.18-1	9.54-1	9.00-1	1.40+0	8.98-1	161
1.05+0	2.91+0	1.03+0	1.20+0	5.14+0	1.10+0	1.35+0	7.12+0	9.94-1	
1.50+0	1.02+1	9.79-1	1.80+0	1.92+1	9.86-1	2.10+0	3.12+1	9.66-1	
2.40+0	4.84+1	9.86-1	2.70+0	7.15+1	1.02+0	3.00+0	1.00+2	1.06+0	
3.30+0	1.25+2	1.02+0	3.60+0	1.49+2	9.59-1				
<b>27 Cobalt</b> <b>Fluorescence yield = 0.373</b>									
6.00-1	1.12-1	1.16+0	7.50-1	3.06-1	1.19+0	9.00-1	6.32-1	1.16+0	160
1.05+0	1.17+0	1.16+0	1.20+0	1.86+0	1.10+0	1.50+0	3.96+0	1.02+0	
1.80+0	7.16+0	9.61-1	2.10+0	1.23+1	9.73-1	2.40+0	1.92+1	9.77-1	
6.00-1	1.39-1	1.44+0	7.50-1	2.98-1	1.16+0	9.00-1	7.75-1	1.42+0	161
1.05+0	1.41+0	1.40+0	1.20+0	2.49+0	1.47+0	1.35+0	3.74+0	1.42+0	

TABLE 4. K-shell x-ray production by helium-3 in target elements from aluminum to silver<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	Ref.
1.50+0	5.51+0	1.42+0	1.80+0	1.00+1	1.34+0	2.10+0	1.54+1	1.22+0	
2.40+0	2.34+1	1.19+0	2.70+0	3.71+1	1.30+0	3.00+0	5.20+1	1.31+0	
3.30+0	6.83+1	1.30+0	3.60+0	8.60+1	1.27+0				
<b>28 Nickel</b>									
									Fluorescence yield = 0.406
6.00-1	5.97-2	8.94-1	7.50-1	1.56-1	8.63-1	9.00-1	3.53-1	9.07-1	161
1.05+0	6.86-1	9.46-1	1.20+0	1.19+0	9.74-1	1.35+0	1.85+0	9.66-1	
1.50+0	2.67+0	9.42-1	1.80+0	5.32+0	9.72-1	2.10+0	9.34+0	1.00+0	
2.40+0	1.24+1	8.49-1	2.70+0	1.94+1	9.07-1	3.00+0	2.56+1	8.60-1	
3.30+0	3.27+1	8.22-1	3.60+0	5.44+1	1.06+0				
<b>29 Copper</b>									
									Fluorescence yield = 0.44
5.25-1	2.60-2	1.06+0	6.00-1	5.00-2	1.07+0	6.75-1	9.00-2	1.12+0	160
7.50-1	1.05-1	8.15-1	9.00-1	3.12-1	1.11+0	1.05+0	5.74-1	1.09+0	
1.20+0	9.35-1	1.05+0	1.50+0	2.12+0	1.02+0	1.80+0	4.03+0	9.93-1	
2.10+0	6.81+0	9.78-1	2.40+0	1.08+1	9.85-1	2.70+0	1.57+1	9.73-1	
3.00+0	2.16+1	9.57-1	3.30+0	2.95+1	9.74-1	3.60+0	3.73+1	9.48-1	
3.90+0	4.81+1	9.68-1	4.20+0	5.83+1	9.51-1	4.50+0	6.72+1	9.06-1	
4.80+0	8.32+1	9.47-1							
<b>32 Germanium</b>									
									Fluorescence yield = 0.535
6.00-1	1.81-2	1.14+0	7.50-1	5.46-2	1.18+0	9.00-1	1.15-1	1.10+0	160
1.05+0	2.31-1	1.14+0	1.20+0	3.95-1	1.13+0	1.50+0	9.03-1	1.08+0	
1.80+0	1.71+0	1.03+0	2.10+0	2.95+0	1.02+0	2.40+0	4.59+0	9.95-1	
6.00-1	1.22-2	7.68-1	7.50-1	3.83-2	8.26-1	9.00-1	8.56-2	8.17-1	161
1.05+0	2.08-1	1.03+0	1.20+0	3.68-1	1.06+0	1.35+0	5.67-1	1.02+0	
1.50+0	7.92-1	9.48-1	1.80+0	1.56+0	9.43-1	2.10+0	2.88+0	9.96-1	
2.40+0	4.14+0	8.97-1	2.70+0	6.53+0	9.48-1	3.00+0	8.40+0	8.60-1	
3.30+0	1.27+1	9.56-1	3.60+0	1.72+1	9.85-1				
<b>34 Selenium</b>									
									Fluorescence yield = 0.589
6.00-1	2.78-3	3.57-1	7.50-1	1.35-2	5.71-1	1.05+0	9.54-2	8.85-1	161
1.20+0	1.55-1	8.22-1	1.35+0	2.83-1	9.32-1	1.50+0	4.45-1	9.68-1	
1.80+0	7.89-1	8.55-1	2.10+0	1.55+0	9.51-1	2.40+0	2.34+0	8.92-1	
2.70+0	3.49+0	8.84-1	3.00+0	5.27+0	9.34-1	3.30+0	7.07+0	9.15-1	
3.60+0	1.02+1	9.97-1							
<b>40 Zirconium</b>									
									Fluorescence yield = 0.73
1.20+0	3.50-2	1.04+0	1.50+0	9.20-2	1.06+0	1.80+0	2.01-1	1.11+0	160
2.10+0	3.58-1	1.09+0	2.40+0	5.34-1	9.86-1	3.00+0	1.21+0	1.00+0	
3.30+0	2.37+0	1.41+0	4.20+0	3.77+0	9.98-1	4.80+0	5.76+0	9.94-1	
<b>41 Niobium</b>									
									Fluorescence yield = 0.74
1.05+0	1.10-2	8.00-1	1.20+0	2.65-2	1.04+0	1.35+0	4.26-2	9.99-1	161
1.50+0	6.33-2	9.49-1	1.80+0	1.39-1	9.93-1	2.40+0	4.06-1	9.63-1	
3.00+0	8.89-1	9.39-1	3.30+0	1.31+0	9.91-1				

TABLE 4. K-shell x-ray production by helium-3 in target elements from aluminum to silver<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	Ref.
<b>46 Palladium                  Fluorescence yield = 0.82</b>									
1.05+0	1.85-3	<b>5.22-1</b>	1.20+0	3.84-3	5.62-1	1.35+0	8.69-3	7.31-1	161
1.50+0	1.51-2	7.91-1	1.80+0	3.51-2	8.44-1	2.10+0	7.64-2	9.85-1	
2.40+0	1.09-1	8.36-1	2.70+0	1.79-1	8.79-1	3.00+0	2.65-1	8.84-1	
3.30+0	3.52-1	8.31-1	3.60+0	5.01-1	8.71-1				
<b>47 Silver                  Fluorescence yield = 0.831</b>									
1.50+0	1.41-2	9.40-1	1.80+0	2.99-2	9.08-1	2.10+0	6.14-2	9.93-1	160
2.40+0	1.00-1	9.58-1	3.00+0	2.26-1	9.37-1	3.30+0	4.48-1	1.32+0	
4.20+0	7.57-1	9.58-1	4.80+0	1.22+0	9.85-1				

<sup>a</sup>

Cross sections and their ratios are printed in a compressed power of 10  
 $-1$   
 notation, e.g. 9.85-1 means  $9.85 \times 10^{-1}$ .

<sup>b</sup>

The ratios shown in **bold** print differ by more than a factor of 2 from  
 the averaged ratios and were -- as described in the text -- rejected.  
 This rejection criterion was applied only to the  $Z_2 > 9$  data.

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
<b>4 Beryllium      Fluorescence yield = 0.00033</b>									
5.00-1	4.06+3	1.43-1	7.50-1	5.43+3	2.26-1	1.00+0	6.29+3	3.20-1	71
1.25+0	6.88+3	4.22-1	1.50+0	7.15+3	5.17-1	1.75+0	7.11+3	5.91-1	
2.00+0	6.82+3	6.40-1							
2.50-1	6.09+3	2.58-1	3.00-1	7.28+3	2.73-1	5.00-1	1.10+4	3.87-1	92
8.00-1	1.10+4	4.76-1	1.00+0	1.07+4	5.44-1	1.20+0	1.02+4	6.04-1	
1.50+0	9.62+3	6.95-1	1.80+0	8.91+3	7.60-1	2.00+0	8.22+3	7.72-1	
4.00-2	1.90+1	2.64-1	5.00-2	3.75+1	1.59-1	6.00-2	6.54+1	1.14-1	119
8.00-2	1.59+2	8.29-2	1.00-1	3.28+2	7.88-2	1.20-1	5.44+2	7.74-2	
<b>6 Carbon      Fluorescence yield = 0.0028</b>									
4.00-2	2.70-1	4.14+0	5.00-2	3.70-1	1.63+0	6.00-2	6.00-1	9.74-1	16
7.00-2	8.40-1	5.92-1	8.00-2	1.30+0	4.49-1				
5.00-2	9.00-1	3.97+0	6.00-2	1.60+0	2.60+0	7.00-2	2.30+0	1.62+0	23
8.00-2	3.40+0	1.17+0	9.00-2	5.40+0	9.97-1	1.00-1	8.50+0	9.01-1	
1.10-1	1.10+1	7.09-1	1.20-1	1.80+1	7.41-1	1.30-1	2.80+1	7.67-1	
1.40-1	3.70+1	7.00-1	1.50-1	5.60+1	7.53-1	1.60-1	8.20+1	8.07-1	
1.70-1	1.00+2	7.3/-1	1.80-1	1.20+2	6.77-1	1.90-1	1.70+2	7.49-1	
2.00-1	1.90+2	6.65-1							
5.00-2	5.60-1	2.47+0	6.00-2	9.50-1	1.54+0	7.00-2	1.60+0	1.13+0	26
8.00-2	2.40+0	8.28-1	9.00-2	3.80+0	7.01-1	1.00-1	5.80+0	6.15-1	
1.10-1	8.90+0	5.74-1	1.20-1	1.20+1	4.94-1	1.30-1	1.70+1	4.66-1	
1.40-1	2.40+1	4.54-1	1.50-1	3.50+1	4.71-1	1.60-1	4.30+1	4.23-1	
1.70-1	6.00+1	4.42-1	1.80-1	7.00+1	3.95-1	1.90-1	8.40+1	3.70-1	
2.00-1	1.10+2	3.85-1							
1.16-1	5.31+0	2.60-1	1.35-1	1.01+1	2.29-1				46
1.50-1	2.26+1	3.04-1	1.75-1	4.13+1	2.66-1	2.00-1	6.62+1	2.32-1	82
2.50-1	1.89+2	2.60-1	3.00-1	3.44+2	2.40-1	4.00-1	1.03+3	2.95-1	
6.00-1	3.09+3	3.67-1	8.00-1	5.24+3	4.19-1	1.00+0	6.89+3	4.58-1	
1.20+0	8.09+3	4.91-1	1.60+0	8.89+3	5.14-1	2.00+0	9.86+3	5.85-1	
2.40+0	9.82+3	6.14-1							

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
<b>7 Nitrogen                  Fluorescence yield = 0.0052</b>									
9.70-2	2.11+0	1.40+0	1.16-1	4.27+0	1.15+0	1.35-1	8.34+0	1.06+0	46
1.25-1	4.65+0	8.66-1	1.50-1	1.06+1	8.03-1	1.75-1	1.90+1	6.83-1	82
2.00-1	3.14+1	6.00-1	2.50-1	8.03+1	5.59-1	3.00-1	1.48+2	4.77-1	
4.00-1	4.86+2	5.31-1	5.00-1	1.02+3	5.48-1	6.00-1	1.74+3	5.74-1	
7.00-1	2.71+3	6.29-1	8.00-1	3.70+3	6.62-1	1.00+0	5.33+3	6.72-1	
1.20+0	6.45+3	6.58-1	1.60+0	8.15+3	6.67-1	2.00+0	9.81+3	7.38-1	
<b>9 Fluorine                  Fluorescence yield = 0.013</b>									
1.00+0	1.67+3	7.79-1	1.70+0	4.49+3	8.19-1	1.80+0	4.82+3	8.19-1	110
4.00-1	7.19+1	6.29-1	6.00-1	2.00+2	3.92-1	8.00-1	4.01+2	3.27-1	116
1.00+0	7.01+2	3.27-1	1.20+0	1.07+3	3.40-1	1.40+0	1.62+3	3.92-1	
1.60+0	1.98+3	3.92-1	1.80+0	2.52+3	4.28-1	2.00+0	3.05+3	4.61-1	
<b>10 Neon                  Fluorescence yield = 0.018</b>									
9.70-2	1.36-1	2.06+0	1.16-1	3.51-1	2.17+0	1.35-1	6.13-1	1.82+0	46
2.00-1	1.38+0	6.57-1	2.50-1	3.66+0	6.39-1	3.00-1	7.10+0	5.58-1	82
4.00-1	2.19+1	5.11-1	5.00-1	5.40+1	5.19-1	6.00-1	1.12+2	5.46-1	
8.00-1	3.09+2	5.77-1	1.00+0	6.17+2	6.05-1	1.20+0	9.87+2	6.16-1	
<b>12 Magnesium                  Fluorescence yield = 0.03</b>									
1.25-1	7.40-2	1.92+0	1.50-1	1.80-1	1.89+0	1.75-1	3.40-1	1.71+0	12
2.00-1	5.90-1	1.59+0							
1.00+0	2.88+2	1.23+0	1.50+0	5.00+2	6.71-1	2.50+0	1.73+3	8.13-1	19
3.00+0	2.39+3	8.58-1	3.50+0	3.15+3	9.37-1	4.00+0	3.55+3	9.21-1	
4.50+0	4.05+3	9.49-1	5.00+0	4.60+3	9.99-1				
<b>13 Aluminum                  Fluorescence yield = 0.039</b>									
1.00-1	6.10-3	1.20+0	1.25-1	3.10-2	1.76+0	1.50-1	8.60-2	1.90+0	12
1.75-1	1.70-1	1.77+0	2.00-1	3.10-1	1.72+0				
2.90+0	2.40+2	1.32-1	3.55+0	3.50+2	1.43-1	3.90+0	4.40+2	1.59-1	15
4.45+0	5.70+2	1.78-1	4.80+0	6.60+2	1.92-1	5.30+0	8.30+2	2.21-1	
6.00-2	2.00-4	1.24+0	7.50-2	7.00-4	8.40-1	8.00-2	1.80-3	1.40+0	17
9.00-2	3.70-3	1.36+0	1.00-1	7.00-3	1.37+0	1.70-1	8.70-2	1.04+0	
1.90-1	1.50-1	1.06+0	2.00-1	2.20-1	1.22+0				
1.00+0	1.34+2	1.07+0	1.50+0	5.90+2	1.37+0	2.00+0	9.61+2	1.09+0	19
2.50+0	1.52+3	1.08+0	3.00+0	1.99+3	1.04+0	3.50+0	2.93+3	1.22+0	
4.00+0	4.10+3	1.44+0	4.50+0	4.81+3	1.49+0	5.00+0	5.08+3	1.42+0	

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
1.80-1	1.20-1	1.09+0	2.00-1	2.00-1	1.11+0	2.20-1	4.00-1	1.43+0	22
2.40-1	7.00-1	1.69+0	2.60-1	8.00-1	1.35+0	2.80-1	1.00+0	1.21+0	
3.00-1	1.60+0	1.44+0	3.20-1	2.00+0	1.36+0				
1.80-1	1.39-1	1.27+0	2.20-1	2.93-1	1.05+0	2.60-1	5.47-1	9.21-1	28
3.00-1	1.06+0	9.53-1	3.40-1	1.73+0	9.07-1				
6.00-2	2.10-4	1.30+0	7.00-2	7.35-4	1.43+0	8.00-2	1.75-3	1.36+0	45
9.00-2	4.06-3	1.50+0	1.00-1	7.65-3	1.50+0	1.20-1	2.10-2	1.48+0	
1.40-1	4.77-2	1.50+0	1.60-1	8.42-2	1.36+0	1.80-1	1.50-1	1.37+0	
2.00-1	2.41-1	1.34+0	4.08-1	3.25+0	7.91-1	5.10-1	6.60+0	6.46-1	
6.12-1	1.25+1	5.96-1	7.14-1	2.30+1	6.10-1	8.16-1	3.78+1	6.14-1	
1.02+0	8.47+1	6.35-1	1.12+0	1.17+2	6.46-1	1.22+0	1.54+2	6.53-1	
1.43+0	2.48+2	6.58-1	1.63+0	3.62+2	6.73-1	1.84+0	4.95+2	6.80-1	
2.04+0	6.46+2	7.01-1	2.24+0	8.20+2	7.26-1	2.35+0	9.15+2	7.37-1	
2.45+0	1.02+3	7.56-1	2.55+0	1.14+3	7.84-1	2.65+0	1.25+3	8.02-1	
2.85+0	1.49+3	8.43-1	2.96+0	1.59+3	8.46-1	3.06+0	1.68+3	8.47-1	
3.16+0	1.76+3	8.45-1	3.26+0	1.85+3	8.48-1	3.67+0	2.18+3	8.49-1	
4.08+0	2.44+3	8.37-1	4.69+0	2.76+3	8.18-1	5.51+0	3.10+3	8.01-1	
5.92+0	3.17+3	7.78-1							
1.00+0	1.00+2	8.00-1	1.50+0	4.41+2	1.02+0	2.00+0	9.42+2	1.07+0	54
2.50+0	1.57+3	1.12+0	3.00+0	2.19+3	1.14+0	3.50+0	2.89+3	1.20+0	
4.00+0	3.52+3	1.23+0	4.50+0	4.18+3	1.29+0	5.00+0	4.83+3	1.35+0	
2.50+0	1.10+3	7.84-1	3.20+0	1.80+3	8.49-1	3.25+0	2.00+3	9.21-1	96
4.00+0	2.39+3	8.38-1	4.90+0	2.60+3	7.40-1				
4.00+0	2.17+3	7.61-1	6.00+0	3.30+3	8.03-1	8.00+0	3.91+3	8.24-1	97
1.00+1	4.15+3	8.22-1	1.20+1	4.28+3	8.33-1	1.60+1	4.12+3	8.19-1	
2.00+1	3.93+3	8.20-1	2.40+1	3.58+3	7.92-1	2.80+1	3.38+3	7.95-1	
3.00+1	3.30+3	7.99-1							
1.00+0	1.13+2	9.04-1	1.80+0	6.64+2	9.61-1				110
<b>14</b>	Silcon		Fluorescence yield = 0.05						
2.90+0	1.40+2	1.12-1	3.55+0	2.00+2	1.13-1	3.90+0	2.50+2	1.23-1	15
4.45+0	3.10+2	1.28-1	4.80+0	3.50+2	1.32-1	5.30+0	4.10+2	1.39-1	
<b>16</b>	Sulfur		Fluorescence yield = 0.078						
1.10+1	4.20+3	1.22+0	1.60+1	4.82+3	1.24+0	2.10+1	4.45+3	1.12+0	109
2.70+1	4.18+3	1.08+0							
<b>17</b>	Chlorine		Fluorescence yield = 0.097						
1.20+1	3.45+3	1.08+0	2.50+1	4.60+3	1.22+0	3.00+1	4.46+3	1.21+0	31
4.00+1	3.69+3	1.07+0							
1.10+1	3.10+3	1.03+0	1.60+1	4.20+3	1.17+0	2.10+1	3.70+3	9.83-1	109
2.70+1	4.45+3	1.19+0							

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$		$E_1$ (MeV)	$\sigma^{\text{Exper}}$		$E_1$ (MeV)	$\sigma^{\text{Exper}}$		$E_1$ (MeV)	$\sigma^{\text{Exper}}$		Ref.
	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{ECPSSR}}$		$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{ECPSSR}}$		$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{ECPSSR}}$		$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{ECPSSR}}$	
8.00-1	1.24+1	1.92+0	1.00+0	2.80+1	1.87+0	1.20+0	4.72+1	1.63+0	1.20+0	4.72+1	1.63+0	123
1.40+0	7.56+1	1.53+0	1.60+0	1.04+2	1.36+0	1.80+0	1.52+2	1.37+0	2.00+0	2.19+2	1.26+0	
2.00+0	2.19+2	1.43+0	2.20+0	2.54+2	1.26+0	2.40+0	2.97+2	1.16+0	2.60+0	4.55+2	1.43+0	
				2.80+0	5.04+2	1.32+0						
<b>18 Argon</b>	Fluorescence yield = 0.118											
3.00-1	4.15-2	6.65-1	5.00-1	4.59-1	7.58-1	8.00-1	3.11+0	7.69-1	8.00-1	3.11+0	7.69-1	82
1.00+0	6.60+0	6.97-1	1.20+0	1.20+1	6.51-1							
<b>19 Potassium</b>	Fluorescence yield = 0.14											
1.10+1	2.54+3	1.16+0	1.60+1	3.20+3	1.12+0	2.10+1	3.69+3	1.17+0	2.70+1	3.78+3	1.14+0	109
<b>20 Calcium</b>	Fluorescence yield = 0.163											
1.50+0	2.00+1	1.16+0	2.00+0	5.00+1	1.10+0	3.00+0	1.80+2	1.16+0	4.00+0	3.70+2	1.13+0	38
4.00+0	3.70+2	1.13+0	5.00+0	6.40+2	1.18+0	6.00+0	8.60+2	1.11+0	7.00+0	1.00+3	9.92-1	
7.00+0	1.00+3	9.92-1	8.00+0	1.30+3	1.05+0	9.00+0	1.40+3	9.66-1	1.00+1	1.60+3	9.72-1	
			1.10+1	1.80+3	9.85-1	1.20+1	1.90+3	9.55-1				
6.00+0	7.24+2	9.36-1	9.00+0	1.48+3	1.02+0	1.20+1	2.14+3	1.08+0	1.50+1	2.62+3	1.10+0	94
			1.80+1	2.97+3	1.12+0							
8.00-1	2.41+0	1.47+0	1.00+0	5.00+0	1.28+0	1.20+0	1.00+1	1.30+0	1.40+0	1.76+1	1.30+0	123
1.40+0	1.76+1	1.30+0	1.60+0	2.90+1	1.34+0	1.80+0	4.27+1	1.32+0	2.00+0	6.15+1	1.35+0	
2.00+0	6.15+1	1.35+0	2.20+0	8.64+1	1.39+0	2.40+0	1.05+2	1.29+0	2.60+0	1.41+2	1.37+0	
			2.80+0	1.70+2	1.33+0							
8.00-1	1.51+0	9.23-1	1.00+0	3.34+0	8.57-1	1.20+0	6.68+0	8.65-1	1.40+0	1.13+1	8.37-1	133
			1.60+0	1.76+1	8.16-1	1.80+0	2.45+1	7.60-1				
2.00+1	3.18+3	1.14+0	2.40+1	3.36+3	1.14+0	2.80+1	3.50+3	1.15+0				136
<b>21 Scandium</b>	Fluorescence yield = 0.188											
8.00-1	1.07+0	1.00+0	1.00+0	2.53+0	9.85-1	1.20+0	4.62+0	9.02-1	1.40+0	8.49+0	9.41-1	133
			1.60+0	1.38+1	9.51-1	1.80+0	2.03+1	9.30-1				
2.00+1	2.85+3	1.15+0	2.40+1	3.21+3	1.20+0	2.80+1	3.32+3	1.19+0				136
<b>22 Titanium</b>	Fluorescence yield = 0.214											
2.90+0	2.40+1	3.39-1	3.90+0	5.30+1	3.24-1	4.80+0	9.10+1	3.29-1	5.30+0	1.16+2	3.34-1	14
1.20+1	1.41+3	1.01+0	2.50+1	2.89+3	1.19+0	3.00+1	3.17+3	1.23+0	4.00+1	3.40+3	1.27+0	31
1.50+0	8.80+0	1.11+0	2.00+0	2.50+1	1.16+0	3.00+0	8.80+1	1.12+0	4.00+0	2.00+2	1.14+0	38
4.00+0	2.00+2	1.14+0	5.00+0	3.30+2	1.08+0	6.00+0	5.70+2	1.25+0	7.00+0	6.60+2	1.07+0	
7.00+0	6.60+2	1.07+0	8.00+0	8.50+2	1.09+0	9.00+0	8.80+2	9.31-1	1.00+1	1.20+3	1.09+0	
			1.10+1	1.40+3	1.12+0	1.20+1	1.70+3	1.22+0				

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
2.00+1	2.40+3	1.10+0							42
5.00-1	1.21-1	1.25+0	6.00-1	2.61-1	1.21+0	7.00-1	4.91-1	1.19+0	67
8.00-1	8.48-1	1.19+0	9.00-1	1.33+0	1.17+0	1.00+0	2.00+0	1.16+0	
1.10+0	2.87+0	1.15+0	1.20+0	3.83+0	1.10+0	1.30+0	5.15+0	1.10+0	
1.40+0	6.57+0	1.07+0	1.50+0	8.42+0	1.07+0	1.60+0	1.03+1	1.04+0	
1.70+0	1.29+1	1.05+0	1.80+0	1.47+1	9.78-1	1.90+0	1.82+1	1.01+0	
2.00+0	2.17+1	1.01+0	2.10+0	2.40+1	9.46-1	2.20+0	2.85+1	9.62-1	
2.30+0	3.22+1	9.41-1	2.40+0	3.70+1	9.42-1				
5.00-1	1.10-1	1.14+0	6.00-1	2.70-1	1.26+0	7.00-1	5.00-1	1.21+0	72
8.00-1	7.00-1	9.83-1	9.00-1	8.50-1	7.45-1	1.00+0	1.30+0	7.53-1	
1.10+0	2.10+0	8.42-1	1.20+0	3.00+0	8.65-1	1.30+0	3.80+0	8.13-1	
1.40+0	5.10+0	8.30-1	1.50+0	6.50+0	8.23-1	1.60+0	8.50+0	8.55-1	
1.70+0	1.00+1	8.12-1	1.80+0	1.20+1	7.99-1	1.90+0	1.50+1	8.29-1	
2.00+0	1.90+1	8.81-1	2.10+0	2.20+1	8.67-1	2.20+0	2.60+1	8.78-1	
2.30+0	2.90+1	8.47-1	2.40+0	3.10+1	7.89-1	2.50+0	3.90+1	8.72-1	
1.00+0	2.63+0	1.52+0	1.10+0	3.37+0	1.35+0	1.30+0	5.43+0	1.16+0	78
1.50+0	7.80+0	9.88-1	1.70+0	1.15+1	9.34-1	1.90+0	1.83+1	1.01+0	
2.10+0	2.28+1	8.98-1	2.20+0	3.06+1	1.03+0	2.30+0	3.49+1	1.02+0	
2.40+0	4.02+1	1.02+0	2.50+0	4.51+1	1.01+0	2.60+0	4.67+1	9.23-1	
2.70+0	5.67+1	9.96-1	2.80+0	6.38+1	1.00+0	2.90+0	7.01+1	9.91-1	
3.00+0	7.75+1	9.90-1	3.10+0	8.45+1	9.79-1	3.20+0	9.08+1	9.60-1	
3.30+0	1.00+2	9.67-1	3.40+0	1.11+2	9.86-1	3.50+0	1.19+2	9.76-1	
3.60+0	1.23+2	9.34-1	3.70+0	1.34+2	9.45-1	3.80+0	1.46+2	9.58-1	
3.90+0	1.58+2	9.66-1	4.00+0	1.69+2	9.64-1	4.10+0	1.75+2	9.37-1	
4.20+0	1.90+2	9.57-1	4.30+0	2.04+2	9.68-1	4.40+0	2.03+2	9.09-1	
2.50+0	5.70+1	1.27+0	3.20+0	1.00+2	1.06+0	3.25+0	1.10+2	1.11+0	96
4.00+0	1.79+2	1.02+0	4.90+0	2.50+2	8.62-1				
1.00+1	1.03+3	9.34-1	1.20+1	1.38+3	9.92-1	1.40+1	1.64+3	9.98-1	105
1.60+1	1.86+3	1.00+0	1.80+1	2.21+3	1.09+0	2.00+1	2.32+3	1.07+0	
2.20+1	2.50+3	1.09+0	2.40+1	2.65+3	1.11+0	2.60+1	2.72+3	1.10+0	
2.80+1	2.80+3	1.11+0	3.00+1	2.96+3	1.15+0				
2.30-1	3.92-4	2.23-1	2.50-1	8.54-4	2.95-1	2.80-1	1.89-3	3.42-1	118
3.00-1	3.21-3	4.00-1	3.50-1	8.62-3	4.82-1	4.00-1	1.93-2	5.62-1	
5.00-1	6.57-2	6.81-1	6.00-1	1.59-1	7.40-1	7.00-1	3.15-1	7.64-1	
8.00-1	5.56-1	7.81-1							
8.00-1	7.80-1	1.09+0	1.00+0	1.81+0	1.05+0	1.20+0	3.34+0	9.63-1	133
1.40+0	5.72+0	9.31-1	1.60+0	9.15+0	9.20-1	1.80+0	1.33+1	8.85-1	
2.00+1	2.36+3	1.08+0	2.40+1	2.72+3	1.14+0	2.80+1	2.88+3	1.14+0	136
7.76-1	6.02-1	9.56-1	9.16-1	1.03+0	8.41-1	1.05+0	1.45+0	6.90-1	146
1.60+0	6.69+0	6.73-1							
1.50+0	4.66+0	5.90-1	2.02+0	1.88+1	8.43-1	2.50+0	3.24+1	7.24-1	148
3.00+0	5.56+1	7.10-1	3.50+0	9.01+1	7.39-1				

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
4.50-1	8.07-2	1.35+0	5.00-1	1.30-1	1.35+0	5.50-1	1.95-1	1.32+0	153
6.00-1	2.78-1	1.29+0	6.50-1	3.84-1	1.27+0	7.00-1	5.14-1	1.25+0	
7.50-1	6.73-1	1.23+0	8.00-1	8.64-1	1.21+0	8.50-1	1.09+0	1.20+0	
9.00-1	1.36+0	1.19+0	1.00+0	2.02+0	1.17+0	1.10+0	2.88+0	1.15+0	
1.20+0	3.97+0	1.14+0	1.30+0	5.31+0	1.14+0	1.40+0	6.93+0	1.13+0	
1.50+0	8.88+0	1.12+0	1.60+0	1.12+1	1.13+0	1.80+0	1.69+1	1.12+0	
2.00+0	2.45+1	1.14+0	2.20+0	3.42+1	1.15+0	2.40+0	4.65+1	1.18+0	
2.60+0	6.17+1	1.22+0	2.80+0	8.03+1	1.26+0	3.00+0	1.03+2	1.32+0	
7.00-1	4.82-1	1.17+0	8.00-1	8.23-1	1.16+0	9.00-1	1.21+0	1.06+0	160
1.00+0	1.83+0	1.06+0	1.20+0	3.57+0	1.03+0	1.40+0	6.29+0	1.02+0	
1.60+0	1.03+1	1.04+0	2.00+0	2.09+1	9.70-1	2.40+0	3.72+1	9.47-1	
<b>23</b>	<b>Vanadium</b>								
1.00+0	1.47+0	1.24+0	1.25+0	2.60+0	9.31-1	1.50+0	4.30+0	7.83-1	54
1.75+0	7.00+0	7.34-1	2.00+0	9.79+0	6.45-1	2.50+0	1.62+1	5.08-1	
3.00+0	2.73+1	4.83-1	3.50+0	4.00+1	<b>4.48-1</b>	4.00+0	5.82+0	<b>4.50-2</b>	
4.50+0	7.59+0	<b>4.29-2</b>							
5.00-1	7.40-2	1.16+0	6.00-1	1.81-1	1.26+0	7.00-1	3.36-1	1.21+0	67
8.00-1	5.83-1	1.21+0	9.00-1	9.47-1	1.22+0	1.00+0	1.39+0	1.17+0	
1.10+0	2.07+0	1.21+0	1.20+0	2.69+0	1.12+0	1.30+0	3.66+0	1.13+0	
1.40+0	4.90+0	1.15+0	1.50+0	6.02+0	1.10+0	1.60+0	7.59+0	1.09+0	
1.70+0	9.34+0	1.08+0	1.80+0	1.13+1	1.07+0	1.90+0	1.37+1	1.08+0	
2.00+0	1.62+1	1.07+0	2.10+0	1.90+1	1.06+0	2.20+0	2.13+1	1.02+0	
2.30+0	2.60+1	1.07+0	2.40+0	2.87+1	1.03+0				
1.00+0	1.21+0	1.02+0	2.20+0	1.97+1	9.39-1	2.30+0	2.26+1	9.30-1	78
2.40+0	2.59+1	9.26-1	2.50+0	2.96+1	9.28-1	2.60+0	2.96+1	8.18-1	
2.70+0	3.70+1	9.07-1	2.80+0	4.15+1	9.08-1	2.90+0	4.54+1	8.90-1	
3.00+0	5.00+1	8.84-1	3.10+0	5.51+1	8.82-1	3.20+0	5.93+1	8.63-1	
3.30+0	6.46+1	8.58-1	3.40+0	6.86+1	8.35-1	3.50+0	7.71+1	8.63-1	
3.60+0	8.37+1	8.64-1	3.70+0	9.01+1	8.61-1	3.80+0	9.70+1	8.61-1	
3.90+0	1.05+2	8.69-1	4.00+0	1.12+2	8.66-1	4.10+0	1.11+2	8.03-1	
4.20+0	1.25+2	8.47-1	4.30+0	1.36+2	8.65-1	4.40+0	1.38+2	8.25-1	
8.00-1	5.39-1	1.11+0	1.00+0	1.28+0	1.08+0	1.20+0	2.42+0	1.01+0	133
1.40+0	4.20+0	9.86-1	1.60+0	7.06+0	1.02+0	1.80+0	1.03+1	9.77-1	
2.00+1	2.04+3	1.06+0	2.40+1	2.38+3	1.11+0	2.80+1	2.68+3	1.17+0	136
<b>24</b>	<b>Chromium</b>								
2.90+0	1.50+1	<b>4.04-1</b>	3.55+0	2.60+1	<b>3.78-1</b>	3.90+0	3.40+1	<b>3.77-1</b>	14
4.80+0	7.10+1	<b>4.48-1</b>	5.30+0	8.10+1	<b>3.99-1</b>				
5.00-1	5.30-2	1.25+0	6.00-1	1.25-1	1.29+0	7.00-1	2.38-1	1.25+0	67
8.00-1	4.25-1	1.28+0	9.00-1	6.61-1	1.22+0	1.00+0	1.03+0	1.25+0	
1.10+0	1.45+0	1.21+0	1.20+0	1.96+0	1.17+0	1.30+0	2.67+0	1.17+0	
1.40+0	3.51+0	1.17+0	1.50+0	4.24+0	1.09+0	1.60+0	5.24+0	1.07+0	
1.70+0	6.97+0	1.14+0	1.80+0	8.28+0	1.10+0	1.90+0	9.92+0	1.09+0	
2.00+0	1.15+1	1.06+0	2.10+0	1.35+1	1.05+0	2.20+0	1.58+1	1.05+0	
2.30+0	1.83+1	1.05+0	2.40+0	1.99+1	9.87-1				

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
9.00-1	1.10+0	2.04+0	1.10+0	2.09+0	1.74+0	1.30+0	4.15+0	1.83+0	80
1.50+0	5.75+0	1.48+0	1.70+0	9.93+0	1.62+0	1.90+0	1.40+1	1.54+0	
2.10+0	2.13+1	1.66+0	2.30+0	2.56+1	1.46+0	2.50+0	3.32+1	1.44+0	
2.70+0	4.70+1	1.59+0	2.90+0	5.40+1	1.45+0	3.10+0	7.46+1	1.63+0	
3.30+0	8.86+1	1.60+0	3.50+0	1.13+2	1.71+0	3.70+0	1.22+2	1.57+0	
3.90+0	1.56+2	1.73+0	4.10+0	1.91+2	1.84+0				
6.00+0	2.91+2	1.07+0	9.00+0	6.75+2	1.09+0	1.20+1	1.17+3	1.20+0	94
1.50+1	1.51+3	1.17+0	1.80+1	1.76+3	1.14+0				
8.00-1	4.02-1	1.21+0	1.00+0	9.17-1	1.12+0	1.20+0	1.80+0	1.07+0	133
1.40+0	3.06+0	1.02+0	1.60+0	4.90+0	9.98-1	1.80+0	6.96+0	9.28-1	
2.00+1	1.86+3	1.10+0	2.40+1	2.18+3	1.14+0	2.80+1	2.33+3	1.12+0	136
1.50+0	4.08+0	1.05+0	2.02+0	1.06+1	9.44-1	2.50+0	2.28+1	9.89-1	148
3.00+0	4.07+1	9.85-1	3.50+0	6.86+1	1.04+0				
<b>25</b>	<b>Manganese</b>		<b>Fluorescence yield = 0.308</b>						
1.00+0	8.14-1	1.41+0	1.10+0	1.02+0	1.21+0	1.30+0	1.66+0	1.03+0	78
1.50+0	2.50+0	9.07-1	1.70+0	3.54+0	8.11-1	1.90+0	6.27+0	9.63-1	
2.20+0	1.08+1	9.94-1	2.30+0	1.23+1	9.73-1	2.40+0	1.35+1	9.24-1	
2.50+0	1.61+1	9.62-1	2.60+0	1.59+1	8.33-1	2.70+0	1.98+1	9.17-1	
2.80+0	2.22+1	9.13-1	2.90+0	2.47+1	9.08-1	3.00+0	2.88+1	9.50-1	
3.10+0	3.01+1	8.96-1	3.20+0	3.19+1	8.59-1	3.30+0	3.53+1	8.65-1	
3.40+0	3.84+1	8.58-1	3.50+0	4.25+1	8.70-1	3.60+0	4.60+1	8.65-1	
3.70+0	4.94+1	8.56-1	3.80+0	5.43+1	8.70-1	3.90+0	5.57+1	8.27-1	
4.00+0	6.20+1	8.56-1	4.10+0	6.74+1	8.67-1	4.20+0	7.19+1	8.63-1	
4.30+0	7.76+1	8.72-1	4.40+0	8.36+1	8.81-1				
8.00-1	2.70-1	1.17+0	1.00+0	6.73-1	1.17+0	1.20+0	1.31+0	1.11+0	133
1.40+0	2.20+0	1.03+0	1.60+0	3.57+0	1.02+0	1.80+0	5.35+0	9.97-1	
<b>26</b>	<b>Iron</b>		<b>Fluorescence yield = 0.34</b>						
2.90+0	1.20+1	6.02-1	3.55+0	2.00+1	5.29-1	3.90+0	2.50+1	4.98-1	14
4.80+0	4.40+1	4.84-1	5.30+0	5.80+1	<b>4.90-1</b>				
3.00+1	2.55+3	1.47+0	4.00+1	2.59+3	1.32+0	5.00+1	2.66+3	1.30+0	24
6.00+1	2.75+3	1.33+0	7.00+1	2.77+3	1.35+0	8.00+1	2.58+3	1.29+0	
2.00+0	7.00+0	1.24+0	3.00+0	2.50+1	1.12+0	4.00+0	6.60+1	1.22+0	38
5.00+0	1.10+2	1.08+0	6.00+0	1.70+2	1.05+0	7.00+0	2.50+2	1.07+0	
8.00+0	3.40+2	1.08+0	9.00+0	4.20+2	1.05+0	1.00+1	5.20+2	1.07+0	
1.10+1	6.30+2	1.09+0	1.20+1	6.90+2	1.04+0				
2.00+1	1.30+3	1.02+0							42
3.00+1	2.18+3	1.26+0							49

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
1.00+0 4.14-1 1.02+0 2.50+0 1.02+1 8.37-1	1.50+0 1.76+0 8.93-1 3.00+0 1.94+1 8.71-1	2.00+0 4.86+0 8.64-1 3.50+0 3.03+1 8.37-1							54
4.00+0 4.31+1 7.96-1 1.00+0 4.53-1 1.12+0 2.40+0 9.86+0 9.29-1 2.70+0 1.41+1 8.94-1 3.00+0 1.93+1 8.67-1 3.30+0 2.65+1 8.79-1 3.60+0 3.59+1 9.10-1 3.90+0 4.77+1 9.50-1 4.20+0 5.66+1 9.07-1	4.50+0 5.68+1 7.47-1 2.20+0 7.48+0 9.51-1 2.50+0 1.10+1 9.03-1 2.80+0 1.59+1 8.94-1 3.10+0 2.15+1 8.69-1 3.40+0 2.99+1 9.04-1 3.70+0 3.89+1 9.07-1 4.00+0 4.94+1 9.13-1 4.30+0 6.00+1 8.98-1	5.00+0 8.41+1 8.29-1 2.30+0 8.66+0 9.44-1 2.60+0 1.17+1 8.41-1 2.90+0 1.79+1 8.98-1 3.20+0 2.34+1 8.56-1 3.50+0 3.33+1 9.20-1 3.80+0 4.14+1 8.91-1 4.10+0 5.21+1 8.95-1 4.40+0 6.12+1 8.58-1							78
3.20+0 3.20+1 1.17+0 4.90+0 8.40+1 8.74-1	3.25+0 3.50+1 1.22+0	4.00+0 6.33+1 1.17+0							96
1.00+1 4.56+2 9.36-1 1.60+1 1.05+3 1.05+0 2.20+1 1.56+3 1.12+0 2.80+1 1.85+3 1.11+0	1.20+1 6.59+2 9.89-1 1.80+1 1.34+3 1.17+0 2.40+1 1.70+3 1.14+0 3.00+1 1.85+3 1.07+0	1.40+1 8.74+2 1.04+0 2.00+1 1.35+3 1.06+0 2.60+1 1.80+3 1.14+0							105
2.80-1 1.32-4 1.78-1 4.00-1 2.96-3 4.87-1 7.00-1 7.96-2 8.83-1	3.00-1 2.84-4 2.47-1 5.00-1 1.31-2 6.86-1 8.00-1 1.49-1 9.28-1	3.50-1 9.70-4 3.35-1 6.00-1 3.63-2 8.06-1							118
8.00-1 2.06-1 1.28+0 1.40+0 1.38+0 9.10-1	1.00+0 4.86-1 1.20+0 1.60+0 2.69+0 1.07+0	1.20+0 1.01+0 1.21+0 1.80+0 4.07+0 1.06+0							133
4.50-1 1.40-2 1.24+0 6.00-1 6.07-2 1.35+0 7.50-1 1.61-1 1.32+0 9.00-1 3.40-1 1.29+0 1.20+0 1.04+0 1.24+0 1.50+0 2.34+0 1.19+0 2.00+0 6.22+0 1.11+0 2.60+0 1.43+1 1.03+0	5.00-1 2.50-2 1.31+0 6.50-1 8.70-2 1.34+0 8.00-1 2.11-1 1.31+0 1.00+0 5.17-1 1.27+0 1.30+0 1.40+0 1.22+0 1.60+0 2.93+0 1.17+0 2.20+0 8.47+0 1.08+0 2.80+0 1.80+1 1.01+0	5.50-1 4.03-2 1.34+0 7.00-1 1.20-1 1.33+0 8.50-1 2.70-1 1.30+0 1.10+0 7.48-1 1.26+0 1.40+0 1.83+0 1.21+0 1.80+0 4.38+0 1.14+0 2.40+0 1.12+1 1.06+0 3.00+0 2.22+1 9.97-1							153
<b>27 Cobalt</b>	<b>Fluorescence yield = 0.373</b>								
2.00+0 6.50+0 1.58+0 3.50+0 3.80+1 1.41+0 5.00+0 1.05+2 1.35+0 8.00+0 2.70+2 1.08+0 1.10+1 5.00+2 1.06+0	2.50+0 1.30+1 1.45+0 4.00+0 5.70+1 1.40+0 6.00+0 1.60+2 1.27+0 9.00+0 3.60+2 1.12+0 1.20+1 6.10+2 1.12+0	3.00+0 2.30+1 1.40+0 4.50+0 7.60+1 1.32+0 7.00+0 2.00+2 1.09+0 1.00+1 4.60+2 1.17+0							38
2.00+1 1.10+3 1.00+0									42
1.00+0 3.99-1 1.38+0 1.50+0 1.47+0 1.04+0 2.20+0 5.31+0 9.23-1 2.50+0 7.83+0 8.74-1 2.80+0 1.11+1 8.46-1 3.10+0 1.48+1 8.07-1 3.40+0 1.97+1 8.00-1	1.10+0 6.19-1 1.46+0 1.70+0 2.12+0 9.35-1 2.30+0 6.00+0 8.93-1 2.60+0 9.69+0 9.47-1 2.90+0 1.23+1 8.35-1 3.20+0 1.62+1 7.98-1 3.50+0 2.16+1 8.00-1	1.30+0 9.75-1 1.19+0 1.90+0 3.39+0 9.95-1 2.40+0 6.80+0 8.74-1 2.70+0 9.86+0 8.49-1 3.00+0 1.36+1 8.25-1 3.30+0 1.79+1 7.98-1 3.60+0 2.31+1 7.84-1							78

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
3.70+0	2.52+1	7.86-1	3.80+0	2.72+1	7.81-1	3.90+0	2.95+1	7.83-1	
4.00+0	3.22+1	7.91-1	4.10+0	3.40+1	7.76-1	4.20+0	3.69+1	7.84-1	
4.30+0	3.92+1	7.77-1	4.40+0	4.35+1	8.07-1				
8.00-1	1.36-1	1.20+0	1.00+0	3.30-1	1.14+0	1.20+0	6.77-1	1.13+0	133
1.40+0	1.22+0	1.12+0	1.60+0	1.91+0	1.05+0	1.80+0	2.81+0	1.00+0	
1.50+0	1.15+0	8.10-1	2.02+0	3.26+0	7.67-1	2.50+0	6.72+0	7.50-1	148
3.00+0	1.23+1	7.46-1	3.20+0	1.55+1	7.63-1	3.50+0	1.98+1	7.33-1	
<b>28 Nickel</b>		<b>Fluorescence yield = 0.406</b>							
2.90+0	8.40+0	7.69-1	3.55+0	1.30+1	6.15-1	3.90+0	1.50+1	5.29-1	14
4.80+0	2.50+1	<b>4.75-1</b>	5.30+0	3.50+1	5.02-1				
3.00+1	1.71+3	1.25+0	4.00+1	1.85+3	1.15+0	5.00+1	2.05+3	1.19+0	24
6.00+1	2.30+3	1.29+0	7.00+1	2.34+3	1.30+0	8.00+1	2.33+3	1.31+0	
2.00+0	2.30+0	7.67-1	3.00+0	9.10+0	7.44-1	4.00+0	2.30+1	7.51-1	38
5.00+0	5.00+1	8.45-1	6.00+0	8.50+1	8.76-1	7.00+0	1.10+2	7.65-1	
8.00+0	1.70+2	8.62-1	9.00+0	2.00+2	7.82-1	1.00+1	2.60+2	8.17-1	
1.10+1	2.90+2	7.59-1	1.20+1	3.30+2	7.37-1				
1.50+0	6.60-1	6.41-1	2.00+0	1.40+0	<b>4.67-1</b>	2.50+0	3.70+0	5.61-1	43
3.00+0	7.40+0	6.05-1	3.50+0	1.10+1	5.45-1	4.00+0	1.80+1	5.88-1	
5.00+1	2.07+3	1.20+0	6.00+1	2.30+3	1.29+0	7.00+1	2.33+3	1.30+0	49
8.00+1	2.21+3	1.24+0	9.00+1	2.23+3	1.27+0	1.00+2	2.12+3	1.24+0	
1.00+0	2.07-1	1.01+0	2.20+0	3.95+0	9.36-1	2.30+0	4.52+0	9.16-1	78
2.40+0	4.97+0	8.67-1	2.50+0	5.80+0	8.79-1	2.60+0	6.66+0	8.82-1	
2.70+0	7.32+0	8.52-1	2.80+0	8.21+0	8.45-1	2.90+0	9.08+0	8.31-1	
3.00+0	1.02+1	8.34-1	3.10+0	1.10+1	8.07-1	3.20+0	1.20+1	7.93-1	
3.30+0	1.34+1	8.02-1	3.40+0	1.46+1	7.94-1	3.50+0	1.61+1	7.97-1	
3.60+0	1.76+1	7.97-1	3.70+0	1.92+1	7.98-1	3.80+0	2.09+1	7.99-1	
3.90+0	2.17+1	7.66-1	4.00+0	2.43+1	7.93-1	4.10+0	2.61+1	7.90-1	
4.20+0	2.81+1	7.91-1	4.30+0	2.97+1	7.79-1	4.40+0	3.24+1	7.94-1	
5.20+0	5.69+1	8.61-1	1.65+1	6.62+2	8.96-1	2.70+1	1.18+3	9.35-1	83
5.00+0	4.00+1	6.76-1							89
4.00+0	2.43+1	7.93-1	6.00+0	8.13+1	8.38-1	8.00+0	1.78+2	9.03-1	97
1.00+1	2.82+2	8.86-1	1.20+1	4.11+2	9.17-1	1.60+1	6.16+2	8.69-1	
2.00+1	8.77+2	9.32-1	2.40+1	1.08+3	9.48-1	2.80+1	1.25+3	9.62-1	
3.00+1	1.30+3	9.51-1							
1.10+1	4.61+2	1.21+0	1.60+1	7.32+2	1.03+0	2.10+1	1.06+3	1.07+0	109
2.70+1	1.32+3	1.05+0							
2.00-1	6.90-6	4.12-1	2.10-1	1.10-5	4.19-1	2.20-1	1.90-5	4.79-1	120
2.40-1	4.70-5	5.72-1	2.60-1	8.50-5	5.54-1	2.80-1	1.50-4	5.66-1	
3.00-1	2.50-4	5.83-1							

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
8.00-1	9.44-2	1.19+0	1.00+0	2.42-1	1.18+0	1.20+0	5.22-1	1.21+0	133
1.40+0	9.94-1	1.26+0	1.60+0	1.61+0	1.23+0	1.80+0	2.38+0	1.17+0	
<b>29 Copper</b> <b>Fluorescence yield = 0.44</b>									
7.00-1	8.87-3	<b>2.87-1</b>	8.00-1	1.45-2	<b>2.56-1</b>	9.00-1	2.78-2	<b>2.93-1</b>	4
1.00+0	4.05-2	<b>2.73-1</b>							
2.90+0	3.60+0	<b>4.39-1</b>	3.55+0	6.40+0	<b>4.01-1</b>	3.90+0	8.70+0	<b>4.05-1</b>	14
4.80+0	1.70+1	<b>4.21-1</b>	5.30+0	2.70+1	5.03-1				
3.00+1	1.39+3	1.15+0	4.00+1	1.72+3	1.19+0	5.00+1	1.86+3	1.17+0	24
6.00+1	2.17+3	1.31+0	7.00+1	2.17+3	1.29+0	8.00+1	2.22+3	1.32+0	
2.00+0	2.30+0	1.04+0	3.00+0	8.60+0	9.37-1	4.00+0	2.30+1	9.89-1	38
5.00+0	4.30+1	9.46-1	6.00+0	7.60+1	1.00+0	7.00+0	1.10+2	9.72-1	
8.00+0	1.50+2	9.57-1	9.00+0	1.90+2	9.26-1	1.00+1	2.40+2	9.33-1	
1.10+1	3.00+2	9.62-1	1.20+1	3.40+2	9.26-1				
3.00+1	1.42+3	1.18+0	4.00+1	1.76+3	1.21+0	5.00+1	2.04+3	1.29+0	49
6.00+1	2.16+3	1.31+0	7.00+1	2.11+3	1.26+0	8.00+1	2.16+3	1.29+0	
9.00+1	2.12+3	1.28+0	1.00+2	2.06+3	1.26+0				
1.00+0	1.40-1	9.43-1	1.50+0	6.56-1	8.69-1	2.00+0	1.94+0	8.74-1	54
2.50+0	4.21+0	8.56-1	3.00+0	8.15+0	8.88-1	3.50+0	1.32+1	8.66-1	
4.00+0	1.94+1	8.34-1	4.50+0	2.77+1	8.31-1	5.00+0	3.68+1	8.09-1	
1.20+1	3.15+2	8.58-1	2.00+1	9.30+2	1.16+0				59
4.00+0	2.11+1	9.07-1							62
2.00+0	2.30+0	1.04+0							68
1.00+0	2.01-1	1.35+0	1.10+0	2.70-1	1.23+0	1.30+0	4.64-1	1.07+0	78
1.50+0	7.09-1	9.39-1	1.70+0	1.02+0	8.40-1	1.90+0	1.85+0	1.01+0	
2.10+0	2.81+0	1.06+0	2.20+0	3.02+0	9.65-1	2.30+0	3.38+0	9.21-1	
2.40+0	3.74+0	8.77-1	2.50+0	4.22+0	8.58-1	2.60+0	4.70+0	8.33-1	
2.70+0	5.45+0	8.49-1	2.80+0	6.12+0	8.42-1	2.90+0	6.84+0	8.35-1	
3.00+0	7.67+0	8.36-1	3.10+0	8.32+0	8.13-1	3.20+0	8.99+0	7.90-1	
3.30+0	9.80+0	7.79-1	3.40+0	1.08+1	7.79-1	3.50+0	1.21+1	7.94-1	
3.60+0	1.35+1	8.09-1	3.70+0	1.46+1	8.02-1	3.80+0	1.59+1	8.02-1	
3.90+0	1.67+1	7.77-1	4.00+0	1.85+1	7.95-1	4.10+0	1.97+1	7.85-1	
4.20+0	2.15+1	7.95-1	4.30+0	2.27+1	7.81-1	4.40+0	2.57+1	8.25-1	
9.00-1	1.60-1	1.69+0	1.00+0	1.60-1	1.08+0	1.20+0	5.10-1	1.62+0	80
1.40+0	9.17-1	1.59+0	1.60+0	1.48+0	1.53+0	1.80+0	2.36+0	1.57+0	
2.00+0	3.30+0	1.49+0	2.20+0	4.65+0	1.49+0	2.40+0	6.25+0	1.47+0	
2.60+0	8.68+0	1.54+0	2.80+0	1.11+1	1.53+0	3.00+0	1.37+1	1.49+0	
3.20+0	1.82+1	1.60+0	3.40+0	2.11+1	1.52+0	3.60+0	2.72+1	1.63+0	
3.80+0	3.22+1	1.62+0	4.00+0	3.69+1	1.59+0				
5.20+0	6.40+1	1.26+0	1.65+1	6.61+2	1.06+0	2.70+1	1.08+3	9.80-1	83

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
6.00+0	7.42+1	9.79-1	9.00+0	1.92+2	9.36-1	1.20+1	3.67+2	1.00+0	94
1.50+1	5.55+2	1.03+0	1.80+1	7.22+2	1.03+0				
7.10-1	3.31-2	1.00+0	9.10-1	9.07-2	9.11-1	1.11+0	1.93-1	8.44-1	99
1.31+0	3.79-1	8.51-1	1.41+0	4.92-1	8.29-1	1.50+0	6.35-1	8.41-1	
1.71+0	1.05+0	8.45-1							
3.00-1	4.83-5	<b>1.84-1</b>	4.00-1	6.19-4	3.65-1	5.00-1	2.81-3	4.76-1	103
6.00-1	9.10-3	6.09-1	7.00-1	2.17-2	7.02-1	8.00-1	4.84-2	8.54-1	
1.00+1	2.36+2	9.18-1	1.20+1	3.65+2	9.94-1	1.40+1	4.90+2	1.02+0	105
1.60+1	6.36+2	1.07+0	1.80+1	8.14+2	1.16+0	2.00+1	8.37+2	1.04+0	
2.20+1	9.75+2	1.08+0	2.40+1	1.09+3	1.10+0	2.60+1	1.18+3	1.10+0	
2.80+1	1.34+3	1.17+0	3.00+1	1.42+3	1.18+0				
1.10+1	3.68+2	1.18+0	1.60+1	7.22+2	1.21+0	2.10+1	9.32+2	1.09+0	109
2.70+1	1.20+3	1.09+0							
3.50-1	2.54-4	3.41-1	4.00-1	7.52-4	4.43-1	4.50-1	1.84-3	5.52-1	118
5.00-1	3.69-3	6.25-1	6.00-1	1.12-2	7.50-1	7.00-1	2.62-2	8.47-1	
8.00-1	4.84-2	8.54-1							
3.20-1	1.58-4	3.83-1	3.60-1	5.83-4	6.54-1	4.00-1	1.12-3	6.60-1	126
4.80-1	3.92-3	8.26-1	5.60-1	9.99-3	9.43-1	6.40-1	2.07-2	1.01+0	
7.20-1	3.69-2	1.05+0	8.00-1	5.81-2	1.02+0	1.00+0	1.42-1	9.57-1	
1.20+0	2.73-1	8.70-1							
8.00-1	6.29-2	1.11+0	1.00+0	1.67-1	1.12+0	1.20+0	3.50-1	1.12+0	133
1.40+0	6.07-1	1.05+0	1.60+0	1.00+0	1.03+0	1.80+0	1.51+0	1.00+0	
8.00-1	7.96-2	1.40+0	9.00-1	1.26-1	1.33+0	1.00+0	2.16-1	1.45+0	134
1.10+0	3.14-1	1.42+0	1.20+0	4.49-1	1.43+0	1.30+0	6.34-1	1.47+0	
1.40+0	7.88-1	1.36+0	1.50+0	1.02+0	1.35+0	1.60+0	1.19+0	1.23+0	
1.70+0	1.42+0	1.17+0	1.80+0	1.70+0	1.13+0				
1.50+0	9.60-1	1.27+0	2.02+0	3.15+0	1.37+0	2.50+0	6.55+0	1.33+0	148
3.00+0	1.23+1	1.34+0	3.50+0	2.03+1	1.33+0				
7.00-1	3.24-2	1.05+0	8.00-1	6.59-2	1.16+0	9.00-1	1.01-1	1.06+0	160
1.00+0	1.72-1	1.16+0	1.20+0	3.64-1	1.16+0	1.40+0	6.59-1	1.14+0	
1.60+0	1.03+0	1.07+0	2.00+0	2.27+0	1.02+0	2.40+0	4.13+0	9.69-1	
<b>30</b>	Zinc	Fluorescence yield = 0.47							
1.16+0	2.17-1	1.09+0	1.36+0	4.03-1	1.07+0	1.55+0	6.60-1	1.05+0	67
1.76+0	1.03+0	1.01+0	1.96+0	1.54+0	1.01+0	2.16+0	2.12+0	9.75-1	
2.36+0	2.96+0	9.91-1	2.46+0	3.55+0	1.02+0				
1.00+0	1.04-1	9.72-1	2.20+0	2.11+0	9.07-1	2.30+0	2.38+0	8.73-1	78
2.40+0	2.68+0	8.45-1	2.50+0	3.05+0	8.32-1	2.60+0	3.39+0	8.06-1	
2.70+0	3.94+0	8.21-1	2.80+0	4.44+0	8.16-1	2.90+0	5.01+0	8.17-1	
3.00+0	5.58+0	8.11-1	3.10+0	6.24+0	8.12-1	3.20+0	6.42+0	7.51-1	
3.30+0	7.11+0	7.51-1	3.40+0	7.74+0	7.40-1	3.50+0	8.44+0	7.35-1	

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
3.60+0	9.42+0	7.48-1	3.70+0	1.04+1	7.55-1	3.80+0	1.13+1	7.54-1	
3.90+0	1.14+1	7.00-1	4.00+0	1.23+1	6.97-1	4.10+0	1.43+1	7.50-1	
4.20+0	1.54+1	7.49-1	4.30+0	1.65+1	7.46-1	4.40+0	1.81+1	7.63-1	
<b>31 Gallium</b> <b>Fluorescence yield = 0.507</b>									
1.29+0	2.03-1	9.08-1	1.49+0	3.66-1	9.22-1	1.69+0	6.11-1	9.46-1	67
1.90+0	9.41-1	9.35-1	2.10+0	1.36+0	9.32-1	2.30+0	1.91+0	9.40-1	
2.41+0	2.21+0	9.20-1							
1.10+1	2.45+2	1.20+0	1.60+1	6.37+2	1.54+0	2.10+1	8.29+2	1.34+0	109
2.70+1	1.18+3	1.41+0							
<b>32 Germanium</b> <b>Fluorescence yield = 0.535</b>									
1.00+0	6.76-2	1.21+0	1.10+0	9.08-2	1.08+0	1.30+0	1.61-1	9.57-1	78
1.50+0	2.62-1	8.77-1	1.70+0	3.89-1	8.00-1	1.90+0	6.30-1	8.47-1	
2.10+0	1.05+0	9.72-1	2.20+0	1.11+0	8.65-1	2.30+0	1.30+0	8.62-1	
2.40+0	1.48+0	8.41-1	2.50+0	1.66+0	8.14-1	2.60+0	1.83+0	7.81-1	
2.70+0	2.17+0	8.10-1	2.80+0	2.42+0	7.94-1	2.90+0	2.73+0	7.93-1	
3.00+0	2.96+0	7.65-1	3.10+0	3.28+0	7.56-1	3.20+0	3.68+0	7.62-1	
3.30+0	4.08+0	7.61-1	3.40+0	4.54+0	7.66-1	3.50+0	5.17+0	7.91-1	
3.60+0	5.56+0	7.75-1	3.70+0	5.91+0	7.53-1	3.80+0	6.49+0	7.57-1	
3.90+0	6.92+0	7.42-1	4.00+0	7.59+0	7.50-1	4.10+0	8.04+0	7.33-1	
4.20+0	8.72+0	7.36-1	4.30+0	9.17+0	7.18-1	4.40+0	1.00+1	7.28-1	
5.20+0	3.47+1	1.51+0	1.65+1	4.47+2	1.24+0	2.70+1	7.75+2	1.08+0	83
8.00-1	4.19-2	<b>2.05+0</b>	9.00-1	5.99-2	1.71+0	1.00+0	8.40-2	1.51+0	134
1.10+0	1.11-1	1.32+0	1.20+0	1.63-1	1.35+0	1.30+0	2.21-1	1.31+0	
1.40+0	2.88-1	1.27+0	1.50+0	3.65-1	1.22+0	1.60+0	4.36-1	1.13+0	
1.70+0	5.72-1	1.18+0	1.80+0	7.06-1	1.17+0				
8.00-1	2.49-2	1.22+0	1.00+0	6.87-2	1.23+0	1.20+0	1.50-1	1.24+0	160
1.40+0	2.76-1	1.22+0	1.60+0	4.53-1	1.18+0	2.00+0	1.03+0	1.14+0	
2.40+0	1.87+0	1.06+0							
<b>33 Arsenic</b> <b>Fluorescence yield = 0.562</b>									
1.36+0	1.13-1	7.61-1	1.57+0	2.21-1	8.37-1	1.77+0	3.54-1	8.41-1	67
1.87+0	4.62-1	8.89-1	1.97+0	5.76-1	9.09-1	2.07+0	6.78-1	8.89-1	
2.17+0	8.66-1	9.53-1	2.37+0	1.19+0	9.46-1	2.47+0	1.40+0	9.59-1	
<b>34 Selenium</b> <b>Fluorescence yield = 0.589</b>									
1.18+0	6.10-2	1.01+0	1.37+0	1.13-1	9.98-1	1.58+0	1.92-1	9.53-1	67
1.78+0	3.04-1	9.45-1	1.98+0	4.95-1	1.02+0				
1.00+0	3.03-2	1.03+0	1.10+0	4.82-2	1.08+0	1.30+0	7.36-2	8.08-1	78
1.50+0	1.21-1	7.40-1	1.70+0	1.86-1	6.92-1	1.90+0	2.95-1	7.14-1	
2.10+0	3.86-1	6.39-1	2.20+0	6.65-1	9.25-1	2.30+0	8.01-1	9.46-1	
2.40+0	8.87-1	8.95-1	2.50+0	9.43-1	8.20-1	2.60+0	1.12+0	8.45-1	
2.70+0	1.30+0	8.57-1	2.80+0	1.48+0	8.56-1	2.90+0	1.62+0	8.28-1	
3.00+0	1.76+0	7.98-1	3.10+0	1.89+0	7.64-1	3.20+0	2.11+0	7.64-1	

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
3.30+0	2.62+0	8.53-1	3.40+0	3.15+0	9.26-1	3.50+0	2.84+0	7.56-1	
3.60+0	3.41+0	8.25-1	3.70+0	3.56+0	7.86-1	3.80+0	3.87+0	7.81-1	
3.90+0	4.26+0	7.88-1	4.00+0	4.92+0	8.37-1	4.10+0	5.08+0	7.97-1	
4.20+0	5.90+0	8.56-1	4.30+0	5.76+0	7.74-1	4.40+0	5.81+0	7.24-1	
1.00+1	8.94+1	1.07+0	1.20+1	1.25+2	9.78-1	1.40+1	1.79+2	1.01+0	105
1.60+1	2.44+2	1.06+0	1.80+1	3.22+2	1.13+0	2.00+1	3.70+2	1.09+0	
2.20+1	4.41+2	1.12+0	2.40+1	5.09+2	1.14+0	2.60+1	5.78+2	1.16+0	
2.80+1	6.20+2	1.14+0	3.00+1	6.79+2	1.15+0				
1.50+0	1.32-1	8.07-1	2.02+0	3.94-1	7.55-1	2.50+0	8.25-1	7.17-1	148
3.00+0	1.56+0	7.08-1	3.20+0	1.99+0	7.20-1	3.50+0	2.67+0	7.10-1	
3.60+0	2.85+0	6.90-1	3.70+0	3.04+0	6.71-1				
<b>35</b>	Bromine		Fluorescence yield = 0.615						
1.10+1	9.68+1	1.15+0	1.60+1	2.42+2	1.28+0	2.10+1	3.76+2	1.22+0	109
2.70+1	4.99+2	1.12+0							
<b>37</b>	Rubidium		Fluorescence yield = 0.667						
1.17+0	2.00-2	8.35-1	1.37+0	3.80-2	8.01-1				67
1.00+0	2.00-2	1.71+0	1.20+0	3.80-2	1.42+0	1.40+0	5.50-2	1.06+0	72
1.60+0	7.30-2	8.07-1	1.80+0	1.10-1	7.58-1	2.00+0	1.90-1	8.67-1	
2.10+0	2.20-1	8.32-1	2.20+0	2.50-1	7.92-1	2.40+0	3.80-1	8.67-1	
3.00+0	9.20-1	9.28-1	4.00+0	2.80+0	1.04+0	5.00+0	6.00+0	1.06+0	
6.00+0	1.20+1	1.20+0	7.00+0	1.70+1	1.07+0	8.00+0	2.30+1	9.81-1	
9.00+0	3.20+1	9.85-1	1.00+1	4.40+1	1.02+0	1.10+1	5.50+1	1.01+0	
1.20+1	7.00+1	1.03+0							
1.50+0	5.49-2	7.91-1	2.02+0	1.85-1	8.12-1	2.50+0	4.10-1	8.03-1	148
3.00+0	8.20-1	8.27-1	3.50+0	1.48+0	8.67-1				
<b>39</b>	Yttrium		Fluorescence yield = 0.71						
1.00+0	4.30-3	6.67-1	1.50+0	3.18-2	7.98-1	2.00+0	1.13-1	8.79-1	54
2.50+0	2.78-1	9.20-1	3.00+0	5.68-1	9.60-1	3.50+0	1.00+0	9.74-1	
3.75+0	1.40+0	1.07+0	4.00+0	1.71+0	1.05+0	4.25+0	2.46+0	1.23+0	
4.50+0	2.56+0	1.05+0							
1.20+1	3.60+1	8.15-1	2.00+1	1.09+2	7.92-1				59
3.00+0	7.80-1	1.32+0	4.00+0	2.10+0	1.29+0	5.00+0	4.20+0	1.22+0	95
6.00+0	7.10+0	1.15+0	8.00+0	1.60+1	1.08+0	1.00+1	3.10+1	1.12+0	
1.20+1	5.30+1	1.20+0							
1.00+1	2.63+1	9.53-1	1.20+1	4.58+1	1.04+0	1.40+1	6.40+1	9.97-1	105
1.60+1	9.24+1	1.06+0	1.80+1	1.21+2	1.08+0	2.00+1	1.49+2	1.08+0	
2.20+1	1.92+2	1.17+0	2.40+1	2.28+2	1.19+0	2.60+1	2.60+2	1.19+0	
2.80+1	2.77+2	1.13+0	3.00+1	3.06+2	1.12+0				

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
<b>40 Zirconium      Fluorescence yield = 0.73</b>									
3.00+1	2.27+2	9.78-1	4.00+1	3.24+2	9.47-1	5.00+1	4.23+2	9.77-1	24
6.00+1	5.51+2	1.09+0	7.00+1	5.99+2	1.07+0	8.00+1	6.42+2	1.06+0	
2.00+1	1.20+2	1.05+0							42
1.50+0	2.20-2	7.22-1	2.02+0	8.20-2	7.94-1	2.50+0	1.90-1	8.09-1	148
3.00+0	3.67-1	7.95-1	3.50+0	6.40-1	7.97-1				
<b>41 Niobium      Fluorescence yield = 0.74</b>									
3.00+1	1.70+2	8.61-1	4.00+1	2.37+2	8.02-1	5.00+1	3.23+2	8.52-1	24
6.00+1	3.96+2	8.87-1	7.00+1	4.75+2	9.50-1	8.00+1	5.20+2	9.61-1	
1.02+0	5.40-3	<b>1.36+0</b>	1.23+0	1.10-2	1.14+0	1.44+0	2.30-2	1.17+0	87
1.66+0	4.30-2	1.20+0	1.87+0	8.50-2	1.45+0	2.08+0	1.00-1	1.11+0	
2.29+0	1.40-1	1.07+0	2.49+0	2.00-1	1.11+0	2.70+0	2.80-1	1.15+0	
6.40-1	1.40-4	4.92-1	8.00-1	1.14-3	1.04+0	9.60-1	2.97-3	1.02+0	126
1.12+0	7.37-3	1.18+0	1.28+0	1.44-2	1.24+0	1.44+0	2.57-2	1.31+0	
1.60+0	4.04-2	1.31+0	1.80+0	7.02-2	1.40+0				
2.02+0	6.20-2	7.75-1	2.50+0	1.46-1	7.98-1	3.00+0	3.00-1	8.31-1	148
3.50+0	5.03-1	7.99-1							
<b>42 Molybdenum      Fluorescence yield = 0.765</b>									
2.90+0	8.20-1	<b>3.27+0</b>	5.30+0	2.10+0	1.01+0				14
2.90+0	7.90-1	<b>3.15+0</b>	3.55+0	1.20+0	<b>2.29+0</b>	3.90+0	1.40+0	1.92+0	15
4.45+0	1.60+0	1.39+0	4.80+0	1.80+0	1.21+0	5.30+0	2.00+0	9.65-1	
1.50+0	1.10-2	6.09-1	2.00+0	3.40-2	5.66-1	2.50+0	6.90-2	<b>4.80-1</b>	43
3.00+0	1.40-1	<b>4.92-1</b>	3.50+0	2.60-1	5.23-1	4.00+0	3.70-1	<b>4.64-1</b>	
1.00+0	2.00-3	7.41-1	1.50+0	1.83-2	1.01+0	2.00+0	5.81-2	9.68-1	54
2.50+0	1.43-1	9.95-1	3.00+0	2.93-1	1.03+0	3.75+0	6.40-1	1.01+0	
4.00+0	8.86-1	1.11+0	4.50+0	1.38+0	1.15+0				
1.02+0	3.30-3	1.11+0	1.23+0	8.00-3	1.08+0	1.44+0	1.70-2	1.13+0	87
1.66+0	3.10-2	1.11+0	1.87+0	4.90-2	1.07+0	2.08+0	7.80-2	1.11+0	
2.29+0	1.10-1	1.07+0	2.49+0	1.60-1	1.13+0				
3.00+0	3.30-1	1.16+0	4.00+0	9.20-1	1.15+0	5.00+0	1.80+0	1.05+0	95
6.00+0	3.40+0	1.10+0	8.00+0	7.60+0	1.00+0	1.00+1	1.50+1	1.04+0	
1.20+1	2.30+1	9.66-1							
1.00+0	2.56-3	9.49-1	1.10+0	3.52-3	8.09-1	1.20+0	5.71-3	8.63-1	134
1.30+0	9.18-3	9.63-1	1.40+0	1.19-2	8.93-1	1.50+0	1.61-2	8.91-1	
1.60+0	2.67-2	1.12+0	1.70+0	3.46-2	1.12+0	1.80+0	3.85-2	9.85-1	
2.02+0	4.90-2	7.84-1	2.50+0	1.13-1	7.86-1	3.00+0	2.32-1	8.16-1	148
3.50+0	4.00-1	8.04-1	3.80+0	5.50-1	8.26-1				

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
<b>45 Rhodium</b> <b>Fluorescence yield = 0.808</b>									
5.20+1	2.10+2	9.01-1							8
3.00+1	9.48+1	9.09-1	4.00+1	1.47+2	8.86-1	5.00+1	2.02+2	9.08-1	24
6.00+1	2.69+2	9.91-1	7.00+1	3.29+2	1.05+0	8.00+1	3.78+2	1.09+0	
1.02+0	1.20-3	9.30-1	1.23+0	3.30-3	9.80-1	1.44+0	6.60-3	9.36-1	87
1.66+0	1.30-2	9.78-1	1.87+0	2.20-2	9.97-1	2.08+0	3.70-2	1.08+0	
2.29+0	4.50-2	8.93-1	2.49+0	7.40-2	1.06+0	2.70+0	9.90-2	1.04+0	
<b>46 Palladium</b> <b>Fluorescence yield = 0.82</b>									
3.00+1	6.26+1	7.03-1	4.00+1	1.02+2	7.13-1	5.00+1	1.53+2	7.87-1	24
6.00+1	2.03+2	8.47-1	7.00+1	2.56+2	9.21-1	8.00+1	2.89+2	9.31-1	
1.50+0	3.98-3	5.98-1	2.02+0	1.99-2	8.25-1	2.50+0	4.99-2	8.79-1	148
3.00+0	1.04-1	9.13-1	3.20+0	1.35-1	9.33-1	3.50+0	1.91-1	9.50-1	
3.60+0	2.11-1	9.47-1							
<b>47 Silver</b> <b>Fluorescence yield = 0.831</b>									
2.70+1	5.60+1	9.05-1	3.40+1	8.40+1	8.87-1	3.90+1	1.01+2	8.49-1	8
4.40+1	1.11+2	7.78-1	5.20+1	1.35+2	7.57-1				
3.00+1	6.52+1	8.60-1	4.00+1	1.02+2	8.25-1	5.00+1	1.41+2	8.30-1	24
6.00+1	2.16+2	1.02+0	7.00+1	2.50+2	1.01+0	8.00+1	2.96+2	1.07+0	
3.00+0	1.00-1	1.09+0	4.00+0	3.10-1	1.18+0	5.00+0	7.70-1	1.35+0	38
6.00+0	1.30+0	1.23+0	7.00+0	2.20+0	1.27+0	8.00+0	3.30+0	1.25+0	
9.00+0	4.60+0	1.21+0	1.00+1	6.30+0	1.21+0	1.10+1	7.80+0	1.14+0	
1.20+1	9.90+0	1.13+0							
1.50+0	2.01-3	<b>3.84-1</b>	2.00+0	9.17-3	4.98-1	2.50+0	2.82-2	6.20-1	54
3.00+0	6.46-2	7.06-1	3.50+0	1.20-1	7.41-1	4.00+0	1.66-1	6.33-1	
4.50+0	2.54-1	6.40-1	5.00+0	4.03-1	7.06-1				
5.20+0	9.10-1	1.39+0	1.65+1	2.14+1	1.05+0	2.70+1	5.11+1	8.26-1	83
8.00-1	2.10-4	<b>1.22+0</b>	1.00+0	5.40-4	8.08-1	1.20+0	1.50-3	8.42-1	87
1.40+0	3.60-3	9.45-1	1.60+0	6.30-3	8.98-1	1.80+0	1.20-2	1.02+0	
2.00+0	2.00-2	1.09+0	2.20+0	2.60-2	9.53-1	2.40+0	4.30-2	1.11+0	
3.00+0	1.00-1	1.09+0	4.00+0	2.70-1	1.03+0	5.00+0	5.90-1	1.03+0	95
6.00+0	1.10+0	1.04+0	8.00+0	2.60+0	9.82-1	1.00+1	5.10+0	9.80-1	
1.20+1	8.30+0	9.45-1							
1.11+0	6.13-4	5.17-1	1.31+0	1.55-3	5.61-1	1.41+0	2.21-3	5.61-1	99
1.50+0	3.14-3	6.00-1	1.71+0	6.25-3	6.63-1				
1.10+1	7.72+0	1.13+0	1.60+1	1.89+1	9.99-1	2.10+1	3.46+1	9.54-1	109
2.70+1	5.97+1	9.65-1							

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
3.00+0	4.80-2	9.83-1	4.00+0	1.50-1	1.06+0	5.00+0	3.20-1	1.03+0	95
6.00+0	5.80-1	1.01+0	8.00+0	1.50+0	1.03+0	1.00+1	2.80+0	9.62-1	
1.20+1	4.60+0	9.25-1							
1.40+0	1.11-4	<b>5.95-2</b>	1.50+0	3.81-4	<b>1.46-1</b>	1.58+0	6.94-4	<b>2.07-1</b>	111
1.60+0	8.84-4	<b>2.49-1</b>	1.62+0	9.36-4	<b>2.50-1</b>	1.64+0	9.97-4	<b>2.52-1</b>	
1.68+0	1.37-3	<b>3.10-1</b>	1.70+0	1.39-3	<b>2.98-1</b>	1.80+0	1.92-3	<b>3.19-1</b>	
1.90+0	3.05-3	<b>3.99-1</b>	2.00+0	3.41-3	<b>3.57-1</b>	2.10+0	4.52-3	<b>3.85-1</b>	
2.20+0	5.30-3	<b>3.72-1</b>	2.30+0	6.56-3	<b>3.83-1</b>	2.40+0	7.59-3	<b>3.73-1</b>	
2.60+0	1.16-2	<b>4.14-1</b>	2.80+0	1.50-2	<b>4.01-1</b>				
9.00+0	2.68+0	1.27+0	1.55+1	1.16+1	1.15+0	1.65+1	1.37+1	1.15+0	128
1.75+1	1.59+1	1.15+0	1.80+1	1.72+1	1.16+0	1.85+1	1.80+1	1.13+0	
1.90+1	1.98+1	1.16+0	1.95+1	2.09+1	1.16+0	2.05+1	2.45+1	1.20+0	
2.15+1	2.65+1	1.16+0	2.25+1	2.90+1	1.14+0	9.00+1	2.35+2	1.08+0	
1.00+2	2.57+2	1.09+0	1.10+2	2.69+2	1.07+0	1.20+2	2.62+2	9.93-1	
1.30+2	3.01+2	1.10+0	1.40+2	3.04+2	1.07+0	1.55+2	3.08+2	1.05+0	
2.40+0	1.11-2	5.45-1	2.60+0	1.59-2	5.68-1	2.80+0	2.16-2	5.77-1	138
2.50+0	1.90-2	7.93-1	3.00+0	3.90-2	7.99-1	3.50+0	7.20-2	8.27-1	148
<b>51</b>	Antimony		Fluorescence yield = 0.87						
1.20+0	4.70-4	7.15-1	1.40+0	1.30-3	8.79-1	1.60+0	2.50-3	8.80-1	87
1.80+0	4.60-3	9.48-1	2.00+0	7.30-3	9.46-1	2.20+0	1.20-2	1.04+0	
2.40+0	1.70-2	1.03+0							
8.00-1	2.03-5	4.01-1	1.00+0	1.38-4	6.07-1	1.20+0	4.46-4	6.78-1	126
1.40+0	1.18-3	7.98-1	1.60+0	2.50-3	8.80-1	1.80+0	4.85-3	1.00+0	
2.00+0	7.59-3	9.84-1	2.20+0	1.17-2	1.01+0	2.40+0	1.70-2	1.03+0	
<b>52</b>	Tellurium		Fluorescence yield = 0.877						
3.00+1	3.00+1	8.70-1	4.00+1	5.43+1	9.04-1	5.00+1	7.90+1	9.16-1	49
6.00+1	1.02+2	9.14-1	7.00+1	1.40+2	1.04+0	8.00+1	1.56+2	1.00+0	
2.20+0	7.97-3	8.47-1	2.40+0	1.02-2	7.55-1	2.60+0	1.45-2	7.77-1	138
2.80+0	1.71-2	6.83-1							
<b>53</b>	Iodine		Fluorescence yield = 0.884						
1.00+0	6.60-5	4.95-1	1.20+0	2.50-4	6.18-1	1.40+0	8.10-4	8.64-1	87
1.60+0	1.70-3	9.27-1	1.80+0	2.60-3	8.18-1	2.00+0	4.30-3	8.44-1	
2.20+0	6.50-3	8.45-1	2.40+0	1.00-2	9.04-1				
2.40+0	1.11-2	1.00+0	2.60+0	1.48-2	9.65-1				138
<b>55</b>	Cesium		Fluorescence yield = 0.897						
1.10+1	8.60-1	5.50-1	1.60+1	3.03+0	6.58-1	2.10+1	7.84+0	8.32-1	109
2.70+1	1.21+1	7.02-1							

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
<b>56 Barium</b> <b>Fluorescence yield = 0.902</b>									
1.40+0	2.50-4	5.19-1	1.60+0	5.00-4	5.14-1	1.80+0	1.00-3	5.76-1	87
2.00+0	1.80-3	6.40-1	2.20+0	3.00-3	6.99-1	2.40+0	4.90-3	7.86-1	
2.40+0	4.23-3	6.79-1	2.60+0	5.71-3	6.57-1				138
<b>58 Cerium</b> <b>Fluorescence yield = 0.912</b>									
5.20+1	5.00+1	1.20+0							8
1.40+0	1.10-4	3.52-1	1.60+0	3.10-4	4.79-1	1.80+0	5.90-4	5.02-1	87
2.00+0	1.20-3	6.22-1	2.20+0	2.00-3	6.74-1	2.40+0	3.00-3	6.92-1	
<b>59 Praseodymium</b> <b>Fluorescence yield = 0.917</b>									
3.00+0	5.60-3	6.12-1	4.00+0	2.10-2	7.65-1	5.00+0	4.80-2	7.87-1	95
6.00+0	9.20-2	8.05-1	8.00+0	2.30-1	7.77-1	1.00+1	4.90-1	8.18-1	
1.20+1	8.10-1	7.76-1							
<b>60 Neodymium</b> <b>Fluorescence yield = 0.921</b>									
2.00+1	2.60+0	6.84-1							42
3.00+1	6.50+0	6.24-1	4.00+1	1.58+1	8.09-1	5.00+1	2.89+1	9.61-1	49
6.00+1	3.37+1	8.23-1	7.00+1	5.01+1	9.65-1	8.00+1	6.61+1	1.06+0	
1.20+1	8.00-1	8.97-1	2.00+1	3.40+0	8.95-1				59
<b>62 Samarium</b> <b>Fluorescence yield = 0.929</b>									
2.00+0	4.40-4	4.65-1	2.50+0	1.30-3	4.96-1	3.00+0	2.10-3	<b>3.73-1</b>	43
3.50+0	3.60-3	<b>3.48-1</b>	4.00+0	5.30-3	<b>3.10-1</b>				
3.00+0	4.30-3	7.64-1	4.00+0	1.50-2	8.77-1	5.00+0	3.40-2	8.90-1	95
6.00+0	6.40-2	8.93-1	8.00+0	1.60-1	8.63-1	1.00+1	3.20-1	8.51-1	
1.20+1	5.10-1	7.77-1							
<b>64 Gadolinium</b> <b>Fluorescence yield = 0.935</b>									
3.00+1	4.80+0	8.03-1	4.00+1	8.80+0	7.66-1	5.00+1	1.76+1	9.70-1	49
6.00+1	2.72+1	1.07+0	7.00+1	3.28+1	1.00+0	8.00+1	3.98+1	9.95-1	
1.20+0	1.30-5	4.23-1	1.40+0	4.40-5	4.94-1	1.60+0	1.00-4	4.96-1	87
1.80+0	1.90-4	4.88-1	2.00+0	3.10-4	4.62-1	2.20+0	4.70-4	<b>4.43-1</b>	
1.33+1	7.47-1	1.12+0	1.52+1	1.12+0	1.14+0	1.83+1	1.83+0	1.10+0	140
2.03+1	2.41+0	1.09+0	2.50+1	4.55+0	1.20+0				
3.49+0	6.54-3	8.66-1	4.54+0	2.02-2	1.00+0	5.40+0	4.93-2	1.33+0	141
6.50+0	7.35-2	1.05+0	8.00+0	1.48-1	1.07+0	9.54+0	2.65-1	1.10+0	
1.10+1	3.81-1	1.01+0	1.30+1	6.68-1	1.07+0	1.51+1	1.01+0	1.06+0	

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
<b>67 Holmium</b>					Fluorescence yield = 0.944				
4.00+1	8.58+0	1.10+0	5.00+1	1.55+1	1.24+0	6.00+1	1.77+1	9.95-1	90
8.00+1	2.94+1	1.02+0		1.10+2	4.25+1	9.53-1			
3.00+0	1.80-3	6.75-1	4.00+0	8.60-3	1.04+0	5.00+0	2.00-2	1.07+0	95
6.00+0	3.50-2	9.97-1	8.00+0	7.70-2	8.48-1	1.00+1	1.70-1	9.25-1	
1.20+1	2.70-1	8.41-1							
9.00+0	1.71-1	1.29+0	1.55+1	8.02-1	1.18+0	1.65+1	9.52-1	1.17+0	128
1.75+1	1.15+0	1.19+0	1.85+1	1.30+0	1.15+0	1.95+1	1.53+0	1.17+0	
2.05+1	1.79+0	1.20+0	2.15+1	1.98+0	1.16+0	2.25+1	2.24+0	1.17+0	
<b>69 Thulium</b>					Fluorescence yield = 0.949				
3.00+1	1.90+0	6.14-1	4.00+1	4.20+0	6.89-1	5.00+1	7.50+0	7.60-1	49
6.00+1	1.18+1	8.35-1	7.00+1	1.72+1	9.22-1	8.00+1	2.00+1	8.62-1	
1.20+1	3.00-1	1.22+0	2.00+1	1.09+0	1.02+0				59
4.00+1	6.18+0	1.01+0	5.00+1	9.94+0	1.01+0	6.00+1	1.43+1	1.01+0	90
8.00+1	2.33+1	1.00+0	1.10+2	3.53+1	9.65-1				
9.00+0	1.31-1	1.29+0	1.55+1	5.83-1	1.12+0	1.65+1	6.89-1	1.10+0	128
1.75+1	7.75-1	1.05+0	1.85+1	9.41-1	1.09+0	1.95+1	1.12+0	1.12+0	
2.05+1	1.29+0	1.12+0	2.15+1	1.54+0	1.18+0	2.25+1	1.68+0	1.14+0	
<b>71 Lutetium</b>					Fluorescence yield = 0.953				
4.00+1	5.04+0	1.05+0	5.00+1	8.71+0	1.12+0	6.00+1	1.24+1	1.10+0	90
8.00+1	2.00+1	1.06+0	1.10+2	3.17+1	1.05+0				
<b>72 Hafnium</b>					Fluorescence yield = 0.955				
3.00+0	1.10-3	8.20-1	4.00+0	4.30-3	9.90-1	5.00+0	1.00-2	1.01+0	95
6.00+0	1.90-2	1.02+0	8.00+0	4.70-2	9.80-1	1.00+1	9.50-2	9.84-1	
1.20+1	1.60-1	9.52-1							
<b>73 Tantalum</b>					Fluorescence yield = 0.957				
5.20+1	5.60+0	8.34-1							8
3.00+1	1.60+0	8.49-1	4.00+1	3.00+0	7.94-1	5.00+1	5.60+0	9.04-1	49
6.00+1	8.50+0	9.45-1	7.00+1	1.38+1	1.15+0	8.00+1	1.40+1	9.19-1	
4.00+1	3.87+0	1.02+0	6.00+1	8.76+0	9.74-1	8.00+1	1.51+1	9.92-1	90
1.10+2	2.45+1	9.90-1							
3.00+0	6.70-4	5.70-1	4.00+0	3.30-3	8.59-1	5.00+0	8.50-3	9.68-1	95
6.00+0	1.60-2	9.67-1	8.00+0	4.00-2	9.39-1	1.00+1	8.20-2	9.58-1	
1.20+1	1.40-1	9.40-1							

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	$E_1$ (MeV)	$\sigma^{\text{Exper}}$ (barn)	$\sigma^{\text{Exper}}$ $\sigma^{\text{ECPSSR}}$	Ref.
<b>74 Tungsten</b> <b>Fluorescence yield = 0.958</b>									
5.20+1	5.40+0	9.00-1							8
4.00+1	3.82+0	1.14+0	5.00+1	7.42+0	1.34+0	6.00+1	9.10+0	1.13+0	90
8.00+1	1.49+1	1.09+0	1.10+2	2.51+1	1.12+0				
3.49+0	1.38-3	6.99-1	4.54+0	4.11-3	7.48-1	5.40+0	9.00-3	8.79-1	141
6.50+0	2.01-2	1.04+0	8.00+0	4.02-2	1.06+0	9.54+0	7.13-2	1.08+0	
1.10+1	1.13-1	1.11+0	1.30+1	1.82-1	1.08+0				
2.20+0	9.15-5	4.08-1	2.40+0	1.61-4	4.52-1	2.60+0	2.54-4	4.80-1	147
2.88+0	4.25-4	4.95-1	3.20+0	6.80-4	4.96-1	3.40+0	8.78-4	4.96-1	
3.60+0	1.07-3	4.77-1	3.80+0	1.36-3	4.88-1	4.00+0	1.67-3	4.90-1	
<b>75 Rhenium</b> <b>Fluorescence yield = 0.959</b>									
4.00+0	3.00-3	9.92-1	5.00+0	7.10-3	1.02+0	6.00+0	1.30-2	9.90-1	95
8.00+0	3.40-2	1.01+0	1.00+1	6.50-2	9.59-1	1.20+1	1.05-1	8.93-1	
<b>78 Platinum</b> <b>Fluorescence yield = 0.963</b>									
5.20+1	3.10+0	8.06-1							8
4.00+1	1.97+0	9.25-1	6.00+1	5.45+0	1.05+0	8.00+1	9.24+0	1.02+0	90
1.10+2	1.55+1	1.01+0							
3.00+0	2.60-4	4.19-1	4.00+0	1.40-3	6.53-1	5.00+0	3.90-3	7.80-1	95
6.00+0	8.70-3	9.18-1	8.00+0	2.50-2	1.03+0	1.00+1	4.70-2	9.66-1	
1.20+1	8.20-2	9.75-1							
<b>79 Gold</b> <b>Fluorescence yield = 0.964</b>									
5.20+1	2.80+0	8.11-1							8
5.00+1	3.10+0	9.77-1	6.00+1	5.30+0	1.13+0	7.00+1	7.60+0	1.19+0	49
8.00+1	9.80+0	1.19+0							
4.00+1	2.04+0	1.07+0	5.00+1	3.58+0	1.13+0	6.00+1	5.24+0	1.12+0	90
8.00+1	8.50+0	1.03+0	1.10+2	1.44+1	1.03+0				
4.00+0	1.30-3	6.78-1	5.00+0	3.50-3	7.79-1	6.00+0	7.30-3	8.56-1	95
8.00+0	1.90-2	8.67-1	1.00+1	4.10-2	9.37-1	1.20+1	6.70-2	8.87-1	
9.00+0	3.73-2	1.18+0	1.65+1	2.35-1	1.24+0	1.75+1	2.70-1	1.21+0	128
1.80+1	2.92-1	1.21+0	1.85+1	2.93-1	1.13+0	1.95+1	3.74-1	1.24+0	
2.05+1	4.04-1	1.17+0	2.15+1	5.09-1	1.30+0	2.25+1	5.40-1	1.22+0	
9.00+1	1.05+1	1.04+0	1.00+2	1.19+1	9.87-1	1.10+2	1.33+1	9.54-1	
1.20+2	1.40+1	8.80-1	1.30+2	1.83+1	1.03+0	1.40+2	1.93+1	9.85-1	
1.55+2	2.06+1	9.29-1							
2.60+0	7.11-5	<b>2.66-1</b>	2.84+0	1.29-4	3.08-1	3.04+0	1.96-4	3.36-1	147
3.29+0	3.08-4	3.67-1	3.60+0	5.62-4	4.53-1	3.80+0	7.63-4	4.90-1	
4.00+0	1.10-3	5.74-1							

TABLE 5. K-shell x-ray production by helium-4 in target elements from beryllium to uranium<sup>a,b</sup>—Continued

$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	$E_1$	$\sigma^{\text{Exper}}$	$\sigma^{\text{Exper}}$	Ref.
(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	(MeV)	(barn)	$\sigma^{\text{ECPSSR}}$	
<b>82 Lead</b> Fluorescence yield = 0.967									
5.20+1	1.90+0	7.52-1							8
1.20+1	5.70-2	1.03+0	2.00+1	2.25-1	9.59-1				59
4.00+1	1.42+0	1.03+0	6.00+1	3.26+0	9.49-1	8.00+1	5.47+0	8.98-1	90
1.10+2	9.68+0	9.16-1							
4.00+0	8.80-4	6.37-1	5.00+0	2.30-3	6.97-1	6.00+0	4.80-3	7.61-1	95
8.00+0	1.40-2	8.63-1	1.00+1	3.00-2	9.28-1	1.20+1	5.00-2	9.00-1	
9.00+0	2.50-2	1.06+0	1.75+1	1.87-1	1.15+0	1.85+1	2.06-1	1.09+0	128
1.95+1	2.44-1	1.11+0	2.05+1	2.88-1	1.15+0	2.15+1	3.16-1	1.11+0	
2.25+1	3.49-1	1.08+0	5.00+1	2.36+0	1.02+0	6.00+1	3.26+0	9.49-1	
7.00+1	4.56+0	9.69-1	8.00+1	5.91+0	9.70-1	9.00+1	7.24+0	9.59-1	
1.00+2	9.78+0	1.08+0	1.10+2	1.11+1	1.05+0	1.20+2	1.19+1	9.88-1	
1.30+2	1.46+1	1.07+0	1.40+2	1.60+1	1.07+0	1.55+2	1.88+1	1.09+0	
<b>83 Bismuth</b> Fluorescence yield = 0.968									
4.00+1	1.32+0	1.06+0	6.00+1	3.04+0	9.80-1	8.00+1	4.87+0	8.81-1	90
1.10+2	8.73+0	9.05-1							
4.00+0	7.60-4	6.12-1	5.00+0	2.10-3	7.04-1	6.00+0	4.60-3	8.03-1	95
8.00+0	1.40-2	9.49-1	1.00+1	2.80-2	9.54-1	1.20+1	4.70-2	9.33-1	
1.75+1	1.30-1	8.84-1	1.85+1	1.74-1	1.02+0	1.95+1	2.59-1	1.31+0	128
2.05+1	2.95-1	1.30+0	2.15+1	2.97-1	1.15+0	2.25+1	3.21-1	1.10+0	
<b>90 Thorium</b> Fluorescence yield = 0.971									
4.00+0	3.20-4	5.39-1	5.00+0	8.90-4	5.85-1	6.00+0	2.10-3	6.97-1	95
8.00+0	6.70-3	8.47-1	1.00+1	1.60-2	1.02+0	1.20+1	2.50-2	9.32-1	
2.60+0	3.00-5	5.29-1							124
4.45+0	4.85-4	5.12-1	5.45+0	1.46-3	6.89-1	6.19+0	2.29-3	6.81-1	141
6.32+0	2.97-3	8.21-1	7.49+0	5.47-3	8.53-1	7.81+0	6.04-3	8.24-1	
8.69+0	1.08-2	1.05+0	9.15+0	1.12-2	9.31-1	9.78+0	1.82-2	1.24+0	
1.16+1	2.94-2	1.21+0	1.17+1	3.06-2	1.22+0	1.21+1	3.02-2	1.09+0	
1.37+1	4.51-2	1.16+0	1.54+1	5.92-2	1.11+0				
<b>92 Uranium</b> Fluorescence yield = 0.972									
4.00+0	2.20-4	4.58-1	5.00+0	6.90-4	5.46-1	6.00+0	1.80-3	7.11-1	95
8.00+0	6.00-3	8.93-1	1.00+1	1.20-2	8.97-1	1.20+1	2.20-2	9.66-1	

<sup>a</sup>

Cross sections and their ratios are printed in a compressed power of 10  
 $-1$   
 notation, e.g. 9.66-1 means  $9.66 \times 10^{-1}$ .

<sup>b</sup>

The ratios shown in bold print differ by more than a factor of 2 from  
 the averaged ratios and were -- as described in the text -- rejected.

This rejection criterion was applied only to the Z2 > 9 data.

**Table 6.** Number of K-shell x-ray production cross sections compiled for each target element (identified in columns by Z2) with source references of Sec. 6.2 (listed in the first column), and tabulated separately for four projectiles: protons, deuterons, helium-3, and helium-4 ions.

Z1 = 1	A1 = 1	Protons
Ref.	Z2 --> 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23	
5		10...
7		5...
9	7 8.	
10	1...	1...
11	25. 21 22.	
12	4 8.	
13	11.	
16	7.	
17	13.	
18	17.	
20		15... 15...
22		13.
23	7.	
25		4... 5...
26	15.	
29	1.	
33	13.	
34	12.	
35	16.	
36		3 3
37	7.	
38		9...
40		7.
44		8...
45	51.	
46	13 13. 7.	8.
47		3 1
48		8.
50	18.	
51	11... 11.	11.
55		9...
57		9...
58	1.	
64		16...
65		1...
66		4... 4...
69		20 20
70	11.	
71	8.	
72		9...
73		13 12
74		1...
75	9 9.	
76		1... 1... 1
77		21...
79		6...
81	19.	
82	13 8... 14.	12...

**Table 6.** Number of K-shell x-ray production cross sections compiled for each target element (identified in columns by Z2) with source references of Sec. 6.2 ( listed in the first column ), and tabulated separately for four projectiles: protons, deuterons, helium-3, and helium-4 ions.

Continued.

Ref.	Z2 -->	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	Protons	
84																						3...	
85																						14	
86																							
88																						1...	
92																							
94																						1...	
97																						6...	
98																						31 31 31 31	
101																						7...	
105																						10...	
106																						13 13 13 13	
108																						10...	
110																							
112																						9...	
113																						11 8	
116																							
117																						15...	
118																						11...	
119																							
121																						8...	
122																						7...	
123																						11 11 11... 12 12...	
132																						14...	
137																						14 14 7	
148																						5...	
149																						1...	
156																							
157																						11 11...	
Ref.	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46
1																						1...	
2																							
3																						8...	
4																							
5																							
6																						15...	
7																						5... 1...	
9																							
10																							
20																							
21																							
27																							
30																							
36	3	3	3	3	3	2																	
38																							
39																							
40																							
44																							
47																							
48																							
52	13	15	15	15	15	15																	

**Table 6.** Number of K-shell x-ray production cross sections compiled for each target element (identified in columns by Z2) with source references of Sec. 6.2 (listed in the first column), and tabulated separately for four projectiles: protons, deuterons, helium-3, and helium-4 ions.

Continued.

Protons

Ref.	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46												
53		16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16	16												
55		10	10	11	9																														
56						18																													
57		9	9	9	14	9	8																												
59						2													2																
61											9	8	8	8	8	8	8	8	8	9	8	8	8												
64												16																							
65													1																						
66						4		4																											
68													7																						
69	18	20	20	19	17	19																													
72														5																					
73	12	12	12	6	10	6																													
74						1																													
76	1	1	1	1	1	1																													
77						21	13												13																
79	6	6																																	
80	25					32																													
84		3	2	3																															
86		9				12	11										11																		
87																		9				9													
89					1																														
94	5	1	1	1	1	5		1	1		1		1		1		1		1																
95																		7	7	7															
97					6																														
98	31	31	31																																
99						15																													
103						8																													
104						3								3																					
105				10		10						10		10				10																	
106	13	13	13																																
108		9	8	6																															
111																6	12	11																	
112						9																													
113	8	8	11	11								10						2																	
114	1		1	1	1	1						1		1				1																	
115				22	22	22	22		21	22																									
117					23																														
118		11		11														9																	
120					21																							6							
121				9																															
122					7	7	7																												
125	1				1	1						1						1	1	1															
126						8													8																
130						6																													
132	14			15	15	15	15																												
135						23		21											20																
137	11			10	7	11	11																												
143					6							6																							
144																				23				23											
148																5		5	5	4														7	

**Table 6.** Number of K-shell x-ray production cross sections compiled for each target element (identified in columns by Z2) with source references of Sec. 6.2 ( listed in the first column ), and tabulated separately for four projectiles: protons, deuterons, helium-3, and helium-4 ions.

Continued.

Protons

Ref.	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46		
149																									
151																									
152	21	...	21	11	...	21																			
Ref.	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69		
1	6	...																							
4	5	...																							
5	14	...																							
6	13	...	12	...																					
30	1	...	1	...													1	...	1	...					
32	27	...																							
38	9	...																							
39	2	...																							
47	3	...	3	3	...																				
48	...					2	...																		
52	15	15	...																						
55	11	...																							
59	...	2	...													2	...						2		
60	...	8	...																						
63	8	8	...	8	8	...				6	7	...													
64	...	11	...																						
77	21	...																							
80	...	17	...																						
86	...	11	11	12	...																				
87	11	...	10	...				10	...	9	...				10	10	10	...	9	9	...				
91	...																				20	...			
94	1	1	5	1	...	5	...																		
95	7	...	7	...											7	...	6	...	6	...	6	...			
99	14	...																							
100	10	...																9	...						
103	5	...																							
104	3	...															3	...							
107	...																	1	...						
111	...	18	16	14	...																				
113	12	...																							
118	8	...																							
120	7	...																							
121	10	...																							
125	1	1	...	1	...	1	...					1	...	1	...	1	...				1	...	1	...	
126	8	...				8	...																		
127	...					18	...																		
129	...	9	...	8	...							6	...												
130	5	...																							
135	19	...	17	...																					
138	...			8	...																				
139	...															9	...								
143	...				6	...	6	6	5	...															
144	...	25	...				26	...	27	...	24	...	25	...											
145	...																		15	...					
148	7	...		4	...																				
151	5	...																							

**Table 6. Number of K-shell x-ray production cross sections compiled for each target element (identified in columns by Z2) with source references of Sec. 6.2 ( listed in the first column ), and tabulated separately for four projectiles: protons, deuterons, helium-3, and helium-4 ions.**

Continued.

Protons

Ref.	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69
154														9	...	10							9
155														6	...	7	...	15	...				14
159														8	...	9	...	9	...	15	...		
Ref.	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92
1	.....	6	.....	1	.....	4	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
2	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	2
5	.....	11	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	20	.....	.....	.....	.....	.....	.....	.....	.....	12
6	.....	3	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
30	.....	1	.....	.....	1	1	.....	1	.....	.....	.....	.....	.....	1	.....	.....	.....	.....	.....	.....	.....	.....	1
41	.....	.....	.....	.....	.....	.....	12	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
59	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	2	.....	.....	.....	.....	.....	.....	.....	.....	.....
91	.....	.....	.....	.....	.....	.....	22	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	15
93	.....	.....	.....	.....	.....	.....	16	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
95	.....	6	6	...	6	.....	6	6	.....	6	6	.....	.....	6	6	.....	.....	.....	.....	.....	.....	.....	4
100	10	.....	9	.....	.....	.....	9	.....	10	.....	.....	.....	.....	9	.....	.....	.....	.....	.....	.....	.....	.....	9
104	.....	.....	.....	.....	.....	.....	3	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
107	.....	.....	1	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	5	.....	1	.....	.....	.....	.....	.....
113	.....	.....	.....	.....	.....	.....	12	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
124	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	12	.....	.....	.....	.....	.....	.....	.....	.....	.....
125	.....	1	.....	.....	1	1	.....	1	1	.....	.....	.....	.....	1	1	.....	.....	.....	.....	.....	.....	.....	.....
139	.....	.....	15	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
142	.....	.....	16	.....	.....	13	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	10	.....	.....	.....	.....	.....	.....
145	.....	.....	.....	6	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
158	.....	15	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
159	.....	14	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
Z1 = 1	.....	A1 = 2	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
Deuterons	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****

Ref.	Z2 -->	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23		
25	.....	.....	.....	.....	.....	.....	4	.....	.....	.....	.....	.....	.....	5	.....	.....	.....	.....	.....	.....	.....		
31	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	3	.....	.....	.....	.....	.....	.....	.....		
45	.....	.....	.....	.....	.....	.....	19	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....		
97	.....	.....	.....	.....	.....	.....	12	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....		
102	.....	.....	.....	.....	.....	3	..	10	2	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....		
119	.....	7	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	8	.....	.....	.....	.....		
121	.....	.....	.....	.....	.....	.....	.....	13	13	13	13	13	13	13	13	13	13	13	13	13	13		
122	.....	.....	.....	.....	.....	.....	.....	13	13	13	13	13	13	13	13	13	13	13	13	13	13		
131	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	4	4	4	4	4	4	4	4		
133	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	6	6	6	6	6	6	6	6	6		
153	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	24	24	24	24	24	24	24	24		
Ref.	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46
4	.....	.....	5	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
97	.....	.....	12	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	10	11	10	10	10	10	10	10	10	
111	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	10	11	10	10	10	10	10	10		
120	.....	.....	14	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....		
121	.....	9	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....		
122	.....	.....	13	13	13	13	13	13	13	13	13	13	13	13	13	13	13	13	13	13	13		
126	.....	.....	10	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	8	.....	.....	.....	.....	.....	.....	.....		

**Table 6.** Number of K-shell x-ray production cross sections compiled for each target element (identified in columns by Z2) with source references of Sec. 6.2 (listed in the first column), and tabulated separately for four projectiles: protons, deuterons, helium-3, and helium-4 ions.

Continued.

	Deuterons																						
Ref.	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46
131	.....	4...	4.....	4.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
133	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	6	
153	.....	23	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
Ref.	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69
111	...	10...	8	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
121	9	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
126	9	.....	9	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
129	.....	8...	7	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
150	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	11	.....	.....	.....	.....	
Ref.	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92
147	.....	.....	15	.....	.....	14	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
158	.....	.....	11	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
Z1 = 2	A1 = 3	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	
Ref.	Z2 -->	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	Helium-3	
12	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	5	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
17	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	12	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
28	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	6	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
45	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	30	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
75	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	7	7	.....	.....	.....	.....	.....	.....	.....	.....	.....	
160	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	160	.....	.....	.....	.....	.....	19	....	.....	.....	.....	
161	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	10...	10	.....	10	.....	10	.....	14...	.....	.....	.....	
Ref.	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46
160	18	.....	9...	19	.....	9	.....	.....	.....	.....	.....	9	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
161	14	.....	14	14	.....	14	14	.....	14	13	.....	13	.....	8	.....	.....	8	.....	.....	11	.....	.....	
Ref.	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69
160	8	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
Z1 = 2	A1 = 4	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	*****	
Ref.	Z2 -->	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	Helium-4	
12	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	4	5	.....	.....	.....	.....	.....	.....	.....	.....	.....	
14	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	14	.....	.....	.....	.....	.....	4	...	.....	.....	.....	
15	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	6	6	.....	.....	.....	.....	.....	.....	.....	.....	.....	
16	.....	.....	5	.....	.....	.....	.....	.....	.....	.....	.....	16	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
17	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	17	.....	.....	8	.....	.....	.....	.....	.....	.....	.....	
19	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	19	.....	8	9	.....	.....	.....	.....	.....	.....	.....	
22	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	22	.....	8	.....	.....	.....	.....	.....	.....	.....	.....	
23	.....	.....	16	.....	.....	.....	.....	.....	.....	.....	.....	23	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
26	.....	.....	16	.....	.....	.....	.....	.....	.....	.....	.....	26	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	
28	.....	.....	.....	.....	.....	.....	.....	.....	.....	5	.....	28	.....	5	.....	.....	.....	.....	.....	.....	.....	.....	
31	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	4	.....	31	.....	4	...	4	...	4	...	4	...	
38	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	12	12	12	12	12	12	12	12	12	12	12	12	

**Table 6.** Number of K-shell x-ray production cross sections compiled for each target element (identified in columns by Z2) with source references of Sec. 6.2 (listed in the first column), and tabulated separately for four projectiles: protons, deuterons, helium-3, and helium-4 ions.

Continued.

Helium-4

Ref.	Z2 -->	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23		
42																					1...		
45																					37...		
46							2	3...			3...												
54																					9...		
67																					10...		
71							7...																
72																					21...		
78																					30 24		
82							13	15...			9...										5...		
92							9...																
94																					5...		
96															5...						5...		
97															10...								
105																					11...		
109															4	4...	4...						
110										3...					2...								
116										9...													
118																					10...		
119							6...																
123															11...	11...							
133																	6	6	6	6			
136																	3	3	3	3			
146																					4...		
148																					5...		
153																					24...		
160																					9...		
Ref.	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46
4							4...																
8																					1...		
14	5...	5...	5	5...																	2...		
15																					6...		
24		6...	6	6...											6	6...					6	6	
38		11	14	11	11...																		
42		1	1...													1...							
43					6...																6...		
49		1...	6	8...																			
54		9...		9...											10...	8...							
59				2...												2...							
62				1...																			
67	20...				8	7...	9	5...			2...												
68					1...																		
72															19...								
78	29	24	29	24	30	24...	30...	30...															
80	17...				17...																		
83					3	3...	3...																
87																	9	8...	9...				
89						1...																	
94		5...				5...																	
95																7...	7...						
96						4...																	

**Table 6.** Number of K-shell x-ray production cross sections compiled for each target element (identified in columns by Z<sub>2</sub>) with source references of Sec. 6.2 (listed in the first column), and tabulated separately for four projectiles: protons, deuterons, helium-3, and helium-4 ions.

Continued.

Helium-4

Ref.	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46
97																							
99																							
103																							
105																							
109																							
118																							
120																							
126																							
133	6	6	6	6	6	6																	
134																							
136	3																						
148	5																						
153																							
160																							
Ref.	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69
-----																							
8	5	.....	1																				
24	6	.....	6																				
38	10																						
42		1																					
43																							
49		8	..	6																			
54	8	.....	8																				
59		2	.....																				
83	3	.....	3																				
87	9	.....	8	7	..	8																	
90																							
94		5	.....																				
95	7	.....	7																				
99	5	.....																					
109	4	.....																					
111	21	..	17																				
126	10	.....	9																				
128		18	.....																				
134	7	.....																					
138	3	..	3	..	4	2	.....	2															
140																							
141																							
148	8	.....	3																				
Ref.	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92
-----																							
8		1	1	.....	1	1	.....	1	1	.....	1												
49		6	.....																				
59																							
90		5	..	4	5	.....	4	5	.....	4	4	.....	4	4	.....	4	4	.....	4	4	.....	4	4
95		7	7	..	6	.....	7	6	.....	6	6	.....	6	6	.....	6	6	.....	6	6	.....	6	6
124																							
128																							
141																							
147																							

Table 7. Contribution (in percentage) of electron capture to ionization according to the ECPSSR theory ( Refs. 15 and 16 ). Collision systems are specified by the target's atomic number Z2 plus projectile's energies per mass and atomic number Z1. Stars appear when the contribution of electron capture to ionization is less than 0.5 %; the numbers in bold print pertain to the systems for which data exist as compiled in Tables 2-5.

Z2	Energy/Mass (in MeV/u) of Hydrogen Ions ( Z1 = 1 )											
	0.01	0.02	0.04	0.08	0.10	0.20	0.40	0.80	1.00	2.00	4.00	8.00
4	<b>25</b>	<b>26</b>	<b>27</b>	<b>26</b>	<b>25</b>	<b>16</b>	<b>7</b>	<b>1</b>	<b>1</b>	*	*	*
5	10	12	13	15	15	13	8	3	2	*	*	*
6	<b>5</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>8</b>	<b>9</b>	<b>7</b>	<b>3</b>	<b>2</b>	*	*	*
7	3	<b>3</b>	<b>4</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>5</b>	<b>3</b>	2	1	*	*
8	2	<b>2</b>	<b>2</b>	<b>3</b>	<b>3</b>	4	4	3	2	1	*	*
9	2	1	1	2	2	<b>2</b>	<b>3</b>	<b>2</b>	<b>2</b>	1	*	*
10	1	1	1	1	1	<b>2</b>	<b>2</b>	<b>2</b>	<b>2</b>	1	*	*
11	1	1	1	1	1	1	1	1	1	1	*	*
12	1	1	1	1	1	1	1	1	1	1	*	*
13	<b>1</b>	*	*	*	*	*	*	1	1	1	*	*
14	1	*	*	*	*	*	*	*	1	1	*	*
15	1	*	*	*	*	*	*	*	1	1	*	*
Z2	Energy/Mass (in MeV/u) of Helium Ions ( Z1 = 2 )											
	0.01	0.02	0.04	0.08	0.10	0.20	0.40	0.80	1.00	2.00	4.00	8.00
4	<b>99</b>	<b>97</b>	<b>89</b>	<b>74</b>	<b>68</b>	<b>48</b>	<b>25</b>	<b>7</b>	5	*	*	*
5	78	80	73	64	61	50	32	13	8	1	*	*
6	<b>45</b>	<b>49</b>	<b>50</b>	<b>47</b>	<b>46</b>	<b>42</b>	<b>32</b>	<b>17</b>	12	2	*	*
7	26	<b>28</b>	<b>30</b>	<b>31</b>	<b>31</b>	<b>32</b>	<b>28</b>	<b>18</b>	14	4	1	*
8	17	17	19	20	21	23	22	16	14	5	1	*
9	11	11	12	13	<b>14</b>	<b>16</b>	<b>17</b>	<b>15</b>	13	6	1	*
10	8	<b>7</b>	<b>8</b>	<b>9</b>	<b>9</b>	<b>11</b>	<b>13</b>	12	11	6	2	*
11	6	5	6	6	7	8	9	10	9	6	2	*
12	5	<b>4</b>	<b>4</b>	<b>4</b>	<b>5</b>	<b>5</b>	<b>7</b>	<b>8</b>	<b>8</b>	6	2	*
13	<b>4</b>	<b>3</b>	<b>3</b>	<b>3</b>	<b>3</b>	<b>4</b>	<b>5</b>	<b>6</b>	<b>6</b>	<b>5</b>	<b>2</b>	<b>1</b>
14	4	3	2	2	3	3	4	<b>5</b>	<b>5</b>	<b>5</b>	2	1
15	3	2	2	2	2	2	3	4	4	4	2	1
16	3	2	2	2	2	2	2	3	3	3	2	1
17	3	2	1	1	1	1	<b>2</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>2</b>	<b>1</b>
18	3	1	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	2	2	2	2	1
19	3	1	1	1	1	1	1	1	1	2	2	1
20	4	1	1	1	1	1	1	1	<b>1</b>	<b>2</b>	<b>2</b>	<b>1</b>
21	5	1	1	1	1	1	1	1	1	1	1	1
22	6	1	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>	<b>1</b>
23	9	1	1	*	*	*	*	*	1	1	1	1
24	14	1	1	*	*	*	*	*	1	1	1	1
25	23	1	1	*	*	*	*	*	*	*	1	1
26	40	1	*	*	*	*	*	*	*	*	1	1
27	64	2	*	*	*	*	*	*	*	*	1	1
28	86	2	*	*	*	*	*	*	*	*	1	1
29	96	3	*	*	*	*	*	*	*	*	1	*
30	99	4	*	*	*	*	*	*	*	*	*	*

## 6. References

### 6.1 Text references

- <sup>1</sup>C. H. Rutledge and R. L. Watson, At. Data Nucl. Data Tables **12**, 195 (1973).
- <sup>2</sup>R. K. Gardner and T. J. Gray, At. Data Nucl. Data Tables **21**, 515 (1978); [Erratum **24**, 281 (1979)].
- <sup>3</sup>T. L. Hardi and R. L. Watson, At. Data Nucl. Data Tables **17**, 107 (1976); R. S. Shokhi and D. Crumpton, *ibid.* **30**, 49 (1984).
- <sup>4</sup>G. Lapicki, Ph.D. thesis, New York University, New York, 1975, see Fig. 1 in this thesis.
- <sup>5</sup>H. Paul, At. Data Nucl. Data Tables **24**, 243 (1979); Nucl. Instrum. Methods **169**, 249 (1980).
- <sup>6</sup>H. Paul, Nucl. Instrum. Methods **192**, 11 (1982).
- <sup>7</sup>H. Paul and W. Obermann, Nucl. Instrum. Methods **214**, 15 (1983).
- <sup>8</sup>H. Paul, Nucl. Instrum. Methods B **3**, 5 (1984) [Erratum: B **5**, 554 (1984)].
- <sup>9</sup>H. Paul, Nucl. Instrum. Methods B **4**, 211 (1984).
- <sup>10</sup>O. Benka and H. Paul, Comm. At. Mol. Phys. **15**, 29 (1984).
- <sup>11</sup>H. Paul and J. Muhr, in *Proceedings of the 2nd Workshop on High-Energy Ion-Atom Collisions, Debrecen 1984*, edited by D. Berenyi and G. Hock (Akademiai Kiado, Budapest, 1985), p. 49.
- <sup>12</sup>H. Paul and J. Muhr, Phys. Rep. **135**, 47 (1986).
- <sup>13</sup>H. Paul, Z. Phys. D **4**, 249 (1987).
- <sup>14</sup>O. Benka, M. Geretschläger, and H. Paul, in *Proceedings of the 3rd Workshop on High-Energy Ion-Atom Collisions, Debrecen, August 1987*, edited by D. Berenyi and G. Hock, *Lecture Notes in Physics 294* (Springer, Berlin, 1988), p. 94.
- <sup>15</sup>W. Brandt and G. Lapicki, Phys. Rev. A **23**, 1717 (1981) for direct ionization.
- <sup>16</sup>G. Lapicki and F. D. McDaniel, Phys. Rev. A **22**, 1896 (1980) (E) **23**, 975 (1981) for electron capture. Also see G. Lapicki, Trans. Nucl. Sci. **28**, 1066 (1981).
- <sup>17</sup>O. Benka and M. Geretschläger (see the data base references 103 and 118 in Sec. 6.2) as well as Polish groups (see Refs. 107, 124, 139, 141, 145, 148, 150, 154, 155, 158, 159, and 161 in Sec. 6.2) report an effective energy that the projectile has after typically traversing through one-half of the target's thickness. These energies were reconverted to the incident projectile energies for the consistency with all other data compiled in Tables 2–5.
- <sup>18</sup>See, for example, Figs. B16–B22 and B38–B42 in the Appendix B of Ref. 12.
- <sup>19</sup>M. O. Krause, J. Phys. Chem. Ref. Data **8**, 307 (1979).
- <sup>20</sup>G. S. Khandelwal, B.-H. Choi, and E. Merzbacher, At. Data **1**, 103 (1969) for direct ionization in the plane-wave Born approximation. These PWBA tables for direct ionization were extended by R. Ricc, G. Basbas, and F. D. McDaniel, At. Data Nucl. Data Tables **20**, 503 (1977).
- <sup>21</sup>V. S. Nikolaev, Zh. Eksp. Teor. Fiz. **51**, 1263 (1966) [Sov. Phys. JETP **24**, 847 (1967)] for electron capture in the OBK approximation.
- <sup>22</sup>M. H. Chen and B. Crasemann, At. Data Nucl. Data Tables **33**, 217 (1985).
- <sup>23</sup>D. D. Cohen and M. Harrigan, At. Data Nucl. Data Tables **33**, 255 (1985).
- <sup>24</sup>J. A. Tanis, S. M. Shafroth, W. W. Jacobs, T. McAbee, and G. Lapicki, Phys. Rev. A **31**, 750 (1985).
- <sup>25</sup>For other critical reviews of the ECPSSR theory see J. M. Hansteen, L. Kocbach, and A. Graue, Phys. Scripta **31**, 63 (1985); J. F. Reading, Nucl. Instrum. Methods A **262**, 160 (1987).
- <sup>26</sup>W. Brandt, R. Laubert, and I. Sellin, Phys. Lett. **21**, 518 (1966); Phys. Rev. **151**, 56 (1966).
- <sup>27</sup>W. Brandt and G. Lapicki, Phys. Rev. A **10**, 474 (1974).
- <sup>28</sup>G. Basbas, W. Brandt, and R. H. Ritchie, Phys. Rev. A **7**, 1971 (1973).
- <sup>29</sup>G. Basbas, W. Brandt, and R. Laubert, Phys. Rev. A **7**, 983 (1973); **17**, 1655 (1978).
- <sup>30</sup>W. Brandt and G. Lapicki, Phys. Rev. A **20**, 465 (1979).
- <sup>31</sup>G. Lapicki, Bull. Am. Phys. Soc. **26**, 1310 (1981).
- <sup>32</sup>Equation (6) of Ref. 30 gives the effective mass for the relativistic electron  $m_K^*$ . Note that, contrary to suggestions of Ref. 23, this mass should not enter into the argument of the energy-loss function  $f_K$  from Eq. (7) of Ref. 15; see G. Lapicki, J. Phys. B **20**, L633 (1987).
- <sup>33</sup>G. Lapicki and A. R. Zander, Phys. Rev. A **23**, 2072 (1981).
- <sup>34</sup>R. Singh and V. Singh, Physica **78**, 343 (1974); **83C**, 200 (1976); A. Langenberg and J. van Eck, J. Phys. B **11**, 1425 (1978); A. Kumar and B. N. Roy, *ibid.* **11**, 1435 (1978); F. F. Komarov and A. P. Novikov, Zh. Tekh. Fiz. **49**, 264 (1979) [Sov. Phys. Tech. Phys. **24**, 155 (1979)]; S. N. Chatterjee, A. Kumar, and B. N. Roy, Physica **122C**, 275 (1983); C. V. Seth, Phys. Rev. A **29**, 1151 (1984).
- <sup>35</sup>J. L. Duggan, P. M. Kocur, J. L. Price, F. D. McDaniel, R. Mehta, and G. Lapicki, Phys. Rev. A **32**, 2088 (1985).
- <sup>36</sup>D. H. Madison and E. Merzbacher, in *Atomic Inner Shell Processes*, edited by B. Crasemann (Academic, New York, 1975), Vol. I, p. 1.
- <sup>37</sup>L. Kocbach, Phys. Norvegica **8**, 187 (1976).
- <sup>38</sup>E. Laegsgaard, J. U. Andersen, and M. Lund, in *Proceedings of the 10th International Conference on the Physics of Electronic and Atomic Collisions, Paris*, edited by G. Watel (North-Holland, Amsterdam, 1978), p. 353; F. Bell, Nucl. Instrum. Methods **192**, 103 (1982).
- <sup>39</sup>E. C. Montenegro and A. G. de Pinho, J. Phys. B **15**, 1521 (1982); E. C. Montenegro and G. B. Baptista, Nucl. Instrum. Methods B **3**, 16 (1984); E. C. Montenegro and G. M. Sigaud, J. Phys. B **18**, 299 (1985).
- <sup>40</sup>K. M. Barfoot, I. V. Mitchell, H. L. Esbach, and W. B. Gilboy, J. Phys. B **15**, L845 (1982).
- <sup>41</sup>D. D. Cohen, J. Phys. B **16**, L415 (1983).
- <sup>42</sup>T. Mukoyama and L. Sarkadi, Phys. Rev. A **23**, 375 (1981); Nucl. Instrum. Methods **179**, 573 (1981); **205**, 341 (1983); **211**, 525 (1983); Phys. Rev. A **25**, 1411 (1982); **28**, 1303 (1983).
- <sup>43</sup>M. H. Chen, B. Crasemann, and H. Mark, Phys. Rev. A **26**, 1243 (1982).
- <sup>44</sup>M. H. Chen, Phys. Rev. A **27**, 2358 (1983); M. H. Chen, B. Crasemann, and H. Mark, *ibid.* **30**, 2082 (1984).
- <sup>45</sup>M. H. Chen and B. Crasemann, Phys. Rev. A **34**, 87 (1986).
- <sup>46</sup>A. Kropf, Nucl. Instrum. Methods **142**, 79 (1977); O. Benka, M. Geretschläger, and H. Paul, *ibid.* **142**, 83 (1977); K. G. Bauer, Q. Fazly, T. Mayer-Kuckuk, H. Mommsen, and P. Schurkes, *ibid.* **148**, 407 (1978); O. Benka, M. Geretschläger, and A. Kropf, *ibid.* **149**, 441 (1978); E. Clayton, *ibid.* **191**, 567 (1981); E. Clayton, D. D. Cohen, and P. Duerden, *ibid.* **191**, 573 (1981); J. L. Campbell, J. A. Cookson, and H. Paul, *ibid.* **212**, 427 (1983); J. A. Cookson and J. L. Campbell, *ibid.* **216**, 489 (1983); B **3**, 185 (1984); G. Linder, *ibid.* B **3**, 130 (1984); D. D. Cohen and E. Clayton, *ibid.* B **22**, 59 (1987); J. L. Campbell, A. Peruj, W. J. Weesdale, and J. A. Cookson, *ibid.* B **30**, 317 (1988). The ECPSSR theory was also recommended for deuteron induced x-ray analysis by K. M. Barfoot, Nucl. Instrum. Methods B **14**, 76 (1986).
- <sup>47</sup>D. D. Cohen and M. Harrigan, Nucl. Instrum. Methods B **15**, 576 (1986) and At. Data Nucl. Data Tables **34**, 393 (1986); T. Hirokawa, F. Ni-shiyama, and Y. Kiso, Nucl. Instrum. Methods B **31**, 525 (1988).
- <sup>48</sup>W. N. Lennard and D. Phillips, Nucl. Instrum. Methods **166**, 151 (1979); R. Mehta, J. L. Duggan, J. L. Price, F. D. McDaniel, and G. Lapicki, Phys. Rev. A **26**, 1883 (1982); W. Maenhaut and H. Raemdonck, Nucl. Instrum. Methods B **1**, 123 (1984).
- <sup>49</sup>W. Jitschin, H. Kleinpoppen, R. Hippeler, and H. O. Lutz, J. Phys. B **12**, 4077 (1979); J. Palinkas, L. Sarkadi, and B. Schlenk, *ibid.* **13**, 3829 (1980); V. V. Sizov and N. M. Kabachnik *ibid.* **16**, 1565 (1983); D. Berenyi, I. Cserny, I. Kadar, A. Kover, S. Ricz, L. Sarkadi, D. Varga, and J. Vegh, *ibid.* **17**, 829 (1984); U. Werner, W. Jitschin, and H. O. Lutz, *ibid.* **18**, 3111 (1984).
- <sup>50</sup>D. D. Cohen and M. Harrigan, At. Data Nucl. Data Tables **34**, 393 (1986); E. Rosato, Nucl. Instrum. Methods B **15**, 591 (1986); D. D. Cohen, *ibid.* **22**, 55 (1987); J. Q. Xu and E. Rosato, Phys. Rev. A **37**, 1946 (1988).
- <sup>51</sup>W. Brandt and G. Basbas, Phys. Rev. A **27**, 578 (1983); (E) **28**, 3142 (1983).
- <sup>52</sup>A. Langerberg and J. van Eck, J. Phys. B **10**, L419 (1977).
- <sup>53</sup>K. Unterseer and N. Kleber, Nucl. Instrum. Methods **192**, 35 (1982); D. J. Land, D. G. Simons, and M. D. Brown, *ibid.* **214**, 35 (1983); B **4**, 239 (1984); L. Kocbach, *ibid.* B **4**, 248 (1984); L. Sarkadi, *ibid.* B **9**, 127 (1985); D. H. Jakubasa-Amundesen, Z. Phys. A **320**, 557 (1985); D. J. Land, Nucl. Instrum. Methods A **240**, 470 (1985) and Bull. Am. Phys. Soc. **31**, 981 (1986).
- <sup>54</sup>J. F. Reading, A. L. Ford, J. S. Smith, and R. L. Becker, in *Invited Talks, the XIIth International Conference on the Physics of Electronic and Atomic Collisions*, edited by J. Eichler, I. V. Hertel, and N. Stolterfoht (Elsevier, Amsterdam, 1984); J. F. Reading and A. L. Ford, Phys. Rev. Lett. **58**, 543 (1987); J. Phys. B **20**, 3747 (1987), applied this method most recently to calculate double, and single, direct ionization of helium by 0.3–40 MeV protons and antiprotons, i.e., in the  $v_1/v_{2K} > 2$  range where the first Born approximation is well justified.
- <sup>55</sup>A. Graue, J. M. Hansteen, R. Gunderson, and L. Kocbach, J. Phys. B **15**, L445 (1982).
- <sup>56</sup>R. Gunderson, J. M. Hansteen, and L. Kocbach, Nucl. Instrum. Methods **192**, 63 (1982); J. M. Hansteen, L. Kocbach, and A. Graue, Phys. Scripta **31**, 63 (1985). See J. M. Hansteen, Ref. 14, p. 39, for the latest status of

- semiclassical calculations.
- <sup>57</sup>J. F. Reading, A. L. Ford, and R. L. Becker, *J. Phys. B* **14**, 1995 (1981); J. F. Reading, A. L. Ford, M. Martir, and R. L. Becker, *Nucl. Instrum. Methods* **192**, 1 (1982).
- <sup>58</sup>D. Trautmann, R. Rosel, and G. Baur, *Nucl. Instrum. Methods* **214**, 21 (1983); A. Jakob, D. Trautmann, R. Rosel, and G. Baur, *ibid. B* **4**, 218 (1984).
- <sup>59</sup>G. Mehler, T. de Reus, U. Muller, J. Reinhardt, B. Muller, W. Greiner, and G. Soff, *Nucl. Instrum. Methods* **A240**, 559 (1985).
- <sup>60</sup>G. Mehler, W. Greiner, and G. Soff, *J. Phys. B* **20**, 2787 (1987).
- <sup>61</sup>L. Kocbach, *Nucl. Instrum. Methods* **B4**, 248 (1984).
- <sup>62</sup>T. Mukoyama and C. D. Lin, *Nucl. Instrum. Methods* **A262**, 15 (1987) and in Ref. 14, p. 84. These authors compare their calculations with an early version of ECPSSR theory, which gives 2%–5% lower values for 0.5–2 MeV protons on copper. As can be seen from Fig. 3 of Ref. 7, the ECPSSR cross section is 10% above the Mukoyama and Lin calculation for 2-MeV protons on copper.
- <sup>63</sup>R. Anholt, W. E. Meyerhof, H. Gould, C. Munger, J. Alonso, P. Thieberger, and H. E. Wegner, *Phys. Rev. A* **32**, 3302 (1985); R. Anholt and H. Gould, *Adv. At. Mol. Phys.* **22**, 315 (1986).
- <sup>64</sup>See, L. Sarkadi, in Ref. 53.
- <sup>65</sup>G. Basbas and D. J. Land, *Phys. Rev. A* **35**, 1003 (1987).
- <sup>66</sup>I. Chadwick, *Philos. Mag.* **25**, 193 (1913); *ibid.* **24**, 594 (1912).
- <sup>67</sup>J. M. Cork, *Phys. Rev.* **59**, 957 (1941).
- <sup>68</sup>G. Lapicki, R. Laubert, and W. Brandt, *Phys. Rev. A* **22**, 1889 (1980); S. Raith, S. Divoux, and B. Gonsior, *Nucl. Instrum. Methods* **B 10/11**, 169 (1985); M. Harrison and D. D. Cohen, *ibid. B* **15**, 581 (1986); T. Papp and B. Schlenk, *J. Phys. B* **20**, 2255 (1987); R. A. Ilkhamov, S. H. Khushmurodov, A. P. Kobzev, J. H. Li, M. Pajek, and R. Sandrik, in Ref. 14, p. 103. See also Refs. 5, 6, 9, 10, 12, and 13.
- <sup>69</sup>G. Lapicki and W. Losonsky, *Phys. Rev. A* **20**, 481 (1979).
- <sup>70</sup>J. M. Hansteen, in Ref. 14, p. 39, argues that the velocity-symmetrized hyperbolic trajectory implies an exact treatment of the Coulomb-deflection effect. However, such symmetrization spoils the unitarity of the scattering matrix; see F. Wolf, R. J. Allen, and H. J. Korsch, *Comm. At. Mol. Phys.* **18**, 107 (1986).
- <sup>71</sup>G. Basbas and G. S. Khandelwal, *Bull. Am. Phys. Soc.* **11**, 307 (1966) and Fig. 3 from the 1973 article of Ref. 29; A. L. Ford, E. Fitchard, and J. F. Reading, *Phys. Rev. A* **16**, 133 (1977); D. J. Land, M. D. Brown, D. Simons, and J. G. Brennan, *Nucl. Instrum. Methods* **192**, 53 (1982); P. Rez, *X-Ray Spectrometry* **13**, 55 (1984).
- <sup>72</sup>L. Sarkadi and T. Mukoyama, *J. Phys. B* **14**, L255 (1981) and *Nucl. Instrum. Methods* **B4**, 296 (1984); K. Finck, W. Jitschin, and H. O. Lutz, *J. Phys. B* **16**, L403 (1983); L. Sarkadi and T. Papp, *Acta Physica Hungarica* **58**, 75 (1985).
- <sup>73</sup>L. Sarkadi, *J. Phys. B* **18**, 2519 and L755 (1986); *Nucl. Instrum. Methods A* **265**, 45 (1987).
- <sup>74</sup>S. Zehendner, G. B. Baptista, R. Donner, E. Justiniano, J. Konrad, H. Schmidt-Bocking, and R. Schuch, *Z. Phys. D* **4**, 243 (1987); L. Sarkadi and T. Mukoyama, *J. Phys. B* **20**, L559 (1987); W. Schadt, H. Schmidt-Bocking, G. Nolte, Z. Roller, A. Skutlartz, M. Wassermann, and G. Zschornack, *Z. Phys. D* **8**, 271 (1988).
- <sup>75</sup>L. Sarkadi and T. Mukoyama, *Phys. Rev. A* **37**, 4540 (1988).
- <sup>13</sup>L. J. Christensen, J. M. Khan, and W. F. Brunner, *Rev. Sci. Instrum.* **38**, 20 (1967).
- <sup>14</sup>P. Komarek, *Acta Phys. Austriaca* **26**, 315 (1967).
- <sup>15</sup>P. Komarek, *Acta Phys. Austriaca* **27**, 369 (1968).
- <sup>16</sup>R. C. Der, T. M. Kavanagh, J. M. Khan, B. P. Curry, and R. J. Fortner, *Phys. Rev. Lett.* **26**, 1731 (1968).
- <sup>17</sup>W. Brandt and R. Laubert, *Phys. Rev.* **178**, 225 (1969).
- <sup>18</sup>R. R. Hart, F. W. Reuter III, H. P. Smith, Jr., and J. M. Khan, *Phys. Rev.* **179**, 4 (1969).
- <sup>19</sup>B. Sellers, F. A. Hanser, and H. H. Wilson, *Phys. Rev.* **182**, 90 (1969).
- <sup>20</sup>G. Bissinger, J. M. Joyce, E. J. Ludwig, W. S. McEver, and S. M. Shafrroth, *Phys. Rev. A* **1**, 841 (1970).
- <sup>21</sup>P. Richard, T. I. Bonner, T. Furuta, I. L. Morgan, and J. R. Rhodes, *Phys. Rev. A* **1**, 1044 (1970).
- <sup>22</sup>P. B. Needham, Jr. and B. D. Sartwell, *Phys. Rev. A* **2**, 27 (1970).
- <sup>23</sup>M. Terasawa, T. Inouye, and H. Kamei, *Jpn. J. Phys.* **29**, 1394 (1970).
- <sup>24</sup>R. L. Watson, C. W. Lewis, and J. B. Natowitz, *Nucl. Phys. A* **154**, 561 (1970).
- <sup>25</sup>K. Shima, I. Makino, and M. Sakisaka, *Jpn. J. Phys.* **30**, 611 (1971).
- <sup>26</sup>M. Terasawa, *Progress Report, Institute of Space and Aeronautical Science*, edited by T. Takayanagi (University of Tokyo, March 1971).
- <sup>27</sup>A. Fahnenius and P. Jauho, *Ann. Acad. Sci. Fenn. Ser. A* **6** 367, 3 (1971).
- <sup>28</sup>K. Shima, I. Makino, and M. Sakisaka, *Jpn. J. Phys.* **31**, 971 (1971).
- <sup>29</sup>P. B. Needham, Jr. and B. D. Sartwell, *Adv. X-Ray Anal.* **14**, 184 (1971).
- <sup>30</sup>O. N. Jarvis, C. Whitehead, and M. Shah, *Phys. Rev. A* **5**, 1198 (1972).
- <sup>31</sup>C. W. Lewis, R. L. Watson, and J. B. Natowitz, *Phys. Rev. A* **5**, 1773 (1972).
- <sup>32</sup>G. Bissinger, S. M. Shafrroth, and A. W. Waltner, *Phys. Rev. A* **5**, 2046 (1972).
- <sup>33</sup>F. W. Reuter III and H. P. Smith, Jr., *J. Appl. Phys.* **43**, 4228 (1972).
- <sup>34</sup>R. G. Musket and W. Bauer, *J. Appl. Phys.* **43**, 4786 (1972).
- <sup>35</sup>M. Terasawa, T. Tamura, and H. Kamada, *Jpn. J. Phys.* **33**, 1420 (1972).
- <sup>36</sup>J. L. Duggan, W. L. Beck, L. Albrecht, L. Munz, and J. D. Spaulding, *Adv. X-Ray Anal.* **15**, 407 (1972).
- <sup>37</sup>G. Bissinger and H. W. Kugel, in *Proceedings of International Conference on Inner Shell Ionization Phenomena, Atlanta 1972*, edited by R. W. Fink, S. T. Manson, M. Palms, and P. V. Rao (U.S. AEC, Oak Ridge, Tn, 1973), p. 993.
- <sup>38</sup>J. Lin, J. L. Duggan, and R. F. Carlton, in *Proceedings of International Conference on Inner Shell Ionization Phenomena, Atlanta 1972*, edited by R. W. Fink, S. T. Manson, M. Palms, and P. V. Rao (U.S. AEC, Oak Ridge, Tn, 1973), p. 998.
- <sup>39</sup>E. Laegsgaard, J. U. Andersen, and L. C. Feldman, in *Proceedings of International Conference on Inner Shell Ionization Phenomena, Atlanta 1972*, edited by R. W. Fink, S. T. Manson, M. Palms, and P. V. Rao (U.S. AEC, Oak Ridge, Tn, 1973), p. 1019.
- <sup>40</sup>L. M. Winters, L. D. Ellsworth, T. Chiao, and J. R. Macdonald, in *Proceedings of International Conference on Inner Shell Ionization Phenomena, Atlanta 1972*, edited by R. W. Fink, S. T. Manson, M. Palms, and P. V. Rao (U.S. AEC, Oak Ridge, Tn, 1973), p. 1069.
- <sup>41</sup>A. W. Waltner, D. M. Peterson, G. Bissinger, A. B. Baskin, C. E. Busch, P. H. Nettles, W. R. Scates, and S. M. Shafrroth, in *Proceedings of International Conference on Inner Shell Ionization Phenomena, Atlanta 1972*, edited by R. W. Fink, S. T. Manson, M. Palms, and P. V. Rao (U.S. AEC, Oak Ridge, Tn, 1973), p. 1080.
- <sup>42</sup>A. Van der Woude, M. J. Saltmarsh, C. A. Ludemann, R. L. Hahn, and E. Eichler, in *Proceedings of International Conference on Inner Shell Ionization Phenomena, Atlanta 1972*, edited by R. W. Fink, S. T. Manson, M. Palms, and P. V. Rao (U.S. AEC, Oak Ridge, Tn, 1973), p. 1388.
- <sup>43</sup>R. H. McKnight, S. T. Thornton, and R. C. Ritter, in *Proceedings of International Conference on Inner Shell Ionization Phenomena, Atlanta 1972*, edited by R. W. Fink, S. T. Manson, M. Palms, and P. V. Rao (U.S. AEC, Oak Ridge, Tn, 1973), p. 1439.
- <sup>44</sup>A. Fahnenius, in *Electrical and Nuclear Technology Publication 3 (Technical Research Center of Finland, Helsinki, 1973)*.
- <sup>45</sup>G. Basbas, W. Brandt, and R. Laubert, *Phys. Rev. A* **7**, 983 (1973).
- <sup>46</sup>K. G. Harrison, H. Tawara, and F. J. De Heer, *Physica* **66**, 6 (1973).
- <sup>47</sup>R. C. Bearse, D. A. Close, J. J. Malanify, and C. J. Umbarger, *Phys. Rev. A* **7**, 1269 (1973).
- <sup>48</sup>L. M. Winters, J. R. Macdonald, M. D. Brown, L. D. Ellsworth, and T. Chiao, *Phys. Rev. A* **7**, 1276 (1973).
- <sup>49</sup>T. L. Hardt and R. L. Watson, *Phys. Rev. A* **7**, 1917 (1973).
- <sup>50</sup>A. Langenberg and J. van Eck, *Phys. Rev. Lett.* **31**, 71 (1973).
- <sup>51</sup>K. Brunner and W. Hink, *Z. Phys.* **262**, 181 (1973).

## 6.2. References to cross-section data compiled in Tables 2–5

- H. W. Lewis, B. E. Simmons, and E. Merzbacher, *Phys. Rev.* **91**, 943 (1953).
- P. R. Bevington and E. M. Bernstein, *Bull. Am. Phys. Soc.* **1**, 198 (1956).
- J. M. Hansteen and S. Messelt, *Nucl. Phys.* **2**, 526 (1957).
- B. Singh, *Phys. Rev.* **107**, 711 (1957).
- E. Merzbacher and H. W. Lewis, in *Handbuch der Physik*, edited by S. Flugge (Springer, Berlin, 1958), Vol. 34, p. 119.
- S. Messelt, *Nucl. Phys.* **5**, 435 (1958).
- R. C. Jopson, H. Mark, and C. D. Swift, *Phys. Rev.* **127**, 1612 (1962). Energies were recalibrated according to the footnote 9 of Ref. 9.
- N. L. Lark, *Bull. Am. Phys. Soc.* **7**, 623 (1962).
- J. M. Khan and D. L. Potter, *Phys. Rev.* **133**, A 890 (1964).
- W. T. Ogier, G. J. Lucas, J. S. Murray, and T. E. Holzer, *Phys. Rev. A* **134**, 1070 (1964).
- J. M. Khan, D. L. Potter, and R. D. Worley, *Phys. Rev.* **139**, A 1735 (1965).
- W. Brandt, R. Laubert, and I. Sellin, *Phys. Rev.* **151**, 56 (1966).

- <sup>52</sup>R. B. Liebert, T. Zabel, D. Miljanic, H. Larson, V. Valkovic, and G. C. Phillips, Phys. Rev. A **8**, 2336 (1973).
- <sup>53</sup>R. D. Lear and T. J. Gray, Phys. Rev. A **8**, 2469 (1973). T. J. Gray, R. D. Lear, R. J. Dexter, F. N. Schwettmann, and K. Wiemer report these data also in Thin Solid Films **19**, 103 (1973).
- <sup>4</sup>R. H. McKnight, S. T. Thornton, and R. R. Karlowicz, Phys. Rev. A **9**, 267 (1974).
- <sup>55</sup>R. Akselsson and T. B. Johansson, Z. Phys. **266**, 245 (1974).
- <sup>56</sup>H. Tawara, K. Ishii, S. Morita, H. Kaji, C. N. Hsu, and T. Shiokawa, Phys. Rev. A **9**, 1617 (1974).
- <sup>57</sup>R. M. Wheeler, R. P. Chaturvedi, and A. R. Zander, in *Proceedings of the 3rd Conference on Applications of Small Accelerator, Denton 1974*, edited by J. L. Duggan and I. L. Morgan (National Technical Information Service, Springfield, Virginia, 1974), Vol. I, p. 387.
- <sup>58</sup>D. Burch, N. Stolterfoht, D. Schneider, H. Wieman, and J. S. Risley, Phys. Rev. Lett. **32**, 1151 (1974).
- <sup>59</sup>F. Folkmann, J. Borggreen, and A. Kjeldgaard, Nucl. Instrum. Methods **119**, 117 (1974).
- <sup>60</sup>K. Ishii, S. Morita, H. Tawara, H. Kaji, and T. Shiokawa, Phys. Rev. A **10**, 774 (1974).
- <sup>61</sup>T. L. Criswell and T. J. Gray, Phys. Rev. A **10**, 1145 (1974).
- <sup>62</sup>R. H. McKnight, S. T. Thornton, and R. R. Karlowicz, Nucl. Instrum. Methods **123**, 1 (1975).
- <sup>63</sup>N. A. Khelil and T. J. Gray, Phys. Rev. A **11**, 893 (1975).
- <sup>64</sup>F. Hopkins, R. Brenn, A. R. Whittemore, J. Karp, and S. K. Bhattacharjee, Phys. Rev. A **11**, 916 (1975).
- <sup>65</sup>S. J. Czuchlewski, J. R. Macdonald, and L. D. Ellsworth, Phys. Rev. A **11**, 1108 (1975).
- <sup>66</sup>F. Hopkins, R. Brenn, A. R. Whittemore, N. Cue, and V. Dutkiewicz, Phys. Rev. A **11**, 1482 (1975).
- <sup>67</sup>F. D. McDaniel, T. J. Gray, and R. K. Gardner, Phys. Rev. A **11**, 1607 (1975).
- <sup>68</sup>W. N. Lennard and I. V. Mitchell, Phys. Rev. A **12**, 1723 (1975).
- <sup>69</sup>F. Bodart, S. Wilk, and G. Deconninck, X Ray Spectrometry **4**, 161 (1975).
- <sup>70</sup>D. Burch, Phys. Rev. A **12**, 2225 (1975).
- <sup>71</sup>K. Kawatsura, K. Ozawa, F. Fujimoto, and M. Terasawa, in *Ion Beam Surface Layer Analysis*, edited by O. Meyer, G. Linker, and F. Käppeler (Plenum, New York, 1976), Vol. 2, p. 719.
- <sup>72</sup>F. D. McDaniel and J. L. Duggan, in *Beam Foil Spectroscopy Collisional and Radiative Processes*, edited by I. A. Sellin and D. J. Pegg (Plenum, New York, 1976), Vol. 2, p. 519.
- <sup>73</sup>V. S. Nikolaev, V. P. Petukhov, E. R. Romanovsky, V. A. Sergeev, I. M. Kruglova, and V. V. Beloshitsky, *9th International Conference on the Physics of Electronic and Atomic Collisions, Seattle 1975*, edited by J. S. Risley and R. Geballe (University of Washington, Washington, 1976), p. 419.
- <sup>74</sup>R. R. Randall, J. A. Bednar, B. Curnette, and C. L. Cocke, Phys. Rev. A **13**, 204 (1976).
- <sup>75</sup>H. Tawara, Y. Hachiya, K. Ishii, and S. Morita, Phys. Rev. A **13**, 572 (1976).
- <sup>76</sup>M. Milazzo and G. Riccobono, Phys. Rev. A **13**, 578 (1976).
- <sup>77</sup>M. R. Khan, D. Crumpton, and P. E. Francois, J. Phys. B **9**, 455 (1976).
- <sup>78</sup>C. G. Soares, R. D. Lear, J. T. Sanders, and H. A. Van Rinsvelt, Phys. Rev. A **13**, 953 (1976).
- <sup>79</sup>A. R. Zander, Y. Chee, J. Walls, and B. Crews, in *Abstracts of Contributed Papers, Second International Conference on Inner Shell Ionization Phenomena, Freiburg 1976*, edited by W. Melhorn (University of Freiburg, Freiburg, 1976), p. 253.
- <sup>80</sup>E. Kolatay, D. Berenyi, I. Kiss, S. Ricsz, G. Hock, and J. Basco, Z. Phys. A **278**, 299 (1976).
- <sup>81</sup>G. Bissinger, J. M. Joyce, and H. W. Kugel, Phys. Rev. A **14**, 1375 (1976).
- <sup>82</sup>A. Langenberg and J. van Eck, J. Phys. B **9**, 2421 (1976).
- <sup>83</sup>T. Badica, C. Ciortea, S. Dima, A. Petrovici, I. Popescu, and V. Neacsu, X-Ray Spectrometry **6**, 90 (1977).
- <sup>84</sup>B. Knauf, G. Presser, and J. Stahler, Z. Phys. A **282**, 25 (1977).
- <sup>85</sup>G. Bissinger, J. M. Joyce, B. L. Doyle, W. W. Jacobs, and S. M. Shafrroth, Phys. Rev. A **16**, 443 (1977).
- <sup>86</sup>M. R. Khan, A. G. Hopkins, D. Crumpton, and P. E. Francois, X-Ray Spectrometry **6**, 140 (1977).
- <sup>87</sup>S. R. Wilson, F. D. McDaniel, J. R. Rowe, and J. L. Duggan, Phys. Rev. A **16**, 903 (1977).
- <sup>88</sup>K. H. Weber and F. Bell, Phys. Rev. A **16**, 1075 (1977).
- <sup>89</sup>H. Schmidt-Bocking, R. Schule, K. E. Steibing, K. Bethge, I. Tserruya, and H. Zekl, J. Phys. B **10**, 2663 (1977).
- <sup>90</sup>G. Deconninck and M. Longree, Phys. Rev. A **16**, 1390 (1977).
- <sup>91</sup>M. Kamiya, K. Ishii, K. Sera, S. Morita, and H. Tawara, Phys. Rev. A **16**, 2295 (1977).
- <sup>92</sup>K. Kawatsura, Ph.D. thesis (University of Kyoto, Kyoto, 1977).
- <sup>93</sup>E. Laegsgaard, J. U. Andersen, and M. Lund, *10th International Conference on the Physics of Electronic and Atomic Collisions, Paris 1977*, edited by G. Watel (North-Holland, Amsterdam, 1978), p. 353.
- <sup>94</sup>G. Bonani, C. Stoller, M. Stockli, M. Suter, and W. Wolfli, Helv. Phys. Acta **51**, 272 (1978).
- <sup>95</sup>R. Anholi, Phys. Rev. A **17**, 983 (1978).
- <sup>96</sup>S. M. Brodskii, S. V. Mamikonyan, and V. I. Filatov, Atomnaya Energiya **44**, 265 (1978) [Sov. Atomic Energy **44**, 300 (1978)].
- <sup>97</sup>G. Basbas, W. Brandt, and R. Laubert, Phys. Rev. A **17**, 1655 (1978).
- <sup>98</sup>J. S. Lopes, A. P. Jesus, G. P. Ferreira, and F. B. Gil, J. Phys. B **11**, 2181 (1978).
- <sup>99</sup>C. Bauer, R. Mann, and W. Rudolph, Z. Phys. A **287**, 27 (1978).
- <sup>100</sup>A. Berinde, C. Deberth, I. Neamu, C. Protop, N. Scinte, V. Zoran, M. Dost, and S. Rohl, J. Phys. B **11**, 2875 (1978).
- <sup>101</sup>K. Shima, Phys. Lett. A **67**, 351 (1978).
- <sup>102</sup>K. Shima, Jpn. J. Appl. Phys. **17**, Supplement 17-2, 350 (1978).
- <sup>103</sup>O. Benka and M. Geretschläger, Z. Phys. A **284**, 29 (1978).
- <sup>104</sup>W. D. Ramsay, M. S. A. L. Al-Ghazi, J. Birchall, and J. S. C. McKay, Phys. Lett. A **69**, 258 (1978).
- <sup>105</sup>M. Poncet and C. Engelmann, Nucl. Instrum. Methods **159**, 455 (1979).
- <sup>106</sup>J. S. Lopes, A. P. Jesus, S. C. Ramos, and G. P. Ferreira, J. Phys. B **12**, 605 (1979).
- <sup>107</sup>A. Celler, J. Kantele, M. Luontama, and J. Zylicz, Nucl. Instrum. Methods **163**, 221 (1979).
- <sup>108</sup>A. R. Zander and M. C. Andrews III, Phys. Rev. A **20**, 1484 (1979).
- <sup>109</sup>T. Badica, C. Ciortea, A. Petrovici, and I. Popescu, X-Ray Spectrometry **8**, 186 (1979).
- <sup>110</sup>W. N. Lennard and D. Phillips, Nucl. Instrum. Methods **166**, 521 (1979).
- <sup>111</sup>C. Magno, M. Milazzo, C. Pizzi, F. Porro, A. Rota, and G. Riccobono, Nuovo Cimento A **54**, 277 (1979).
- <sup>112</sup>K. M. Barfoot, I. V. Mitchell, H. L. Eschbach, and W. B. Gilboy, Nucl. Instrum. Methods **168**, 131 (1980).
- <sup>113</sup>E. Laegsgaard, J. U. Andersen, and F. Hogedal, Nucl. Instrum. Methods **169**, 293 (1980).
- <sup>114</sup>M. Dost, Nucl. Instrum. Methods **169**, 305 (1980).
- <sup>115</sup>J. S. Lopes, A. P. Jesus, and S. C. Ramos, Nucl. Instrum. Methods **169**, 311 (1980).
- <sup>116</sup>K. Kawatsura, A. Ootuka, K. Ozawa, F. Fujimoto, K. Komaki, and M. Terasawa, Nucl. Instrum. Methods **170**, 265 (1980).
- <sup>117</sup>K. Sera, K. Ishii, M. Kamiya, A. Kuwako, and S. Morita, Phys. Rev. A **21**, 1412 (1980).
- <sup>118</sup>O. Benka and M. Geretschläger, J. Phys. B **13**, 3223 (1980).
- <sup>119</sup>T. Scharnagl and W. Hink, J. Phys. B **13**, 4021 (1980).
- <sup>120</sup>G. Lapicki, R. Laubert, and W. Brandt, Phys. Rev. A **22**, 1889 (1980).
- <sup>121</sup>A. P. Jesus and J. S. Lopes, *Inner-Shell and X-Ray Physics of Atoms and Solids*, edited by D. J. Fabian, H. Kleinpoppen, and L. M. Watson (Plenum, New York, 1981), p. 21.
- <sup>122</sup>Z. Szokefalvi-Nagy and I. Demeter, Nucl. Instrum. Methods **181**, 1 (1981).
- <sup>123</sup>L. Avaldi, M. Milazzo, A. Rota, and G. Riccobono, J. Phys. B **14**, 2223 (1981).
- <sup>124</sup>P. Hornshøj, Z. Zelazny, M. Jaskola, L. Zemlo, A. Celler, and J. Szerypo, J. Phys. B **14**, 2391 (1981).
- <sup>125</sup>M. Dost, S. Hoppenau, J. Kising, S. Rohl, and P. Schorn, Phys. Rev. A **24**, 693 (1981).
- <sup>126</sup>R. K. Rice, F. D. McDaniel, G. Basbas, and J. L. Duggan, Phys. Rev. A **24**, 758 (1981).
- <sup>127</sup>P. Cuzzocrea, E. Perillo, E. Rosato, G. Spadaccini, N. De Cesare, and M. Vigilante, Lett. Nuovo Cimento **32**, 33 (1981).
- <sup>128</sup>M. Dost, S. Hoppenau, S. Rohl, and W. A. Schönfeldt, J. Phys. B **14**, 3153 (1981).
- <sup>129</sup>A. P. Jesus and J. S. Lopes, Nucl. Instrum. Methods **192**, 25 (1982).
- <sup>130</sup>J. U. Andersen, E. Laegsgaard, and M. Lund, Nucl. Instrum. Methods **192**, 79 (1982).
- <sup>131</sup>K. M. Barfoot, I. V. Mitchell, H. L. Eschbach, and W. B. Gilboy, J. Phys. B **15**, L845 (1982).
- <sup>132</sup>M. D. Brown, D. G. Simons, D. J. Land, and J. G. Brennan, Phys. Rev. A **25**, 2935 (1982).
- <sup>133</sup>J. S. Lopes, A. P. Jesus, and M. F. Da Silva, J. Phys. B **15**, 1749 (1982).

- <sup>134</sup>D. Bhattacharya, A. Roy, S. K. Bhattacherjee, and S. K. Mitra, *J. Phys. B* **15**, 3047 (1982).
- <sup>135</sup>D. Bhattacharya and S. K. Mitra, *Pramana* **19**, 399 (1982).
- <sup>136</sup>K. Ishii, M. Sebata, M. Kamiya, A. Kuwako, S. Morita, Y. Awaya, and T. Tonuma, *Jpn. J. Phys.* **51**, 4021 (1982).
- <sup>137</sup>M. D. Brown, D. G. Simons, D. J. Land, and J. G. Brennan, *IEEE Trans. Nucl. Sci.* **30**, 957 (1983).
- <sup>138</sup>L. Avaldi, M. Milazzo, G. Trivia, and I. V. Mitchell, *J. Phys. B* **16**, 1957 (1983).
- <sup>139</sup>M. Gocłowski, M. Jaskola, J. Szerypo, P. Hornshoj, and Z. Zelazny, *J. Phys. B* **16**, 3571 (1983).
- <sup>140</sup>Z. Sujkowski, D. Chmielewska, and M. N. Harakeh, *Nucl. Instrum. Methods* **219**, 111 (1984).
- <sup>141</sup>Z. Zelazny and P. Hornshoj, *J. Phys. B* **17**, 1867 (1984).
- <sup>142</sup>N. V. De Castro Faria, F. L. Freire, Jr., E. C. Montenegro, A. G. De Pinho, and E. F. Da Silveira, *J. Phys. B* **17**, 2307 (1984).
- <sup>143</sup>L. Avaldi, I. V. Mitchell, and H. L. Eschbach, *Nucl. Instrum. Methods B* **3**, 21 (1984).
- <sup>144</sup>S. Divoux, B. Raith, and B. Gonsior, *Nucl. Instrum. Methods B* **3**, 27 (1984).
- <sup>145</sup>M. Pfuetzner, J. Szerypo, Z. Zelazny, J. Zylicz, M. Gocłowski, M. Jaskola, L. Zemlo, and P. Hornshoj, *Nucl. Instrum. Methods B* **3**, 33 (1984).
- <sup>146</sup>W. N. Lennard, J. S. Foster, H. Geissel, K. M. Barfoot, and D. Phillips, *Nucl. Instrum. Methods B* **4**, 262 (1984).
- <sup>147</sup>F. L. Freire, Jr., E. C. Montenegro, A. G. De Pinho, and G. M. Sigaud, *J. Phys. B* **18**, 313 (1985).
- <sup>148</sup>E. Braziewicz, J. Braziewicz, M. Pajek, G. N. Osetynski, and J. Płoskonka, *J. Phys. B* **19**, 1471 (1986).
- <sup>149</sup>M. Geretschläger and O. Benka, *Phys. Rev. A* **34**, 866 (1986).
- <sup>150</sup>F. M. El-Ashry, M. Gocłowski, L. Glowacka, M. Jaskola, Z. Zelazny, and J. Szerypo, *J. Phys. B* **19**, 2311 (1986).
- <sup>151</sup>H. Xu, C. Ren, J. Tang, and F. Yang, in announcement of the 10th International CODATA Conference, Ottawa, 1986. Data obtained by private communication from H. Xu of Fudan University (1986). J. Li, H. Xu, C. Ren, and F. Lu, *Nucl. Instrum. Methods B* **30**, 16 (1988), report  $\sigma_{KX}^{\text{Expt}} = 42.9 b$  for 2-MeV protons on copper. This cross section agrees with  $(45.5 \pm 3.2) b$  from Xu's 1986 communication; it is in excellent agreement with the empirical "reference"  $(42.2 \pm 0.8) b$  from Ref. 12 of Sec. 6.2 and it lies 1% below  $\sigma_{KX}^{\text{ECPSSR}} = 43.4 b$ .
- <sup>152</sup>S. O. Olabanji and J. M. Calvert, *Nucl. Instrum. Methods A* **251**, 354 (1986).
- <sup>153</sup>E. C. Montenegro, A. G. De Pinho, and G. M. Sigaud, *J. Phys. B* **19**, 3287 (1986).
- <sup>154</sup>Z. Zelazny, M. Pfuetzner, J. Szerypo, M. Jaskola, and M. Gocłowski, *J. Phys. B* **19**, 4185 (1986).
- <sup>155</sup>F. M. El-Ashry, L. Glowacka, M. Jaskola, M. Pfuetzner, J. Szerypo, Z. Zelazny, G. M. Osetynski, and M. Pajek, *Nucl. Instrum. Methods B* **22**, 82 (1987). Only the data that have not been tabulated in prior references are taken from this article.
- <sup>156</sup>S. O. Olabanji and B. G. Martinsson, *Nucl. Instrum. Methods B* **24/25**, 81 (1987).
- <sup>157</sup>R. P. Bhalla, F. D. McDaniel, and G. Lapicki, *Phys. Rev. A* **35**, 3655 (1987).
- <sup>158</sup>M. Pfuetzner, J. Szerypo, Z. Zelazny, F. M. El-Ashry, M. Gocłowski, L. Glowacka, D. Trautmann, and M. Jaskola, *J. Phys. B* **20**, 3453 (1987).
- <sup>159</sup>J. Szerypo, M. Pfuetzner, W. Kretschmer, R. Schmitt, W. Schuster, A. Bienkowski, L. Glowacka, D. Trautmann, and M. Jaskola, *J. Phys. B* **20**, 5475 (1987).
- <sup>160</sup>D. G. Simons, J. L. Price, Jr., and D. J. Land, *15th International Conference on the Physics of Electronic and Atomic Collisions, Brighton 1987*, edited by J. Geddes, H. B. Gilbody, A. E. Kingston, C. J. Latimer, and H. J. R. Walters (Queen's University, Belfast, 1987), p. 610.
- <sup>161</sup>R. A. Ilkhamov, S. H. Khusmurodov, A. P. Kobzev, J. H. Li, M. Pajek, and R. Sandrik, in *Proceedings of the 3rd Workshop on High-Energy Ion-Atom Collisions, Debrecen 1987*, edited by D. Berenyi and G. Hock, *Lecture Notes in Physics* **294** (Springer, Berlin, 1988), p. 103.

## 6.3. Author index for the data base references in Section 6.2

- Akselsson,R. 55  
 Al-Ghazi,M.S.A.L. 104  
 Albrecht,L. 36  
 Andersen,J.U. 39,93,113,130  
 Andrews,M.C.,III 108  
 Anholt,R. 95  
 Aivaldi,L. 123,138,143  
 Awaya,Y. 136  
 Badica,T. 83,109  
 Barfoot,K.M. 112,131,146  
 Basbas,G. 45,97,126  
 Basco,J. 80  
 Baskin,A.B. 41  
 Bauer,C. 99  
 Bauer,W. 34  
 Bearse,R.C. 47  
 Beck,W.L. 36  
 Bednar,J.A. 74  
 Bell,F. 88  
 Beloshitsky,V.V. 73  
 Benka,O. 103,118,149  
 Berenyi,D. 80  
 Berinde,A. 100  
 Bernstein,E.M. 2  
 Bethge,K. 89  
 Bevington,P.R. 2  
 Bhalla,R.P. 157  
 Bhattacharya,D. 134,135  
 Bhattacherjee,S.K. 64,134  
 Bienkowski,A. 159  
 Birchall,J. 104  
 Bissinger,G. 20,32,37,41,81,85  
 Bodart,F. 69  
 Bonani,G. 94  
 Bonner,T.I. 21  
 Borggreen,J. 59  
 Brandt,W. 12,17,45,97,120  
 Braziewicz,E. 148  
 Braziewicz,J. 148  
 Brenn,R. 64,66  
 Brennan,J.G. 132,137  
 Brodskii,S.M. 96  
 Brown,M.D. 48,132,137  
 Brunner,K. 51  
 Brunner,W.F. 13  
 Burch,D. 58,70  
 Busch,C.E. 41  
 Calvert,J.M. 152  
 Carlton,R.F. 38  
 Celler,A. 107,124  
 Chaturvedi,R.P. 57  
 Chee,Y. 79  
 Chiao,T. 40,48  
 Chmielewska,D. 140  
 Christensen,L.J. 13  
 Ciortea,C. 83,109  
 Close,D.A. 47  
 Cocke,C.L. 74  
 Crews,B. 79  
 Criswell,T.L. 61  
 Crumpton,D. 77,86  
 Cue,N. 66  
 Curnette,B. 74  
 Curry,B.P. 16  
 Cuzzocrea,P. 127  
 Czuchlewski,S.J. 65  
 Da Silva,M.F. 133  
 Da Silviera,E.F. 142  
 De Castro Faria,N.V. 142  
 De Cesare,N. 127  
 De Heer,F.J. 46  
 De Pinho,A.G. 142,147,153  
 Deberth,C. 100  
 Deconninck,G. 69,90  
 Demeter,I. 122  
 Der,R.C. 16  
 Dima,S. 83  
 Divoux,S. 144  
 Dost,M. 100,114,125,128  
 Doyle,B.L. 85  
 Duggan,J.L. 36,38,72,87,126  
 Dutkiewicz,V. 66  
 Eichler,E. 42  
 El-Ashry,F.M. 150,155,158  
 Ellsworth,L.D. 40,48,65  
 Engelmann,C. 105  
 Eschbach,H.L. 112,131,143  
 Fahlenius,A. 27,44  
 Feldman,L.C. 39  
 Ferreira,G.P. 98,106  
 Filatov,V.I. 96  
 Folkmann,F. 59  
 Fortner,R.J. 16  
 Foster,J.S. 146  
 Francois,P.E. 77,86  
 Freire,F.L.,Jr. 142,147  
 Fujimoto,F. 71,116  
 Furuta,T. 21  
 Gardner,R.K. 67  
 Geissel,H. 146  
 Geretschlager,M. 103,118,149  
 Gil,F.B. 98

## 6.3. Author index for the data base references in Sec. 6.2 -- Continued

- Gilboy,W.B. 112,131  
 Glowacka,L. 150,155,158,159  
 Goclawski,M. 139,145,150,154,158  
 Gonsior,B. 144  
 Gray,T.J. 53,61,63,67  
 Hachiya,Y. 75  
 Hahn,R.L. 42  
 Hanser,F.A. 19  
 Hansteen,J.M. 3  
 Harakeh,M.N. 140  
 Hardt,T.L. 49  
 Harrison,K.G. 46  
 Hart,R.R. 18  
 Hink,W. 51,119  
 Hock,G. 80  
 Hogedal,F. 113  
 Holzer,T.E. 10  
 Hopkins,A.G. 86  
 Hopkins,F. 64,66  
 Hoppenau,S. 125,128  
 Hornshoj,P. 124,139,141,145  
 Hsu,C.N. 56  
 Ilkhamov,R.A. 161  
 Inouye,T. 23  
 Ishii,K. 56,60,75,91,117,136  
 Jacobs,W.W. 85  
 Jarvis,O.N. 30  
 Jaskola,M. 124,139,145,150,154,155,158,159  
 Jauho,P. 27  
 Jesus,A.P. 98,106,115,121,129,133  
 Johansson,T.B. 55  
 Jopson,R.C. 7  
 Joyce,J.M. 20,81,85  
 Kaji,H. 56,60  
 Kamada,H. 35  
 Kamei,H. 23  
 Kamiya,M. 91,117,136  
 Kantele,J. 107  
 Karlowicz,R.R. 54,62  
 Karp,J. 64  
 Kavanagh,T.M. 16  
 Kawatsura,K. 71,92,116  
 Khan,J.M. 9,11,13,16,18  
 Khan,M.R. 77,86  
 Khelil,N.A. 63  
 Khusmurodov,S.H. 161  
 Kising,J. 125  
 Kiss,I. 80  
 Kjeldgaard,A. 59  
 Knaf,B. 84  
 Kobzev,A.P. 161  
 Kolatay,E. 80  
 Komaki,K. 116  
 Komarek,P. 14,15  
 Kretschmer,W. 159  
 Kruglova,I.M. 73  
 Kugel,H.W. 37,81  
 Kuwako,A. 117,136  
 Laegsgaard,E. 39,93,113,130  
 Land,D.J. 132,137,160  
 Langenberg,A. 50,82  
 Lapicki,G. 120,157  
 Lark,N.L. 8  
 Larson,H. 52  
 Laubert,R. 12,17,45,97,120  
 Lear,R.D. 53,78  
 Lennard,W.N. 68,110,146  
 Lewis,C.W. 24,31  
 Lewis,H.W. 1,5  
 Li,J.H. 161  
 Liebert,R.B. 52  
 Lin,J. 38  
 Longree,M. 90  
 Lopes,J.S. 98,106,115,121,129,133  
 Lucas,G.J. 10  
 Ludemann,C.A. 42  
 Ludwig,E.J. 20  
 Lund,M. 93,130  
 Luontama,M. 107  
 Macdonald,J.R. 40,48,65  
 Magno,C. 111  
 Makino,I. 25,28  
 Malanify,J.J. 47  
 Mamikonyan,S.V. 96  
 Mann,R. 99  
 Mark,H. 7  
 Martinsson,B.G. 156  
 McDaniel,F.D. 67,72,87,126,157  
 McEver,W.S. 20  
 McKay,J.S.C. 104  
 McKnight,R.H. 43,54,62  
 Merzbacher,E. 1,5  
 Messelt,S. 3,6  
 Milazzo,M. 76,111,123,138  
 Miljanic,D. 52  
 Mitchell,I.V. 68,112,131,138,143  
 Mitra,S.K. 134,135  
 Montenegro,E.C. 142,147,153  
 Morgan,I.L. 21  
 Morita,S. 56,60,75,91,117,136  
 Munz,L. 36  
 Murray,J.S. 10

## 6.3. Author index for the data base references in Sec. 6.2 -- Continued

- Musket,R.G. 34
- Natowitz,J.B. 24,31
- Neacsu,V. 83
- Neamu,I. 100
- Needham,P.B.,Jr. 22,29
- Nettles,P.H. 41
- Nikolaev,V.S. 73
- Ogier,W.T. 10
- Olabanji,S.O. 152,156
- Ootuka,A. 116
- Osetynski,G.M. 148,155
- Ozawa,K. 71,116
- Pajek,M. 148,155,161
- Perillo,E. 127
- Peterson,D.M. 41
- Petrovici,A. 83,109
- Petukhov,V.P. 73
- Pfuetzner,M. 145,154,155,158,159
- Phillips,D. 110,146
- Phillips,G.C. 52
- Pizzi,C. 111
- Ploskonka,J. 148
- Poncet,M. 105
- Popescu,I. 83,109
- Porro,F. 111
- Potter,D.L. 9,11
- Presser,G. 84
- Price,J.L.,Jr. 160
- Protop,C. 100
- Raith,B. 144
- Ramos,S.C. 106,115
- Ramsay,W.D. 104
- Randall,R.R. 74
- Ren,C. 151
- Reuter,F.W.,III 18,33
- Rhodes,J.R. 21
- Riccobono,G. 76,111,123
- Rice,R.K. 126
- Richard,P. 21
- Ricz,S. 80
- Risley,J.S. 58
- Ritter,R.C. 43
- Rohl,S. 100,125,128
- Romanovsky,E.R. 73
- Rosato,E. 127
- Rota,A. 111,123
- Rowe,J.R. 87
- Roy,A. 134
- Rudolph,W. 99
- Sakisaka,M. 25,28
- Saltmarsh,M.J. 42
- Sanders,J.T. 78
- Sandrik,R. 161
- Sartwell,B.D. 22,29
- Scates,W.R. 41
- Scharnagl,T. 119
- Schmidt-Bocking,H. 89
- Schmitt,R. 159
- Schneider,D. 58
- Schonfeldt,W.A. 128
- Schorn,P. 125
- Schule,R. 89
- Schuster,W. 159
- Scintei,N. 100
- Sebata,M. 136
- Sellers,B. 19
- Sellin,I. 12
- Sera,K. 91,117
- Sergeev,V.A. 73
- Shafrroth,S.M. 20,32,41,85
- Shah,M. 30
- Shima,K. 25,28,101,102
- Shiokawa,T. 56,60
- Sigaud,G.M. 147,153
- Simmons,B.E. 1
- Simons,D.G. 132,137,160
- Singh,B. 4
- Smith,H.P.,Jr. 18,33
- Soares,C.G. 78
- Spadaccini,G. 127
- Spaulding,J.D. 36
- Stahler,J. 84
- Steibing,K.E. 89
- Stockli,M. 94
- Stoller,C. 94
- Stolterfoht,N. 58
- Sujkowski,Z. 140
- Suter,M. 94
- Swift,C.D. 7
- Szerypo,J. 124,139,145,150,154,155,158,159
- Szokefalvi-Nagy,Z. 122
- Tamura,T. 35
- Tang,J. 151
- Tawara,H. 46,56,60,75,91
- Terasawa,M. 23,26,35,71,116
- Thornton,S.T. 43,54,62
- Tonuma,T. 136
- Trautmann,D. 158,159
- Trivia,G. 138
- Tserruya,I. 89

### **6.3. Author index for the data base references in Sec. 6.2 -- Continued**

- Umbarger,C.J. 47  
Valkovic,V. 52  
Van der Woude,A. 42  
Van Eck,J. 50,82  
Van Rinsveld,H.A. 78  
Vigilante,M. 127  
  
Walls,J. 79  
Waltner,A.W. 32,41  
Watson,R.L. 24,31,49  
Weber,K.H. 88  
Wheeler,R.M. 57  
Whitehead,C. 30  
Whittemore,A.R. 64,66  
Wieman,H. 58  
Wilk,S. 69  
  
Wilson,H.H. 19  
Wilson,S.R. 87  
Winters,L.M. 40,48  
Wolfli,W. 94  
Worley,R.D. 11  
  
Xu,H. 151  
  
Yang,F. 151  
  
Zabel,T. 52  
Zander,A.R. 57,79,108  
Zekl,H. 89  
Zelazny,Z. 124,139,141,145,150,154,155,158  
Zemlo,L. 124,145  
Zoran,V. 100  
Zylicz,J. 107,145