International Equations for the Pressure Along the Melting and Along the Sublimation Curve of Ordinary Water Substance

Cite as: Journal of Physical and Chemical Reference Data **23**, 515 (1994); https://doi.org/10.1063/1.555947 Submitted: 10 October 1992 . Published Online: 15 October 2009

Wolfgang Wagner, A. Saul, and A. Pruss



ARTICLES YOU MAY BE INTERESTED IN

The IAPWS Formulation 1995 for the Thermodynamic Properties of Ordinary Water Substance for General and Scientific Use

Journal of Physical and Chemical Reference Data **31**, 387 (2002); https://doi.org/10.1063/1.1461829

New Equations for the Sublimation Pressure and Melting Pressure of H₂O Ice Ih

Journal of Physical and Chemical Reference Data **40**, 043103 (2011); https://doi.org/10.1063/1.3657937

A Fundamental Equation for Water Covering the Range from the Melting Line to 1273 K at Pressures up to 25000 MPa

Journal of Physical and Chemical Reference Data 18, 1537 (1989); https://doi.org/10.1063/1.555836





Journal of Physical and Chemical Reference Data **23**, 515 (1994); https://doi.org/10.1063/1.555947 © 1994 American Institute of Physics for the National Institute of Standards and Technology.

International Equations for the Pressure along the Melting and along the Sublimation Curve of Ordinary Water Substance

W. Wagner^a, A. Saul^b, and A. Pruß

Institut für Thermo- und Fluiddynamik, Ruhr-Universität Bochum, D-44780 Bochum, Germany

Received October 10, 1992; revised manuscript received November 12, 1993

In order to define the phase boundary between the solid phase (ice) and the fluid phase (liquid and gas) of ordinary (light) water substance in pressure-temperature coordinates, correlation equations for the pressure along the melting curve of the various modifications of ice as well as for the pressure along the sublimation curve are presented. The five equations for the melting pressure of the ice phases ice I, ice III, ice V, ice VI, and ice VII, which only contain one to three fitted coefficients, cover the pressure range from the "normal" triple point to 20000 MPa. In this entire range the equations represent the selected measurements of the melting pressure within their experimental uncertainty. The 2-coefficient equation for the sublimation pressure covers the temperature range from 190 K to the triple point (273.16 K). The equations form the basis of the revised release on the pressure along the melting and sublimation curves of ordinary water substance. A verbatim copy of this release is presented in the Appendix of this paper.

Key words: correlation equations; melting line; sublimation line; water.

Contents

Nomenclature 516 Introduction 516 1. 2. Melting Curve 516 3. Sublimation Curve 517 4. Estimates of Uncertainties and Computerprogram Verification. 518 5. Acknowledgment 525 References 525 6. Appendix. IAPWS Revised Release on the Pressure along the Melting and the Sublimation Curves of Ordinary Water Substance 526 List of Tables

1.	values for the triple points of the solid phases	
	which coexist with the liquid	519
2.	Selected data sets for the melting pressure p_m and estimated total experimental uncertainties in	
	pressure	519
3.	Equations for the melting pressure of the various	
	ice phases of ordinary water $(\theta = T/T_n)$ and	
	$\pi = p_{\rm m}/p_{\rm n}$)	519
4.	Selected data sets for the sublimation pressure	
	p_{subl} and their estimated total experimental uncer-	
	tainty in pressure	520

^aTo whom correspondence should be addressed.

^bPresent address: Ruhrgas AG, D-45138 Essen, Germany.

©1994 by the U.S. Secretary of Commerce on behalf of the United States. This copyright is assigned to the American Institute of Physics and the American Chemical Society.

Reprints available from ACS; see Reprints List at back of issue.

5.	Estimated uncertainties of the calculated melting	
	pressures, Eqs. (2.4) to (2.8) and the sublimation	
	pressure, Eq. (3.5)	520
6.	Pressure values calculated from Eqs. (2.4) to (2.8)	
	and (3.5) at a selected temperature	520

List of Figures

1.	The phase-boundary curves of water in a pT dia-	501
2.	Percentage deviation of the experimental melting pressures $p_{m,expt}$ from the values $p_{m,calc}$ calculated from Eq. (2.4) for ice II Eq. (2.5) for ice III Eq.	521
	(2.6) for ice V, Eq. (2.7) for ice VI, and Eq. (2.8) for ice VII	522
3.	Percentage deviation $\Delta p_{subl} = (p_{subl,expt} - p_{subl,calc})/p_{subl,expt}$ and absolute deviations of the experimen- tal sublimation pressures $p_{subl,expt}$ from values $p_{subl,calc}$ calculated from Eq. (3.3); the absolute de- viation diagram only covers the lower tempera- ture range from 190 K to 240 K. Values calcu- lated from Wexler's equation ¹⁸ are plotted for	
	comparison	523
4.	Percentage deviation $\Delta p_{subl} = (p_{subl,expt} - p_{subl,calc})/p_{subl,expt}$ and absolute deviations of the experimen- tal sublimation pressures $p_{subl,expt}$ from values $p_{subl,calc}$ calculated from Eq. (3.5); the absolute deviation diagram only covers the lower tempera- ture range from 190 K to 240 K. Values calculated from Wexler's equation ¹⁸ are plotted for	
	comparison	524

Nomenclature

Symbol	Description
а	Adjustable coefficient
C_p	Isobaric specific heat capacity
Ġ	Gas
h	Specific enthalpy
i	Serial number
Ι	Maximum number of terms in an equation
L	Liquid
n	Exponent
р	Pressure
S	Solid
t	Exponent
Т	Absolute temperature
v	Specific volume

Greek

π	Reduced	pressure	$(\pi =$	$p/p_{\rm n}$)
---	---------	----------	----------	-----------------

 θ Reduced temperature ($\theta = T/T_n$)

Superscripts

+ Denotes the state of the solid phase on the melting-pressure curve and on the sublimation-pressure curve, respectively.

Denotes the state of the liquid phase on the vapor-pressure curve.

" Denotes the state of the gas phase on the vapor-pressure curve and on the sublimation-pressure curve, respectively. gen General; full bank of terms.

Subscripts

calc	Calcu	lated
------	-------	-------

expt Experimental

- *i* Index
- m Denotes the melting pressure
- n Denotes a quantity used to normalize another quantity, e.g. $\pi = p_m/p_n$
- s Denotes the vapor pressure
- subl Denotes the sublimation pressure
- t At a triple point

1. Introduction

The International Association for the Properties of Water and Steam (IAPWS) provides internationally accepted formulations for the properties of light and heavy steam, water and selected aqueous solutions for scientific and industrial applications. Besides publications on other properties there are the IAPWS approved equation of state, called IAPS-84,¹ covering the fluid range of ordinary water substance (H₂O), and special correlation equations for the gas-liquid saturation properties of ordinary water substance.^{2, 3} In this context, it is the purpose of this paper to define the locus of the phase boundary in pressure-temperature coordinates between the fluid phases of water (liquid and gas) and solid water (ice). Based on the available data sources correlation equations for the sublimation pressure as well as for the melting pressures of the various modifications of ice (ice l, ice III, ice V, ice VI, and ice VII) are presented.

Figure 1 shows the phase-boundary curves of water in a pT diagram. The curves for which correlation equations are given in this paper are drawn by bold lines. The equation for the vapor pressure p_s is described in the paper of Wagner *et al.*² (ITS-90) which is a revision of the article of Saul and Wagner³ (IPTS-68) with regard to the temperature scale.

2. Melting Curves

One of the anomalies of water substance is the variety of modifications of the solid. Experimental melting-pressure data exist for the species ice I, ice III, ice V to ice VII along the solid-liquid phase boundaries. There is no set of equations which allows the calculation of the melting pressure for the whole phase boundary covered by experimental data. There are only correlation equations developed by Pistorius *et al.*⁴ and Mishima and Endo⁵ for the part of the melting curve where ice VII coexists with the liquid. Both authors use the Simon equation as well as a correlation equation in the form of a power series for the phase transition ice VII-liquid. The Simon equation

$$\frac{p_{\rm m}}{p_{\rm t}} = 1 + a \left[1 - \left(\frac{T}{T_{\rm t}}\right)^n \right] \tag{2.1}$$

is characterized by its simplicity; temperature can be obtained directly for a given pressure and it is very suitable for extrapolation purposes. However, neither Pistorius *et al.*⁴ nor Mishima and Endo⁵ were able to correlate their own measurements within the experimental uncertainty by the Simon equation. To yield a satisfactory representation of the experimental data both authors fitted power series to their own data.

The triple-point temperatures and the corresponding pressures where two solid phases coexist with the liquid are given in Table 1.

The melting curve is mainly described by the measurements of Bridgman⁶⁻⁸ obtained in the period 1912 to 1942. As the reference points for the calibration of pressure indicators were changed after Bridgman had taken his measurements, we corrected all pressures in accordance with the advice given by Babb.⁹ Moreover, all temperature values of the measurements were converted into the 1990 International Temperature Scale (ITS-90).^{10, 11}

Besides the Bridgman data, Table 2 lists the selected data sets for the melting curve of water. The recalculated data of Mishima and Endo⁵ for the melting pressure of ice VII were not used because they are considered to be inconsistent with the data of the other authors. Based on the selected data we have developed equations for the melting pressure of the phase transitions ice I-L, ice III-L, ice V-L, ice VI-L, and ice VII-L (ice II and ice IV are modifications which do not coexist with the liquid). Each of the equations was produced from a general functional formulation called "bank of terms" by applying the optimization method developed by Setzmann and Wagner.¹⁵ The following two expressions formed the bank of terms for all correlation equations of the melting pressure:

$$\frac{p_{\rm m}^{\rm gen}}{p_{\rm n}} = 1 + \sum_{i=1}^{I} a_i \left[1 - \left(\frac{T}{T_{\rm n}}\right)^{t_i} \right]$$
(2.2)

$$\ln\left[\frac{p_{\rm m}^{\rm gen}}{p_{\rm n}}\right] = \sum_{i=1}^{l} a_i \left[1 - \left(\frac{T}{T_{\rm n}}\right)^{t_i}\right]$$
(2.3)

where p_n and T_n are suitably chosen normalizing values for pressure and temperatures; see below. Eq. (2.2) corresponds to a general bank of terms of the Simon type. It was found that this type of equation provides a suitable functional form for the melting-pressure equations of ice I, ice III, ice V, and ice VI. The new equations (2.4) to (2.7) (see Table 3) represent all the experimental data within the estimated experimental uncertainty. Equations which were established on the basis of Eq. (2.3) behaved quite similarly (except for ice I) but did not improve the representation of the experimental data. For the ice VII region, however, the new general functional form defined by Eq. (2.3) is superior to the Simon equation and allowed the development of a melting-pressure equation for ice VII that represents all selected data within the experimental uncertainty given in Table 2. However, within the experimental uncertainty, the data of Holzapfel and Franck¹⁴ and of Pistorius et al.⁴ show systematic deviations of about -4%. The reason for this behavior is based on the fact that, just for fitting the equation, we shifted these data by about +4% to bring them in agreement with the much more accurate data of Bridgman,^{7,8} and of Bendeliani and Vereschagin¹³ for temperatures below 475 K. This "manipulation" was to stabilize the fitting process; of course, the comparison shown in Fig. 2 was carried out with the original, unshifted data.

To yield thermodynamic consistency at the triple points, with the exception of Eq. (2.8), each of the melting-pressure equations was fixed at the lower triple point (that one which corresponds to the lower pressure, see Table 1) by the choice of the functional form of the equation and at the upper triple point by a constraint. However, since for the phase equilibrium ice VII-liquid only one triple point is known, Eq. (2.8) was only fixed at the triple point ice VII-L.

At the "normal" triple point solid-liquid-gas (S-L-G, S = ice I) thermodynamic consistency with respect to the internationally recommended vapor-pressure equation^{2,3} [$p_s(T_t) = p_m(T_t)$] was achieved by calculating the reduced pressure π of the melting-pressure equation (2.4) with the p_s value obtained from the vapor-pressure equation at the triple point, namely $p_s(T_t) = 611.657$ Pa. Furthermore, the slope of the melting-pressure equation for ice I at the S-L-G triple point was fitted to the value determined as follows: At the S-L-G triple point the volume difference between the saturated liquid and the melting solid, $v'-v^+ = -0.090 \text{ cm}^3/\text{g}$, is known from the measurements of Bridgman.⁶ If one uses the value 333.66 J/g calculated by Jancso *et al.*¹⁶ for the enthalpy of melting namely, $h'-h^+ = 333.66 \text{ J/g}$, the slope of the melting curve ice I-L at the triple point S-L-G can be determined from the Clausius-Clapeyron equation as $dp_m/dT = -13.57 \text{ MPa/K}$.

As a result of these calculations the correlation equations for the melting pressure of the various ice phases together with the corresponding temperature and pressure ranges are given in Table 3 as Eqs. (2.4) to (2.8). Fig. 2 shows that the melting-pressure equations for ice I, ice III, ice IV, and ice V represent the experimental data within about $\pm 3\%$, which corresponds to the estimated experimental uncertainty. Along the melting curve of ice VII the uncertainty of the experimental data rises to $\pm 10\%$. Therefore, Eq. (2.8) was not improved by increasing the number of coefficients.

3. Sublimation Curves

Several correlation equations for the sublimation pressure are known. The two latest versions are the five-coefficient equation of Jancso *et al.*¹⁶ and the six-coefficient equation of Wexler, ¹⁸ which are based on temperatures according to the IPTS-68. These two equations cover the temperature range between 173 K and 273 K.

There are only a few experimental studies of the sublimation pressure of water. Information about the two data sets of Jancso *et al.*¹⁶ and of Douslin and Osborn ¹⁹ is summarized in Table 4. The values of the experimental sublimation pressures given in the paper of Jancso *et al.*¹⁶ had to be corrected because the capacitance manometer was calibrated by measuring the known pressure on the sublimation curve at 273.15 K which corresponded to a pressure at the triple point (273.16 K) of 611.283 Pa. Since, however, this value differs from the triple-point pressure which is internationally accepted at present, namely 611.657 Pa measured by Guildner *et al.*, ¹⁷ we corrected the sublimation pressures of Jancso et al.¹⁶ correspondingly.

To ensure thermodynamic consistency at the ice I-L-G triple point with respect to the vapor-pressure equation² and to the melting-pressure equation for ice I $[p_s(T_t) = p_{subl}(T_t) = p_m(T_t)]$, we fixed the sublimation-pressure equation at the triple-point pressure of 611.657 Pa by the choice of the functional structure. The slope of the sublimation pressure curve at the triple point was determined by the Clausius-Clapeyron equation

$$\frac{\mathrm{d}p_{\rm subl}}{\mathrm{d}T} = \frac{1}{T} \frac{h'' - h^+}{v'' - v^+} \tag{3.1}$$

In order to evaluate this equation at the ice I-L-G triple point, it can be written in the following form:

$$\frac{\mathrm{d}p_{\mathrm{subl}}}{\mathrm{d}T} = \frac{1}{T} \frac{(h'' - h') + (h' - h^{+})}{(v'' - v') + (v' - v^{+})} \,. \tag{3.1a}$$

The values for the enthalpy of vaporization h''-h'=2500.3 kJ/kg and for the difference between the volume of the saturated vapor and liquid v''-v'=206.030 m³/kg were obtained from the internationally recommended equations for the saturation line.^{2,3} The values for $h'-h^+$ as well as for $v'-v^+$ have already been given in Sec. 2. The resulting slope of the sublimation-pressure curve at the ice I-L-G triple point is determined as:

$$\frac{\mathrm{d}p_{\text{subl}}}{\mathrm{d}T} = 50.36 \frac{\mathrm{Pa}}{\mathrm{K}} . \tag{3.2}$$

$$T=273.16 \mathrm{K}$$

Based on a bank of terms similar to Eq. (2.3) and taking into account the selected data and the determined slope of the sublimation pressure at the triple point, the optimization procedure of Setzmann and Wagner¹⁵ yielded the following equation:

$$\ln\left[\frac{p_{\text{subl}}}{p_{\text{t}}}\right] = a_1(1-\theta^{-1}) + a_2(1-\theta^{-3}) + a_3(1-\theta^{-5})$$
(3.3)

with $\theta = T/T_t$, $T_t = 273.16$ K, and $p_t = 611.657$ Pa. The coefficients a_1 to a_3 are not given for the following reason. Although Fig. 3 shows that Eq. (3.3) represents the measured sublimation pressures (see Table 4) in the entire temperature range 190 K $\leq T \leq 273.16$ K within the scattering of the data, Eq. (3.3) yields values for the derivatives which are not sufficiently accurate. The good representation of the experimental sublimation pressures, especially at temperatures below 240 K, leads to incorrect derivatives dp_{subl}/dT and d^2p_{subl}/dT^2 . Rabinovitch²⁰ found this out when calculating the molar isobaric heat capacity c_p^+ of the solid ice I on the sublimation pressure curve from the relation

$$c_{p}^{+} = c_{p}^{+} -T \cdot \frac{d^{2} p_{\text{subl}}}{dT^{2}} (v^{\prime \prime} - v^{+})$$
$$-2T \cdot \frac{dp_{\text{subl}}}{dT} \left[\left(\frac{\partial v^{\prime \prime}}{\partial T} \right)_{p} - \left(\frac{\partial v^{+}}{\partial T} \right)_{p} \right]$$
$$-T \cdot \left(\frac{dp_{\text{subl}}}{dT} \right)^{2} \left[\left(\frac{\partial v^{\prime \prime}}{\partial p} \right)_{T} - \left(\frac{\partial v^{+}}{\partial p} \right)_{T} \right], \qquad (3.4)$$

which can be derived from Eq. (3.1). When Rabinovitch determined the needed derivatives of the sublimation pressure from the IPTS-68 version of Eq. (3.3),²¹ he obtained physically unreasonable values for c_p^+ , and for $T \leq 240$ K he found even negative values. The reason for this misguided behavior is not based on incorrect values for $c_p^{"}$, $v^{"}$, v^+ and the derivatives of v^+ and $v^{"}$ (there are correlation equations in literature which are sufficiently accurate) but on the values of the derivatives dp_{subl}/dT and d^2p_{subl}/dT .² which were not sufficiently accurate.

The reason for obtaining incorrect derivatives from Eq. (3.3) is that the experimental sublimation pressures

especially at lower temperatures must have much larger experimental uncertainties than estimated by Jancso *et al.*¹⁶ and by Douslin and Osborn¹⁹, see Table 4.

Thus, the new sublimation-pressure equation could not be developed by using the experimental sublimation pressures as the only input data. In order to extend the data base for the new sublimation-pressure equation, we calculated dp_{subl}/dT and d^2p_{subl}/dT^2 from Eq. (3.1) where the enthalpy of sublimation $(h''-h^+)$ and the specific volume v'' of the gas on the desublimation line were determined from correlation equations given by Wexler.¹⁸ Then, based on the bank of terms similar to Eq. (2.3) and taking into account the experimental sublimation pressures with a low weight and the calculated derivatives dp_{subl}/dT and d^2p_{subl}/dT^2 with relatively high weights, the optimization procedure of Setzmann and Wagner¹⁵ iteratively used in combination with a nonlinear least-squares procedure yielded the following equation:

$$\ln\left[\frac{p_{\text{subi}}}{p_{\text{t}}}\right] = a_1(1-\theta^{-1.5}) + a_2(1-\theta^{-1.25})$$
(3.5)

with $\theta = T/T_t$, $T_t = 273.16$ K, $p_t = 611.657$ Pa, $a_1 =$ -13.9281690, and $a_2 = 34.7078238$. Eq. (3.5) is valid for the temperature range 190 K $\leq T \leq$ 273.16 K. Fig. 4 shows that for temperatures above 245 K, Eq. (3.5) represents the experimental sublimation pressures within their scattering. The lower, absolute deviation diagram of Fig. 4 illustrates the systematic deviations between the experimental data and the values calculated from Eq. (3.5); these deviations are "necessary" to obtain the correct derivatives from Eq. (3.5). The very good agreement between Eq. (3.5) and Wexler's six-coefficient equation is not surprising because a part of the input data (the derivatives) used to establish Eq. (3.5) were calculated from Wexler's auxiliary equations given in Ref. 18. In this way, Eq. (3.5) is partly based on Wexler's concept to establish a sublimation-pressure equation based on theoretical considerations. We mainly used Wexler's "physically correct" derivatives of the sublimation pressure instead of adhering too closely to uncertain experimental data. As well as Wexler's equation,¹⁸ Eq. (3.5) also yields physically meaningful c_p^+ values when applying Eq. (3.4).

All in all, Eq. (3.5) represents the sublimation pressure and the corresponding derivatives of the sublimation-pressure curve of water as well as Wexler's six-coefficient equation but this quality is achieved with only two fitted coefficients and Eq. (3.5) corresponds to ITS-90.

4. Estimates of Uncertainties and Computer-program Verification

The estimated uncertanties of the melting and sublimation pressures calculated from Eqs. (2.4) to (2.8) and (3.5) are listed in Table 5. Based on the relatively high uncertainties of the experimental melting pressures, the derivatives of Eqs. (2.4) to (2.8) could be subjected to larger errors.

To assist the user in computer-program verification, Table 6 lists values for the pressures calculated from each of the six equations at one temperature within the range of validity.

PRESSURE ALONG MELTING AND SUBLIMATION CURVES OF WATER

Table 1. Values for the triple points of the solid phases which coexist with the liquid (L)

Coexisting phases	T _i /K	p _i / MPa	Author
ice I – ice III – L	251.165	209.9	Bridgman ⁶
ice III – ice V – L	256.164	350.1	Bridgman ⁶
ice V – ice VI – L	273.31	632.4	Bridgman ⁶
ice VI – ice VII – L	355	2216	Bridgman ⁷

Table 2. Selected data sets for the melting pressure p_m and estimated total experimental uncertainties in pressure

Authors	Year	Temp. range T/K	Number of data ^a	Ice phase	Uncertainty
Bridgman ⁶ Henderson & Speedy ¹²	1912 1987	251 - 273 258 - 274	5 9 (10)	I	± 3% ± 0.5 MPa
Bridgman ⁶	1912	251 - 256	4	III	± 3%
Bridgman ⁶	1912	256 - 273	7	v	± 3%
Bridgman ⁶ Bridgman ⁷ Bridgman ⁸ Bendeliani & Vereschagin ¹³	1912 1937 1942 1969	260 - 334 325 - 355 298 298 + 331	13 6 1 1 (2)	VI	± 3% ± 3% ± 3% ± 5%
Bridgman ⁷ Bridgman ⁸ Holzapfel & Franck ¹⁴ Pistorius <i>et al.</i> ⁴ Bendeliani & Vereschagin ¹³	1937 1942 1969 1963 1969	355 - 465 355 - 448 350 - 620 473 - 700 378 - 619	10 3 20 14 7	VII	± 3% ± 3% ± 10% ± 10% ± 5%

^aIf the number of selected data does not correspond to the total number of data, the latter is given in brackets.

Table 3. Equations for the melting pressure of the various ice phases of ordinary water $(\theta = T/T_n \text{ and } \pi = p_m/p_n)$

Ice phase	Temp. range T/K	Pressure range p / MPa	<i>T</i> _n / K	p _n / MPa	Equation	
I	273.16 - 251.165	0.000611657	273.16	0.000611657	$\pi - 1 - 0.626000 \times 10^{6} (1 - \theta^{-3})$ + 0.197135 × 10 ⁶ (1 - $\theta^{21.2}$)	(2.4)
III	251.165 - 256.164	209.9 - 350.1	251.165	209,9	$\pi = 1 - 0.295252 (1 - \theta^{60})$	(2.5)
v	256.164 - 273.31	350.1 - 632.4	256.164	350.1	$\pi = 1 - 1.18721 (1 - \theta^8)$	(2.6)
VI	273.31 - 355	632.4 - 2216	273.31	632.4	$\pi = 1 - 1.07476 (1 - \theta^{4.6})$	(2.7)
VII	355 -715 ^a	2216 - 19910 ^a	355	2216	$\ln (\pi) = 0.173683 \times 10^{1} (1 - \theta^{-1})$	• •
					$-0.544606 \times 10^{-1} (1-\theta^{5}) \\ +0.806106 \times 10^{-7} (1-\theta^{22})$	(2.8)

^aThese values do not characterize the end of the melting curve for ice VII but correspond to the highest temperature and pressure for which measurements exist.

WAGNER, SAUL, AND PRUSS

Table 4. Selected data sets for the sublimation pressure psubl and their estimated total experimental uncertainties in pressure

Authors	Temp.range T/K	Number of data	Uncertainty ^a
Douslin & Osborn ¹⁹	242 - 271	8	$\pm 3 \times 10^{-3} p (\pm 3 \times 10^{-4} p)$
Jancso et al. ¹⁶	194 — 271	77	$T \ge 230 \text{ K: } \pm 5 \times 10^{-3} p \ (\pm (3-4) \times 10^{-4} \text{Pa})$ $T < 230 \text{ K: } \pm 0.2 \text{ Pa} \ (\pm (3-4) \times 10^{-5} \text{Pa})$

^aUncertainty values given in brackets were estimated by ourselves.

Table 5. Estimated uncertainties of the calculated melting pressures, Eqs. (2.4 to 2.8), and the sublimation pressure, Eq. (3.3). For a comment on the uncertainty of the derivatives of the Eqs. (2.4) to (2.8) see Scc. 4.

Table 6. Pressure values calculated from Eqs. (2.4) to (2.8) and (3.3) at a selected temperature

Equation	Equilibrium	Percentage uncertainty
(2.4)	ice I – liquid	≤ ± 3
(2.5)	ice III – liquid	$\leq \pm 3$
(2.6)	ice V – liquid	≤ ± 3
(2.7)	ice VI – liquid	$\leq \pm 3$
(2.8)	ice VII – liquid	≤ ± 7
(3.3)	ice I – gas	$T < 250 \text{ K}$: $\leq \pm 0.5$
		$T \ge 250 \text{ K}$: $\le \pm 0.1$

Equation	Equilibrium	T/\mathbf{K}	p/MPa
(2.4)	ice I – liquid	260.0	139.382
(2.5)	ice III - liquid	254.0	269.456
(2.6)	ice V – liquid	265.0	479.640
(2.7)	ice VI – liquid	320.0	1356.76
(2.8)	ice VII – liquid	550.0	6308.71
(3.3)	ice I – gas	230.0	0.0000089465



FIG. 1. The phase-boundary curves of water in a pT diagram.



FIG. 2. Percentage deviation of the experimental melting pressures p_{m.expl} from the values p_{m.calc} calculated from Eq. (2,4) for ice I, Eq. (2.5) for ice III, Eq. (2.6) for ice V, Eq. (2.7) for ice VI, and Eq. (2.8) for ice VII.

J. Phys. Chem. Ref. Data, Vol. 23, No. 3, 1994



FIG. 3. Percentage deviation $\Delta p_{subl.expt} - p_{subl.expt}$ and absolute deviations of the experimental sublimation pressures $p_{subl.expt}$ from values $p_{subl.expt}$ calculated from Eq. (3.3); the absolute deviation diagram only covers the lower temperature range from 190 K to 240 K. Values calculated from Wexler's equation¹⁸ are plotted for comparison.



FIG. 4. Percentage deviation $\Delta p_{subl.expl} = (p_{subl.expl} - p_{subl.expl}) p_{subl.expl}$ and absolute deviations of the experimental sublimation pressures $p_{subl.expl}$ from values $p_{subl.expl}$ calculated from Eq. (3.5); the absolute deviation diagram only covers the lower temperature range from 190 K to 240 K. Values calculated from Wexler's equation¹⁸ are plotted for comparison.

524

5. Acknowledgments

The authors are grateful to the Deutsche Forschungsgemeinschaft for financial support of this project. We would also like to thank the members of the Working Group "Thermophysical Properties of Water and Steam" of IAPWS for fruitful discussions. The comments of A. A. Aleksandrov, J. R. Cooper, J. M. H. Levelt Sengers, O. Sifner, H. Sato, K. Watanabe and E. Whalley had a significant impact on the final form of this paper. We would especially like to thank one of the reviewers and also Prof. V. A. Rabinovitch for their important hints with regard to the consistency test of the sublimation-pressure equation via the c_p^{+} calculation.

6. References

- ¹J. Kestin and J. V. Sengers, J. Phys. Chem. Ref. Data 15, 305 (1986).
- ²W. Wagner and A. Pruss, J. Phys. Chem. Ref. Data 22, 783 (1993).
- ³A. Saul and W. Wagner, J. Phys. Chem. Ref. Data 16, 893 (1987).

⁴C. W. F. T. Pistorius, M. C. Pistorius, J. P. Blakey, and L. J. Admiraal, J. Chem. Phys. 38, 600 (1963).

⁵O. Mishima and S. Endo, J. Chem. Phys. 68, 4417 (1978).

- ⁶P. W. Bridgman, Proc. Am. Acad. Arts Sci. 47, 441 (1912).
- ⁷P. W. Bridgman, J. Chem. Phys. 5, 964 (1937).
- ⁸P. W. Bridgman, Proc. Am. Acad. Arts Sci 74, 419 (1942).
- ⁸S. E. Babb, Some notes concerning Bridgman's manganin pressure scale, High pressure measurements, p. 115, Editor A.A. Giardanini and E.C. Lloyd, Butterworth, Washington D.C. (1963).
- ¹⁰H. Preston-Thomas, Metrologia 27, 3 (1990).
- ¹¹Supplementary Information for the ITS-90. International Bureau of Weights and Measures. Pavillon de Breteuil, F-02312 Sevres, France, 1990.
- ¹²S. J. Henderson and R. J. Speedy, J. Phys. Chem. 91, 3069 (1987).
- ¹³N. A. Bendeliani and L. F. Vereschagin, Zh. Fiz. Chim. 43, 1631 (1969), in Russian.
- ¹⁴W. B. Holzapfel and E. U. Franck, High Temp.-High Press. 1, 675 (1969).
- ¹⁵U. Setzmann and W. Wagner, Int. J. Thermophys. 10, 1103 (1989).
- ¹⁶G. Jancso, J. Pupezin, and W. van Hook, J. Phys. Chem. 74, 2984 (1970).
 ¹⁷L. A. Guildner, D. P. Johnson, and F. E. Jones, J. Res. Natl. Bur. Stand. Sect. A, 80, 505 (1986).
- ¹⁸A. Wexler, J. Res. Natl. Bur. Stand. Sect. A, **81**, 5 (1977).
- ¹⁹D. R. Douslin and A. Osborn, J. Sci. Instrum. 42, 369 (1965).
- ²⁰V. A. Rabinovitch, Presentation at the 1992 IAPWS meeting in St. Petersburg.
- ²¹Release on the Pressure along the Melting and along the Sublimation Curves of Ordinary Water Substance. International Association on the Properties of Water and Steam (IAPWS), September 1989.

WAGNER, SAUL, AND PRUSS

Appendix

The International Association for the Properties of Water and Steam Milan, Italy September 1993

Release on the Pressure along the Melting and the Sublimation Curves of Ordinary Water Substance

Unrestricted publication allowed in all countries.

Issued by the International Association for the Properties of Water and Steam

> President: J. R. Cooper Queen Mary and Westfield College London E1 4NS England

Executive Secretary: Dr. R. B. Dooley Electric Power Research Institute Palo Alto, California 94304 USA

This release is a revision of the edition of 1989 and contains 4 numbered pages.

This release has been authorized by the International Association for the Properties of Water and Steam (IAPWS) at its meeting in Milan, Italy, 12 - 18 September 1993, for issue by its Secretariat. The members of IAPWS are: Canada, Czech Republic, Denmark, the Federal Republic of Germany, France, Japan, Russia, the United Kingdom, and the United States of America, and the associate members are: Argentina and Italy.

IAPWS issued in 1989 a Release on the Pressure along the Melting and Sublimation Curves of Ordinary Water Substance based on the IPTS-68 temperature scale. This revised release has been prepared with the temperatures of the triple points and the coefficients in the correlation equations changed to correspond to the ITS-90. To obtain the values for the new coefficients, the equations have been refitted using the same input data but with temperatures converted to ITS-90.

Compared with the IPTS-68 temperatures in the original release, one more decimal place is given here to the converted ITS-90 temperatures. This ensures that any recalculation to the original IPTS-68 temperature values produces the same figures as given in the original source after rounding to the same number of decimal places. This increase by one decimal in the converted ITS-90 temperatures does not imply that these values have been redetermined or are more accurate than previously stated on IPTS-68. In the second revision, Eq. (6) has been replaced by a simpler equation which better represents the derivatives of the sublimation pressure. Moreover, the parameter in Eq. (4) has been adjusted for a better representation of the ice VI - ice VII - liquid triple point.

Further details about the equations presented in this release can be found in an article "International Equations for the Pressure along the Melting and the Sublimation Curves of Ordinary Water Substance" by W. Wagner, A. Saul, and A. Pru β , to be published in the Journal of Physical and Chemical Reference Data.

Further information about this release and other releases issued by IAPWS can be obtained from the Executive Secretary of IAPWS.

Equations for the Pressure along the Melting and the Sublimation Curves of Ordinary Water Substance

1 Nomenclature

Thermodynamic quantities:

- p Pressure
- T = Temperature
- $\theta = T/T_n$
- $\pi = p_m/p_n$

Subscripts:

- m Denotes a value on the melting curve
- n Denotes a quantity used to obtain a dimensionless variable
- sublDenotes a value on the sublimation curvetDenotes a value at a triple point
- Note: T denotes absolute Temperature on the International Temperature Scale of 1990.

PRESSURE ALONG MELTING AND SUBLIMATION CURVES OF WATER

2 Melting Pressure

2.1 Melting pressure of ice I (temperature range from 273.16 to 251.165 K)

$$\pi = 1 - 0.626000 \times 10^{6} (1 - \theta^{-3}) + 0.197135 \times 10^{6} (1 - \theta^{21.2})$$
(1)

with $T_n = 273.16$ K and $p_n = 0.000611657$ MPa.

2.2 Melting pressure of ice III (temperature range from 251.165 to 256.164 K)

 $\pi = 1 - 0.295252 (1 - \theta^{60}) \tag{2}$

with $T_n = 251.165$ K and $p_n = 209.9$ MPa.

2.3 Melting pressure of ice V (temperature range from 256.164 to 273.31 K)

 $\pi = 1 - 1.18721 (1 - \theta^8) \tag{3}$

with $T_n = 256.164$ K and $p_n = 350.1$ MPa.

2.4 Melting pressure of ice VI (temperature range from 273.31 to 355 K)

 $\pi = 1 - 1.07476 (1 - \theta^{4.6}) \tag{4}$

with $T_n = 273.31$ K and $p_n = 632.4$ MPa.

2.5 Melting pressure of ice VII (temperature range from 355 to 715 K)

 $\ln(\pi) = 0.173683 \times 10^{1} (1 - \theta^{-1}) - 0.544606 \times 10^{-1}$ $(1 - \theta^{5}) + 0.806106 \times 10^{-7} (1 - \theta^{22})$ (5)

with $T_n = 355$ K and $p_n = 2216$ MPa.

Note: The upper temperature of the range of Eq. (5) corresponds to the highest temperature for which measurements exist and not the end of the melting curve of ice VII.

Eqs. (1) to (5) are constrained to fit the experimental values T_t and p_t of the relevant triple points given in Table 1.

TABLE 1. Values for the triple points of the solid phases which coexist with the liquid

$T_{\rm t}/{\rm K}$	pt / MPa
251.165	209.9
256.164	350.1
273.31	632.4
355	2216
	<i>T</i> ₁ /K 251.165 256.164 273.31 355

3 Sublimation pressure

$$\ln\left[\frac{p_{\text{subl}}}{p_{\text{n}}}\right] = a_{1}(1-\theta^{-1.5}) + a_{2}(1-\theta^{-1.25})$$
(6)

with $a_1 = -13.9281690$ $T_n = 273.16 \text{ K} = T_t$ $a_2 = 34.7078238$ $p_n = 611.657 \text{ Pa} = p_t$

•Note: Eq. (6) is valid from 190 to 273.16 K.

4 Range of validity of the equations

IAPWS endorses the validity of the equations presented in this release for each of the saturation lines. Eqs. (1) to (4) are valid for the entire range of the solid-liquid equilibrium. Eqs. (5) to (6) only cover the ranges of the solid-liquid or solid-vapor equilibrium, respectively, as indicated.

5 Estimates of uncertainty

The estimated uncertainties of the melting and sublimation pressures calculated from Eqs. (1) to (6) are listed in Table 2. Based on the relatively high uncertainties of the experimental melting pressures, the derivatives of Eqs. (1) to (5) could be subjected to larger errors.

Equation	Equilibrium	Percentage uncertainty
(1)	ice I – liquid	≤ ± 3
(2)	ice III - liquid	$\leq \pm 3$
(3)	ice V – liquid	$\leq \pm 3$
(4)	ice VI - liquid	≤ ± 3
(5)	ice VII – liquid	≤ ± 7
(6)	ice I – gas	$T < 250 \text{ K}$: $\leq \pm 0.5$
		$T \ge 250 \text{ K}$: $\le \pm 0.1$

TABLE 2. Estimated uncertainties of the calculated pressures

6 Computer-program verification

To assist the user in computer-program verification, Table 3 lists values for the pressures calculated from each of the six equations at one temperature within the range of validity.

TABLE 3. Pressure calculated from Eqs. (1) to (6) at a selected temperature

Equation	Equilibrium	T/K	ρ/MPa
(1)	ice I – liquid	260.0	139.382
(2) .	ice III – liquid	254.0	269.456
(3)	ice V - liquid	265.0	479.640
(4)	ice VI - liquid	320.0	1356.76
(5)	ice VII – liquid	550.0	6308.71
(6)	ice I – gas	230.0	0.0000089465